

A particularly unstable film

Jeffrey F. Morris[†]

Levich Institute and Department of Chemical Engineering, CUNY City College of New York, New York, NY 10031, USA

Analysis of the classic problem of shallow film flow on an inclined plane is revisited for a Brownian suspension. The particle phase, considered in a two-fluid model, is predicted to cause pronounced changes to the instability characteristics of the flow. These are due to an indirect effect of the non-Newtonian rheology, with the normal stresses causing migration and a viscosity stratification which strongly alters the base state from its Newtonian counterpart. Both the short- and long-wavelength inertial stability boundaries are altered, and, more strikingly, an instability at zero Reynolds number is found.

Key words: thin films, particle/fluid flow, instability

1. Introduction

Dhas & Roy (2022) (DR) recently presented a stability analysis for flow of a Brownian suspension down an inclined plane, as illustrated by figure 1(a). The flow is considered in the shallow limit, with the flow length l and film depth h_0 satisfying $h_0/l \ll 1$, allowing the simplifications of lubrication and boundary layer analyses. Here, h_0 is the uniform base state thickness seen in figure 1(b). The work by DR predicts not only destabilizing effects of the particles to the classical (Benjamin 1957; Yih 1963) inertial instabilities of a film, but also a previously unknown instability at vanishing inertia.

An issue that features prominently in the work of DR is shear-induced particle migration, which has long been a theme in suspension studies (Leighton & Acrivos 1987), and is associated with the normal stresses attributable to the particle phase (Morris & Boulay 1999). Due to variation in particle volume fraction ϕ caused by migration, the base state for the stability analysis is altered from the half-parabola velocity profile of the pure liquid. Migration in the analogous pressure-driven channel flow has been studied for Brownian suspensions (Frank *et al.* 2003), and its effects are more pronounced for larger Péclet number, $Pe_p = \dot{\gamma} a^2 / D_0$, with $D_0 = kT / 6\pi\mu_f a$ being the Stokes–Einstein–Sutherland diffusivity of a particle of radius a in a liquid of viscosity μ_f with shear rate $\dot{\gamma}$. Analysis of suspension film stability involves a large parameter space, and this makes for a daunting problem. Specifically, the film Reynolds number $Re = \rho h_0 u_0 / \mu_f$ (here u_0 is the mean axial velocity of the film) along with ϕ and Pe_p are, at a minimum, necessary for the fully developed base state, while a/h_0 affects the rate of development of this state.

[†] Email address for correspondence: morris@ccny.cuny.edu




Figure 1. (a) A particle-laden film and (b) the predicted development of the height and ϕ profile for $\phi_0 = 0.3$ at $Pe_p = 1$, from a uniformly concentrated film at $x = 0$. The axial coordinate x is scaled by h_i , and the final thickness in (b) is the base-state h_0 . From Dhas & Roy (2022).

One touchstone for this study in prior analysis is the stability of a flowing film on a heated substrate (Goussis & Kelly 1985). The base flow studied by DR resembles this flow as both exhibit a viscosity stratification, with a lower viscosity near the plane wall. For the suspension, this is a result of migration towards the free surface and viscosity dependence on ϕ of the typical form $\mu(\phi) = \mu_f(1 - \phi/\phi_m)^{-2}$, with maximum packing fraction ϕ_m . For the Brownian suspensions at $Pe_p \leq 10$ described here, relatively modest variations in ϕ are seen, as indicated by figure 1(b) at $Pe_p = 1$, yet the variation has major implications for film stability. This suggests that the influence on non-colloidal suspensions, where $Pe_p \rightarrow \infty$ and migration is stronger, will be more pronounced.

2. Overview

2.1. Migration, base flow and surface corrugation

A distinguishing feature of the suspension film is that the axially invariant base state requires a significant length to allow migration towards the free surface to develop. An important basic result of DR is that while details may vary, the primary consequences of migration are unaffected by the specific model chosen. The authors have implemented Brownian versions of the diffusive flux model (Khoshnood & Jalali 2012) and the suspension balance model, in which particle flux is driven by the particle stress Σ^P as $J \sim \nabla \cdot \Sigma^P$ (Frank *et al.* 2003). Full development of the flow is achieved when the particle flux normal to the inclined plane vanishes, $J_y = 0$.

Figure 1(b) shows predictions from DR at $Pe_p = 1$ and an initially uniform concentration $\phi_0 = 0.3$. As the flow proceeds axially, the film thins while ϕ increases near the free surface, in agreement with experiments at higher Pe_p (Timberlake & Morris 2005). The film thins because the near-wall viscosity becomes smaller than in the uniform- ϕ film. At larger ϕ_0 , the normal stresses driving migration increase, and the development length decreases. By contrast, the extent of migration and hence the development length both increase with Pe_p ; this is simply understood by noting that when $Pe_p \rightarrow 0$, there is no migration because Brownian motion maintains a uniform ϕ .

Surface corrugations are seen in film flows of suspensions. As shown by figure 2, features large relative to the particle size, and with broad wavenumber content, were found by Timberlake & Morris (2005). A postulate of DR is that these corrugations represent the nonlinear saturation of waves predicted by the current linear stability analysis. These structures found at $Re < 0.01$ may stem from the non-inertial instability of § 2.3.

A particularly unstable film




Figure 2. Surface corrugations in a film of $h_0 = 2.1$ mm, of neutrally buoyant $180 \mu\text{m}$ diameter spheres at uniform $\phi_0 = 0.3$ on a plane inclined at $\alpha = 34.6^\circ$ from horizontal. Lighting and camera orientation are shown at left, with flow from top to bottom (at $Re < 0.01$) of the imaged region, which is 3 cm tall and 2.4 cm wide. Adapted from Timberlake & Morris (2005).

2.2. Stability analysis and modulation of the inertial instability

The coupled equations and associated boundary conditions describing the flow and particle conservation accounting for migration are analysed for the fully developed film. The linear stability analysis follows a standard approach: the variables are expressed as $X = X_b + \hat{X} \exp(ik(x - ct))$, with X_b the base flow value and \hat{X} the infinitesimal magnitude of the perturbation. The equations are then linearized with respect to the perturbation to yield a set of modified Orr–Sommerfeld equations. These are similar to forms found in other viscosity-stratified free-surface flows (Goussis & Kelly 1985), but this set is unique owing to normal stress and particle flux terms specific to suspensions. Detailed guidance to the dominant destabilizing effects is provided by DR.

The addition of particles increases the suspension viscosity and thus an increase in the threshold Re , based on the pure fluid viscosity, for instability is seen with increasing ϕ . This is seen in figures 3(a) and 3(b), where we see the neutral stability curves for the long-wave (surface, low- Re) and the short-wave (shear, higher- Re) modes, respectively; dotted black arrows show the increase in Re for the low- Pe_p , and hence relatively uniform- ϕ , state for both instabilities. The red arrows show the change in the neutral stability conditions with a change from $Pe_p = 0.1$ to $Pe_p = 1$, where the base state is significantly more inhomogeneous due to shear-induced migration (see figure 1b). We see that very long-wave disturbances are predicted to grow at $Re < 1$. As the authors note, this implies that the surface mode is destabilized at Re below the value needed for instability of the pure liquid. This is a striking and exciting result, as it presages a zero-inertia instability seen in the long-wave surface mode. The shear mode is predicted to occur at much larger Re .

The basis for destabilization was explored through systematic removal of terms in the governing Orr–Sommerfeld equations, as elaborated by DR in discussion of their figure 13 and table 3. The authors find that the dominant destabilizing effect, which causes the large majority of the shift associated with each red arrow in figure 3, results from a coupling of the y -directed variation of base-state volume fraction and velocity profiles to the perturbation in ϕ . Base-state stratification is thus the source of the dominant destabilizing effect due to the particles, and this stratification is more pronounced as Pe_p increases.

2.3. Inertialess ($Re = 0$) instability

By taking the limit of $Re \rightarrow 0$, DR show that there is instability in the absence of inertia. This novel prediction occurs above a critical Péclet number of $Pe_p \approx 0.7$. The maximum




Figure 3. Neutral stability curves for the surface (a) and shear (b) modes for $\phi_0 = 0.1$ (black curves) and $\phi_0 = 0.3$ (blue curves) at $Pe_p = 0.1$ (dashed curves) and $Pe_p = 1$ (solid curves). Dotted black arrows highlight the change with increasing ϕ ; solid red arrows show the change due to increase of Pe_p and stronger stratification. Adapted from Dhas & Roy (2022).

growth rate at $Re = 0$ is smaller than at $Re = 20$, but the two are comparable above the critical Pe_p . Because the particle concentration is the sole destabilizing influence, the envelope of instability is larger – extending to shorter wavelengths – for $\phi_0 = 0.3$ than for $\phi_0 = 0.1$. A succinct physical explanation for this instability is not immediately clear. The presence of a viscosity gradient at the ‘critical layer’ where the flow speed equals the wave speed, through its effect on the curvature of the base flow profile as developed by Craik (1969), appears to play a central role.

3. Summary and perspective

The free-surface suspension, flow stability analysis of DR sheds light in a number of directions. At a practical level, the set of instabilities explored by DR may play roles in flows of coatings, crystallizing magma and mud. The alteration of the low- Re surface mode, and appearance of a zero- Re surface instability, are predictions that warrant experimental study. In a broader context, the work of DR serves as a foundation for a more general exploration of instability of suspension flows, providing guidance to various issues that will need to be considered in other flows.

In the instabilities described by DR for the film, a central role is played by particle migration towards the free surface. This induces a continuous viscosity stratification, with increase towards the free surface in the neutrally buoyant case studied here. Establishing this coupling of migration, a basic rheophysical behaviour in suspensions, to the flow stability is a very satisfying finding. In their consideration of film flow, the authors have pointed to the importance of the critical layer, where the flow speed matches the wave speed. The finding of a lower bound on Pe_p to see the zero-inertia instability is presumably based on a need for sufficient stratification due to migration; the influence of varying viscosity at the critical layer seems to be a key physical issue (Craik 1969). The film flow was studied by DR for a neutrally buoyant suspension with stratification solely due to migration, but immediately suggests study of the more general case of particles which rise or settle. These cases could exhibit a wide range of ϕ profiles in a non-colloidal ($Pe_p \rightarrow \infty$) suspension, and this may facilitate experimental study of the role of stratification. In their work, DR chose not to consider larger ϕ in order to avoid a jammed condition at the interface, but the implications of such a ‘frozen’ layer seem intriguing. Further issues of interest include the role of finite-amplitude perturbations, including the natural fluctuations of ϕ , and – as suggested by DR – the nonlinear development of the linearly unstable modes.

A particularly unstable film

Flows which yield to stability analysis often offer significant new understanding. While the pioneering examination of stability by DR considered a shallow suspension film, the work opens the way to deep insight.

Declaration of interest. The author reports no conflict of interest.

REFERENCES

- BENJAMIN, T.B. 1957 Wave formation in laminar flow down an inclined plane. *J. Fluid Mech.* **2** (6), 554–573.
- CRAIK, A.D.D. 1969 The stability of plane Couette flow with viscosity stratification. *J. Fluid Mech.* **36** (4), 685–693.
- DHAS, D.J. & ROY, A. 2022 Stability of gravity-driven particle-laden flows—roles of shear-induced migration and normal stresses. *J. Fluid Mech.* **938**, A29.
- FRANK, M., ANDERSON, D., WEEKS, E.R. & MORRIS, J.F. 2003 Particle migration in pressure-driven flow of a Brownian suspension. *J. Fluid Mech.* **493**, 363–378.
- GOUSSIS, D. & KELLY, R.E. 1985 Effects of viscosity variation on the stability of film flow down heated or cooled inclined surfaces: long-wavelength analysis. *Phys. Fluids* **28**, 3207–3214.
- KHOSHNOOD, A. & JALALI, M.A. 2012 Long-lived and unstable modes of Brownian suspensions in microchannels. *J. Fluid Mech.* **701**, 407–418.
- LEIGHTON, D.T. & ACRIVOS, A. 1987 The shear-induced migration of particles in concentrated suspensions. *J. Fluid Mech.* **181**, 415–439.
- MORRIS, J.F. & BOULAY, F. 1999 Curvilinear flows of noncolloidal suspensions: the role of normal stresses. *J. Rheol.* **43** (5), 1213–1237.
- TIMBERLAKE, B.D. & MORRIS, J.F. 2005 Particle migration and free-surface topography in inclined plane flow of a suspension. *J. Fluid Mech.* **538**, 309–341.
- YIH, C.-S. 1963 Stability of liquid flow down an inclined plane. *Phys. Fluids* **6** (3), 321–334.