

G. Okugawa, K. Takase, K. Nobuhara, Y. Saito, T. Kinoshita. *Department of Neuropsychiatry, Kansai Medical University, Osaka, Japan*

**Objective:** The caudate nucleus is involved in cognitive function. Schizophrenic patients showed cognitive dysfunction. It has been reported that volume reduction of the caudate nucleus was associated with cognitive impairment in schizophrenic patients. Because treatment with olanzapine improves cognitive dysfunction in schizophrenia, olanzapine may affect the caudate nucleus volume in patients with schizophrenia. We measured volumes of grey and white matter in the caudate nucleus of schizophrenic patients.

**Methods:** Ten schizophrenic patients and ten healthy subjects were examined magnetic resonance imaging. Ten patients were scanned at the time of pre-treatment and post-treatment with olanzapine. MR data analysis was performed using BRAINS software in order to measure grey and white matter volume of the caudate nucleus.

**Results:** Schizophrenic patients had reduced volume of grey and white matter of the caudate nucleus compared with healthy subjects. The average duration of treatment with olanzapine was 186 days in schizophrenic patients. The volume of grey and white matter in the caudate nucleus at the time of post-treatment was significant larger than that at the time of pre-treatment with olanzapine in patients with schizophrenia. There was no significant difference between the volume of grey matter of the caudate nucleus at the post-treatment with olanzapine and that of healthy subjects.

**Conclusion:** Schizophrenic patients had reduced volume of the caudate nucleus. Treatment with olanzapine may improve volume reduction of grey matter of the caudate nucleus in schizophrenic patients.

## P104

Once-year experience with aripiprazole in acute care units. Recommendations for use

J. Orta<sup>1</sup>, R. Dueñas<sup>2</sup>, A. Arévalo<sup>3</sup>. <sup>1</sup> *Unidad de Agudos, Hospital Sant Joan de Déu-SSM, Sant Boi, Barcelona, Spain* <sup>2</sup> *Unidad de Agudos, Hospital Benito Menni, Sant Boi, Barcelona, Spain* <sup>3</sup> *Unidad de Agudos, Centro Neuropsiquiátrico Sagrado Corazón, Martorell, Barcelona, Spain*

**Background:** In the last year, a new antipsychotic (AP) was approved in Spain for treatment of schizophrenia: aripiprazole. Objectives: 23 clinical psychiatrists of Acute Units throughout Spain have constituted a regular work group (PSIQ-A) for the purpose of sharing clinical experiences and examining topics of interest to our daily clinical practice.

**Methods:** In periodic meetings, members of PSIQ-A have made a compilation of different approaches to distinct clinical situations that hospitalized schizophrenic patients may present: approach in Emergency Room to try to reach a consensus, specifically with respect to aripiprazole use in each situations.

**Results:** Usually recommended dosage with predominance of positive symptoms: 25-30 mg/day (generally more than 15 mg/day) and with predominance of negative symptoms: smaller doses are sufficient and effective. Because of the demands of Acute Care rapid changes are chosen (about 1 week), except with clozapine and depot preparations (2 weeks) with full doses of aripiprazole in 1-3 days and tapering off of previous AP. The initial, temporary association of drugs with a more sedative profile is frequent (BZD, levomepromazine, quetiapine). Some benefits have been: Reduction of psychotic anxiety; possibility of improving insight; excellent tolerance, even

at high doses; response in negative-residual patients: more activity, more eagerness to do things, “evident improvement in the most chronic patients”.

**Conclusion:** Aripiprazole is a new and interesting drug in the approach to the phase of decompensation and admission of patients with schizophrenia, with good tolerability in major areas of patient concern and feeling well-being and improving insight.

## P105

First psychotic episode - a descriptive study

A. Marques, S. Pereira, M. Martins. *Department of Psychiatry, Centro Hospitalar de Vila Nova de Gaia, Vila Nova de Gaia, Portugal*

**Background and aim:** The importance of early recognition and treatment of the first psychotic episode is well documented in literature. This study aims to describe and analyze the sociodemographic and clinical characteristics of a sample of patients admitted in a psychiatric ward for their first psychotic episode.

**Methods:** Data from 48 patients was retrospectively analyzed using a specific clinical protocol. Inclusion criteria were admission with a first psychotic episode during January 2003 to June 2005. Patients with primary affective and organic disorder were excluded. ACCESS was used for statistic analysis.

**Results:** Patients were aged 19-56, mainly of the masculine gender (77%), single (68%), living with own family (89%), and receiving any kind of social support (13%).

Main diagnoses were Schizophrenia (54%); Persistent Delusional Disorders (17%); Acute and Transitory Psychotic Disorders (29%).

Age of onset was 28 years (median) for males and 36 years for females. Onset was

insidious for 44% of the patients and the Time Disease Untreated (TDU) mean-2,2 month; median 18,8 month, witch is similar with literature data. Ten percent were involuntarily admitted and 84% were taking oral atypical antipsychotic with total compliance for 33% and partial for 25% of patients.

Only 23% of the patients or their families were attending therapeutic groups.

**Conclusion:** The results of our study in part agree with the data from the literature on the other hand they reflect the characteristics of our healthcare system and population, and can provide ways to improve care.

## P106

Antipsychotic treatment and the need for hospitalization: advantages of long-acting neuroleptics

M. Perez Garcia, M. Paramo Fernandez, V. Prado Robles, J. Alonso San Gregorio, J. Perez Perez, I. Tortajada Bonasselt. *Department of Psychiatry, Hospital de Conxo, Santiago de Compostela, A Coruña, Spain*

**Introduction:** At present, the need of antipsychotic treatments for the improvement of the condition of people with psychotic disorders is unquestionable. Despite the current availability of highly effective drugs with few secondary effects, the main cause behind hospitalization is still the lack of compliance.

**Objectives:** Analysis of the determining variables behind the need for hospitalization and the influence of the types of antipsychotic treatments.