

## RIEMANNIAN HOMOGENEOUS FOLIATIONS WITHOUT HOLONOMY

ROBERT A. BLUMENTHAL

### § 1. Introduction

Let  $M$  be a compact connected  $C^\infty$  manifold with a smooth Riemannian foliation  $\mathcal{F}$ . That is,  $(M, \mathcal{F})$  admits a bundle-like metric in the sense of [7]. In [4] it is shown that if all leaves of  $\mathcal{F}$  are closed without holonomy, then the space of leaves  $M/\mathcal{F}$  of the foliation is a manifold and the natural projection  $M \rightarrow M/\mathcal{F}$  is a locally trivial fibre space. In the present work we show that for certain of the Riemannian homogeneous foliations, holonomy is the only obstruction to the foliation being a fibration.

Let  $G/K$  be a simply connected, even dimensional, positively curved Riemannian homogeneous space of a compact connected Lie group  $G$  and let  $\mathcal{F}$  be a homogeneous  $G/K$ -foliation of a compact connected manifold  $M$ . For example,  $\mathcal{F}$  is a codimension  $2q$  elliptic (i.e., homogeneous  $SO(2q+1)/SO(2q) \cong S^{2q}$ -) foliation. Then  $\mathcal{F}$  is cohomologically a fibration in the sense that the base-like cohomology algebra of the foliated manifold  $(M, \mathcal{F})$  is isomorphic to the de Rham cohomology algebra of  $G/K$  [3]. The main result of this paper asserts that if  $\mathcal{F}$  has no holonomy, then it is actually a fibration.

(1.1) **THEOREM.** *If  $\mathcal{F}$  is without holonomy, then  $M$  fibres over  $G/K$  with the leaves of  $\mathcal{F}$  as fibres.*

### § 2. Riemannian Homogeneous Foliations

In this section we prove (1.1) and use its proof to elucidate further properties of Riemannian homogeneous foliations.

Let  $G/K$  be a connected homogeneous space on which  $G$  acts effectively and let  $\mathcal{F}$  be a homogeneous  $G/K$ -foliation of a connected manifold  $M$ . That is,  $\mathcal{F}$  is defined by a  $G/K$ -cocycle  $\{(U_\alpha, f_\alpha, \lambda_{g_{\alpha\beta}})\}_{\alpha, \beta \in A}$  where

---

Received November 12, 1979.

- i)  $\{U_\alpha\}_{\alpha \in A}$  is an open cover of  $M$
- ii)  $f_\alpha : U_\alpha \rightarrow G/K$  is a submersion defining  $\mathcal{F}|_{U_\alpha}$
- iii)  $f_\alpha = \lambda_{g_{\alpha\beta}} \circ f_\beta$  on  $U_\alpha \cap U_\beta$  where  $g_{\alpha\beta} \in G$  and  $\lambda_{g_{\alpha\beta}}$  denotes the diffeomorphism of  $G/K$  which sends  $aK$  into  $g_{\alpha\beta}aK$ .

To prove the theorem, we recall a construction from [2] which gives a useful realization of the holonomy group of a leaf of  $\mathcal{F}$ .

Let  $P = \{[\lambda_g \circ f_\alpha]_x : x \in U_\alpha, \alpha \in A, g \in G\}$ , where  $[\lambda_g \circ f_\alpha]_x$  denotes the germ of  $\lambda_g \circ f_\alpha$  at  $x$  and let  $\pi : P \rightarrow M$  be defined by  $\pi([\lambda_g \circ f_\alpha]_x) = x$ . Then  $P$  possesses a well-defined topology and differentiable structure such that  $\pi : P \rightarrow M$  is a smooth regular covering. Moreover, the evaluation map  $F : P \rightarrow G/K$  defined by  $F([\lambda_g \circ f_\alpha]_x) = \lambda_g(f_\alpha(x))$  is a smooth submersion constant along the leaves of  $\pi^{-1}(\mathcal{F})$  where  $\pi^{-1}(\mathcal{F})$  denotes the foliation of  $P$  whose leaves are the connected components of the inverse images under  $\pi$  of the leaves of  $\mathcal{F}$ . By choosing a connected component of  $P$ , we may assume that  $P$  is connected. This connected regular covering gives rise to a homomorphism  $\Phi : \pi_1(M, x_0) \rightarrow G$  such that  $\Gamma = \text{image}(\Phi)$  is the group of covering transformations and such that the diagram

$$\begin{array}{ccc}
 P & \xrightarrow{F} & G/K \\
 \gamma \downarrow & & \downarrow \lambda_\gamma \\
 P & \xrightarrow{F} & G/K
 \end{array}$$

is commutative for each  $\gamma \in \Gamma$ .

Let  $L$  be a leaf of  $\mathcal{F}$  and choose a leaf  $L'$  of  $\pi^{-1}(\mathcal{F})$  which projects to  $L$ . Then the holonomy group of  $L$  is isomorphic to  $\Gamma_{L'} = \{\gamma \in \Gamma : \gamma(L') = L'\}$  and thus  $\pi|_{L'} : L' \rightarrow L$  is a regular covering with the holonomy group of  $L$  as its group of covering transformations.

If  $K$  is compact, then  $M$  admits a bundle-like metric (whence  $\mathcal{F}$  is called a Riemannian homogeneous foliation) such that the lifted metric on  $P$  is bundle-like with respect to the foliation defined by the submersion  $F$ . Thus if  $M$  is also compact, we have that  $F : P \rightarrow G/K$  is a locally trivial fibre space [5]. (We remark that since an isometry of a connected Riemannian manifold is determined by its value and differential at a point, this construction remains valid for a foliation defined by an  $N$ -cocycle  $\{(U_\alpha, f_\alpha, g_{\alpha\beta})\}_{\alpha, \beta \in A}$  where  $N$  is a connected Riemannian manifold and each  $g_{\alpha\beta}$  extends to an isometry of  $N$ .)

Assume now that  $M, \mathcal{F}$ , and  $G/K$  satisfy the hypotheses of (1.1). Fix

an orientation on  $G/K$  invariant under the action of  $G$ . Let  $\gamma \in \Gamma$ . Then  $\lambda_\gamma$  is an orientation-preserving isometry of  $G/K$  and hence has a fixed point  $y$  [6]. It is here that we have used the assumption in Theorem (1.1) that  $G/K$  is an even dimensional, positively curved Riemannian homogeneous space of the compact connected Lie group  $G$ . Since  $F: P \rightarrow G/K$  is a fibration over a simply connected manifold, we have that the space of leaves of  $\pi^{-1}(\mathcal{F})$  is diffeomorphic to  $G/K$  and hence there exists a unique leaf  $L'_0$  of  $\pi^{-1}(\mathcal{F})$  such that  $F(L'_0) = y$ . Now  $F(\gamma(L'_0)) = \lambda_\gamma(F(L'_0)) = \lambda_\gamma(y) = y = F(L'_0)$  and hence  $\gamma(L'_0) = L'_0$ . Since  $\Gamma_{L'_0}$  is isomorphic to the holonomy group of the leaf  $L_0 = \pi(L'_0)$  of  $\mathcal{F}$ , it follows that  $\gamma$  is the identity transformation. Hence  $\Gamma$  is trivial and so  $M$  is diffeomorphic to  $P$  whence  $F: M \rightarrow G/K$  is a fibration with the leaves of  $\mathcal{F}$  as fibres.

(2.1) COROLLARY. *Let  $G/K$  be a simply connected, even dimensional, positively curved Riemannian homogeneous space of a compact connected Lie group  $G$  and let  $\mathcal{F}$  be a one-dimensional homogeneous  $G/K$ -foliation of a compact connected manifold  $M$ . Then  $\mathcal{F}$  has a compact leaf and  $\pi_1(M)$  has polynomial growth of degree  $\leq 1$ .*

*Proof.* If  $\mathcal{F}$  is without holonomy, then by (1.1) the leaves of  $\mathcal{F}$  are the fibres of a fibration  $M \rightarrow G/K$  and hence all the leaves are circles. Since  $G/K$  is simply connected, the exact homotopy sequence of this fibration gives a surjection  $\pi_1(S^1) \rightarrow \pi_1(M)$  whence  $\pi_1(M)$  has polynomial growth of degree  $\leq 1$ . If  $\mathcal{F}$  has a leaf  $L$  whose holonomy group is non-trivial, then  $L$  is diffeomorphic to  $S^1$ . Moreover, since  $L$  is compact, the image of its fundamental group in  $\pi_1(M)$  is a subgroup of finite index [2] and hence  $\pi_1(M)$  has polynomial growth of degree  $\leq 1$  [1].

(2.2) PROPOSITION. *Let  $\mathcal{F}$  be a codimension 2 transversely Euclidean (homogeneous  $SO(2) \cdot \mathbb{R}^2 / SO(2) \cong \mathbb{R}^2$ -) foliation of a compact connected manifold  $M$ . If  $\mathcal{F}$  is without holonomy, then  $M$  fibres over  $T^2$ .*

*Proof.* In this case  $\Gamma$  is a subgroup of  $SO(2) \cdot \mathbb{R}^2$ , the group of rigid motions of the plane. Since  $\mathcal{F}$  is without holonomy, it follows that  $\Gamma$  acts freely on  $\mathbb{R}^2$  and hence is a group of translations. If  $(x, y)$  denotes the standard coordinates on  $\mathbb{R}^2$ , then  $dx$  and  $dy$  are linearly independent translation-invariant one-forms and hence there exist smooth linearly independent one-forms  $\omega_1$  and  $\omega_2$  on  $M$  such that  $\pi^*\omega_1 = F^*(dx)$ ,  $\pi^*\omega_2 = F^*(dy)$ . Moreover,  $\omega_1$  and  $\omega_2$  are closed and so  $M$  fibres over  $T^2$  [8].

We now apply the above construction to study the existence of compact leaves for a class of Riemannian homogeneous foliations. Let  $G/K$  be a compact simply connected Riemannian symmetric space with nonzero Euler characteristic where  $G = I_0(G/K)$ .

(2.3) PROPOSITION. *Let  $M$  be a compact manifold with solvable fundamental group. Then every homogeneous  $G/K$ -foliation of  $M$  has a compact leaf.*

*Proof.* The image  $\Gamma$  of the homomorphism  $\Phi : \pi_1(M, x_0) \rightarrow G$  is a solvable subgroup of the compact Lie group  $G$  and so its closure is a compact solvable Lie subgroup of  $G$ . Let  $H$  be the connected component of the identity of  $\bar{\Gamma}$ . Then  $H$  is a toral subgroup. Since the Euler characteristic of  $G/K$  is nonzero,  $G$  and  $K$  have the same rank [9] and hence  $H$  is contained in some conjugate of  $K$ . Thus there exists a point  $y$  of  $G/K$  fixed under the action of  $H$  and hence, since  $H$  is a subgroup of  $\Gamma$  of finite index, the orbit of  $y$  under the action of  $\Gamma$  is a finite set of points  $y = y_1, y_2, \dots, y_r$ . For each  $i = 1, \dots, r$  let  $L_i$  be the unique leaf of  $\pi^{-1}(\mathcal{F})$  such that  $F(L_i) = y_i$ . Then  $L_1, \dots, L_r$  all project via  $\pi$  to the same leaf  $L$  of  $\mathcal{F}$  and, since  $\pi^{-1}(L) = \cup_{i=1}^r L_i$  is a closed subset of  $P$ , it follows that  $L$  is compact.

Recall that by a codimension  $q$  elliptic foliation of a manifold  $M$  we mean a homogeneous  $G/K$ -foliation of  $M$  where  $G = SO(q+1)$ ,  $K = SO(q)$ ,  $G/K \cong S^q$ .

(2.4) COROLLARY. *Every codimension  $2q$  elliptic foliation of a compact manifold with solvable fundamental group has a compact leaf.*

#### REFERENCES

- [ 1 ] H. Bass, The degree of polynomial growth of finitely generated nilpotent groups, Proc. London Math. Soc., (3) **25** (1972), 603–614.
- [ 2 ] R. Blumenthal, Transversely homogeneous foliations, Annales de l'Institut Fourier **29** (1979), 143–158.
- [ 3 ] —, The base-like cohomology of a class of transversely homogeneous foliations, Bulletin des Sciences Mathématiques, **104** (1980), 301–303.
- [ 4 ] R. Hermann, On the differential geometry of foliations, Annals of Math., (3) **72** (1960), 445–457.
- [ 5 ] —, A sufficient condition that a mapping of Riemannian manifolds be a fibre bundle, Proc. Amer. Math. Soc., **11** (1960), 236–242.
- [ 6 ] S. Kobayashi, Transformation groups in differential geometry, Ergebnisse der Mathematik und ihrer Grenzgebiete (70) Springer-Verlag, Berlin, 1972.

- [ 7 ] B. Reinhart, Foliated manifolds with bundle-like metrics, *Annals of Math.*, (1) **69** (1959), 119–132.
- [ 8 ] D. Tischler, On fibering certain foliated manifolds over  $S^1$ , *Topology* **9** (1970), 153–154.
- [ 9 ] J. Wolf, *Spaces of constant curvature*, McGraw-Hill, New York, 1967.

*Université des Sciences et Techniques de Lille I*  
*U.E.R. de Mathématiques Pures et Appliquées*  
*B.P. 36-59650 Villeneuve d'Ascq*  
*France*

*Department of Mathematics*  
*St. Louis University*  
*St. Louis, Missouri 63103*  
*U.S.A.*