

Short Communication

Preserving pathogens for wildlife conservation: a case for action on amphibian declines

JAMIE VOYLES, SCOTT D. CASHINS, ERICA BREE ROSENBLUM
and ROBERT PUSCHENDORF

Abstract Infectious disease is an important driver in biological systems but its importance in conservation has historically been underestimated. Recently, however, researchers have increasingly recognized the impact of diseases on wildlife populations and have grappled with disease-related conservation challenges. For example, the phenomenon of worldwide amphibian declines caused by the fungal disease chytridiomycosis has contributed to the creation of a global Amphibian Conservation Action Plan. The sense of urgency in the protection of amphibians and mitigation of the effects of chytridiomycosis is well-warranted but determining the best way to respond to chytridiomycosis is challenging. Current conservation strategies focus on the preservation of the amphibian hosts, their habitats and their genetic materials. However, we suggest that to confront disease threats fully, particularly in the case of amphibian declines, insight into host–pathogen coevolution may be critical and we must therefore also preserve the pathogen for basic disease research. Here we outline priority targets for virulence research and urge researchers and managers to isolate and archive the pathogen *Batrachochytrium dendrobatidis* to ensure viable long-term amphibian conservation.

Keywords Amphibian declines, *Batrachochytrium dendrobatidis*, chytridiomycosis, pathogen preservation, wildlife disease

Parasites can regulate host population densities, alter community dynamics and affect entire ecosystems (Scott, 1988; Daszak et al., 2000; Hudson et al., 2002; de Castro & Bolker, 2005; Whiles et al., 2006). Disease has obvious consequences for already-threatened animal populations (e.g. black-footed ferrets, Thorne & Williams, 1988)

JAMIE VOYLES (Corresponding author) School of Public Health, Tropical Medicine and Rehabilitation Sciences, Amphibian Disease Ecology Group, James Cook University, Townsville, Queensland 4811, Australia. E-mail jamie.voyles@gmail.com

SCOTT D. CASHINS and ROBERT PUSCHENDORF School of Marine and Tropical Biology, Amphibian Disease Ecology Group, James Cook University, Townsville, Australia.

ERICA BREE ROSENBLUM Department of Biological Sciences, University of Idaho, Moscow, USA.

Received 25 November 2008. Revision requested 22 December 2008.
Accepted 23 January 2009.

but it can also pose a significant threat to otherwise healthy populations (Scott, 1988; Daszak et al., 2000; de Castro & Bolker, 2005). Many amphibians, for example, are experiencing unprecedented declines because of the disease chytridiomycosis, caused by the fungal pathogen *Batrachochytrium dendrobatidis* (Berger et al., 1998; Lips et al., 2006; Mendelson et al., 2006; Gascon et al., 2007). The relatively recent discovery of *B. dendrobatidis* was precipitated by amphibian mass mortality events and disappearances in pristine locations around the world (Berger et al., 1998; Lips et al., 2006; Mendelson et al., 2006; Gascon et al., 2007). Although *B. dendrobatidis* is confined to the superficial layers of epidermis, infected frogs can die rapidly because of the physiological importance of amphibian skin (Berger et al., 1998; Voyles et al., 2007). Current *B. dendrobatidis* distribution patterns generated by predictive models are based on the occurrence of the pathogen on the amphibians (Ron, 2005; Puschendorf et al., 2009). It is unclear whether *B. dendrobatidis* can persist independently of amphibian hosts, in an alternate life stage (Di Rosa et al., 2007), by saprobic growth (Mitchell et al., 2008) or in non-amphibian carriers. In spite of these unresolved possibilities, the loss of biodiversity (Lips et al., 2006; Mendelson et al., 2006), subsequent ecosystem disturbance (Whiles et al., 2006) and the risk for naïve amphibian populations justify immediate conservation efforts.

Management of infectious diseases is not easy; traditional conservation strategies, such as managing reservoir hosts, culling or vaccination, are not practical options where host management is not feasible or where vaccination is unavailable (Hudson et al., 2002). For amphibian conservation, one remediation prescribed by the Amphibian Conservation Action Plan and currently underway is the ‘rescue’ of amphibians from their natural habitats into ex situ breeding programmes before the disease emerges (Mendelson et al., 2006; Gascon et al., 2007). This strategy is controversial (Pounds et al., 2007) and many scientists recognize the conservation crux: re-establishment of amphibians into habitats where the pathogen is now endemic (Pounds et al., 2007; Schlaepfer, 2007). Understanding host–pathogen interactions and coevolution may be critical for long-term conservation. However, with the current focus on preserving vulnerable amphibian hosts and their genetic material (Gascon et al., 2007), the importance of pathogen collection and preservation is being

overlooked. Chytridiomycosis is an important example of how access to pathogen isolates will be invaluable for disease research and conservation efforts.

Determinants of mortality from chytridiomycosis are multiple and complex. Pivotal factors include environmental conditions, population dynamics, host resistance and pathogen virulence. To date, most studies have focused on environmental factors (Woodhams et al., 2003; Berger et al., 2004) and host resistance (Rollins-Smith et al., 2002; Woodhams et al., 2007), whereas differential pathogen virulence has been largely neglected. Preliminary evidence suggests that *B. dendrobatidis* virulence may differ among isolates (Berger et al., 2005; Retallick & Miera, 2007; Fisher et al., 2009) but the underlying reasons for the variation are unknown. Furthermore, reduced mortality has been observed in populations after surviving initial emergence (Morgan et al., 2007) and different populations, even within species, can experience differential susceptibility to chytridiomycosis (Briggs et al., 2005). This variation in susceptibility to *B. dendrobatidis* suggests the possibility of increased host resistance, decreased parasite virulence or both.

Determining the factors that influence disease development will lead to a greater understanding of the disease and thus direct more effective conservation action. For example, we may be able to distinguish a novel pathogen from a pathogen that has increased in virulence (Daszak et al., 2000; Rachowicz et al., 2005), assess potential disease risk (Kuiken et al., 2005), track pathogen movements (Kuiken et al., 2005; Kang et al., 2006) or incorporate disease resistance into breeding programmes (Schlaepfer, 2007). Substantial advancements in virulence research are achievable but depend on access to pathogen isolates from diverse spatio-temporal origins. We suggest three priority targets for pathogen collection and cryopreservation: (1) isolates from geographically and taxonomically diverse hosts, (2) isolates preserved during and following pathogen emergence, and (3) isolates collected at finer scales from populations exhibiting differential responses to infection. Such isolates will be a key resource for laboratory experiments and genetic studies aiming to identify pathogenicity factors (Rosenblum et al., 2008) in chytridiomycosis or for any disease that threatens wildlife populations.

In the case of amphibian declines due to chytridiomycosis, the ability to confront the disease threat fully will hinge on coordinating multiple mitigation strategies including careful screening for the pathogen in naïve populations (Skerratt et al., 2008; Cashins et al., in press) and in the international amphibian trade (Australian Government Department of the Environment and Heritage, 2006; Gascon et al., 2007), identifying and preserving vulnerable amphibians (Mendelson et al., 2006; Gascon et al., 2007) and, most importantly, advancing basic disease research (Australian Government Department of the Environment and Heritage, 2006; Mendelson et al., 2006; Gascon et al.,

2007). Understanding pathogen virulence is important to this endeavour and we urge managers and researchers to participate in a collaborative effort to archive *B. dendrobatidis*. Established methodologies for isolating (Longcore et al., 1999) and cryo-archiving (Boyle et al., 2003) *B. dendrobatidis* are available in multiple languages at the Global *Bd* Banking Project website (2009). In developing this website as an international forum, we aim to share the necessary resources to preserve *B. dendrobatidis* and to collate information on global isolates into a single database to facilitate virulence research. Confronting disease threats and implementing effective conservation action will require a better understanding of the disease, the host and the pathogen.

References

- AUSTRALIAN GOVERNMENT DEPARTMENT OF THE ENVIRONMENT AND HERITAGE (2006) *Threat Abatement Plan: Infection of Amphibians With Chytrid Fungus Resulting in Chytridiomycosis*. Canberra, Commonwealth of Australia. <http://www.environment.gov.au/biodiversity/threatened/publications/tap/chytrid> [accessed 9 November 2008].
- BERGER, L., MARANTELLI, G., SKERRATT, L.F. & SPEARE, R. (2005) Virulence of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* varies with the strain. *Diseases of Aquatic Organisms*, 68, 47–50.
- BERGER, L., SPEARE, R., DASZAK, P., GREEN, D.E., CUNNINGHAM, A.A., GOGGIN, C.L. et al. (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences of the USA*, 95, 9031.
- BERGER, L., SPEARE, R., HINES, H.B., MARANTELLI, G., HYATT, A.D., McDONALD, K.R. et al. (2004) Effect of season and temperature on mortality in amphibians due to chytridiomycosis. *Australian Veterinary Journal*, 82, 434–439.
- BOYLE, D.G., HYATT, A.D., DASZAK, P., BERGER, L., LONGCORE, J.E., PORTER, D. et al. (2003) Cryo-archiving of *Batrachochytrium dendrobatidis* and other chytridiomycetes. *Diseases of Aquatic Organisms*, 56, 59–64.
- BRIGGS, C.J., VREDENBURG, V.T., KNAPP, R.A. & RACHOWICZ, L.J. (2005) Investigating the population-level effects of chytridiomycosis: an emerging infectious disease of amphibians. *Ecology*, 86, 3149–3159.
- CASHINS, S.D., SKERRATT, L.F. & ALFORD, R.A. (in press) Effect of sample collection techniques on sensitivity of real time PCR assay for detecting the amphibian pathogen *Batrachochytrium dendrobatidis*. *Veterinary Microbiology*.
- DASZAK, P., CUNNINGHAM, A.A. & HYATT, A.D. (2000) Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science*, 287, 443–449.
- DE CASTRO, F. & BOLKER, B. (2005) Mechanisms of disease-induced extinction. *Ecology Letters*, 8, 117–126.
- DI ROSA, I., SIMONCELLI, F., FAGOTTI, A. & PASCOLINI, R. (2007) Ecology: the proximate cause of frog declines? *Nature*, 447, E4–E5.
- FISHER, M.C., BOSCH, J., YIN, Z., STEAD, D.A., WALKER, J., SELWAY, L. et al. (2009) Proteomic and phenotypic profiling of the amphibian pathogen *Batrachochytrium dendrobatidis* shows that genotype is linked to virulence. *Molecular Ecology*, 18, 415–429.

- GASCON, C., COLLINS, J.P., MOORE, R.D., CHURCH, D.R., MCKAY, J.E. & MENDELSON, J.R. (2007) *Amphibian Conservation Action Plan, Proceedings: IUCN/SSC Amphibian Conservation Summit 2005*. IUCN, Gland, Switzerland.
- GLOBAL *Bd* BANKING PROJECT (2009) [Http://www.bdbank.org](http://www.bdbank.org) [accessed 6 August 2009].
- HUDSON, P.J., RIZZOLI, A. & BRYAN, G. (2002) *The Ecology of Wildlife Diseases*. Oxford University Press, Oxford, UK.
- KANG, S., BLAIR, J.E., GEISER, D.M., KHANG, C.H., PARK, S.Y., GAHEGAN, M. et al. (2006) Plant pathogen culture collections: it takes a village to preserve these resources vital to the advancement of agricultural security and plant pathology. *Phytopathology*, 96, 920–925.
- KUINEN, T., LEIGHTON, F.A., FOUCHIER, R.A., LEDUC, J.W., PEIRIS, J.S., SCHUDEL, A. et al. (2005) Pathogen surveillance in animals. *Science*, 309, 1680–1681.
- LIPS, K.R., BREM, F., BRENES, R., REEVE, J.D., ALFORD, R.A., VOYLES, J. et al. (2006) Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proceedings of the National Academy of Sciences of the USA*, 103, 3165–3170.
- LONGCORE, J.E., PESSIER, A.P. & NICHOLS, D.K. (1999) *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia*, 91, 219–227.
- MENDELSON, III, J.R., LIPS, K.R., GAGLIARDO, R.W., RABB, G.B., COLLINS, J.P., DIFFENDORFER, J.E. et al. (2006) Confronting amphibian declines and extinctions. *Science*, 313, 48.
- MITCHELL, K.M., CHURCHER, T.S., GARNER, T.W.J. & FISHER, M.C. (2008) Persistence of the emerging pathogen *Batrachochytrium dendrobatidis* outside the amphibian host greatly increases the probability of host extinction. *Proceedings of the Royal Society B*, 275, 329–334.
- MORGAN, J.A.T., VREDENBURG, V.T., RACHOWICZ, L.J., KNAPP, R.A., STICE, M.J., TUNSTALL, T. et al. (2007) Population genetics of the frog-killing fungus *Batrachochytrium dendrobatidis*. *Proceedings of the National Academy of Sciences of the USA*, 104, 13845.
- POUNDS, J.A., CARNAVAL, A.C., PUSCHENDORF, R., HADDAD, C.F.B. & MASTERS, K.L. (2007) Action on amphibian extinctions: going beyond the reductive. *Science* (E-Letter, 28 August 2007), 313.
- PUSCHENDORF, R., CARNAVAL, A.C., VANDERWAL, J., ZUMBADO-ULATE, H., CHAVES, G., BOLAÑOS, F. & ALFORD, R.A. (2009) Distribution models for the amphibian chytrid *Batrachochytrium dendrobatidis* in Costa Rica: proposing climatic refuges as a conservation tool. *Diversity and Distributions*, 15, 401–408.
- RACHOWICZ, L.J., HERO, J.M., ALFORD, R.A., TAYLOR, J.W., MORGAN, J.A.T., VREDENBURG, V.T. et al. (2005) The novel and endemic pathogen hypotheses: competing explanations for the origin of emerging infectious diseases of wildlife. *Conservation Biology*, 19, 1441–1448.
- RETALLICK, R.W.R. & MIERA, V. (2007) Strain differences in the amphibian chytrid *Batrachochytrium dendrobatidis* and non-permanent, sub-lethal effects of infection. *Diseases of Aquatic Organisms*, 75, 201–207.
- ROLLINS-SMITH, L.A., DOERSAM, J.K., LONGCORE, J.E., TAYLOR, S.K., SHAMBLIN, J.C., CAREY, C. & ZASLOFF, M.A. (2002) Antimicrobial peptide defenses against pathogens associated with global amphibian declines. *Developmental and Comparative Immunology*, 26, 63–72.
- RON, S. (2005) Predicting the distribution of the amphibian pathogen *Batrachochytrium dendrobatidis* in the New World. *Biotropica*, 37, 209–221.
- ROSENBLUM, E.B., STAJICH, J.E., MADDOX, N. & EISEN, M.B. (2008) Global gene expression profiles for life stages of the deadly amphibian pathogen *Batrachochytrium dendrobatidis*. *Proceedings of the National Academy of Sciences of the USA*, 105, 17034.
- SCHLAEPFER, M.A. (2007) Is there a role for evolutionary management in addressing the threat of chytridiomycosis? *Science* (E-Letter, 23 March 2007), 313.
- SCOTT, M.E. (1988) The impact of infection and disease on animal populations: implications for conservation biology. *Conservation Biology*, 2, 40–56.
- SKERRATT, L.F., BERGER, L., HINES, H.B., McDONALD, K.R., MENDEZ, D. & SPEARE, R. (2008) Survey protocol for detecting chytridiomycosis in all Australian frog populations. *Ecohealth*, 80, 85–94.
- THORNE, E. & WILLIAMS, E.S. (1988) Disease and endangered species: the black-footed ferret as a recent example. *Conservation Biology*, 2, 66–74.
- VOYLES, J., BERGER, L., YOUNG, S., SPEARE, R., WEBB, R., WARNER, J. et al. (2007) Electrolyte depletion and osmotic imbalance in amphibians with chytridiomycosis. *Diseases of Aquatic Organisms*, 77, 113–118.
- WHILES, M.R., LIPS, K.R., PRINGLE, C.M., KILHAM, S.S., BIXBY, R.J., BRENES, R. et al. (2006) The effects of amphibian population declines on the structure and function of Neotropical stream ecosystems. *Frontiers in Ecology and the Environment*, 4, 27–34.
- WOODHAMS, D.C., ALFORD, R.A. & MARANTELLI, G. (2003) Emerging disease of amphibians cured by elevated body temperature. *Diseases of Aquatic Organisms*, 55, 65–67.
- WOODHAMS, D.C., ARDIPRADJA, K., ALFORD, R.A., MARANTELLI, G., REINERT, L.K. & ROLLINS-SMITH, L.A. (2007) Resistance to chytridiomycosis varies among amphibian species and is correlated with skin peptide defenses. *Animal Conservation*, 10, 409–417.

Biographical sketches

JAMIE VOYLES is interested in ecosystem health and the evolution of virulence in host-pathogen systems, focusing on pathogenesis of chytridiomycosis and virulence of *Batrachochytrium dendrobatidis* in amphibians. SCOTT D. CASHINS studies disease ecology, focusing on transmission and seasonal dynamics of *Batrachochytrium dendrobatidis* in tadpoles in rainforest habitats of North Queensland, Australia. ERICA B. ROSENBLUM's research interests centre on the mechanisms of adaptive evolution, using functional genomics tools to elucidate the genetic basis of pathogenicity of chytridiomycosis. ROBERT PUSCHENDORF is interested in disease ecology and the influence of environmental factors on host-pathogen interactions, especially on dynamics of *Batrachochytrium dendrobatidis* across habitats in Costa Rica and Australia.