



COMPOSITIO MATHEMATICA

Martin boundaries of the duals of free unitary quantum groups

Sara Malacarne and Sergey Neshveyev

Compositio Math. **155** (2019), 1171–1193.

[doi:10.1112/S0010437X19007322](https://doi.org/10.1112/S0010437X19007322)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Martin boundaries of the duals of free unitary quantum groups

Sara Malacarne and Sergey Neshveyev

ABSTRACT

Given a free unitary quantum group $G = A_u(F)$, with F not a unitary 2×2 matrix, we show that the Martin boundary of the dual of G with respect to any G - \hat{G} -invariant, irreducible, finite-range quantum random walk coincides with the topological boundary defined by Vaes and Vander Venet. This can be thought of as a quantum analogue of the fact that the Martin boundary of a free group coincides with the space of ends of its Cayley tree.

Introduction

The study of harmonic functions on trees has a long history. In the early 1960s Dynkin and Malyutov [DM61] considered nearest-neighbor random walks on free groups and obtained an analogue of the Poisson formula for them by identifying the Martin boundary of such a group with the space of ends of its Cayley graph. This result was then generalized by a number of authors, among others by Cartier [Car72], who considered nearest-neighbor, but not necessarily homogeneous, random walks on trees. Finite-range random walks were subsequently studied by Derriennic [Der75] in the homogeneous case and by Picardello and Woess [PW87] in general. In both cases the result was the same as before: the Martin boundary of a tree coincides with its space of ends. This was later generalized to hyperbolic graphs by Ancona [Anc88] who considered finite-range random walks on such graphs and showed that (under mild assumptions) the corresponding Martin boundaries coincide with the Gromov boundaries. A related result was also obtained by Kaimanovich [Kai00] who studied only bounded harmonic functions on hyperbolic groups but went beyond finite-range random walks.

The natural quantum analogues of free groups are the duals of free quantum groups of Van Daele and Wang [VDW96]. An interpretation of these duals as vertices of some quantum trees was proposed by Vergnioux [Ver05], and an analogue of the end compactification was defined by Vaes and Vergnioux [VV07] in the free orthogonal case and by Vaes and Vander Venet in the free unitary case [VVV10]. The duals of free orthogonal quantum groups behave in many respects like the group \mathbb{Z} , so the corresponding tree has valency 2, but the quantum boundary is less trivial than this analogy might suggest; for example, for the dual of $SU_q(2)$ ($q \neq \pm 1$) it is the Podleś quantum sphere S_q^2 . An identification of the quantum Poisson and Martin boundaries with the quantum spaces of ends was obtained in [Izu02, NT04, VV07]; see also the earlier paper by Biane [Bia94] on the dual of $SU(2)$, which to a great extent motivated the development of the quantum theory.

Received 2 July 2018, accepted in final form 21 December 2018, published online 31 May 2019.

2010 Mathematics Subject Classification 20G42, 46L65, 60J50 (primary).

Keywords: quantum group, Martin boundary, Gromov boundary, random walks on trees.

The research leading to these results received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement no. 307663.

This journal is © Foundation Compositio Mathematica 2019.

The duals of free unitary quantum groups behave more like the free group with two generators. In this case Vaes and Vander Venet showed [VVV10] that the Poisson boundary is isomorphic to the quantum boundary they defined, or more precisely, that the von Neumann algebra of bounded harmonic functions is isomorphic to the von Neumann algebra generated by the C^* -algebra of continuous functions on the boundary in the GNS-representation defined by a particular state. Classically this would mean a computation of the Martin boundary as a measure space rather than as a topological space, where the measure is defined as the hitting distribution of the random walk. The goal of the present paper is to show that the Martin boundary of the dual of a free unitary quantum group does coincide with the boundary defined in [VVV10] topologically, that is, we have an equality of C^* -algebras.

It should be added that a precise relation between quantum Poisson and Martin boundaries has not been established. It was shown in [NT04] that for the dual of $SU_q(2)$ the computation of the Martin boundary allows one to easily identify the Poisson boundary as well. This was also recently confirmed in [Jor17] by a different argument, and the same is true for the duals of free orthogonal quantum groups. But a general result of this sort is not available yet, so we cannot claim that our result automatically covers the computation of the Poisson boundary in [VVV10].

If one wants to compute the Martin boundary of the dual a free unitary quantum group $G = A_u(F)$, it is natural to try to follow the strategy used in [VV07] in the free orthogonal case. However, this requires more precise information on the Green functions of particular classical random walks on trees than seems to be available. We propose a different approach, which also reduces the computation to a classical problem (and works equally well in the free orthogonal case). The idea is that by considering the spectral subspaces of $\ell^\infty(\hat{G})$ with respect to the adjoint action of G , we see that a quantum random walk is determined by a collection of operators on the ℓ^2 -spaces of branches of the classical tree $\text{Irr}(G)$. It turns out that the matrix coefficients of any two such operators corresponding to the common branch of definition are exponentially small perturbations of each other (with respect to the distance to the root of $\text{Irr}(G)$). Our main technical result roughly says that the corresponding classical Green functions all have the same asymptotics at infinity. While this is probably not surprising, we point out that the tree $\text{Irr}(G)$ has exponential growth, so it is not immediately obvious that even an exponentially small perturbation is enough to compensate for this growth.

1. Preliminaries

1.1 Compact and discrete quantum groups

In this section we recall the basic notions related to compact and discrete quantum groups; see, for example, [NT13] for more details.

A *compact quantum group* G is given by a unital C^* -algebra $C(G)$ and a unital $*$ -homomorphism

$$\Delta: C(G) \rightarrow C(G) \otimes C(G)$$

such that $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$ and the spaces $(C(G) \otimes 1)\Delta(C(G))$ and $(1 \otimes C(G))\Delta(C(G))$ are dense in $C(G) \otimes C(G)$.

A *finite-dimensional unitary representation* of G is a pair (U, H_U) consisting of a finite-dimensional Hilbert space H_U and a unitary element $U \in B(H_U) \otimes C(G)$ such that $\Delta(U) = U_{12}U_{13}$. The tensor product of two representations U and V is defined by

$$U \otimes V = U_{13}V_{23} \in B(H_U) \otimes B(H_V) \otimes C(G).$$

With this tensor product, the finite-dimensional unitary representations of G form a C^* -tensor category $\text{Rep } G$.

The space $\mathbb{C}[G]$ of matrix coefficients of all finite-dimensional unitary representations of G is a Hopf $*$ -algebra with comultiplication Δ and antipode S such that $(\iota \otimes S)(U) = U^*$ for all $U \in \text{Rep } G$. The dual space $\mathcal{U}(G) = \mathbb{C}[G]^*$ is a $*$ -algebra with multiplication dual to Δ , so that $\omega\eta = (\omega \otimes \eta)\Delta$, and involution $\omega^* = \bar{\omega}S$, where $\bar{\omega}(a) = \overline{\omega(a^*)}$. This is by definition the algebra of all (unbounded) functions on the *dual discrete quantum group* $\Gamma = \hat{G}$.

More concretely, $\mathcal{U}(G)$ can be described as follows. For every $U \in \text{Rep } G$ we have a $*$ -representation $\pi_U: \mathcal{U}(G) \rightarrow B(H_U)$ defined by $\pi_U(\omega) = (\iota \otimes \omega)(U)$. Consider the set $I = \text{Irr}(G)$ of isomorphism classes of irreducible unitary representations of G . For every $s \in I$ choose a representative U_s . We write π_s, H_s , etc., instead of π_{U_s}, H_{U_s} . Then the collection of representations $(\pi_s)_{s \in I}$ defines a $*$ -isomorphism $\mathcal{U}(G) \cong \prod_{s \in I} B(H_s)$. The subalgebra $\ell^\infty(\Gamma) \subset \mathcal{U}(G)$ corresponding to $\ell^\infty\text{-}\bigoplus_{s \in I} B(H_s)$ under this isomorphism is, by definition, the algebra of bounded functions on Γ . We will not distinguish between the algebras $\ell^\infty(\Gamma)$ and $\ell^\infty\text{-}\bigoplus_{s \in I} B(H_s)$. We will also consider the subalgebras

$$c_0(\Gamma) = c_0\text{-}\bigoplus_{s \in I} B(H_s) \quad \text{and} \quad c_c(\Gamma) = \bigoplus_{s \in I} B(H_s)$$

of functions vanishing at infinity and of finitely supported functions, respectively.

By duality the product on $\mathbb{C}[G]$ defines a comultiplication

$$\hat{\Delta}: \mathcal{U}(G) \rightarrow \mathcal{U}(G \times G) := (\mathbb{C}[G] \otimes \mathbb{C}[G])^* \cong \prod_{s,t \in I} B(H_s) \otimes B(H_t).$$

By restriction it gives a map $\hat{\Delta}: \ell^\infty(\Gamma) \rightarrow \ell^\infty(\Gamma) \bar{\otimes} \ell^\infty(\Gamma)$.

The antipode S on $\mathbb{C}[G]$ defines an antipode \hat{S} on $\mathcal{U}(G)$ by $\hat{S}(\omega) = \omega S$. There exists a unique element $\rho \in \mathcal{U}(G)$, called the *Woronowicz character*, such that ρ is positive, invertible, $\hat{S}^2 = \text{Ad } \rho$ and $\text{Tr}(\pi_U(\rho)) = \text{Tr}(\pi_U(\rho^{-1}))$ for all $U \in \text{Rep } G$. It also has the property $\hat{\Delta}(\rho) = \rho \otimes \rho$ justifying the name ‘character’. The number $\text{Tr}(\pi_U(\rho))$ is called the *quantum dimension* of U and denoted by $\text{dim}_q(U)$. We will write $\text{dim}_q(s)$ instead of $\text{dim}_q(U_s)$.

The element ρ allows one to define a canonical duality functor on $\text{Rep } G$. Namely, given a finite-dimensional unitary representation U of G , the *conjugate representation* is defined by

$$\bar{U} = (j(\pi_U(\rho^{1/2})) \otimes 1)(j \otimes \iota)(U^*)(j(\pi_U(\rho^{-1/2})) \otimes 1) \in B(\bar{H}_U) \otimes C(G),$$

where \bar{H}_U is the complex conjugate Hilbert space to H_U and j is the canonical $*$ -anti-isomorphism $B(H_U) \rightarrow B(\bar{H}_U)$, $j(T)\bar{\xi} = \overline{T^*\xi}$. Note that the element $U^c = (j \otimes \iota)(U^*)$ already satisfies $(\iota \otimes \Delta)(U^c) = U_{12}^c U_{13}^c$, and the additional conjugation by $j(\pi_U(\rho^{1/2})) \otimes 1$ is needed only to make this element unitary.

The object \bar{U} is conjugate to U in $\text{Rep } G$, meaning that there exist morphisms $R_U: \mathbb{1} \rightarrow \bar{U} \otimes U$ and $\bar{R}_U: \mathbb{1} \rightarrow U \otimes \bar{U}$, where $\mathbb{1}$ is the trivial representation of G on the one-dimensional space \mathbb{C} , solving the *conjugate equations*: the compositions

$$U \xrightarrow{\iota \otimes R_U} U \otimes \bar{U} \otimes U \xrightarrow{\bar{R}_U^* \otimes \iota} U \quad \text{and} \quad \bar{U} \xrightarrow{\iota \otimes \bar{R}_U} \bar{U} \otimes U \otimes \bar{U} \xrightarrow{R_U^* \otimes \iota} \bar{U}$$

are the identity morphisms. Using the Woronowicz character ρ , we can define such morphisms by

$$R_U(1) = \sum_i \bar{\xi}_i \otimes \pi_U(\rho^{-1/2})\xi_i, \quad \bar{R}_U(1) = \sum_i \pi_U(\rho^{1/2})\xi_i \otimes \bar{\xi}_i,$$

where $\{\xi_i\}_i$ is an orthonormal basis in H_U . The pair (R_U, \bar{R}_U) is *standard*, meaning that $\|R_U\| = \|\bar{R}_U\|$ and the product $\|R_U\| \|\bar{R}_U\|$ is minimal among all possible solutions of the conjugate equations for U and \bar{U} . Note that $\|R_U\| = \|\bar{R}_U\| = \dim_q(U)^{1/2}$.

1.2 Quantum random walks

We now recall the main concepts of the boundary theory of quantum random walks; see [Izu02, NT04].

For every $s \in I = \text{Irr}(G)$, consider the state ϕ_s on $B(H_s)$ defined by

$$\phi_s = \frac{\text{Tr}(\cdot \pi_s(\rho^{-1}))}{\dim_q(s)}.$$

This is a unique state invariant under the left action of G on $B(H_s)$ given by

$$B(H_s) \rightarrow C(G) \otimes B(H_s), \quad T \mapsto (U_s)_{21}^*(1 \otimes T)(U_s)_{21}.$$

Taken together, these actions define the *left adjoint action* of G on $\ell^\infty(\Gamma)$.

Consider now a probability measure μ on I and define a normal state ϕ_μ on $\ell^\infty(\Gamma)$ by

$$\phi_\mu = \sum_{s \in I} \mu(s) \phi_s.$$

Consider the corresponding convolution Markov operator P_μ on $\ell^\infty(\Gamma)$:

$$P_\mu = (\phi_\mu \otimes \iota) \hat{\Delta}.$$

This operator commutes with the left adjoint action of G and the right action by translations of Γ on $\ell^\infty(\Gamma)$, which is defined by $\hat{\Delta}$. We therefore say that P_μ defines a *G-Γ-invariant quantum random walk* on Γ .

By the G -equivariance, the operator P_μ preserves the center $\ell^\infty(I)$ of $\ell^\infty(\Gamma)$, hence it defines a classical random walk on the set I with transition probabilities $p_\mu(s, t)$ such that

$$P_\mu(I_t)I_s = p_\mu(s, t)I_s, \tag{1}$$

where I_s denotes the unit of $B(H_s) \subset \ell^\infty(\Gamma)$. Explicitly, the transition probabilities are given by

$$p_\mu(s, t) = \sum_{r \in I} \mu(r) m_{rs}^t \frac{\dim_q(t)}{\dim_q(r) \dim_q(s)}, \tag{2}$$

where m_{rs}^t is the multiplicity of U_t in $U_r \otimes U_s$. We will only be interested in the case where this classical random walk is *irreducible* and *transient*. The irreducibility assumption means that, given s, t , there exist $s_0 = s, s_1, \dots, s_n = t$ such that $p_\mu(s_0, s_1) \cdots p_\mu(s_{n-1}, s_n) > 0$. Equivalently, every representation U_t is a subrepresentation of $U_{r_1} \otimes \cdots \otimes U_{r_n}$ for some $r_1, \dots, r_n \in \text{supp } \mu$. In this case we also say that the measure μ and the state ϕ_μ are *generating*. The assumption of transience means that the sum of all such products $p_\mu(s_0, s_1) \cdots p_\mu(s_{n-1}, s_n)$ (with $s_0 = s$ and $s_n = t$) is finite for all $s, t \in I$. In this case we also say that ϕ_μ and μ are *transient*. Similarly to [DG73], it can be shown that, assuming irreducibility, the random walk is transient if the quantum dimension function is non-amenable; in particular, when Γ is either non-amenable or non-Kac (recall that the latter means that $\rho \neq 1$). We will prove a related result more useful for our purposes in Lemma 4.1 below.

Assuming irreducibility and transience, the *Green kernel* is defined as the completely positive map

$$G_\mu: c_c(\Gamma) \rightarrow \ell^\infty(\Gamma), \quad G_\mu(a) = \sum_{n=0}^\infty P_\mu^n(a),$$

and then the *Martin kernel* is defined as the map

$$K_\mu: c_c(\Gamma) \rightarrow \ell^\infty(\Gamma), \quad K_\mu(a) = G_\mu(I_e)^{-1}G_\mu(a),$$

where $e \in I$ is the point corresponding to the trivial representation $\mathbf{1}$. We remark that the element $G_\mu(I_e)$ is a strictly positive scalar in every matrix block $B(H_s)$, so $G_\mu(I_e)^{-1}$ is a well-defined, but possibly unbounded, function on I . The elements $K_\mu(a) = G_\mu(I_e)^{-1}G_\mu(a)$ are, however, bounded. Furthermore, if $a \in c_c(\Gamma)$ is positive and non-zero, then $K_\mu(a)$ is a positive invertible element of $\ell^\infty(\Gamma)$.

We have an involution $s \mapsto \bar{s}$ on I defined by $\bar{U}_s \cong U_{\bar{s}}$. For every probability measure μ on I , we define a new probability measure $\check{\mu}$ on I by $\check{\mu}(s) = \mu(\bar{s})$. The corresponding transition probabilities satisfy

$$p_{\check{\mu}}(s, t) = \frac{\dim_q(t)^2}{\dim_q(s)^2} p_\mu(t, s). \tag{3}$$

It follows that $\check{\mu}$ is generating and transient if and only if μ has the same properties.

Finally, given a generating transient probability measure μ , the *Martin compactification* of Γ is defined as the C*-subalgebra $C(\bar{\Gamma}_{M,\mu})$ of $\ell^\infty(\Gamma)$ generated by $K_{\check{\mu}}(c_c(\Gamma))$ and $c_c(\Gamma)$. This C*-algebra is invariant under the left adjoint action of G and the right action of Γ by translations. The *Martin boundary* is defined by $C(\partial\Gamma_{M,\mu}) = C(\bar{\Gamma}_{M,\mu})/c_0(\Gamma)$.

1.3 Module categories and categorical random walks

To every C*-algebra A equipped with a left action of G one can associate the category of equivariant finitely generated Hilbert modules. We will need this only for unital C*-subalgebras $A \subset \ell^\infty(\Gamma)$, equipped with the action given by the restriction of the adjoint action, in which case this category has the following concrete description [NY14, § 4].

For $U, V \in \text{Rep } G$, consider the subspace

$$\begin{aligned} \mathcal{D}_\Gamma(U, V) &:= \ell^\infty\text{-}\bigoplus_{s \in I} \text{Hom}_G(H_s \otimes H_U, H_s \otimes H_V) \\ &\subset \ell^\infty\text{-}\bigoplus_{s \in I} B(H_s \otimes H_U, H_s \otimes H_V) = \ell^\infty(\Gamma) \otimes B(H_U, H_V), \end{aligned}$$

and define $\mathcal{D}_A(U, V)$ as the intersection of this space with $A \otimes B(H_U, H_V)$. We can then define a new category \mathcal{D}_A with the same objects as in $\text{Rep } G$, but with the morphism spaces $\mathcal{D}_A(U, V)$. (To be more precise, we also need to complete \mathcal{D}_A with respect to subobjects, but whether we do this or not is not going to play any role in this paper.) The category $\text{Rep } G$ can be considered as a (non-full) subcategory of \mathcal{D}_A , if we identify a morphism $T: H_U \rightarrow H_V$ in $\text{Rep } G$ with the collection of morphisms $\iota_s \otimes T: H_s \otimes H_U \rightarrow H_s \otimes H_V$. Furthermore, the functors of tensoring on the right by objects of $\text{Rep } G$ define on \mathcal{D}_A the structure of a right $(\text{Rep } G)$ -module category.

The morphism spaces $\mathcal{D}_A(U, V)$ are completely determined by the spaces of the form $\mathcal{D}_A(\mathbf{1}, U)$ thanks to the Frobenius reciprocity isomorphisms

$$\mathcal{D}_A(U, V) \cong \mathcal{D}_A(\mathbf{1}, V \otimes \bar{U}), \quad T \mapsto (T \otimes \iota_{\bar{U}})\bar{R}_U. \tag{4}$$

By decomposing $V \otimes \bar{U}$ into irreducibles we see that \mathcal{D}_A is completely determined already by the spaces $\mathcal{D}_A(\mathbf{1}, U_s)$, $s \in I$.

The G - C^* -algebra A can be easily reconstructed from the morphism spaces $\mathcal{D}_A(U, V)$: the elements of the form $(\iota \otimes \omega)(T)$, where $T \in \mathcal{D}_A(U, V) \subset \ell^\infty(\Gamma) \otimes B(H_U, H_V)$ and $\omega \in B(H_U, H_V)^*$, span a dense $*$ -subalgebra of A . By the previous paragraph it is clear that it suffices to take $T \in \mathcal{D}_A(\mathbb{1}, U_s)$, $s \in I$. For each $s \in I$, the linear span of elements of the form $(\iota \otimes \omega)(T)$, with $T \in \mathcal{D}_A(\mathbb{1}, U_s)$ and $\omega \in B(H_e, H_e \otimes H_s)^* = H_s^*$, is nothing other than the spectral subspace of A corresponding to \bar{U}_s .

Assume now that we are given a probability measure μ on I and consider the corresponding Markov operator P_μ on $\ell^\infty(\Gamma)$. Since P_μ commutes with left adjoint action of G , the operator $P_\mu \otimes \iota$ is well defined on $\mathcal{D}_\Gamma(U, V) \subset \ell^\infty(\Gamma) \otimes B(H_U, H_V)$. We will denote this operator by P_μ again. This operator can be described in purely categorical terms as follows.

The space $\mathcal{D}_\Gamma(U, V)$ can be identified with the space $\text{Nat}_b(\iota \otimes U, \iota \otimes V)$ of bounded natural transformations between the functors $\iota \otimes U$ and $\iota \otimes V$ on $\text{Rep } G$, that is, uniformly bounded collections $(\eta_W)_W$ of morphisms $\eta_W: W \otimes U \rightarrow W \otimes V$ that are natural in W , since any such collection is completely determined by the morphisms $\eta_s = \eta_{U_s}$. For every $s \in I$, we have a contraction P_s on $\text{Nat}_b(\iota \otimes U, \iota \otimes V)$ defined by

$$P_s(\eta)_W = (\text{tr}_s \otimes \iota)(\eta_{U_s \otimes W}).$$

where $\text{tr}_s \otimes \iota$ denotes the *normalized categorical partial trace* $\text{Mor}(U_s \otimes X, U_s \otimes Y) \rightarrow \text{Mor}(X, Y)$ defined by

$$(\text{tr}_s \otimes \iota)(T) = \dim_q(s)^{-1} (R_s^* \otimes \iota_Y)(\iota_{\bar{s}} \otimes T)(R_s \otimes \iota_X),$$

and where (R_s, \bar{R}_s) is a standard solution of the conjugate equations for U_s and $U_{\bar{s}}$. Then

$$P_\mu = \sum_{s \in I} \mu(s) P_s.$$

This leads to the following categorical description of the Martin compactification $C(\bar{\Gamma}_{M, \mu})$ in [Jor17]; see also [NY17] for the initial motivation and [DRVV10] for a precursor of this picture.

Consider the subspace $\text{Nat}_c(\iota \otimes U, \iota \otimes V) \subset \text{Nat}_b(\iota \otimes U, \iota \otimes V)$ consisting of all natural transformations η such that $\eta_s = 0$ for all but finitely many s . In other words, we consider the subspace

$$\mathcal{D}_{\Gamma, c}(U, V) := \bigoplus_{s \in I} \text{Hom}_G(H_s \otimes H_U, H_s \otimes H_V) \subset \ell^\infty - \bigoplus_{s \in I} \text{Hom}_G(H_s \otimes H_U, H_s \otimes H_V).$$

We have well-defined maps

$$K_\mu: \mathcal{D}_{\Gamma, c}(U, V) \rightarrow \mathcal{D}_\Gamma(U, V), \quad K_\mu(\eta) = G_\mu(I_e)^{-1} G_\mu(\eta),$$

where $G_\mu(\eta) = \sum_{n=0}^\infty P_\mu^n(\eta)$. Then $\mathcal{D}_{C(\bar{\Gamma}_{M, \mu})}$ is the smallest C^* -subcategory of \mathcal{D}_Γ containing $\text{Rep } G$, $\mathcal{D}_{\Gamma, c}$ and all morphisms of the form $K_{\bar{\mu}}(\eta)$, where η is a morphism in $\mathcal{D}_{\Gamma, c}$. Again using (4), we see that it suffices to take $\eta \in \mathcal{D}_{\Gamma, c}(\mathbb{1}, U_s)$, $s \in I$.

Note that the previous paragraph does not really add any new information on $C(\bar{\Gamma}_{M, \mu})$. What we are basically saying is that in computing $K_{\bar{\mu}}: c_c(\Gamma) \rightarrow \ell^\infty(\Gamma)$ it suffices to consider elements a lying in spectral subspaces of $c_c(\Gamma)$ (with respect to the left adjoint action of G) corresponding to U_s , and then $K_{\bar{\mu}}(a)$ is again an element of the spectral subspace of $\ell^\infty(\Gamma)$ corresponding to U_s . But this is clear, since $c_c(\Gamma)$ is the direct sum of its spectral subspaces and the operator P_μ is G -equivariant. Nevertheless, working with the spaces $\mathcal{D}_\Gamma(\mathbb{1}, U_s)$ is more convenient than with the spectral subspaces, and the interpretation of elements of $\mathcal{D}_\Gamma(\mathbb{1}, U_s)$ as natural transformations between certain functors makes some computations more transparent.

1.4 Free unitary quantum groups

In this section we recall some properties of free unitary quantum groups introduced in [VDW96] and studied in detail in [Ban97].

Fix a natural number $n \geq 2$ and a matrix $F \in GL_n(\mathbb{C})$ such that $\text{Tr}(F^*F) = \text{Tr}((F^*F)^{-1})$. The compact *free unitary quantum group* $A_u(F)$ is defined as the universal unital C^* -algebra with generators u_{ij} , $1 \leq i, j \leq n$, such that the matrices $U = (u_{ij})_{i,j}$ and FU^cF^{-1} are unitary, where $U^c = (u_{ij}^*)_{i,j}$, equipped with the comultiplication

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

Therefore U defines a unitary representation of the quantum group $A_u(F)$ on the n -dimensional Hilbert space, called the *fundamental representation*.

The set I of isomorphism classes of irreducible representations of $A_u(F)$ is the free monoid on letters α and β , with α corresponding to U . The involution $s \mapsto \bar{s}$ is the anti-automorphism of the monoid defined by $\bar{\alpha} = \beta$ and $\bar{\beta} = \alpha$.

From now on we will use the conventions of [VVV10] in that we write x instead of U_x for $x \in I$ whenever convenient. The fusion rules for the representations of $A_u(F)$ are given by

$$x \otimes y \cong \bigoplus_{z \in I: x=x_0z, y=\bar{z}y_0} x_0y_0.$$

Therefore if the last letter of x is the same as the first letter of y , then $U_x \otimes U_y$ is irreducible and isomorphic to U_{xy} . In this case we write $xy = x \otimes y$.

The Woronowicz character ρ of $A_u(F)$ is determined by the property

$$\pi_U(\rho) = (F^*F)^t \quad (\text{the transpose of } F^*F).$$

Clearly, $\dim_q(\alpha) = \text{Tr}(F^*F) \geq n$, and the equality holds if and only if F is unitary. Let $q \in (0, 1]$ be such that

$$\text{Tr}(F^*F) = q + q^{-1}. \tag{5}$$

Then $q = 1$ if and only if F is a unitary 2×2 matrix.

An element $x \in I$ is said to be *indecomposable* if there are only trivial decompositions $x = y \otimes z$, that is, we must have $y = e$ or $z = e$. Equivalently, x is an alternating product of α and β . Every word $x \in I$ can be written as $x_1 \otimes \dots \otimes x_n$, where x_1, \dots, x_n are indecomposable. Then

$$\dim_q(x) = \prod_{i=1}^n [|x_i| + 1]_q, \tag{6}$$

where $|x_i|$ denotes the length of x_i and $[n]_q = (q^n - q^{-n}) / (q - q^{-1})$.

2. Topological boundary of the dual of $A_u(F)$

Consider a free unitary quantum group $G = A_u(F)$, with F not a unitary 2×2 matrix. Recall that this assumption means that the number $q \in (0, 1]$ defined by (5) is strictly less than 1. Denote by Γ the dual discrete quantum group.

As we discussed in § 1.4, the set I of isomorphism classes of irreducible representations of G is a free monoid on letters α and β . The empty word is denoted by e . Consider the tree with

vertex set I such that different elements x and y of I are connected by an edge if and only if one of them is obtained from the other by adding (or removing) one letter on the left. Denote by \bar{I} the corresponding end compactification of I . The elements of \bar{I} are words in α and β that are either finite or infinite on the left, and the boundary $\partial I = \bar{I} \setminus I$ is the set of infinite words. The algebra $C(\bar{I})$ of continuous functions on \bar{I} can be identified with the algebra of functions $f \in \ell^\infty(I)$ such that

$$|f(yx) - f(x)| \rightarrow 0 \text{ as } x \rightarrow \infty, \text{ uniformly in } y \in I.$$

In [VVV10], Vaes and Vander Venet extended this construction to $\ell^\infty(\Gamma)$ as follows. (More precisely, they consider words infinite on the right, while in order to be consistent with our conventions, we consider words infinite on the left.)

For all $x, y \in I$, fix an isometry $V(xy, x \otimes y) \in \text{Mor}(xy, x \otimes y)$. Define ucp maps

$$\psi_{yx,x}: B(H_x) \rightarrow B(H_{yx}) \quad T \mapsto V(yx, y \otimes x)^*(1 \otimes T)V(yx, y \otimes x).$$

They do not depend on any choices. Define

$$\mathcal{B} = \{a \in \ell^\infty(\Gamma) : \|a_{yx} - \psi_{yx,x}(a_x)\| \rightarrow 0 \text{ as } x \rightarrow \infty, \text{ uniformly in } y \in I\}. \tag{7}$$

By [VVV10, Theorem 3.2], this is a unital C^* -subalgebra of $\ell^\infty(\Gamma)$ containing $c_0(\Gamma)$. It can therefore be considered as the algebra of continuous functions on a compactification of Γ . The (non-commutative) algebra of continuous functions on the corresponding boundary is defined by $\mathcal{B}_\infty = \mathcal{B}/c_0(\Gamma)$. The left adjoint action of G and the right action by translations of Γ on $\ell^\infty(\Gamma)$ define actions of G and Γ on \mathcal{B} and \mathcal{B}_∞ .

If we view \mathcal{B} only as a G - C^* -algebra, then from our discussion in § 1.3 we immediately get the following result.

PROPOSITION 2.1. *Consider the module category $\mathcal{D}_{\mathcal{B}}$ associated with the left action of G on \mathcal{B} . Then, for all $U, V \in \text{Rep } G$, the morphism space $\mathcal{D}_{\mathcal{B}}(U, V)$ consists of all elements*

$$T \in \ell^\infty\text{-}\bigoplus_{x \in I} \text{Hom}_G(H_x \otimes H_U, H_x \otimes H_V)$$

such that $\|T_{yx} - (V(yx, y \otimes x)^* \otimes \iota_V)(\iota_y \otimes T_x)(V(yx, y \otimes x) \otimes \iota_U)\| \rightarrow 0$ as $x \rightarrow \infty$, uniformly in $y \in I$.

In order to get a better understanding of the morphism spaces $\mathcal{D}_{\mathcal{B}}(U, V)$ we will now make a particular choice of representations U_x and isometries $V(z, x \otimes y) \in \text{Mor}(z, x \otimes y)$ for all $z \prec x \otimes y$; cf. [VV07, Appendix].

Fix representatives U_α and U_β of α and β . All other representatives U_x we construct as follows. If $x = x_1 \cdots x_n \in I$, with $x_i \in \{\alpha, \beta\}$ for all i , then, since $\text{Mor}(x, x_1 \otimes \cdots \otimes x_n)$ is one-dimensional, $H_{x_1} \otimes \cdots \otimes H_{x_n}$ contains a unique invariant subspace on which $U_{x_1} \otimes \cdots \otimes U_{x_n}$ is irreducible of class x . We take this subspace as H_x and define U_x as the restriction of $U_{x_1} \otimes \cdots \otimes U_{x_n}$ to H_x . Denote by p_x the projection in $B(H_{x_1} \otimes \cdots \otimes H_{x_n})$ with image H_x .

Note that the fusion rules and our choice of representatives imply that H_{xy} is a subspace of $H_x \otimes H_y$. We take the embedding map $H_{xy} \rightarrow H_x \otimes H_y$ as $V(xy, x \otimes y)$. Then $V(xy, x \otimes y)^*$ coincides with the projection $p_{xy}^{x \otimes y} := p_{xy}|_{H_x \otimes H_y}$.

Next, viewing U_β as the dual of U_α , fix a standard solution $(R_\alpha, \bar{R}_\alpha)$ of the conjugate equations for U_α . Put $R_\beta = \bar{R}_\alpha$ and $\bar{R}_\beta = R_\alpha$. Assume now that $x = x_1 \cdots x_n$ and $y = y_1 \cdots y_m$,

with $x_i, y_j \in \{\alpha, \beta\}$ for all i and j , and assume that z is a subrepresentation of $x \otimes y$, so $z = x_1 \cdots x_{n-k} y_{k+1} \cdots y_m$ for some k and $x_{n-i} = \bar{y}_{i+1}$ for $i = 0, \dots, k - 1$. We then define an operator $\tilde{V}(z, x \otimes y): H_z \rightarrow H_x \otimes H_y$ by

$$\tilde{V}(z, x \otimes y) = (p_x \otimes p_y)(\iota^{\otimes(n-1)} \otimes \bar{R}_{x_{n-k+1}} \otimes \iota^{\otimes(m-1)}) \dots (\iota^{\otimes(n-k)} \otimes \bar{R}_{x_n} \otimes \iota^{\otimes(m-k)}).$$

This can also be described as follows. Our fixed standard solutions $(R_\alpha, \bar{R}_\alpha)$ and $(R_\beta, \bar{R}_\beta) = (\bar{R}_\alpha, R_\alpha)$ of the conjugate equations allow us to construct standard solutions of the conjugate equations for tensor products of U_α and U_β . By restriction we then get standard solutions (R_x, \bar{R}_x) of the conjugate equations for all $x \in I$. By construction, we have $(R_{\bar{x}}, \bar{R}_{\bar{x}}) = (\bar{R}_x, R_x)$ and

$$R_{xy} = (p_{\bar{y}\bar{x}}^{\otimes \bar{x}} \otimes p_{xy}^{\otimes y})(\iota_{\bar{y}} \otimes R_x \otimes \iota_y)R_y, \quad \bar{R}_{xy} = (p_{xy}^{\otimes y} \otimes p_{\bar{y}\bar{x}}^{\otimes \bar{x}})(\iota_x \otimes \bar{R}_y \otimes \iota_{\bar{x}})\bar{R}_x. \tag{8}$$

Then the morphism $\tilde{V}(z, x \otimes y)$ for $x = sv, y = \bar{v}t$ and $z = st$ is given by

$$\tilde{V}(st, sv \otimes \bar{v}t) = (p_{sv}^{s \otimes v} \otimes p_{\bar{v}t}^{\bar{v} \otimes t})(\iota_s \otimes \bar{R}_v \otimes \iota_t)V(st, s \otimes t). \tag{9}$$

Since $\tilde{V}(st, sv \otimes \bar{v}t)$ is an element of $\text{Mor}(st, sv \otimes \bar{v}t)$, it must be isometric up to a scalar factor.

LEMMA 2.2. *There is a constant $c > 0$ depending only on q such that*

$$\dim_q(v)^{1/2} \geq \|\tilde{V}(st, sv \otimes \bar{v}t)\| \geq c \dim_q(v)^{1/2}$$

for all $v, s, t \in I$.

Proof. The first inequality is an immediate consequence of (9), since $\|\bar{R}_v\| = \dim_q(v)^{1/2}$.

The second inequality can be deduced from an analogous result for $\text{SU}_{-q}(2)$ using arguments similar to those in [VVV10, Van08]. Namely, we proceed as follows.

We first reduce the proof to a particular case. If $v = v_1 \otimes v_2$, then, using that $\bar{R}_v = (\iota_{v_1} \otimes \bar{R}_{v_2} \otimes \iota_{\bar{v}_1})\bar{R}_{v_1}$ by (8) and that $p_{sv}^{s \otimes v} = p_{sv_1}^{s \otimes v_1} \otimes \iota_{v_2}$, $p_{\bar{v}t}^{\bar{v} \otimes t} = \iota_{\bar{v}_2} \otimes p_{\bar{v}_1 t}^{\bar{v}_1 \otimes t}$, we see that

$$\|\tilde{V}(st, sv \otimes \bar{v}t)\| = \dim_q(v_2)^{1/2} \|\tilde{V}(st, sv_1 \otimes \bar{v}_1 t)\|.$$

We may therefore assume that v is indecomposable. We also assume that $v \neq e$, since $\tilde{V}(st, s \otimes t) = V(st, s \otimes t)$ is isometric by definition.

Next, write sv as $s_1 \otimes s_2 v$, with $s_2 v$ indecomposable. Assume first that $s_2 = e$. Then

$$\tilde{V}(st, sv \otimes \bar{v}t) = (\iota_s \otimes \iota_v \otimes p_{\bar{v}t}^{\bar{v} \otimes t})(\iota_s \otimes \bar{R}_v \otimes \iota_t)V(st, s \otimes t),$$

hence

$$\tilde{V}(st, sv \otimes \bar{v}t)^* \tilde{V}(st, sv \otimes \bar{v}t) = V(st, s \otimes t)^*(\iota_s \otimes (\text{Tr}_{\bar{v}} \otimes \iota)(p_{\bar{v}t}^{\bar{v} \otimes t}))V(st, s \otimes t),$$

where $\text{Tr}_{\bar{v}}$ denotes the categorical trace. The morphism $(\text{Tr}_{\bar{v}} \otimes \iota)(p_{\bar{v}t}^{\bar{v} \otimes t})$ is a scalar multiple of ι_t , and, applying Tr_t , we see that this scalar is $\dim_q(\bar{v}t)/\dim_q(t)$. It follows that

$$\|\tilde{V}(st, sv \otimes \bar{v}t)\| = \frac{\dim_q(\bar{v}t)^{1/2}}{\dim_q(t)^{1/2}}.$$

From (6) it is easy to see that this quantity is not smaller than $c \dim_q(v)^{1/2}$ for a constant c depending only on q ; cf. [VV07, (5)].

Assume now that $s_2 \neq e$. Then $st = s_1 \otimes s_2t$ and $\tilde{V}(st, sv \otimes \bar{v}t) = \iota_{s_1} \otimes \tilde{V}(s_2t, s_2v \otimes \bar{v}t)$. We see that in this case the computation reduces to the case where $s_1 = e$, that is, we may assume that sv is indecomposable. In a similar way we reduce the computation to the case where $\bar{v}t$ is indecomposable as well, so we assume that all three elements sv , $v\bar{v}$ and $\bar{v}t$ are indecomposable (and $v \neq e$). In other words, $sv\bar{v}t$ equals $\alpha\beta\alpha \cdots$ or $\beta\alpha\beta \cdots$. But in this case the computation of the norm of $\tilde{V}(st, sv \otimes \bar{v}t)$ is equivalent to a similar computation for $SU_{-q}(2)$ (see [Van08, Lemma 8.7.2]), which gives

$$\|\tilde{V}(st, sv \otimes \bar{v}t)\| = \begin{bmatrix} |s| + |t| + |v| + 1 \\ |v| \end{bmatrix}_q^{1/2} \begin{bmatrix} |s| + |v| \\ |v| \end{bmatrix}_q^{-1/2} \begin{bmatrix} |t| + |v| \\ |v| \end{bmatrix}_q^{-1/2}$$

(see [VV07, (7.3)]), where $\begin{bmatrix} n \\ k \end{bmatrix}_q$ denote the q -binomial coefficients. Using that

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = q^{-k(n-k)} \prod_{i=0}^{k-1} \frac{1 - q^{2(n-i)}}{1 - q^{2(k-i)}},$$

we easily deduce that $\|\tilde{V}(st, sv \otimes \bar{v}t)\| \geq c[|v| + 1]_q^{1/2} = c \dim_q(v)^{1/2}$ for a constant $c > 0$ depending only on q ; cf. [VV07, Lemma 8.5]. □

In particular, the morphisms $\tilde{V}(z, x \otimes y)$ are non-zero, so we can define isometries

$$V(z, x \otimes y) = \frac{\tilde{V}(z, x \otimes y)}{\|\tilde{V}(z, x \otimes y)\|}.$$

The following result will play a crucial role in our computations. Note that it does not depend on the particular choice of isometries $V(z, x \otimes y)$.

PROPOSITION 2.3 ([VVV10, Lemma A.1], [Van08, Lemma 8.7.3]). *There is a constant C depending only on q such that*

$$\begin{aligned} \|(\iota_u \otimes V(z, x \otimes y))p_{uz}^{u \otimes z} - (p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y))\| &\leq Cq^{(|z|+|x|-|y|)/2}, \\ \|(V(z, x \otimes y) \otimes \iota_u)p_{zu}^{z \otimes u} - (\iota_y \otimes p_{yu}^{y \otimes u})(V(z, x \otimes y) \otimes \iota_u)\| &\leq Cq^{(|z|+|y|-|x|)/2}, \end{aligned}$$

for all $u, x, y, z \in I$ such that $z \prec x \otimes y$.

For our particular choice of $V(z, x \otimes y)$ this implies the following estimates. (In fact, it is not difficult to see that these estimates are equivalent to the ones above, but with different constants.)

COROLLARY 2.4. *There is a constant C depending only on q such that*

$$\begin{aligned} \|V(uz, ux \otimes y)p_{uz}^{u \otimes z} - (p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y))\| &\leq Cq^{(|z|+|x|-|y|)/2}, \\ \|V(zu, x \otimes yu)p_{zu}^{z \otimes u} - (\iota_x \otimes p_{yu}^{y \otimes u})(V(z, x \otimes y) \otimes \iota_u)\| &\leq Cq^{(|z|+|y|-|x|)/2}, \end{aligned}$$

for all $u, x, y, z \in I$ such that $z \prec x \otimes y$.

Proof. By construction we have

$$\tilde{V}(uz, ux \otimes y)p_{uz}^{u \otimes z} = (p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes \tilde{V}(z, x \otimes y))p_{uz}^{u \otimes z},$$

hence

$$\|\tilde{V}(z, x \otimes y)\|^{-1} \tilde{V}(uz, ux \otimes y) p_{uz}^{u \otimes z} = (p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y)) p_{uz}^{u \otimes z},$$

so in particular

$$\|\tilde{V}(z, x \otimes y)\|^{-1} \|\tilde{V}(uz, ux \otimes y)\| = \|(p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y)) p_{uz}^{u \otimes z}\|.$$

Therefore in order to prove the first inequality in the formulation it suffices to show that the last norm is close to 1 up to $C'q^{(|z|+|x|-|y|)/2}$ for some constant C' , and that

$$(p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y)) p_{uz}^{u \otimes z}$$

is close to $(p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(z, x \otimes y))$ up to $C'q^{(|z|+|x|-|y|)/2}$. But this is indeed the case by the previous proposition, since $(\iota_u \otimes V(z, x \otimes y)) p_{uz}^{u \otimes z}$ is an isometry on H_{uz} . The second inequality is proved in a similar way. \square

We can now get a description of the morphism spaces $\mathcal{D}_{\mathcal{B}}(e, y)$. If y is not of the form $\bar{z}z$, $z \in I$, then $\text{Mor}(x, x \otimes y) = 0$ for all x , so $\mathcal{D}_{\mathcal{B}}(e, y) = 0$. If $y = \bar{z}z$, then $x \prec x \otimes y$ precisely for x of the form uz , $u \in I$. Denote by Ω_y the set of all such x . It forms a branch of the tree I and its closure $\bar{\Omega}_y$ in \bar{I} is a clopen subset.

COROLLARY 2.5. Fix $y \in I$ of the form $\bar{z}z$. Define an element $T = (T_x)_x \in \ell^\infty - \bigoplus_{x \in I} \text{Mor}(x, x \otimes y)$ by

$$T_x = \begin{cases} V(x, x \otimes y) & \text{if } x \in \Omega_y, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\mathcal{D}_{\mathcal{B}}(e, y) = C(\bar{\Omega}_y)T$.

Proof. By Proposition 2.1, in order to show that $T \in \mathcal{D}_{\mathcal{B}}(e, y)$ we have to prove that

$$\|T_{ux} - (p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes T_x)V(ux, u \otimes x)\| \rightarrow 0$$

as $x \rightarrow \infty$, uniformly in $u \in I$. But this is true by the previous corollary. Next, if $S \in \mathcal{D}_{\mathcal{B}}(e, y)$, then $S = fT$ for a function $f \in \ell^\infty(\Omega_y)$ such that $|f(ux) - f(x)| \rightarrow 0$ as $x \rightarrow \infty$ in Ω_y , uniformly in $u \in I$. But this means precisely that $f \in C(\bar{\Omega}_y)$. \square

Later we will also need the following estimate.

COROLLARY 2.6. There is a constant C depending only on q such that

$$\|(V(ux, uv \otimes \bar{v}x) \otimes \iota_y)V(ux, ux \otimes y) - (\iota_{uv} \otimes V(\bar{v}x, \bar{v}x \otimes y))V(ux, uv \otimes \bar{v}x)\| \leq Cq^{|x|-|y|/2}$$

for all $u, v, x, y \in I$ such that $x \prec x \otimes y$.

Proof. Using Corollary 2.4, we see that up to an operator of norm $\leq Cq^{|x|-|y|/2}$ the morphism

$$(V(ux, uv \otimes \bar{v}x) \otimes \iota_y)V(ux, ux \otimes y) p_{ux}^{u \otimes x}$$

equals $(V(ux, uv \otimes \bar{v}x) \otimes \iota_y)(p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(x, x \otimes y)) p_{ux}^{u \otimes x}$. By (9), the latter operator equals

$$\|\tilde{V}(ux, uv \otimes \bar{v}x)\|^{-1} (p_{uv}^{u \otimes v} \otimes p_{\bar{v}x}^{\bar{v} \otimes x} \otimes \iota_y)(\iota_u \otimes \bar{R}_v \otimes \iota_x \otimes \iota_y)(p_{ux}^{u \otimes x} \otimes \iota_y)(\iota_u \otimes V(x, x \otimes y)) p_{ux}^{u \otimes x}.$$

By Proposition 2.3, this, in turn, equals

$$\begin{aligned} & \|\tilde{V}(ux, uv \otimes \bar{v}x)\|^{-1} (p_{uv}^{u \otimes v} \otimes p_{\bar{v}x}^{\bar{v} \otimes x} \otimes \iota_y) (\iota_u \otimes \bar{R}_v \otimes \iota_x \otimes \iota_y) (\iota_u \otimes V(x, x \otimes y)) p_{ux}^{u \otimes x} \\ &= \|\tilde{V}(ux, uv \otimes \bar{v}x)\|^{-1} (p_{uv}^{u \otimes v} \otimes p_{\bar{v}x}^{\bar{v} \otimes x} \otimes \iota_y) (\iota_u \otimes \bar{R}_v \otimes V(x, x \otimes y)) p_{ux}^{u \otimes x} \end{aligned} \tag{10}$$

up to an operator of norm not greater than

$$C \frac{\|\bar{R}_v\|}{\|\tilde{V}(ux, uv \otimes \bar{v}x)\|} q^{|x|-|y|/2} = C \frac{\dim_q(v)^{1/2}}{\|\tilde{V}(ux, uv \otimes \bar{v}x)\|} q^{|x|-|y|/2}.$$

The last quantity is not greater than $Cc^{-1}q^{|x|-|y|/2}$, where c is the constant from Lemma 2.2.

In a similar way one checks that the right-hand side of (10) equals

$$(\iota_{uv} \otimes V(\bar{v}x, \bar{v}x \otimes y)) V(ux, uv \otimes \bar{v}x) p_{ux}^{u \otimes x}$$

up to an operator of norm not greater than $Cq^{|x|-|y|/2} + Cc^{-1}q^{|x|-|y|/2}$, where C is from Proposition 2.3 and c is from Lemma 2.2. □

3. Identification of the Martin boundary

The following theorem is our main result.

THEOREM 3.1. *Consider a free unitary quantum group $G = A_u(F)$, with F not a unitary 2×2 matrix, and a generating finitely supported probability measure μ on $I = \text{Irr}(G)$. Then the Martin compactification $C(\bar{\Gamma}_{M,\mu})$ of the discrete quantum group $\Gamma = \hat{G}$ with respect to the quantum random walk defined by μ coincides with the compactification \mathcal{B} defined in (7). It follows that the Martin boundary $C(\partial\Gamma_{M,\mu})$ coincides with \mathcal{B}_∞ .*

In order to simplify the notation, let us denote the Martin compactification $C(\bar{\Gamma}_{M,\mu})$ by \mathcal{M} . Since both \mathcal{M} and \mathcal{B} are G - C^* -subalgebras of $\ell^\infty(\Gamma)$, with the actions of G coming from the left adjoint action of G on $\ell^\infty(\Gamma)$, by the discussion in § 1.3 in order to show that they coincide it suffices to check that $\mathcal{D}_{\mathcal{M}}(e, y) = \mathcal{D}_{\mathcal{B}}(e, y)$ for all $y \in I$.

Consider first $y = e$. Then, by definition, $\mathcal{D}_{\mathcal{B}}(e, e)$ coincides with $C(\bar{I})$ considered as a subalgebra of the center $\ell^\infty(I)$ of $\ell^\infty(\Gamma)$. On the other hand, $\mathcal{D}_{\mathcal{M}}(e, e)$ contains the algebra of continuous functions on the Martin compactification of I defined by the Markov operator with transition probabilities $p_\mu(s, t)$ defined by (1). More precisely, denoting by P this Markov operator and by G_P the corresponding classical Green kernel,

$$G_P(s, t) = \sum_{n=0}^{\infty} p_\mu^{(n)}(s, t) = \delta_{st} + \sum_{n=1}^{\infty} \sum_{s=s_0, s_1, \dots, s_n=t} p_\mu(s_0, s_1) \cdots p_\mu(s_{n-1}, s_n),$$

in view of (3) we have

$$G_{\bar{\mu}}(I_s)I_t = \frac{\dim_q(s)^2}{\dim_q(t)^2} G_P(s, t).$$

Therefore, denoting by K_P the classical Martin kernel, $K_P(s, t) = G_P(s, t)/G_P(e, t)$, we see that the function $K_P(s, \cdot) \in \ell^\infty(I)$ differs from $K_{\bar{\mu}}(I_s)$ only by a scalar factor:

$$K_{\bar{\mu}}(I_s) = \dim_q(s)^2 K_P(s, \cdot).$$

Hence the C^* -algebra generated by $c_c(I) \subset c_c(\Gamma)$ and the elements $K_{\tilde{\mu}}(I_s)$, $s \in I$, coincides with the algebra of continuous functions on the Martin compactification of I . (Note that this does not yet exclude the possibility that $\mathcal{D}_{\mathcal{M}}(e, e) = \mathcal{M} \cap \ell^\infty(I)$ is a strictly larger algebra.) By a result of Picardello and Woess [PW87] (see also [Woe00, Corollary 26.14] or [Woe00, Theorem 27.1]), the Martin compactification of I coincides with the end compactification \bar{I} . It is checked in [VVV10, § 2] that this result can indeed be applied in our case, since the transition probabilities $p_\mu(s, t)$ satisfy the following properties.

- The spectral radius $\lim_n p_\mu^{(n)}(s, s)^{1/n}$ of the random walk is strictly less than 1 (see also Remark 4.2 below).
- The random walk has bounded range: there is $S \in \mathbb{N}$ such that $p_\mu(s, t) = 0$ whenever $d(s, t) > S$,
- The random walk is uniformly irreducible (in the sense used in [Woe00]): there are $\varepsilon_0 > 0$ and $K \in \mathbb{N}$ such that for any $s, t \in I$ with $d(s, t) = 1$ we have $p_\mu^{(k)}(s, t) \geq \varepsilon_0$ for some $k \leq K$.

We thus have

$$\mathcal{D}_{\mathcal{B}}(e, e) = C(\bar{I}) = C^*(c_c(I), K_{\tilde{\mu}}(c_c(I))) \subset \mathcal{D}_{\mathcal{M}}(e, e). \tag{11}$$

Assume next $y = \bar{z}z$ for some $z \neq e$. Then, by Corollary 2.5, we have $\mathcal{D}_{\mathcal{B}}(e, y) = C(\bar{\Omega}_y)T$, where $T_x = V(x, x \otimes y)$ for $x \in \Omega_y$. For every $t \in \Omega_y$ define $\eta^{(t)} \in \mathcal{D}_{\Gamma, c}(e, y) = \text{Nat}_c(\iota, \iota \otimes y)$ by

$$\eta_s^{(t)} = \begin{cases} V(t, t \otimes y) & \text{if } s = t, \\ 0 & \text{otherwise.} \end{cases} \tag{12}$$

Define numbers $\check{q}_\mu(s, t)$ for $s, t \in \Omega_y$ by

$$P_{\tilde{\mu}}(\eta^{(t)})_s = \check{q}_\mu(s, t)V(s, s \otimes y).$$

Motivated by (3), we then put

$$q_\mu(s, t) = \frac{\dim_q(t)^2}{\dim_q(s)^2} \check{q}_\mu(t, s).$$

LEMMA 3.2. *The numbers $q_\mu(s, t)$ are real and $|q_\mu(s, t)| \leq p_\mu(s, t)$ for all $s, t \in \Omega_y$.*

Proof. Equivalently, we have to show that the numbers $\check{q}_\mu(s, t)$ are real and $|\check{q}_\mu(s, t)| \leq p_{\tilde{\mu}}(s, t)$.

Since $\tilde{\mu}$ is a convex combination of point masses, it suffices to consider $\tilde{\mu} = \delta_u$. Fix $s, t \in \Omega_y$. We may assume that $t \prec u \otimes s$, as otherwise $\check{q}_\mu(s, t) = p_{\tilde{\mu}}(s, t) = 0$. Then, with $\eta^{(t)}$ given by (12),

$$P_{\tilde{\mu}}(\eta^{(t)})_s = (\text{tr}_u \otimes \iota)(\eta_{u \otimes s}^{(t)}) = (\text{tr}_u \otimes \iota)((V(t, u \otimes s) \otimes \iota_y)V(t, t \otimes y)V(t, u \otimes s)^*),$$

and hence

$$\check{q}_\mu(s, t) = \text{tr}_{u \otimes s}((\iota_u \otimes V(s, s \otimes y)^*)(V(t, u \otimes s) \otimes \iota_y)V(t, t \otimes y)V(t, u \otimes s)^*). \tag{13}$$

Somewhat informally, the statement that $\check{q}_\mu(s, t)$ is real follows now from the fact that in order to compute the above trace we need only to use that R_α and \bar{R}_α solve the conjugate equations and satisfy $R_\alpha^* R_\alpha = q + q^{-1} = \bar{R}_\alpha^* \bar{R}_\alpha$, and all these relations involve only real numbers. This can

be formalized, for example, as follows. We can choose orthonormal bases $(e_i)_i$ of H_α and $(f_i)_i$ of H_β such that

$$R_\alpha(1) = \sum_i \lambda_i f_i \otimes e_i \quad \text{and} \quad \bar{R}_\alpha(1) = \sum_i \lambda_i^{-1} e_i \otimes f_i$$

for some $\lambda_i > 0$ (with $\sum_i \lambda_i^2 = \sum_i \lambda_i^{-2} = q + q^{-1}$). Then the real linear spans of these bases define real forms of H_α and H_β . Taking tensor products of these forms, we get a real form of $H_{x_1} \otimes \cdots \otimes H_{x_n}$ for every word $x = x_1 \cdots x_n$ in α and β . Since H_x is the orthogonal complement of the images of operators of the form $\iota \otimes \cdots \otimes \iota \otimes R_\alpha \otimes \iota \otimes \cdots \otimes \iota$ and $\iota \otimes \cdots \otimes \iota \otimes \bar{R}_\alpha \otimes \iota \otimes \cdots \otimes \iota$, which respect the real forms of the tensor products, we then also get a real form of H_x . Now, in computing the right-hand side of (13) we may work only with the real forms of all the spaces involved, hence the result must be real.

In order to prove the second statement of the lemma, note that we also have $|\check{q}_\mu(s, t)| = \|P_{\check{\mu}}(\eta^{(t)})_s\|$. Consider the map $P_{\check{\mu}}$ on $\text{Nat}_b(\iota \otimes (e \oplus y), \iota \otimes (e \oplus y)) \cong \ell^\infty\text{-}\bigoplus_x \text{End}(x \otimes (e \oplus y))$. Then, by restricting it to the block corresponding to t and by projecting the image onto the block corresponding to s , we get a completely positive map $P_{st}: \text{End}(t \otimes (e \oplus y)) \rightarrow \text{End}(s \otimes (e \oplus y))$. By definition, this map, being divided by $p_{\check{\mu}}(s, t)$, becomes unital. Hence $\|P_{st}\| = p_{\check{\mu}}(s, t)$. Viewing $V(t, t \otimes y)$ as the element

$$\begin{pmatrix} 0 & 0 \\ V(t, t \otimes y) & 0 \end{pmatrix} \in \text{End}(t \otimes (e \oplus y)),$$

we conclude that $\|P_{\check{\mu}}(\eta^{(t)})_s\| \leq \|P_{st}\| = p_{\check{\mu}}(s, t)$. □

Denote by Q the matrix $(q_\mu(s, t))_{s, t \in \Omega_y}$. We can define the ‘Green kernel’

$$G_Q(s, t) = \delta_{st} + \sum_{n=1}^\infty \sum_{s=s_0, s_1, \dots, s_n=t} q_\mu(s_0, s_1) \cdots q_\mu(s_{n-1}, s_n).$$

Then

$$G_{\check{\mu}}(\eta^{(s)})_t = \frac{\dim_q(s)^2}{\dim_q(t)^2} G_Q(s, t) V(t, t \otimes y).$$

Therefore if we let $K_Q(s, t) = G_Q(s, t)/G_P(e, t)$, then

$$K_{\check{\mu}}(\eta^{(s)}) = \dim_q(s)^2 K_Q(s, \cdot) T, \tag{14}$$

with T as in Corollary 2.5. We thus need to understand the behavior of the functions $K_Q(s, \cdot)$ at infinity. As we will see, this behavior is not that different from that of $K_P(s, \cdot)$. The starting point is the following estimate.

LEMMA 3.3. *There is a constant C (depending on q , $|y|$ and the support of μ) such that*

$$|q_\mu(s, t) - p_\mu(s, t)| \leq Cq^{|s|}$$

for all $s, t \in \Omega_y$.

Proof. Recall that $p_\mu(s, t) = 0$ if $d(s, t) > S$, and then $q_\mu(s, t) = 0$ as well, so we could equally well use $q^{|t|}$ instead of $q^{|s|}$ in the formulation. Using (6), it is also easy to see that the ratios $\dim_q(s)/\dim_q(t)$ are bounded when $d(s, t) \leq S$. It follows that the statement of the lemma is equivalent to the existence of C' such that $|\check{q}_\mu(s, t) - p_{\check{\mu}}(s, t)| \leq C'q^{|s|}$.

Similarly to the proof of the previous lemma, we may assume that $\check{\mu} = \delta_u$ and consider $s, t \in \Omega_y$ such that $t \prec u \otimes s$. We may also assume that $|s| - |u| \geq |y|/2$, as there are only finitely many pairs (s, t) as above not satisfying this condition. Then $u = u_0v, s = \bar{v}s_0$ and $t = u_0s_0$ for some $s_0, u_0, v \in I$. Since $|s_0| = |s| - |v| \geq |s| - |u| \geq |y|/2$, we have $s_0 \prec s_0 \otimes y$. By (13) we have

$$\check{q}_\mu(s, t) = \text{tr}_{u \otimes s}((\iota_{u_0v} \otimes V(\bar{v}s_0, \bar{v}s_0 \otimes y)^*)(V(u_0s_0, u_0v \otimes \bar{v}s_0) \otimes \iota_y)V(u_0s_0, u_0s_0 \otimes y)V(t, u \otimes s)^*).$$

By Corollary 2.6 the last expression is close to

$$\text{tr}_{u \otimes s}(V(u_0s_0, u_0v \otimes \bar{v}s_0)V(t, u \otimes s)^*) = \text{tr}_{u \otimes s}(V(t, u \otimes s)V(t, u \otimes s)^*) = \frac{\dim_q(t)}{\dim_q(u) \dim_q(s)}$$

up to $Cq^{|s_0|-|y|/2} \leq Cq^{|s|-|u|-|y|/2}$. By (2), the quantity on the right is exactly $p_{\check{\mu}}(s, t)$. □

From this we will deduce in the next section the following result.

PROPOSITION 3.4. *For every $s \in \Omega_y$, the function $K_Q(s, \cdot) = G_Q(s, \cdot)/G_P(e, \cdot)$ on Ω_y extends to a continuous function on $\bar{\Omega}_y \subset \bar{I}$.*

By (14), this already shows that

$$K_{\check{\mu}}(\mathcal{D}_{\Gamma,c}(e, y)) \subset C(\bar{\Omega}_y)T = \mathcal{D}_{\mathcal{B}}(e, y). \tag{15}$$

As opposed to the functions $K_P(s, \cdot)$, it may happen that the functions $K_Q(s, \cdot)$ vanish at some points of the boundary of Ω_y . Nevertheless we have the following result, whose proof we also postpone till the next section.

PROPOSITION 3.5. *For every point $t_\infty \in \bar{\Omega}_y \cap \partial I$, we have $K_Q(s, t_\infty) > 0$ for all s sufficiently close to t_∞ .*

It is now easy to complete the proof of the equality $\mathcal{M} = \mathcal{B}$.

Proof of Theorem 3.1. By (11), (15) and our discussion at the end of § 1.3, we can already conclude that $K_{\check{\mu}}(c_c(\Gamma)) \subset \mathcal{B}$. Hence $\mathcal{M} \subset \mathcal{B}$, since we also have $c_0(\Gamma) \subset \mathcal{B}$.

Furthermore, $\mathcal{D}_{\mathcal{M}}(e, y)$ is a bimodule over $\mathcal{D}_{\mathcal{M}}(e, e)$ and by (11) we know that $\mathcal{D}_{\mathcal{M}}(e, e) \supset C(\bar{I})$. Using Proposition 3.5, for every function $f \in C(\bar{\Omega}_y)$ we can find $s_1, \dots, s_n \in \Omega_y$ and $g_1, \dots, g_n \in C(\bar{\Omega}_y)$ such that $f(t_\infty) = \sum_i g_i(t_\infty)K_Q(s_i, t_\infty)$ for all $t_\infty \in \bar{\Omega}_y \cap \partial I$. In view of (14) this shows that, with T as in Corollary 2.5, we have

$$\mathcal{D}_{\mathcal{B}}(e, y) = C(\bar{\Omega}_y)T \subset C(\bar{I})K_{\check{\mu}}(\mathcal{D}_{\Gamma,c}(e, y)) + c_0(I)T. \tag{16}$$

Hence $\mathcal{D}_{\mathcal{B}}(e, y) \subset \mathcal{D}_{\mathcal{M}}(e, y)$ and therefore $\mathcal{B} \subset \mathcal{M}$. □

Remark 3.6. The equality $\mathcal{M} = \mathcal{B}$ and (16) imply that the space $C(\bar{I})K_{\check{\mu}}(c_c(\Gamma)) + c_0(\Gamma)$ is dense in $\mathcal{B} = C(\bar{\Gamma}_{M,\mu})$.

4. Estimates of the Green kernels

In this section we will prove Propositions 3.4 and 3.5. We thus fix a generating probability measure μ , a point $y \in I$ of the form $\bar{z}z$, $z \neq e$, and consider the set $\Omega_y \subset I$ of words of the form uz . In order to simplify the notation we denote the transition probabilities $p_\mu(s, t)$ by $p(s, t)$.

In the previous section we introduced the numbers $q_\mu(s, t)$, $s, t \in \Omega_y$, which we now denote simply by $q(s, t)$. The only information we will need about them is that, by Lemmas 3.2 and 3.3, they are real and

$$|q(s, t)| \leq p(s, t) \quad \text{and} \quad |q(s, t) - p(s, t)| \leq Cq^{|s|} \quad \text{for all } s, t \in \Omega_y. \tag{17}$$

Denote by P and Q the matrices $(p(s, t))_{s, t \in I}$ and $(q(s, t))_{s, t \in \Omega_y}$.

The space $c_c(I)$ can be completed to two different Hilbert spaces, depending on the inner product we choose. Denote by $\ell^2(I)$ the completion of $c_c(I)$ with respect to the inner product (\cdot, \cdot) corresponding to the counting measure: $(f, g) = \sum_{x \in I} f(x)\overline{g(x)}$. Denote by $\ell^2(I, m)$ the completion of $c_c(I)$ with respect to the inner product corresponding to the ‘Haar measure’ m given by $m(x) = \dim_q(x)^2$: $(f, g)_m = \sum_{x \in I} m(x) f(x)\overline{g(x)}$. For any $f \in c_c(I)$ and $s \in I$, we have

$$(f, \delta_s) = m(s)^{-1}(f, \delta_s)_m. \tag{18}$$

LEMMA 4.1. *The operator P on $c_c(I)$ extends to a bounded operator on $\ell^2(I, m)$ of norm $\lambda < 1$. The operator Q on $c_c(\Omega_y)$ extends to a bounded operator on $\ell^2(\Omega_y, m)$ of norm not greater than λ .*

Proof. Denote by \dim_{\min} the dimension function on the finite-dimensional representations of our quantum group $G = A_u(F)$ obtained by letting $q = 1$ in formula (6).

As can be easily seen from (2), the unitary operator $\ell^2(I, m) \rightarrow \ell^2(I)$, $\delta_s \mapsto \dim_q(s)\delta_s$, transforms the operator P into the operator

$$\sum_{r \in I} \frac{\mu(r)}{\dim_q(r)} \lambda_r,$$

where λ_r is the operator of multiplication by r on the fusion algebra $\mathbb{C}[I]$ of G (which we identify with $c_c(I)$ as a space). By [NT13, Lemma 2.7.3], the operator λ_r on $\ell^2(I)$ is bounded, of norm not greater than $\dim_{\min}(r)$. Therefore

$$\lambda \leq \sum_{r \in I} \mu(r) \frac{\dim_{\min}(r)}{\dim_q(r)} < 1,$$

since $\dim_{\min}(r) < \dim_q(r)$ for all $r \neq e$.

The second statement of the lemma follows from (17), since we get

$$|(Qf, g)_m| \leq (P|f|, |g|)_m$$

for all $f, g \in c_c(\Omega_y)$. □

Remark 4.2. Note that by (18) we have

$$p^{(n)}(s, t) = (P^n \delta_t, \delta_s) = m(s)^{-1}(P^n \delta_t, \delta_s)_m. \tag{19}$$

From this we see that the spectral radius of the random walk defined by P is not greater than λ . This (already mentioned) fact that the spectral radius of the random walk is strictly less than one is actually enough for the results below, but since the inequality $\|P\| < 1$ slightly simplifies the arguments, we will use it.

For $x \in I$, let us introduce the following subset of I :

$$\Delta_x = \{ux : u \in I\}.$$

We will have to consider the substochastic matrices $P_{\Delta_x} = (p(s, t))_{s, t \in \Delta_x}$ and the corresponding Green kernels, which we denote by G_{P, Δ_x} . For $x \in \Omega_y$, we similarly define Q_{Δ_x} and G_{Q, Δ_x} .

The Green kernels G_{P, Δ_x} satisfy the following uniform version of Harnack’s inequality.

LEMMA 4.3. *There exists $\delta \in (0, 1)$ such that*

$$G_{P, \Delta_x}(s, t) \leq \delta^{-d(s, v)} G_{P, \Delta_x}(v, t) \quad \text{and} \quad G_{P, \Delta_x}(s, t) \leq \delta^{-d(t, v)} G_{P, \Delta_x}(s, v)$$

for all $x \in I$ and $v, s, t \in \Delta_x$.

Proof. As we already mentioned in the previous section, the random walk defined by P is uniformly irreducible. In fact, the following slightly stronger property (also called uniform irreducibility in [PW87]) is shown in [VVV10, §2]. There exist $K \in \mathbb{N}$ and $\delta_0 > 0$ such that:

- $p(s, t) \geq \delta_0$ whenever $p(s, t) > 0$;
- for any $s, t \in I$ with $d(s, t) = 1$ we have $p^{(k)}(s, t) > 0$ for some $k \leq K$.

In particular, given $s, t \in I$ with $d(s, t) = 1$, there are points $x_0 = s, x_1, \dots, x_k = t, k \leq K$, such that $p(x_i, x_{i+1}) \geq \delta_0$ for all $i = 0, \dots, k - 1$. But then we also have $p(x_i x, x_{i+1} x) \geq \delta_0$ for any $x \in I$. That is, the same constants K and δ_0 work for P_{Δ_x} . This implies the lemma; specifically, we can take $\delta = \delta_0^K$. □

The most important property of the Green kernels of random walks on trees (shared also by random walks on hyperbolic graphs, where it is much more difficult to prove [Anc88]) is their almost multiplicativity along geodesics. We will need the following uniform version of this property, cf. [INO08, Proposition 2.1].

For $s, t \in I$, we denote by $[s, t]$ the unique geodesic segment between s and t .

LEMMA 4.4. *There exists a constant $C_1 > 0$ such that*

$$\frac{1}{C_1} G_{P, \Delta_x}(s, v) G_{P, \Delta_x}(v, t) \leq G_{P, \Delta_x}(s, t) \leq C_1 G_{P, \Delta_x}(s, v) G_{P, \Delta_x}(v, t)$$

for all $x \in I, s, t \in \Delta_x$ and $v \in [s, t]$.

Proof. The first inequality is true, with $C_1 = (1 - \rho)^{-1}$, for any irreducible random walk with spectral radius $\rho < 1$. Namely, we have

$$\frac{G_{P, \Delta_x}(s, v)}{G_{P, \Delta_x}(v, v)} G_{P, \Delta_x}(v, t) \leq G_{P, \Delta_x}(s, t),$$

since the expression on the left is the sum of $p(s_0, s_1) \cdots p(s_{n-1}, s_n)$ over all paths $(s = s_0, s_1, \dots, s_n = t)$ in Δ_x that pass through v . As $G_{P, \Delta_x}(v, v) \leq G_P(v, v) \leq (1 - \lambda)^{-1}$ by Remark 4.2, we can take $C_1 = (1 - \lambda)^{-1}$.

For the second inequality, take $x \in I, s, t \in \Delta_x$ and $v \in [s, t]$. Then any path from s to t in Δ_x contributing to $G_{P, \Delta_x}(s, t)$ must pass through the set $A = \Delta_x \cap B_S(v)$, where $B_S(v)$ is

the open ball of radius S with center v . (Recall that $S \in \mathbb{N}$ is such that $p(u, w) = 0$ whenever $d(u, w) > S$.) It follows that

$$G_{P,\Delta_x}(s, t) \leq \sum_{u \in A} G_{P,\Delta_x}(s, u)G_{P,\Delta_x}(u, t).$$

By Lemma 4.3, for every $u \in A$, we have

$$G_{P,\Delta_x}(s, u)G_{P,\Delta_x}(u, t) \leq \delta^{-2(S-1)}G_{P,\Delta_x}(s, v)G_{P,\Delta_x}(v, t).$$

Since the number of vertices in $B_S(v)$ is not greater than $3 \cdot 2^{S-1} - 2$, we thus see that for the second inequality in the formulation of the lemma we can take $C_1 = 3(2/\delta^2)^{S-1}$. \square

We are now ready to prove the following key estimate.

LEMMA 4.5. *There is a constant C_2 such that*

$$|G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| \leq C_2q^{|x|}G_{P,\Delta_x}(s, t)$$

for all $x \in \Omega_y$ and $s, t \in \Delta_x$.

Proof. By (19) we have $G_P(s, t) = m(s)^{-1}((1 - P)^{-1}\delta_t, \delta_s)_m$. Similarly,

$$G_{P,\Delta_x}(s, t) = m(s)^{-1}((1 - P_{\Delta_x})^{-1}\delta_t, \delta_s)_m \quad \text{and} \quad G_{Q,\Delta_x}(s, t) = m(s)^{-1}((1 - Q_{\Delta_x})^{-1}\delta_t, \delta_s)_m.$$

Define functions $\tilde{\delta}_u = \delta_u/\sqrt{m(u)}$ for $u \in I$. Then we can write

$$G_{P,\Delta_x}(s, t) = \frac{\sqrt{m(t)}}{\sqrt{m(s)}}((1 - P_{\Delta_x})^{-1}\tilde{\delta}_t, \tilde{\delta}_s)_m \quad \text{and} \quad G_{Q,\Delta_x}(s, t) = \frac{\sqrt{m(t)}}{\sqrt{m(s)}}((1 - Q_{\Delta_x})^{-1}\tilde{\delta}_t, \tilde{\delta}_s)_m.$$

From this we get

$$|G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| = \frac{\sqrt{m(t)}}{\sqrt{m(s)}}|((1 - Q_{\Delta_x})^{-1}(Q_{\Delta_x} - P_{\Delta_x})(1 - P_{\Delta_x})^{-1}\tilde{\delta}_t, \tilde{\delta}_s)_m|.$$

Since $\{\tilde{\delta}_u\}_{u \in \Delta_x}$ is an orthonormal basis in $\ell^2(\Delta_x, m)$, the scalar product in the expression above equals

$$\sum_{u, v \in \Delta_x} ((1 - P_{\Delta_x})^{-1}\tilde{\delta}_t, \tilde{\delta}_v)_m((Q_{\Delta_x} - P_{\Delta_x})\tilde{\delta}_v, \tilde{\delta}_u)_m((1 - Q_{\Delta_x})^{-1}\tilde{\delta}_u, \tilde{\delta}_s)_m.$$

It follows that

$$\begin{aligned} & |G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| \\ & \leq \frac{\sqrt{m(t)}}{\sqrt{m(s)}} \sum_{u, v \in \Delta_x} |((1 - P_{\Delta_x})^{-1}\tilde{\delta}_t, \tilde{\delta}_v)_m((Q_{\Delta_x} - P_{\Delta_x})\tilde{\delta}_v, \tilde{\delta}_u)_m((1 - Q_{\Delta_x})^{-1}\tilde{\delta}_u, \tilde{\delta}_s)_m| \\ & = \sum_{u, v \in \Delta_x} G_{P,\Delta_x}(v, t) |q(u, v) - p(u, v)| |G_{Q,\Delta_x}(s, u)|. \end{aligned}$$

By (17) we have $|G_{Q,\Delta_x}(s, u)| \leq G_{P,\Delta_x}(s, u)$ and $|q(u, v) - p(u, v)| \leq Cq^{|u|}$. Since we also have $q(u, v) = p(u, v) = 0$ if $d(u, v) > S$, we conclude that

$$|G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| \leq C \sum_{\substack{u, v \in \Delta_x : \\ d(u, v) \leq S}} q^{|u|}G_{P,\Delta_x}(s, u)G_{P,\Delta_x}(v, t).$$




FIGURE 1. Possible positions of the closest point on the geodesic.

Applying Lemma 4.3, we then get

$$|G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| \leq 3 \left(\frac{2}{\delta}\right)^S C \sum_{u \in \Delta_x} q^{|u|} G_{P,\Delta_x}(s, u) G_{P,\Delta_x}(u, t),$$

where we used again that every closed ball of radius S contains not more than $3 \cdot 2^S - 2$ vertices.

In order to estimate the above expression, for every $u \in \Delta_x$ denote by u' the point on the geodesic $[s, t]$ closest to u . Then $u' \in [u, s] \cap [u, t]$, so by Lemma 4.4 we have

$$\begin{aligned} G_{P,\Delta_x}(s, u) G_{P,\Delta_x}(u, t) &\leq C_1^2 G_{P,\Delta_x}(s, u') G_{P,\Delta_x}(u', u) G_{P,\Delta_x}(u, u') G_{P,\Delta_x}(u', t) \\ &\leq C_1^3 G_{P,\Delta_x}(s, t) G_{P,\Delta_x}(u', u) G_{P,\Delta_x}(u, u'), \end{aligned} \tag{20}$$

and hence

$$|G_{Q,\Delta_x}(s, t) - G_{P,\Delta_x}(s, t)| \leq 3 \left(\frac{2}{\delta}\right)^S C C_1^3 G_{P,\Delta_x}(s, t) \sum_{u \in \Delta_x} q^{|u|} G_{P,\Delta_x}(u', u) G_{P,\Delta_x}(u, u'). \tag{21}$$

It remains to estimate the sum in the above expression on the right. Observe that if $u' \neq x'$ (where $x' \in [s, t]$ is the vertex closest to x), then

$$|u| \geq |u'| = |x'| + d(u', x') \geq |x| + d(u', x').$$

If $u' = x'$, then the inequality $|u| \geq |x| + d(u', x')$ is obvious. See Figure 1 illustrating these two cases.

It follows that

$$\begin{aligned} \sum_{u \in \Delta_x} q^{|u|} G_{P,\Delta_x}(u', u) G_{P,\Delta_x}(u, u') &\leq q^{|x|} \sum_{u \in \Delta_x} q^{d(u', x')} G_{P,\Delta_x}(u', u) G_{P,\Delta_x}(u, u') \\ &\leq q^{|x|} \sum_{u \in \Delta_x} \sum_{w \in [s, t]} q^{d(w, x')} G_{P,\Delta_x}(w, u) G_{P,\Delta_x}(u, w). \end{aligned} \tag{22}$$

Next, for every $w \in [s, t]$, we have

$$\begin{aligned} \sum_{u \in \Delta_x} G_{P, \Delta_x}(w, u)G_{P, \Delta_x}(u, w) &= \sum_{u \in \Delta_w} ((1 - P_{\Delta_x})^{-1}\tilde{\delta}_u, \tilde{\delta}_w)_m ((1 - P_{\Delta_x})^{-1}\tilde{\delta}_w, \tilde{\delta}_u)_m \\ &= ((1 - P_{\Delta_x})^{-2}\tilde{\delta}_w, \tilde{\delta}_w)_m \leq (1 - \lambda)^{-2}. \end{aligned} \tag{23}$$

We also have

$$\sum_{w \in [s, t]} q^{d(w, x')} < 2(1 + q + q^2 + \dots) = 2(1 - q)^{-1}. \tag{24}$$

Combining (22)–(24), we get from (21) that

$$|G_{Q, \Delta_x}(s, t) - G_{P, \Delta_x}(s, t)| \leq 6 \left(\frac{2}{\delta}\right)^S \frac{C C_1^3}{(1 - \lambda)^2(1 - q)} q^{|x|} G_{P, \Delta_x}(s, t),$$

and we are done. □

Proof of Proposition 3.4. Fix $s \in \Omega_y$ and $t_\infty \in \bar{\Omega}_y \cap \partial I$. We have to show that as $t \rightarrow t_\infty$ the numbers $K_Q(s, t) = G_Q(s, t)/G_P(e, t)$ converge to a finite limit.

Fix $\varepsilon > 0$. By Lemma 4.5 there exists $x \in \Omega_y$ such that $s \notin \Delta_x$, $t_\infty \in \bar{\Delta}_x \cap \partial I$ and

$$|G_{Q, \Delta_x}(u, t) - G_{P, \Delta_x}(u, t)| \leq \varepsilon G_{P, \Delta_x}(u, t) \quad \text{for all } u, t \in \Delta_x.$$

Take $t \in \Delta_x$. Then $x \in [s, t]$, so, similarly to the proof of Lemma 4.4, every path from s to t in I contributing to $G_P(s, t)$ must pass through the set $A = \Delta_x \cap B_S(x)$ and there is a well-defined moment when it leaves $I \setminus \Delta_x$ and enters A for the last time. It follows that if for $u \in A$ we put

$$M_P(s, u) = \sum_{n=1}^{\infty} \sum_{\substack{s=s_0, s_1, \dots, s_n=u: \\ s_{n-1} \notin \Delta_x}} p(s_0, s_1) \cdots p(s_{n-1}, s_n),$$

then

$$G_P(s, t) = \sum_{u \in A} M_P(s, u)G_{P, \Delta_x}(u, t). \tag{25}$$

We can similarly define $M_Q(s, u)$ and get a decomposition of $G_Q(s, t)$. Then, for any $t, t' \in \Delta_x$, we have

$$\begin{aligned} |K_Q(s, t) - K_Q(s, t')| &= \left| \sum_{u \in A} \frac{M_Q(s, u)G_{Q, \Delta_x}(u, t)}{G_P(e, t)} - \sum_{u \in A} \frac{M_Q(s, u)G_{Q, \Delta_x}(u, t')}{G_P(e, t')} \right| \\ &\leq \varepsilon \sum_{u \in A} \frac{|M_Q(s, u)|G_{P, \Delta_x}(u, t)}{G_P(e, t)} + \varepsilon \sum_{u \in A} \frac{|M_Q(s, u)|G_{P, \Delta_x}(u, t')}{G_P(e, t')} \\ &\quad + \sum_{u \in A} |M_Q(s, u)| \left| \frac{G_{P, \Delta_x}(u, t)}{G_P(e, t)} - \frac{G_{P, \Delta_x}(u, t')}{G_P(e, t')} \right| \\ &\leq \varepsilon(K_P(s, t) + K_P(s, t')) + \sum_{u \in A} |M_Q(s, u)| \left| \frac{G_{P, \Delta_x}(u, t)}{G_P(e, t)} - \frac{G_{P, \Delta_x}(u, t')}{G_P(e, t')} \right|, \end{aligned}$$

where we used that $|M_Q(s, u)| \leq M_P(s, u)$ by (17). If we could prove that the sum above converges to zero as $t, t' \rightarrow t_\infty$, then we would get

$$\limsup_{t, t' \rightarrow t_\infty} |K_Q(s, t) - K_Q(s, t')| \leq 2\varepsilon K_P(s, t_\infty),$$

and since ε could be taken arbitrarily small, we would be able to conclude that the numbers $K_Q(s, t)$ converge to a finite limit as $t \rightarrow t_\infty$.

Therefore it remains to show that for every $u \in A$ we have

$$\lim_{t, t' \rightarrow t_\infty} \left| \frac{G_{P, \Delta_x}(u, t)}{G_P(e, t)} - \frac{G_{P, \Delta_x}(u, t')}{G_P(e, t')} \right| = 0.$$

Again using (25), but now applied to $G_P(e, t)$, we can rewrite this as

$$\lim_{t, t' \rightarrow t_\infty} \left| \left(\sum_{v \in A} M_P(e, v) \frac{G_{P, \Delta_x}(v, t)}{G_{P, \Delta_x}(u, t)} \right)^{-1} - \left(\sum_{v \in A} M_P(e, v) \frac{G_{P, \Delta_x}(v, t')}{G_{P, \Delta_x}(u, t')} \right)^{-1} \right| = 0.$$

In order to prove this, it suffices to show that for all $u, v \in A$ the function $G_{P, \Delta_x}(v, \cdot)/G_{P, \Delta_x}(u, \cdot)$ on Δ_x extends to a non-vanishing continuous function on $\bar{\Delta}_x \subset \bar{I}$. But this is indeed true, since this function is nothing else than the Martin kernel of the random walk on the tree Δ_x defined by the matrix P_{Δ_x} . Although this matrix is only substochastic, it still satisfies the assumptions of [Woe00, Corollary 26.14] or [Woe00, Theorem 27.1], as we essentially discussed in the proof of Lemma 4.3. So the same arguments as in [Woe00] show that the Martin compactification of Δ_x with respect to P_{Δ_x} coincides with the end compactification, which is $\bar{\Delta}_x$. \square

Proof of Proposition 3.5. Fix $t_\infty \in \bar{\Omega}_y \cap \partial I$. By Lemma 4.5 we can find $x \in \Omega_y$ such that $t_\infty \in \bar{\Delta}_x$ and

$$G_{Q, \Delta_x}(s, t) \geq \frac{1}{2} G_{P, \Delta_x}(s, t) \quad \text{for all } s, t \in \Delta_x.$$

Suppose next that we can find $u \in \Delta_x$ such that $t_\infty \in \bar{\Delta}_u$ and

$$G_P(s, t) - G_{P, \Delta_x}(s, t) \leq \frac{1}{4} G_P(s, t) \quad \text{for all } s, t \in \Delta_u. \tag{26}$$

Then, by (17),

$$|G_Q(s, t) - G_{Q, \Delta_x}(s, t)| \leq G_P(s, t) - G_{P, \Delta_x}(s, t) \leq \frac{1}{4} G_P(s, t) \leq \frac{1}{3} G_{P, \Delta_x}(s, t) \leq \frac{2}{3} G_{Q, \Delta_x}(s, t),$$

whence

$$G_Q(s, t) \geq \frac{1}{3} G_{Q, \Delta_x}(s, t) \geq \frac{1}{6} G_{P, \Delta_x}(s, t) \geq \frac{1}{8} G_P(s, t),$$

and therefore

$$K_Q(s, t_\infty) \geq \frac{1}{8} K_P(s, t_\infty) > 0 \quad \text{for all } s \in \Delta_u.$$

Therefore it remains to find u such that (26) is satisfied.

For $s, t \in \Delta_x$, similarly to the proof of Lemma 4.4, the paths from s to t contributing to $G_P(s, t) - G_{P, \Delta_x}(s, t)$ must pass through the set $A = B_S(x) \cap \Delta_x$ of cardinality $2^S - 1$. It follows that

$$G_P(s, t) - G_{P, \Delta_x}(s, t) \leq \sum_{v \in A} G_P(s, v) G_P(v, t),$$

and, using Lemma 4.3, we get

$$G_P(s, t) - G_{P, \Delta_x}(s, t) \leq \left(\frac{2}{\delta^2} \right)^S G_P(s, x) G_P(x, t).$$

Now take any $u \in \Delta_x$ and assume that $s, t \in \Delta_u$. Let v be the point on $[s, t]$ closest to u , or equivalently, to x . Then $v \in [s, x] \cap [t, x]$, so, similarly to (20), we have

$$G_P(s, x) G_P(x, t) \leq C_1^3 G_P(s, t) G_P(x, v) G_P(v, x).$$

Hence

$$G_P(s, t) - G_{P, \Delta_x}(s, t) \leq \left(\frac{2}{\delta^2}\right)^S C_1^3 G_P(s, t) G_P(x, v) G_P(v, x).$$

Therefore for (26) to be satisfied it suffices to have

$$G_P(x, v) G_P(v, x) \leq \frac{1}{4C_1^3} \left(\frac{\delta^2}{2}\right)^S \quad (27)$$

for all $v \in \Delta_u$. But, similarly to (23),

$$\sum_{v \in I} G_P(x, v) G_P(v, x) \leq (1 - \lambda)^{-2} < \infty,$$

so (27) is true for all but finitely many v in I . \square

Remark 4.6. Essentially the same proof shows that actually $K_Q(s, t_\infty)/K_P(s, t_\infty) \rightarrow 1$ as $s \rightarrow t_\infty$.

ACKNOWLEDGEMENTS

Part of the research leading to this article was carried out during the authors' visit to the Texas A&M University. The authors are grateful to Michael Brannan, Ken Dykema, Guoliang Yu and the staff of the university for their hospitality.

REFERENCES

- Anc88 A. Ancona, *Positive harmonic functions and hyperbolicity*, in *Potential theory – surveys and problems (Prague, 1987)*, Lecture Notes in Mathematics, vol. 1344 (Springer, Berlin, 1988), 1–23.
- Ban97 T. Banica, *Le groupe quantique compact libre $U(n)$* , *Comm. Math. Phys.* **190** (1997), 143–172.
- Bia94 Ph. Biane, *Théorème de Ney-Spitzer sur le dual de $SU(2)$* , *Trans. Amer. Math. Soc.* **345** (1994), 179–194.
- Car72 P. Cartier, *Fonctions harmoniques sur un arbre*, in *Symposia Mathematica*, Convegno di Calcolo delle Probabilità, INDAM, Rome, 1971 vol. IX (Academic Press, London, 1972), 203–270.
- DRVV10 A. De Rijdt and N. Vander Vennet, *Actions of monoidally equivalent compact quantum groups and applications to probabilistic boundaries*, *Ann. Inst. Fourier (Grenoble)* **60** (2010), 169–216.
- Der75 Y. Derriennic, *Marche aléatoire sur le groupe libre et frontière de Martin*, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **32** (1975), 261–276.
- DG73 Y. Derriennic and Y. Guivarc'h, *Théorème de renouvellement pour les groupes non moyennables*, *C. R. Acad. Sci. Paris Sér. A-B* **277** (1973), A613–A615.
- DM61 E. B. Dynkin and M. B. Maljutov, *Random walk on groups with a finite number of generators*, *Dokl. Akad. Nauk SSSR* **137** (1961), 1042–1045 (in Russian).
- Izu02 M. Izumi, *Non-commutative Poisson boundaries and compact quantum group actions*, *Adv. Math.* **169** (2002), 1–57.
- INO08 M. Izumi, S. Neshveyev and R. Okayasu, *The ratio set of the harmonic measure of a random walk on a hyperbolic group*, *Israel J. Math.* **163** (2008), 285–316.
- Jor17 B. P. A. Jordans, *Convergence to the boundary for random walks on discrete quantum groups and monoidal categories*, *Münster J. Math.* **10** (2017), 287–365.

- Kai00 V. A. Kaimanovich, *The Poisson formula for groups with hyperbolic properties*, Ann. of Math. (2) **152** (2000), 659–692.
- NT04 S. Neshveyev and L. Tuset, *The Martin boundary of a discrete quantum group*, J. Reine Angew. Math. **568** (2004), 23–70.
- NT13 S. Neshveyev and L. Tuset, *Compact quantum groups and their representation categories*, Cours Spécialisés, vol. 20 (Société Mathématique de France, Paris, 2013).
- NY14 S. Neshveyev and M. Yamashita, *Categorical duality for Yetter-Drinfeld algebras*, Doc. Math. **19** (2014), 1105–1139.
- NY17 S. Neshveyev and M. Yamashita, *Poisson boundaries of monoidal categories*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), 927–972.
- PW87 M. A. Picardello and W. Woess, *Martin boundaries of random walks: ends of trees and groups*, Trans. Amer. Math. Soc. **302** (1987), 185–205.
- VVV10 S. Vaes and N. Vander Vennet, *Poisson boundary of the discrete quantum group $\hat{A}_u(F)$* , Compos. Math. **146** (2010), 1073–1095.
- VV07 S. Vaes and R. Vergnioux, *The boundary of universal discrete quantum groups, exactness, and factoriality*, Duke Math. J. **140** (2007), 35–84.
- VDW96 A. Van Daele and S. Wang, *Universal quantum groups*, Internat. J. Math. **7** (1996), 255–263.
- Van08 N. Vander Vennet, *Probabilistic boundaries of discrete quantum groups*, PhD thesis, Leuven (2008).
- Ver05 R. Vergnioux, *Orientation of quantum Cayley trees and applications*, J. Reine Angew. Math. **580** (2005), 101–138.
- Woe00 W. Woess, *Random walks on infinite graphs and groups*, Cambridge Tracts in Mathematics, vol. 138 (Cambridge University Press, Cambridge, 2000).

Sara Malacarne saramal@math.uio.no

Department of Mathematics, University of Oslo, P.O. Box 1053 Blindern,
NO-0316 Oslo, Norway

Sergey Neshveyev sergeyn@math.uio.no

Department of Mathematics, University of Oslo, P.O. Box 1053 Blindern,
NO-0316 Oslo, Norway