

# The Reconstruction Property in Banach Spaces and a Perturbation Theorem

Peter G. Casazza and Ole Christensen

*Abstract.* Perturbation theory is a fundamental tool in Banach space theory. However, the applications of the classical results are limited by the fact that they force the perturbed sequence to be equivalent to the given sequence. We will develop a more general perturbation theory that does not force equivalence of the sequences.

## 1 Introduction

Perturbation theory is a very important tool in several areas of mathematics. It began with the fundamental perturbation result by Paley and Wiener [6], stating that a sequence that is sufficiently close to an orthonormal basis in a Hilbert space automatically forms a basis, that is, the *reconstruction property* is preserved. Since then, a number of variations and generalizations of this perturbation theorem have appeared, e.g., to the setting of Banach spaces (see Singer [7, pp. 84–109]). All of these generalizations have in common that a perturbation  $\{g_i\}_{i \in I}$  of a sequence  $\{f_i\}_{i \in I}$  in a Banach space  $X$  must be equivalent to  $\{f_i\}_{i \in I}$ , that is, there exists a bounded and invertible operator  $T$  on  $X$  such that  $Tf_i = g_i$  for all  $i \in I$ . This puts severe restrictions on applications of the theory. In this paper we will present a more general perturbation theory for reconstruction families in Banach spaces; it is strong enough to capture existing results, but does not force the involved sequences to be equivalent.

## 2 The Reconstruction Property

We first give a formal definition of the reconstruction property.

**Definition 2.1** Let  $X$  be a separable Banach space. A sequence  $\{f_i^*\}_{i \in I} \subset X^*$  has the *reconstruction property* for  $X$  with respect to a sequence  $\{f_i\}_{i \in I} \subset X$  if

$$(2.1) \quad f = \sum_{i \in I} f_i^*(f) f_i, \quad \text{for all } f \in X.$$

In short, we will also say that the pair  $\{f_i, f_i^*\}_{i \in I}$  has the reconstruction property for  $X$ .

It is important for our applications that  $\{f_i\}_{i \in I}$  and  $\{f_i^*\}_{i \in I}$  come from  $X$  and  $X^*$  in Definition 2.1. For example, if  $f_i^* \in \ell_\infty$  and  $\{f_i^*\}_{i \in I}$  is unitarily equivalent to the

---

Received by the editors September 30, 2005.  
The first author was supported by NSF DMS 0405376  
AMS subject classification: 42C15.  
©Canadian Mathematical Society 2008.

unit vector basis of  $\ell_2$ , then this sequence clearly has a “reconstruction” property with respect to its own predual (i.e., expansions with respect to the orthonormal basis) but this family cannot have the reconstruction property with respect to  $\ell_1$ , which is the pre-dual of  $\ell_\infty$ . We refer the reader to [2] for a generalization of the reconstruction property.

*Remark 2.2.* If  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for a Banach space  $X$ , then  $X$  has the bounded approximation property [1, p. 286]. In fact, the sequence of finite rank operators  $T_n: X \rightarrow X, T_n f = \sum_{i=1}^n f_i^*(f) f_i$  has the property that  $T_n f \rightarrow f$  in norm for all  $f \in X$ . Therefore,  $X$  is isomorphic to a complemented subspace of a Banach space with a basis [1, p. 290]. Conversely, if  $X$  has the bounded approximation property, then there exists a Banach space  $X \subset Y$  with a basis  $\{f_i, f_i^*\}$  and a projection  $P$  of  $Y$  onto  $X$ . Now,  $\{P f_i, P f_i^*\}$  has the reconstruction property for  $X$ . For information on the bounded approximation property, see [1, pp. 271–316].

We observe that the reconstruction property (2.1) is stronger than the assumption that  $\{f_i\}_{i \in I}$  spans the space  $X$ .

**Proposition 2.3** *There exists a Banach space  $X$  with the following properties:*

- (i) *there is a sequence  $\{f_i\}_{i=1}^\infty$  such that each  $f \in X$  has an expansion  $f = \sum_{i=1}^\infty a_i f_i$ ;*
- (ii)  *$X$  does not have the reconstruction property with respect to any pair  $\{h_i, h_i^*\}_{i \in I}$ .*

**Proof** Let  $X$  be a separable Banach space failing the bounded approximation property (see [1, Ch. 7]). Then  $X$  does not have the reconstruction property with respect to any family  $\{h_i, h_i^*\}_{i \in I}$ . Let  $T: \ell_1 \rightarrow X$  be a quotient map. If  $\{e_i\}_{i=1}^\infty$  is the unit vector basis of  $\ell_1$ , let  $f_i = T e_i$ . If  $f \in X$ , then there is a  $g \in \ell_1$  so that  $T g = f$ . Since  $g = \sum_{i=1}^\infty g(i) e_i$ , we have that

$$f = T g = \sum_{i=1}^\infty g(i) T e_i = \sum_{i=1}^\infty g(i) f_i. \quad \blacksquare$$

Given that  $f_i \in X$  satisfies (i) in Proposition 2.3, it would be interesting to find further conditions which guarantee the existence of  $f_i^* \in X^*$  so that  $\{f_i, f_i^*\}$  has the reconstruction property for  $X$ . This however is a very deep question, and we do not know the answer even for Hilbert spaces.

### 3 A Perturbation Theorem

For our main perturbation result we will need several standard results from Banach space theory, and we state them in the following lemma. For notation, if  $X$  is a Banach space we write  $B_X$  for the unit ball of  $X$ .

**Lemma 3.1** *Let  $X, Y$  be Banach spaces and  $T: X \rightarrow Y$  be a bounded linear operator.*

- (i) *If  $T$  is an isomorphism onto  $Y$ , then  $\|T f\| \geq A \|f\|$  for all  $f \in X$  if and only if  $A B_Y \subset T(B_X)$ .*
- (ii) *If  $T$  is an isomorphism onto  $Y$  which satisfies estimates of the form*

$$A \|f\| \leq \|T f\| \leq C \|f\|$$

for all  $f \in X$ , then for all  $g \in Y^*$  we have

$$A\|g\| \leq \|T^*g\| \leq C\|g\|.$$

(iii) If  $T$  is bounded, linear, and surjective, and  $AB_Y \subset T(B_X)$ , then  $T^*$  is an isomorphism (but not necessarily surjective), satisfying for all  $g \in Y^*$  that

$$A\|g\| \leq \|T^*g\| \leq \|T\| \|g\|.$$

The result below is a Banach space version of the Paley–Wiener theorem for frames in Hilbert space [3].

**Theorem 3.2** Suppose that  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . Let  $X_d$  be a sequence space which has the unit vectors  $\{e_i\}_{i=1}^\infty$  as a basis. Assume that

$$T\{c_i\}_{i=1}^\infty := \sum_{i=1}^\infty c_i f_i$$

defines a bounded linear operator from  $X_d$  into  $X$ . Assume further that the operator  $R: X \rightarrow X_d$  given by  $Rf = \{f_i^*(f)\}_{i=1}^\infty$  is a bounded operator. Let  $\{g_i\}$  be a sequence in  $X$  for which there exist constants  $\lambda, \mu > 0$  such that  $\lambda + \mu\|R\| < 1$  and

$$(3.1) \quad \left\| \sum_{i=1}^\infty c_i (f_i - g_i) \right\| \leq \lambda \left\| \sum_{i=1}^\infty c_i f_i \right\| + \mu \| \{c_i\}_{i=1}^\infty \|_{X_d},$$

for all finitely non-zero scalar sequences  $\{c_i\}_{i=1}^\infty$ . Then there are functionals  $\{g_i^*\}_{i=1}^\infty \subset X^*$  so that  $\{g_i, g_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ .

Moreover,  $U: X_d \rightarrow X$  given by  $U\{c_i\}_{i=1}^\infty = \sum_{i=1}^\infty c_i g_i$  is a bounded, linear, and surjective operator, and

$$(3.2) \quad \frac{1}{\|R\|} (1 - (\lambda + \mu\|R\|)) \|f\| \leq \|U^*f\| \leq \|T\| \left( 1 + \lambda + \frac{\mu}{\|T\|} \right) \|f\|$$

for all  $f \in X^*$ . Finally, if the unit vectors form an unconditional basis for  $X_d$ , then the series  $\sum_{i=1}^\infty c_i g_i$  converges unconditionally for all  $\{c_i\}_{i=1}^\infty \in X_d$ .

**Proof** For all finite sequences  $\{c_i\}_{i=1}^n$  we have

$$\left\| \sum_{i=1}^n c_i g_i \right\| \leq \left\| \sum_{i=1}^n c_i (f_i - g_i) \right\| + \left\| \sum_{i=1}^n c_i f_i \right\| \leq (1 + \lambda) \left\| \sum_{i=1}^n c_i f_i \right\| + \mu \| \{c_i\}_{i=1}^n \|_{X_d}.$$

It follows that for all  $\{c_i\}_{i=1}^\infty \in X_d$  and all  $n > m$  in  $\mathbb{N}$ ,

$$(3.3) \quad \left\| \sum_{i=1}^n c_i g_i - \sum_{i=1}^m c_i g_i \right\| = \left\| \sum_{i=m+1}^n c_i g_i \right\| \leq (1 + \lambda) \left\| \sum_{i=m+1}^n c_i f_i \right\| + \mu \left\| \sum_{i=m+1}^n c_i e_i \right\|_{X_d}.$$

Since  $\sum_{i=1}^\infty c_i e_i$  converges by the fact that  $\{c_i\}_{i=1}^\infty \in X_d$  and the unit vectors form a basis for  $X_d$ , it follows that  $\sum_{i=1}^\infty c_i f_i$  converges by our assumption that  $T$  is a bounded operator. Now it follows from (3.3) that  $\sum_{i=1}^\infty c_i g_i$  converges in  $X$  (unconditionally if the unit vectors form an unconditional basis for  $X_d$ ). If we define  $U: X_d \rightarrow X$  by  $U\{c_i\}_{i=1}^\infty = \sum_{i=1}^\infty c_i g_i$ , we have

$$\begin{aligned} \|U\{c_i\}_{i=1}^\infty\| &\leq (1 + \lambda)\|T\{c_i\}_{i=1}^\infty\| + \mu\|\{c_i\}_{i=1}^\infty\|_{X_d} \\ &\leq ((1 + \lambda)\|T\| + \mu)\|\{c_i\}_{i=1}^\infty\|_{X_d}. \end{aligned}$$

Hence,  $\|U\| = \|U^*\| \leq ((1 + \lambda)\|T\| + \mu)$ , which verifies the right-hand side of (3.2). Next, define an operator  $L: X \rightarrow X$  by  $Lf = \sum_{i=1}^\infty f_i^*(f)g_i$ . For any  $f \in X$  we have

$$\begin{aligned} \|(I - L)f\| &= \|f - Lf\| = \left\| \sum_{i=1}^\infty f_i^*(f)f_i - \sum_{i=1}^\infty f_i^*(f)g_i \right\| = \left\| \sum_{i=1}^\infty f_i^*(f)(f_i - g_i) \right\| \\ &\leq \lambda \left\| \sum_{i=1}^\infty f_i^*(f)f_i \right\| + \mu\|\{f_i^*(f)\}_{i=1}^\infty\|_{X_d} = \lambda\|f\| + \mu\|Rf\|_{X_d} \\ &\leq \lambda\|f\| + \mu\|R\|\|f\| = (\lambda + \mu\|R\|)\|f\|. \end{aligned}$$

Since  $\lambda + \mu\|R\| < 1$ , it follows that  $L$  is an invertible operator on  $X$ . Now let  $g_i^* = (L^{-1})^* f_i^*$  for all  $i \in \mathbb{N}$ . If  $f \in X$ , we have

$$\sum_{i=1}^\infty g_i^*(f)g_i = \sum_{i=1}^\infty [(L^{-1})^* f_i^*](f)g_i = \sum_{i=1}^\infty f_i^*(L^{-1}f)g_i = LL^{-1}f = f.$$

So  $\{g_i, g_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ .

In order to prove the left-hand side of (3.2), we note that for  $f \in X$

$$\begin{aligned} \|Lf\| &\geq \|f\| - \|(I - L)(f)\| \\ &\geq \|f\| - (\lambda + \mu\|R\|)\|f\| = (1 - (\lambda + \mu\|R\|))\|f\|. \end{aligned}$$

Now consider  $U: R(X) \rightarrow X$  as defined above. If  $f \in B_X$ , then

$$\left\| \left\{ \frac{1}{\|R\|} f_i^*(f) \right\}_{i=1}^\infty \right\|_{X_d} = \frac{1}{\|R\|} \|R(f)\|_{X_d} \leq \frac{1}{\|R\|} \|R\|\|f\| = 1.$$

So,  $\left\{ \frac{1}{\|R\|} f_i^*(f) \right\}_{i=1}^\infty \in B_{X_d}$ . Also,

$$\left\| U \left\{ \frac{1}{\|R\|} f_i^*(f) \right\}_{i=1}^\infty \right\| = \frac{1}{\|R\|} \|Lf\| \geq \frac{1}{\|R\|} (1 - (\lambda + \mu\|R\|)) \|f\|.$$

By Lemma 3.1(i),  $U(B_{X_d}) \supset U(B_{R(X)}) \supset \frac{1}{\|R\|}(1 - (\lambda + \mu\|R\|))B_X$ . By Lemma 3.1(iii), we have  $\|U^* f\| \geq \frac{1}{\|R\|}(1 - (\lambda + \mu\|R\|))\|f\|$ . ■

We now consider some applications of Theorem 3.2.

**Example 3.3** Choose  $\lambda, \mu > 0$  so that  $\lambda + \mu\|R\| < 1$ , ( $T, R$  as in Theorem 3.2). Choose any bounded linear operator  $L: X_d \rightarrow X$  with  $\|L\| \leq \|R\|$ . For  $i \in \mathbb{N}$  let

$$g_i = (1 - \lambda)f_i + \mu Le_i,$$

where  $\{e_i\}_{i=1}^\infty$  is a basis of  $X_d$ . Then for all  $\{c_i\}_{i=1}^\infty \in X_d$  we have

$$\begin{aligned} \left\| \sum_{i=1}^{\infty} c_i(f_i - g_i) \right\| &= \left\| \lambda \sum_{i=1}^{\infty} c_i f_i + \mu \sum_{i=1}^{\infty} c_i Le_i \right\| \\ &\leq \lambda \left\| \sum_{i=1}^{\infty} c_i f_i \right\| + \mu \left\| \sum_{i=1}^{\infty} c_i Le_i \right\| \\ &\leq \lambda \left\| \sum_{i=1}^{\infty} c_i f_i \right\| + \mu \|L\| \|\{c_i\}_{i=1}^\infty\|_{X_d} \\ &\leq \lambda \left\| \sum_{i=1}^{\infty} c_i f_i \right\| + \mu \|R\| \|\{c_i\}_{i=1}^\infty\|_{X_d} \end{aligned}$$

So the hypotheses of Theorem 3.2 are satisfied.

Another natural application of Theorem 3.2 is to take a Banach space  $X_d$  with a basis  $\{g_i, g_i^*\}$  and let  $P$  be a projection on  $X_d$ . Letting  $X = P(X_d)$ ,  $f_i = P(g_i)$ ,  $f_i^* = P(g_i^*)$ ,  $T = P$ , and with  $R$  the injection of  $X$  into  $X_d$ , we can apply the theorem.

An important aspect of Theorem 3.2 is that it does not require the perturbed family  $\{g_i\}$  to be equivalent to the original reconstruction sequence  $\{f_i\}$ . We will give an example of this below. Recall that two sequences  $\{f_i\}_{i \in I}$ ,  $\{g_i\}_{i \in I}$  in a Banach space are *equivalent* if the mapping  $T: f_i \mapsto g_i$  can be extended to a well-defined bounded linear map of  $\overline{\text{span}}\{f_i\}$  onto  $\overline{\text{span}}\{g_i\}$ .

**Proposition 3.4** *There is a Banach space  $X$  and a pair  $\{f_i, f_i^*\}_{i=1}^\infty$  having the reconstruction property for  $X$ , a sequence space  $X_d$  with an unconditional basis  $\{e_i\}_{i=1}^\infty$  so that the operators  $T, R$  in Theorem 3.2 exist and there is a sequence  $\{g_i\}_{i=1}^\infty$  in  $X$  satisfying the perturbation criterion (3.1), but  $\{g_i\}_{i=1}^\infty$  is not equivalent to  $\{f_i\}_{i=1}^\infty$ .*

**Proof** Let  $P$  be a non-trivial projection on  $\ell_p$  onto a subspace for any  $1 \leq p < \infty$ . Let  $X_d = \ell_p$ ,  $X = P(X_d)$ . With the notation in Theorem 3.2, let  $T = P$  and  $R$  be the injection of  $X$  into  $X_d$ . Since  $X$  is isomorphic to  $\ell_p$  (see [5]), there is an isomorphism  $L: X_d \rightarrow X$ . Now choose  $\lambda, \mu > 0$  so that  $\lambda + \mu \max\{\|L\|, \|R\|\} < 1$ . Let  $\{e_i\}_{i=1}^\infty$  be the unit vector basis of  $\ell_p$ , let  $f_i = Pe_i$  for all  $i = 1, 2, \dots$ , and let  $g_i = (1 - \lambda)f_i + \mu Le_i$ . For all finitely non-zero sequences  $\{c_i\}_{i=1}^\infty$  we have

$$\begin{aligned} \left\| \sum_{i=1}^{\infty} c_i(f_i - g_i) \right\| &\leq \lambda \left\| \sum_{i=1}^{\infty} c_i f_i \right\| + \mu \left\| \sum_{i=1}^{\infty} c_i Le_i \right\| \\ &\leq \lambda \left\| \sum_{i=1}^{\infty} c_i f_i \right\| + \mu \|L\| \|\{c_i\}_{i=1}^\infty\|_{X_d}. \end{aligned}$$

So  $\{g_i\}_{i=1}^\infty$  is a perturbation of  $\{f_i\}_{i=1}^\infty$ . If we choose any  $\{0\} \neq \{c_i\}_{i=1}^\infty \in X_d$  so that  $\sum_{i=1}^\infty c_i p e_i = \sum_{i=1}^\infty c_i f_i = 0$ , then

$$\sum_{i=1}^\infty c_i g_i = (1 - \lambda) \sum_{i=1}^\infty c_i f_i + \mu \sum_{i=1}^\infty c_i L e_i = \mu \sum_{i=1}^\infty c_i L e_i.$$

Since  $\{L e_i\}_{i=1}^\infty$  is a basis for  $X$ , it follows that  $\sum_{i=1}^\infty c_i L e_i \neq 0$ , and so  $\{f_i\}_{i=1}^\infty$  is not equivalent to  $\{g_i\}_{i=1}^\infty$ . ■

We will now show that the conclusion in Theorem 3.2 can be obtained under weaker assumptions. Let us discuss why this is important. In Theorem 3.2, it is easily checked that the operator  $RT$  is a projection of  $X_d$  onto  $R(X)$ . This is a rather strong restriction on the application of the result. As we saw earlier, the very existence of a reconstruction family implies that  $X$  is isomorphic to a complemented subspace of a Banach space with a basis. However, the space with a basis may not be the space  $X_d$  above. The next result has the advantage that it does not require that  $X$  be isomorphic to a complemented subspace of  $X_d$ , but just that it embed into  $X_d$ . The proof follows line by line the proof of Theorem 3.2 using  $R^{-1}$  in place of  $T$ .

**Theorem 3.5** *Suppose  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for a Banach space  $X$ . Let  $X_d$  be a sequence space which has the unit vectors as a basis. Assume the operator  $R: X \rightarrow X_d$  given by  $Rf = \{f_i^*(f)\}_{i=1}^\infty$  is a (not necessarily surjective) isomorphism. Let  $\{g_i\}_{i=1}^\infty$  be a sequence in  $X$  for which there exist constants  $\lambda, \mu > 0$  such that  $\lambda + \mu\|R\| < 1$  and*

$$\left\| \sum_{i=1}^\infty c_i (f_i - g_i) \right\| \leq \lambda \left\| \sum_{i=1}^\infty c_i f_i \right\| + \mu \|\{c_i\}_{i=1}^\infty\|_{X_d},$$

for all finitely non-zero scalar sequences  $\{c_i\}_{i=1}^\infty$  taken from  $\{f_i^*(f)\}_{i=1}^\infty$  for any  $f \in X$ . Then there are functionals  $\{g_i^*\}_{i=1}^\infty \subset X^*$  so that  $\{g_i, g_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ .

Moreover,  $U: R(X) \rightarrow X$  given by  $U\{c_i\}_{i=1}^\infty = \sum_{i=1}^\infty c_i g_i$  is an isomorphism satisfying for all  $f \in X^*$ ,

$$\frac{1}{\|R\|} (1 - (\lambda + \mu\|R\|)) \|f\| \leq \|U^* f\| \leq \|T\| ((1 + \lambda)\|R^{-1}\| + \mu) \|f\|.$$

### 4 The Reconstruction Property Revisited

We will now consider some theoretical consequences of the reconstruction property and related examples.

**Proposition 4.1** *Suppose that  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . Then for all  $g \in X^*$  we have that the sequence*

$$\left\{ \sum_{i=1}^n g(f_i) f_i^* \right\}_{n=1}^\infty$$

converges to  $g \in X^*$  in the  $\omega^*$ -topology.

**Proof** For any  $f \in X$  we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[ \left( \sum_{i=1}^n g(f_i) f_i^* \right) (f) \right] &= \lim_{n \rightarrow \infty} \sum_{i=1}^n g(f_i) f_i^*(f) = \lim_{n \rightarrow \infty} g \left( \sum_{i=1}^n f_i^*(f) f_i \right) \\ &= g \left( \lim_{n \rightarrow \infty} \sum_{i=1}^n f_i^*(f) f_i \right) = g(f). \end{aligned}$$

This proves the proposition. ■

In the case that  $X$  is reflexive, the convergence in Proposition 4.1 becomes weak convergence. It is natural to ask whether we also obtain convergence in norm in this case. Unfortunately, this fails. Even in a Hilbert space, having the reconstruction property with respect to  $\{f_i, f_i^*\}$  does not imply the reconstruction property for  $\{f_i^*, f_i\}$ :

**Example 4.2** Let  $H$  be a separable Hilbert space. There are vectors  $f_i^*, f_i \in H$  so that for every  $f \in H$  we have  $f = \sum_{i=1}^\infty f_i^*(f) f_i$  but we do not have that  $f = \sum_{i=1}^\infty f(f_i) f_i^*$  for all  $f \in H$ .

To see this, let  $\{e_i\}_{i=1}^\infty$  be an orthonormal basis for  $H$ , and define the vectors  $\{f_i\}_{i=1}^\infty$  and  $\{f_i^*\}_{i=1}^\infty$  by

$$f_{2i} = e_i, f_{2i-1} = e_1, \quad f_1^* = e_1, \quad f_{2i}^* = e_i, \quad f_{2i+1}^* = e_{i+1} - e_i.$$

Now, for all  $f \in H$ ,

$$\begin{aligned} \sum_{i=1}^\infty f_{2i}^*(f) f_{2i} &= f, \\ \sum_{i=1}^n f_{2i+1}^*(f) f_{2i+1} &= \langle e_1, f \rangle e_1 + \left( \sum_{i=1}^n \langle e_{i+1} - e_i, f \rangle \right) e_1 = \langle e_{n+1}, f \rangle e_1. \end{aligned}$$

Since  $\lim_{n \rightarrow \infty} \langle e_{n+1}, f \rangle = 0$ , we have that  $\sum_{i=1}^\infty f_{2i+1}^*(f) f_{2i+1} = 0$ . Hence, for all  $f \in H$ ,  $f = \sum_{i=1}^\infty f_i^*(f) f_i$ . On the other hand, if  $f = e_1$ , then

$$\begin{aligned} \sum_{i=1}^{2n+1} f(f_i) f_i^* &= \langle e_1, f_2 \rangle f_2^* + \sum_{i=0}^n \langle e_1, f_{2i+1} \rangle f_{2i+1}^* \\ &= e_1 + \sum_{i=0}^n \langle e_1, e_1 \rangle (e_{i+1} - e_i) = e_1 + e_{n+1}. \end{aligned}$$

It follows that  $\sum_{i=1}^\infty f(f_i) f_i^*$  does not converge in  $H$ .

The next proposition shows that we can get the reconstruction property with respect to  $X^*$  if the reconstruction property for  $X$  holds with unconditional convergence.

**Proposition 4.3** Assume that  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$  and that the series  $\sum_{i=1}^\infty f_i^*(f) f_i (= f)$  converges unconditionally for all  $f \in X$ . Then the following are equivalent.

- (i) For all  $g \in X^*$  we have  $g = \sum_{i=1}^\infty g(f_i) f_i^*$ .
- (ii)  $c_0$  does not embed into  $X^*$ .

**Proof** (i)  $\Rightarrow$  (ii): By (i),  $X^*$  is separable and so  $c_0$  cannot embed into  $X^*$  [5].

(ii)  $\Rightarrow$  (i): For  $E \subset \mathbb{N}$  finite, define  $T_E f = \sum_{i \in E} g_i(f) f_i$ . The family  $\{T_E\}$  is a family of finite rank bounded linear operators on  $X$  which are pointwise bounded because of the unconditional convergence of  $\sum_{i=1}^\infty f_i^*(f) f_i$ . By the uniform boundedness principle, this family is uniformly bounded, i.e.,

$$\sup_{E \subset \mathbb{N}} \|T_E\| = K < \infty.$$

If  $E, F$  are finite subsets of  $\mathbb{N}$ , then

$$\begin{aligned} \left\| \sum_{i \in E} g(f_i) f_i^* - \sum_{i \in F} g(f_i) f_i^* \right\| &= \sup_{\|f\|=1} \left| \sum_{i \in E} g(f_i) f_i^*(f) - \sum_{i \in F} g(f_i) f_i^*(f) \right| \\ &\leq \sup_{\|f\|=1} \left( \left| g\left(\sum_{i \in E} f_i^*(f) f_i\right) \right| + \sum_{i \in F} \left| g\left(\sum_{i \in F} f_i^*(f) f_i\right) \right| \right) \\ &\leq \|g\| \left( \sup_{\|f\|=1} \left\| \sum_{i \in E} f_i^*(f) f_i \right\| + \sup_{\|f\|=1} \left\| \sum_{i \in F} f_i^*(f) f_i \right\| \right) \\ &\leq 2K \|g\| \|f\|. \end{aligned}$$

By [4, Theorem 6, p. 44], it follows that  $\sum_{i=1}^\infty g(f_i) f_i^*$  is weakly unconditionally Cauchy. Since  $c_0$  does not embed into  $X^*$ , by [4, Theorem 8, p. 45], we have that

$$\sum_{i=1}^\infty g(f_i) f_i^*$$

is unconditionally convergent in  $X^*$ . Since this series converges weakly to  $g$  by Proposition 4.1, we have that  $g = \sum_{i=1}^\infty g(f_i) f_i^*$  and the series converges unconditionally. ■

Recall that we say a subspace  $Y \subset X^*$  norms  $X$  if there is a constant  $A > 0$  so that for all  $f \in X$  we have

$$A \|f\| \leq \sup_{\|g\|=1, g \in Y} |g(f)|.$$

**Proposition 4.4** Assume  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . Then  $\overline{\text{span}}\{f_i^*\}_{i=1}^\infty$  norms  $X$ .

**Proof** For all  $f \in X$  we have  $f = \sum_i f_i^*(f) f_i$ . It follows that the finite rank operators  $T_n(f) = \sum_{i=1}^n f_i^*(f) f_i$ , are pointwise bounded on  $X$ . By the uniform boundedness principle, there exists a constant  $K$  so that for all  $n$  and all  $f \in X$ ,

$$\left\| \sum_{i=1}^n f_i^*(f) f_i \right\| \leq K \|f\|.$$



Now, for every  $g \in X^*$  we have,

$$\begin{aligned} \left\| \sum_{i=1}^n g(f_i) f_i^* \right\| &= \sup_{\|f\|=1} \left| \left( \sum_{i=1}^n g(f_i) f_i^* \right) (f) \right| = \sup_{\|f\|=1} \left| \sum_{i=1}^n g(f_i) f_i^*(f) \right| \\ &= \sup_{\|f\|=1} \left| g \left( \sum_{i=1}^n f_i^*(f) f_i \right) \right| \\ &\leq \|g\| \sup_{\|f\|=1} \left\| \sum_{i=1}^n f_i^*(f) f_i \right\| \\ &\leq K \|g\| \|f\|. \end{aligned}$$

Now fix  $f \in X$   $\epsilon > 0$  and choose  $g \in X^*$  so that  $\|g\| = 1$  and  $\|f\| \leq (1 + \epsilon)|g(f)|$ . We have (for  $n$  sufficiently large),

$$\begin{aligned} \|f\| &\leq (1 + \epsilon)|g(f)| \leq (1 + \epsilon) \left| \sum_{i=1}^{\infty} f_i^*(f) g(f_i) \right| \\ &\leq (1 + \epsilon)^2 \left| \sum_{i=1}^n f_i^*(f) g(f_i) \right| \leq (1 + \epsilon)^2 \left| \left( \sum_{i=1}^n g(f_i) f_i^* \right) (f) \right|. \end{aligned}$$

Since  $\sum_{i=1}^n g(f_i) f_i^*$  is in the closed linear span of the  $\{f_i^*\}_{i=1}^{\infty}$  and since the norms of these vectors are uniformly bounded, it follows that the space  $\overline{\text{span}}\{f_i^*\}_{i=1}^{\infty}$  norms  $X$ . ■

**Proposition 4.5** *If  $\{f_i, f_i^*\}_{i=1}^{\infty}$  has the reconstruction property for  $X$  and  $\{f_i^*\}_{i=1}^{\infty}$  is equivalent to  $\{g_i^*\}_{i=1}^{\infty}$  in  $X^*$ , then  $\{g_i^*\}_{i=1}^{\infty}$  has the reconstruction property with respect to some Banach space  $Y$  and elements  $\{g_i\}_{i=1}^{\infty}$  in  $Y$ .*

**Proof** Given the isomorphism  $Tg_i^* = f_i^*$ , let  $g_i = T^* f_i$ . Note that  $f_i$  is a linear functional on the closed linear span of the  $\{f_i^*\}$  in an obvious way; but, its norm as a linear functional may not be the same as its norm as an element of  $X$ . However, by Proposition 4.4, these norms are equivalent.

Now for  $f \in X$

$$\begin{aligned} \sum_{i=1}^{\infty} g_i^*(f) g_i &= T^* \sum_{i=1}^{\infty} (T^{-1} f_i^*)(f) f_i \\ &= T^* \sum_{i=1}^{\infty} f_i^*((T^{-1})^* f) f_i = T^*(T^{-1})^* f = f. \end{aligned}$$

So  $\{g_i, g_i^*\}_{i=1}^{\infty}$  has the reconstruction property for  $X$ . ■

We can strengthen the results in the particular case of a reflexive Banach space.

**Theorem 4.6** Let  $X$  be reflexive. Assume  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . Let  $\{g_i^*\}_{i=1}^\infty$  be elements of  $X^*$ . Assume there is a  $0 < \lambda < 1$  so that for all  $n \in \mathbb{N}$  and all sequences of scalars  $\{a_i\}_{i=1}^n$  we have

$$\left\| \sum_{i=1}^n a_i(f_i^* - g_i^*) \right\| \leq \lambda \left\| \sum_{i=1}^n a_i f_i^* \right\|.$$

Then there are vectors  $\{g_i\}_{i=1}^\infty \subset X$  so that for all  $f \in X$   $f = \sum_{i=1}^\infty g_i^*(f)g_i$ .

**Proof** We first observe that  $\overline{\text{span}}\{f_i^*\}_{i=1}^\infty = X^*$ . In fact, if this were not the case, the reflexivity of  $X$  would imply the existence of an element  $f \in X^{**} = X$  so that  $f(g) = 0$  for all  $g \in \overline{\text{span}}\{f_i^*\}_{i=1}^\infty$ ; but then  $\sum_{i=1}^\infty f_i^*(f)f_i = 0$ , which contradicts the assumption that  $\{f_i, f_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . Now define  $T: X^* \rightarrow X^*$  by  $T(f_i^*) = g_i^*$ . Since  $\{g_i^*\}_{i=1}^\infty$  is a perturbation of  $\{f_i^*\}_{i=1}^\infty$ , this is a well-defined operator on  $X^*$ . But the perturbation condition implies that  $\|I - T\| \leq \lambda < 1$ . Hence,  $T$  is an invertible operator on  $X^*$ , and

$$\sum_{i=1}^\infty g_i^*(f)(T^*)^{-1}f_i = (T^*)^{-1} \sum_{i=1}^\infty f_i^*(T^*f)f_i = (T^*)^{-1}T^*f = f.$$

So  $\{(T^*)^{-1}f_i, g_i^*\}_{i=1}^\infty$  has the reconstruction property for  $X$ . ■

Unfortunately, besides the reflexive case, a perturbation of a family with the reconstruction property need not have the reconstruction property.

**Example 4.7** Let  $X = c_0$  so  $X^* = \ell_1$ . Let  $\{e_i\}$  (respectively,  $\{e_i^*\}$ ) be the unit vector basis of  $X$  (respectively,  $X^*$ ). Define  $f_i^* = e_i^*$ ,  $f_i = e_i$ , for all  $i = 1, 2, 3, \dots$ . Also, let  $g_1^* = e_1^*$ ,  $g_i^* = \frac{1}{2}e_1^* + e_i^*$ , for all  $i \geq 2$ . Of course,  $\{f_i^*\}_{i=1}^\infty$  has the reconstruction property with respect to  $\{f_i\}_{i=1}^\infty$ . Also, for all  $n \in \mathbb{N}$  and all families of scalars  $\{a_i\}_{i=1}^n$  we have

$$\begin{aligned} \left\| \sum_{i=1}^n a_i(f_i^* - g_i^*) \right\| &= \left\| \sum_{i=1}^n a_i e_i^* - \left( \sum_{i=1}^n a_i e_i^* + \frac{1}{2} \sum_{i=2}^n a_i e_1^* \right) \right\| \\ &= \frac{1}{2} \left| \sum_{i=2}^n a_i \right| \leq \frac{1}{2} \sum_{i=1}^n |a_i| = \frac{1}{2} \left\| \sum_{i=1}^n a_i f_i^* \right\|. \end{aligned}$$

So  $\{g_i^*\}_{i=1}^\infty$  is a perturbation of  $\{f_i^*\}_{i=1}^\infty$  and hence is a basic sequence in  $\ell_1$  which is equivalent to the unit vector basis of  $\ell_1$ . But also,  $e_i^* = g_i^* - \frac{1}{2}g_1^*$  for all  $i = 2, 3, \dots$ . It follows that  $\{g_i^*\}_{i=1}^\infty$  is actually a basis for  $\ell_1$  equivalent to the unit vector basis.

We proceed, by way of contradiction, to show that this family  $\{g_i^*\}$  does not have the reconstruction property with respect to any sequence of vectors in  $c_0$ . So, assume  $\{g_i\}_{i=1}^\infty \subset c_0$  satisfies that  $f = \sum_{i=1}^\infty g_i^*(f)g_i$ , for all  $f \in c_0$ . Then for all  $j \geq 2$  we have  $e_j = \sum_{i=1}^\infty g_i^*(e_j)g_i = g_j$ . Also,

$$e_1 = \sum_{i=1}^\infty g_i^*(e_1)g_i = g_1 + \sum_{i=2}^\infty \frac{1}{2}g_i = g_1 + \sum_{i=2}^\infty \frac{1}{2}e_i.$$

Hence,  $g_1 = e_1 - \frac{1}{2} \sum_{i=2}^{\infty} e_i$ . It follows that  $g_1 \notin c_0$ , contradicting our assumption. So  $\{g_i^*\}_{i=1}^{\infty}$  does not have the reconstruction property with respect to any sequence of vectors in  $c_0$ .

## References

- [1] P. G. Casazza, *Approximation properties*. In: Handbook on the Geometry of Banach Spaces. Vol. 1. Elsevier, Amsterdam, 2001, pp. 271–316.
- [2] P. G. Casazza, O. Christensen, and D. T. Stoeva, *Frame expansions in separable Banach spaces*. J. Math. Anal. Appl. **307**(2005), no. 2, 710–723.
- [3] O. Christensen, *A Paley-Wiener theorem for frames*. Proc. Amer. Math. Soc. **123**(1995), no. 7, 2199–2202.
- [4] J. Diestel, *Sequences and Series in Banach Spaces*. Graduate Texts in Mathematics 92, Springer-Verlag, New York, 1984.
- [5] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces. I. Sequence Spaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete 92, Springer-Verlag, Berlin, 1977.
- [6] R. E. A. C. Paley and N. Wiener, *Fourier transforms in the complex domain*. Reprint of the 1934 original. American Mathematical Society Colloquium Publications 19. American Mathematical Society, Providence, RI, 1987.
- [7] I. Singer, *Bases in Banach Spaces*. Die Grundlehren der Mathematischen Wissenschaften 154, Springer-Verlag, New York, 1970.

*Department of Mathematics, University of Missouri, Columbia, Mo 65211, U.S.A.*  
*e-mail:* pete@casazza.math.missouri.edu

*Technical University of Denmark, Department of Mathematics, 2800 Lyngby, Denmark*  
*e-mail:* Ole.Christensen@mat.dtu.dk