

Mechanisms of intestinal phosphate transport in small ruminants

BY B. SCHRÖDER¹, H. KÄPPNER¹, K. FAILING¹, E. PFEFFER²
AND G. BREVES¹

¹ Department of Veterinary Physiology, Justus-Liebig-University, Giessen, Germany

² Department of Animal Nutrition, Rheinische Friedrich-Wilhelms-University, Bonn, Germany

(Received 25 April 1994 – Revised 5 January 1995 – Accepted 28 February 1995)

In order to study the localization and mechanisms of intestinal phosphate transport in sheep and goats, unidirectional inorganic phosphate (P_i) flux rates across isolated stripped epithelial tissues were measured *in vitro* by applying the Ussing-chamber technique. In the first experiment the tissues were obtained from animals which had been kept on an adequate dietary P supply. In the second experiment the animals had either been kept on an adequate Ca and P supply or were Ca- and/or P-depleted. Significant net P_i absorption was measured in all segments of the small intestine and in the proximal colon of sheep and in the duodenum and jejunum of goats. Since the experiments were carried out in the absence of any electrochemical gradient, this clearly indicates the presence of active mechanisms for P_i transport in the intestinal tract of small ruminants. In sheep jejunum, reduction of mucosal Na concentration to 1.8 mM or serosal application of ouabain (0.1 mM) resulted in significant decreases of net P_i absorption of the same order of magnitude, indicating that about 65% of active P_i transport in sheep jejunum is mediated by a Na-dependent active transport mechanism. The mechanism for the remaining Na⁺-independent active P_i transport has not yet been identified. Dietary P depletion caused hypophosphataemia and induced a significant stimulation of net P_i absorption in goat duodenum and jejunum. This increase was independent of dietary Ca supply and was not associated with increased plasma calcitriol concentrations. This suggests substantial differences in hormonal regulation of P_i transport in small ruminants in comparison with single-stomached species.

Intestinal phosphate absorption: Phosphorus: Calcium: Ruminant

In ruminants large amounts of inorganic phosphate (P_i) enter the gastrointestinal tract by salivary secretion, and this endogenous secretion is well balanced by net P_i absorption from the intestinal tract. Thus the overall absorption rate from the intestinal tract is substantially higher compared with single-stomached animals (for review see Breves & Schröder, 1991). Although it is well known that in ruminants the upper small intestine is the major site for net P_i absorption, little is known about the epithelial transport processes involved and their hormonal regulation. This is in contrast to many single-stomached species where a calcitriol-stimulated secondary active Na⁺- P_i -cotransport system has been identified as the predominant mechanism for active P_i absorption (Bruce *et al.* 1966/67; Pfeffer *et al.* 1970; Grace *et al.* 1974; Berner *et al.* 1976; Murer & Hildmann, 1981; Brandis *et al.* 1987; Danisi *et al.* 1988; Shirazi-Beechey *et al.* 1988).

For sheep a carrier-mediated mechanism has been assumed for mucosal P_i uptake from *in vivo* studies (Care *et al.* 1980; Scott *et al.* 1984). From a recent *in vitro* study with isolated brush border membrane vesicles from sheep, evidence was obtained for active P_i uptake, which was driven by a transmembrane proton gradient and which was stimulated in response to dietary P depletion (Shirazi-Beechey *et al.* 1991). However, it has to be excluded that this adaptation was controlled by changes in plasma calcitriol since it is well known from different studies that in ruminants neither calcitriol nor its production rate is changed

in response to dietary P depletion as has been described for single-stomached species (Breves *et al.* 1985; Maunder *et al.* 1986; Schröder *et al.* 1990).

In order to get more detailed knowledge about intestinal P_i transport mechanisms in sheep and goats, Ussing chambers were used for studying transepithelial P_i flux rates in the absence of electrochemical gradients. In the first experiment the intestinal tissues were obtained from animals which had been kept on an adequate P supply. In the second experiment the tissues were taken from growing goats which had either been kept on an adequate dietary Ca and P supply or were Ca- and/or P-depleted.

MATERIALS AND METHODS

Animals and feeding

In Expt 1, intestinal tissues were obtained from adult sheep and goats weighing between 10 and 65 kg, which had been kept on an adequate dietary Ca and P supply. In Expt 2 the tissues were taken from male growing goats of 4–5 months of age. Nine weeks before slaughter animals were allotted to four different dietary regimens with either adequate or reduced Ca and/or P contents (Table 1). From all animals of Expt 2, blood samples were taken the day before slaughter and plasma was stored at -70° . Intestinal segments were taken within 3 min after exsanguination, from either the descending part of the duodenum, mid jejunum, terminal ileum or proximal colon. They were immediately rinsed with ice-cold physiological saline (9 g NaCl/l) and kept in a glucose-containing buffer solution at 4° , being continuously gassed with O_2 - CO_2 (95:5 v/v).

Incubation of epithelial tissues and calculation of P_i flux rates

In each experiment intestinal segments were opened along the mesenteric line and fixed on a Pyrex plate with the mucosal surface uppermost. Epithelial tissues were stripped off the muscle layers and mounted in computer-controlled Ussing chambers (Mussler Microclamp, Aachen, Germany) with an exposed area of 100 mm². The tissues were incubated on both sides with 13 ml of an isotonic buffer solution (pH 7.4) containing (mmol/l) NaCl 125.4, KCl 5.4, $CaCl_2$ 1.2, $NaHCO_3$ 21, Na_2HPO_4 0.3, NaH_2PO_4 1.2 and indomethacin 0.01. In addition, the serosal solution contained 10 mM glucose, which for osmotic reasons was compensated on the mucosal side by 10 mM-mannitol. In some experiments choline and K salts were used to reduce Na concentrations from 148.2 mM to 1.8 mM. In order to check the functional viability of the tissues at the start and the end of flux measurements maximal increases of short-circuit currents (I_{sc}) were determined in a number of experiments in response to either theophylline (10 mM serosal) or glucose (10 mM mucosal) or L-alanine (5 mM mucosal).

For calculations of unidirectional P_i flux rates (J_{ms} : flux from mucosal to serosal side, J_{sm} : flux in the opposite direction), 185 KBq [³²P]orthophosphate (370 MBq/ml, Amersham Buchler, Braunschweig, Germany) were added to one side of the tissue. Samples (0.1 ml) were taken from the labelled side 20 min later and subsequently in at least three 10 min intervals from the unlabelled side (0.5 ml) of the Ussing chamber. All samples taken from the unlabelled side were replaced by equal volumes of isosmotic bathing fluid. Net fluxes (J_{net}) were calculated as differences between J_{ms} and J_{sm} of paired tissues whose conductances did not differ by more than 25%. In another series of experiments flux measurements were done before and after the addition of arsenate (mucosal) or ouabain (serosal) to the bathing solution. Radioactivity measurements were done in a Packard Tricarb liquid scintillation counter with counting accuracies > 95%. For further experimental details see Schröder *et al.* (1991, 1993a). Unidirectional P_i -flux rates were calculated using standard equations (Schultz & Zalusky, 1964).

Table 1. *Dietary calcium and phosphorus contents of rations fed to growing goats*

Grouping*		Feed (g/kg dry matter)	
Ca	P	Ca	P
+	+	11.1	4.6
+	-	10.1	2.0
-	+	4.2	4.5
-	-	3.9	1.9

+, adequate mineral content; -, reduced mineral content.

Analytical procedures

Plasma Ca and P_i were measured using test kits provided by Boehringer (Mannheim, Germany).

Plasma parathyroid hormone (PTH) was determined using a commercial PTH assay system, which detects the midregional 44–68 amino acid sequence fragment of the entire human PTH polypeptide (Immunodiagnostik, Bensheim, Germany). This assay system was used due to lack of appropriate kits to measure intact PTH or PTH fragments in goat plasma. The detection limit for measuring the hPTH(44–68) immunoreactive fraction in goat plasma was < 16 pM. Calcitriol concentrations in goat plasma were measured according to the method of Hollis (1986) involving single cartridge extraction and separation of calcitriol from plasma (Bond Elut C18-OH; ICT, Bad Homburg, Germany) and subsequent determination of calcitriol using a competitive protein-binding assay with a preparation of calf thymus cytosol (Reinhardt & Horst, 1990). To assess sufficient analytical recovery of calcitriol the extraction and separation procedure was modified as follows: to 2 ml goat plasma 10 µl [³H]calcitriol in ethanol (67 Bq, specific activity 6.36 TBq/mmol; Amersham Buchler) was added. After vortex-mixing the sample was incubated for 60 min at room temperature. Acetonitrile extraction of plasma vitamin D metabolites from the sample was performed twice for 30 min each on a shaker. After centrifugation and dilution of combined supernatant fractions with 1 volume of 0.4 mM-K₂HPO₄ (pH 10.5) the entire extract was applied to a prewashed C18-OH cartridge (2 ml hexane, 2 ml isopropanol, 2 ml water). Separation of calcitriol from the plasma extract was performed by washing the cartridge successively with 5 ml water, 5 ml methanol–water (70:30 v/v), 5 ml hexane–methylene chloride (90:10 v/v), 5 ml hexane–isopropanol (99:1 v/v), and finally elution of the calcitriol-containing fraction with 6 ml hexane–isopropanol (94:6 v/v). All solvent elutions of the cartridges were performed under reduced pressure using an Adsorbex sample preparation unit (Merck, Darmstadt, Germany). Analytical recovery of [³H]calcitriol through the entire extraction and purification procedure was 48.1 (SD 15.9)%, (*n* 32). The sensitivity of the nonequilibrium competitive receptor binding assay using the specific vitamin D receptor from calf thymus, defined as 2 SD from the mean for data from the zero sample, was 2 pg/tube. Bound tracer was 50% displaced at approximately 10 pg/tube. In the absence of calcitriol standard, approximately 25% of the total [³H]calcitriol was bound to the thymus receptor.

Other chemicals

The solvents were purchased from Merck (Darmstadt, Germany). The chemicals for preparing the buffer solutions, arsenate, ouabain, theophylline and L-alanine were obtained from Sigma Chemie GmbH (Deisenhofen, Germany). All chemicals were of analytical

grade. Unlabelled calcitriol was kindly provided by Hoffman-La Roche (Basel, Switzerland).

Statistics

All calculations were performed using the BMDP-87 software (Dixon, 1987). Univariate statistics such as arithmetic mean, standard deviation (SD) and standard error of the mean (SE) of the appropriate number of animals were computed for each variable by the BMDP1D program. In general, P values < 0.05 were taken to be statistically significant. According to the design of the experiments assessment of consecutive experimental periods in a given preparation (same animal, same segment) were based on paired statistics. In contrast, comparisons of preparations from different localizations were based on unpaired statistics because in most of the experiments the individual intestinal segments were derived from different animals. Paired and unpaired Student's t tests were used for statistical analysis (BMDP3D). One-way and two-way ANOVA were performed using the BMDP2V routine. For a significant result, individual mean values were checked in pairs for significant differences using the conventional Scheffé test (Sachs, 1992).

RESULTS

Experiment 1

Viability and electrical properties of intestinal epithelia from goats and sheep. In both species no significant changes of short-circuit currents (I_{sc}) and tissue conductances (G_T) occurred during the experimental period of at least 2 h, indicating constant conditions during the experiments. In addition, in a number of experiments tissues were either treated with theophylline (sheep duodenum) or glucose (duodenum of growing goats) or L-alanine (jejunum of growing goats) at the end of the flux measurements in order to check tissue viability (Fig. 1). Theophylline was used to increase second messenger concentrations, i.e. cyclic nucleotides, in the enterocytes. Glucose and L-alanine were both used in order to increase electrogenic Na^+ transport which resulted in increase of I_{sc} . In response to these compounds no significant differences in maximal increases of I_{sc} were detected in comparison with respective initial treatments after mounting the tissues (results not shown).

In the duodenum of adult goats I_{sc} values were positive and significantly higher ($P < 0.001$, paired t test) compared with those for jejunal preparations (Table 2). The negative jejunal I_{sc} may have resulted from either net absorption of anions or net secretion of cations. G_T values in goat duodenum were significantly lower than in the jejunum and in all intestinal segments of sheep. In sheep, G_T decreased from the duodenum to the proximal colon. In sheep all I_{sc} were positive and significantly higher in the ileum and proximal colon compared with the upper parts of the small intestines (Table 2).

Intestinal P_i flux rates in goats and sheep. In both species and in all segments of the intestinal tract J_{ms} exceeded J_{sm} under short-circuit-current conditions resulting in significant net flux rates from the mucosal to the serosal compartment (Table 3). J_{net} values in goat jejunum were significantly higher than in the duodenum. In sheep, J_{net} slightly increased within the distal parts of the small intestines.

Effects of arsenate, sodium substitution and ouabain on jejunal net P_i flux rates in sheep. When arsenate was added at concentrations of 2, 5 or 7 mM to the mucosal bathing solution, J_{net} of P_i decreased. Maximal inhibition of about 65% was achieved by 5 and 7 mM-arsenate (Fig. 2). A similar inhibition of J_{net} was observed either when mucosal Na was reduced from 148.2 mM to 1.8 mM (Fig. 3) or after serosal application of 0.1 mM-




Fig. 1. Maximal increases of short-circuit current (ΔI_{sc}) across small-intestinal epithelia from sheep and growing goats, in response to different substances. Theophylline (10 mM in serosal bathing solution), glucose (10 mM in mucosal fluid) and L-alanine (5 mM in mucosal fluid) were added at the end of unidirectional flux measurements. For osmotic reasons equimolar amounts of mannitol were given to the untreated compartment. Values are means with their standard errors represented by vertical bars. For details of procedures, see p. 636.

Table 2. *Electrical properties of stripped epithelia from goats and sheep*
(Mean values with their standard errors)

Species	Segment	n	I_{sc} ($\mu\text{eq}/\text{cm}^2$ per h)		G_T (mS/cm^2)	
			Mean	SE	Mean	SE
Goat	Duodenum	13	0.41	0.05	13.5	1.1
	Jejunum	13	-1.04	0.29	28.1	1.2
Sheep	Duodenum	9	0.45	0.07	30.3	3.4
	Jejunum	30	0.29	0.11	26.4	1.5
	Ileum	12	1.93	0.42	26.6	2.9
	Proximal colon	6	1.55	0.26	22.0	2.1

I_{sc} , short-circuit current; G_T , tissue conductance.

ouabain (Fig. 4). In each of these experiments remaining J_{net} values were still significantly different from zero.

Experiment 2

Variables of calcium and phosphorus homeostasis in goats. P depletion resulted in hypophosphataemia and hypercalcaemia and these changes were not related to the level of Ca intake (Table 4). In Ca-depleted goats no significant changes in plasma Ca concentrations could be detected. These animals, however, became hyperphosphataemic when they were kept on an adequate P supply. In both Ca-depleted groups calcitriol concentrations significantly increased in comparison with control animals. Plasma PTH levels were only higher in Ca depletion without P depletion as indicated by a significant interaction between dietary Ca and P conditions (Table 4).

P_i flux rates and electrical properties of duodenal and jejunal tissues as affected by calcium and/or phosphorus depletion. In both duodenal and jejunal tissues J_{ms} increased significantly with low P intake irrespective of the level of Ca intake. Since J_{sm} remained unaffected this

Table 3. Unidirectional and net flux rate of inorganic phosphate in intestinal segments from goats and sheep*

(Mean values with their standard errors)

Species	Segment	n	J_{ms} (nmol/cm ² per h)		J_{sm} (nmol/cm ² per h)		J_{net} (nmol/cm ² per h)		P
			Mean	SE	Mean	SE	Mean	SE	
Goat	Duodenum	13	33.9	4.2	12.4	2.3	21.5	2.8	< 0.001
	Jejunum	13	114.8	14.7	40.7	11.3	74.1	15.0	< 0.001
Sheep	Duodenum	9	69.1	12.0	43.2	6.7	25.9	6.8	< 0.01
	Jejunum	30	87.7	7.5	34.0	3.3	53.8	7.2	< 0.001
	Ileum	12	131.1	17.1	42.8	7.2	88.2	19.2	< 0.001
	Proximal colon	6	52.6	4.5	29.3	3.8	23.3	6.8	< 0.05

J_{ms} , mucosal to serosal flux rate; J_{sm} , serosal to mucosal flux rate; J_{net} , net flux rate ($J_{net} = J_{ms} - J_{sm}$); P, statistical significance of difference from zero of mean J_{net} .

* For details of procedures, see p. 636.




Fig. 2. Inhibition of net flux rates (J_{net}) of inorganic phosphate across jejunal epithelia from sheep in the presence of mucosal arsenate. Values are means with their standard errors represented by vertical bars; (▨), basal; (□), arsenate-treated. Values for inhibition as a percentage of basal J_{net} were: 2 mM -27.0, 5 mM -66.8, 7 mM -61.2. * Mean values were significantly different from control, $P < 0.05$. For details of procedures, see p. 636.

change resulted in significant increases of J_{net} in both intestinal segments (Tables 5 and 6). Electrical properties of duodenal tissues were not affected by dietary treatment whereas in jejunal tissues P depletion induced increases in negative I_{sc} compared with both adequately P-fed groups (Tables 5 and 6). In all these experiments no statistically significant interactions could be observed.

DISCUSSION

Viability of the tissues during experiments

In vitro transepithelial flux measurements across intact intestinal epithelia with the Ussing-chamber system require sufficient integrity and viability of the tissues during the experimental period. Functional viability of the epithelia over at least 2 h has been




Fig. 3. Change in the net flux rate (J_{net}) of inorganic phosphate across jejunal epithelia from sheep in response to reducing the sodium concentration in the mucosal bathing solution from 144 mM (high [Na⁺]) to 1.8 mM (low [Na⁺]). Values are means with their standard errors represented by vertical bars for four animals for four experiments. Inhibition as a percentage of basal J_{net} was -61.9. * Mean value was significantly different from high [Na⁺], $P < 0.05$. For details of procedures, see p. 636.




Fig. 4. Inhibition of net flux rates (J_{net}) of inorganic phosphate across jejunal epithelia from sheep in the presence of serosal ouabain (0.1 mM). Values are means with their standard errors represented by vertical bars, for six experiments for six animals; (▨), basal; (□), ouabain-treated. Inhibition as a percentage of basal J_{net} was -63.3. ** Mean value was significantly different from basal, $P < 0.01$. For details of procedures, see p. 636.

demonstrated by measuring short-circuit currents (electrogenic net ion transport) and tissue conductances (representing trans- and paracellular pathways). Neither electrical variable changed significantly during the experiments. Furthermore, addition of organic substrates such as glucose and L-alanine to the tissues or treatment with the secretagogue theophylline provoked similar increases of I_{sc} at the start and the end of the experiments. Since the experiments were performed in the absence of any organic substrates in the luminal compartment, electrogenic Na⁺/substrate-cotransport was kept at a low level

Table 4. Concentrations of calcium, inorganic phosphate, parathyroid hormone, and calcitriol in plasma from growing goats as affected by long-term dietary calcium and/or phosphorus depletion*

(Mean values with their standard errors for eight animals per dietary group)

Grouping		Ca (mmol/l)		P _i (mmol/l)		PTH (pmol/l)		Calcitriol (pmol/l)	
Ca	P	Mean	SE	Mean	SE	Mean	SE	Mean	SE
+	+	2.73	0.20	2.22	0.40	95	8	107	17
+	-	3.02	0.13	0.61	0.17	85	9	102	19
-	+	2.70	0.19	2.78	0.20	148	18	233	26
-	-	2.95	0.18	0.87	0.50	99	4	224	25
Source of variation		<i>P</i> values of two-way ANOVA							
Ca		NS		< 0.01		< 0.01		< 0.001	
P		< 0.001		< 0.001		< 0.01		NS	
Ca × P interaction		NS		NS		< 0.05		NS	

P_i, inorganic phosphate; PTH, parathyroid hormone.

* For details of treatments, see Table 1.

Table 5. Unidirectional and net flux rates of inorganic phosphate, and electrical properties of duodenal preparations of growing goats as affected by calcium and/or phosphorus depletion*

(Mean values with their standard errors)

Grouping		<i>n</i>	J _{ms} (nmol/ cm ² per h)		J _{sm} (nmol/ cm ² per h)		J _{net} (nmol/ cm ² per h)		I _{sc} (μeq/ cm ² per h)		G _T (mS/cm ²)	
Ca	P		Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
+	+	8	44.2	12.8	8.7	2.3	35.5	11.9	0.51	0.07	12.4	1.2
+	-	7	95.4	36.2	10.1	2.8	85.4	33.4	0.52	0.08	14.3	1.8
-	+	8	34.7	7.9	7.7	1.5	27.0	7.1	0.50	0.05	13.2	1.3
-	-	7	86.5	18.6	11.3	1.7	75.3	18.2	0.73	0.03	14.8	1.2
Source of variation		<i>P</i> values of two-way ANOVA										
Ca		NS		NS		NS		NS		NS		
P		< 0.05		NS		< 0.05		NS		NS		
Ca × P interaction		NS		NS		NS		NS		NS		

J_{ms}, mucosal to serosal flux rate; J_{sm}, serosal to mucosal flux rate; J_{net}, net flux rate (J_{net} = J_{ms} - J_{sm}); I_{sc}, short-circuit current; G_T, tissue conductance.

* For details of treatments, see Table 1.

during the experiments. This procedure gave the opportunity to test Na⁺/substrate-cotransport capacity of the epithelia at the end of the flux measurements as indicated by increases of the I_{sc} after addition of respective substrates (Ferrante *et al.* 1988). In order to prove the tissue's capability to stimulate electrolyte transport by second messengers, theophylline was added to the serosal buffer solution. This secretagogue is known to increase cytosolic cAMP and cGMP concentrations and intracellular free Ca levels (Schröder *et al.* 1991). Tissue viability and integrity could be confirmed by these functional

Table 6. Unidirectional and net flux rates of inorganic phosphate, and electrical properties of jejunal preparations of growing goats as affected by calcium and/or phosphorus depletion*

(Mean values with their standard errors)

Grouping		J_{ms} (nmol/ cm ² per h)	J_{sm} (nmol/ cm ² per h)	J_{net} (nmol/ cm ² per h)	I_{sc} (μ eq/ cm ² per h)	G_T (mS/cm ²)						
Ca	P	n	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
+	+	8	86.1	12.2	34.5	3.2	51.6	11.3	-0.81	0.23	27.8	2.0
+	-	7	171.6	38.1	39.0	6.6	132.6	36.2	-1.35	0.24	28.9	1.4
-	+	8	122.2	26.5	30.8	5.8	91.5	24.7	-0.31	0.25	26.1	2.8
-	-	7	156.2	31.9	32.1	5.9	124.1	32.6	-1.01	0.44	29.5	1.7
Source of variation						P values of two-way ANOVA						
Ca			NS		NS		NS		NS		NS	
P			< 0.05		NS		< 0.05		< 0.05		NS	
Ca \times P interaction			NS		NS		NS		NS		NS	

J_{ms} , mucosal to serosal flux rate; J_{sm} , serosal to mucosal flux rate; J_{net} , net flux rate ($J_{net} = J_{ms} - J_{sm}$); I_{sc} , short-circuit current; G_T , tissue conductance.

* For details of treatments, see Table 1.

tests since the epithelial response was identical at both start and end of the flux measurements. It could further be confirmed by the constancy of the successive P_i flux rates in individual Ussing chambers.

Active P_i absorption in the small intestine and the proximal colon

In both goats and sheep, J_{ms} exceeded J_{sm} in all intestinal segments tested in the present study, thus resulting in significant net P_i absorption (Table 3). These J_{net} clearly indicate the presence of active mechanisms for P_i transport in the small intestine and the proximal colon since they were measured in the absence of any transepithelial electrochemical gradients. These findings are the first experimental proof for actively driven P_i transport obtained from intact intestinal epithelia of ruminants, thus confirming the assumptions made by Care *et al.* (1980), Scott *et al.* (1984) and Scharrer (1985) who have postulated the involvement of carrier-mediated mechanisms in the intestinal transport of P_i in sheep. In the small intestine, J_{net} of P_i increased from oral to aboral. The reasons for this are not yet fully understood. Physiological as well as experimental conditions may be involved. Different P_i transport capacities along the intestinal tract, i.e. different V_{max} values of the brush border Na^+/P_i cotransport system, can cause different flux rates in respective intestinal segments as has been described for pigs and rabbits (Brandis *et al.* 1987; Danisi & Murer, 1991).

In the present study the physiological pH value of the extracellular fluid (7.4) was adjusted in both mucosal and serosal solutions. Thus, it has to be taken into account that a mucosal pH of 7.4 was only equivalent to *in vivo* conditions using tissues from the mid jejunum. From experiments with sheep a pH range of 3.5 to 5.2 has been reported for the luminal fluid in the upper small intestine (1-3 m from the pylorus) and 7.8 to 8.2 in the terminal ileum (Ben-Ghedalia *et al.* 1975). However, the main principle to detect active transport mechanisms with the Ussing-chamber technique implies the absence of any electrochemical gradient which also excludes the application of transepithelial pH gradients. Therefore, the

present study was performed under iso-pH conditions. Nevertheless, it still has to be proved for ruminants how luminal pH reflects cell surface pH. It has been shown for rats and pigs that the mucosal surface is isolated from changes in the pH of luminal contents by a zone of moderate pH ('acid microclimate') (Daniel *et al.* 1989; McEwan *et al.* 1990). In conclusion, it cannot be excluded for duodenal tissues that factors such as the ratio between mono- and divalent P_i and proton gradients present under *in vivo* conditions may have been underestimated.

Sodium dependency of P_i transport

Since it is well established that in ruminants, as in single-stomached species, the small intestine is the major site for net P_i absorption, and with respect to physiological pH conditions, tissues from the mid jejunum were used for further characterizing the properties of active P_i transport in sheep. It could be demonstrated that almost 0.66 of active P_i transport depended on the presence of Na since the reduction of mucosal Na concentration from 148.2 mM to 1.8 mM reduced J_{net} by about 60% (Fig. 3). Similar findings were obtained when ouabain was added to the serosal compartment at a concentration of 0.1 mM (Fig. 4). This concentration had been found to be sufficient for complete inhibition of basolateral Na^+K^+ -ATPase (EC 3.6.1.3) activity (Blaustein, 1993). In further studies arsenate was added to the mucosal side of the intact tissues resulting in a significant decrease of J_{net} of about 65% in comparison with controls. Maximal inhibition was found at a concentration of 5 mM. Adjusting arsenate to 7 mM did not further increase the degree of inhibition (Fig. 2). Arsenate, an anion analogous in structure to the phosphate anion, has been used as a competitive inhibitor of the renal and intestinal Na^+ -coupled P_i cotransport system in *in vitro* P_i uptake measurements with isolated brush border membrane vesicles (BBMV) from single-stomached species (Berner *et al.* 1976; Hoffmann *et al.* 1976; Danisi *et al.* 1984; Quamme, 1985; Shirazi-Beechey *et al.* 1988). This inhibitory effect has been confirmed by recent studies with intestinal BBMV from goats where Na^+ -dependent P_i uptake could be inhibited completely in the presence of arsenate (Schröder *et al.* 1993b). However, the fact that the level of inhibition by arsenate was similar to the effects in response to the reduction of Na in the mucosal buffer solution, or in response to addition of ouabain in the serosal compartment, may be a coincidence, since arsenate may also have a variety of other effects on living cells, i.e. intermediary metabolism, which can limit the suitability of this compound for studies of active P_i transport system (Szczepanska-Konkel *et al.* 1986).

Although from the inhibition and substitution experiments the Na-dependency of active intestinal P_i transport in sheep has been shown, it may not be concluded that this mechanism is identical with the secondary active Na^+/P_i cotransport which has been shown in different single-stomached species (for review see Breves & Schröder, 1991) and sheep ileum (Schneider *et al.* 1985a, b). The observed effect of Na^+ could also be due to the functioning of a Na^+/H^+ antiporter which may be present in the luminal membrane. Na^+ -driven H^+ extrusion could provide protons which could be used to be coupled to the movement of P_i into the cell via a H^+/P_i cotransporter as has been shown for sheep duodenum (Shirazi-Beechey *et al.* 1989, 1991). A similar mechanism has also been demonstrated for the internal mitochondrial membrane of eukaryotic cells (Wehrle & Pedersen, 1989). Hence the exclusion of Na^+ from the luminal compartment would lead to the reduction of activity of the Na^+/H^+ antiporter and this could consequently reduce the rates of P_i transport via the H^+/P_i cotransporter. The addition of ouabain to the serosal compartment inhibits the activity of basolateral Na^+K^+ -ATPase, thus leading to increases of the cytosolic Na^+ concentration and reduction of the Na^+ gradient across the luminal membrane, which again could reduce the activity of the Na^+/H^+ antiporter.

Independent of to what extent and how Na^+ may be involved in active P_i -transport, it has to be stated that in the present study a proportion (about 30%) of total J_{net} of P_i remained unaffected by any mucosal or serosal manipulations of Na^+ -dependent P_i transport. From this the existence of further Na^+ -independent mechanisms for luminal P_i -uptake may be surmised.

Active P_i absorption as affected by dietary calcium and/or phosphorus depletion

Studies in single-stomached species have shown that active P_i absorption can be regulated in response to changes of dietary P intake, and that this effect is mediated by calcitriol (Fox & Care, 1978; Peterlik & Wasserman, 1980; Jungbluth & Binswanger, 1989). Therefore, in the second experiment tissues were taken from goats which had been kept on Ca- and/or P-deficient diets. Significant changes in different variables of Ca and P homeostasis were observed (Table 4). Independent of level of Ca intake P depletion resulted in hypophosphataemia and hypercalcaemia. Increases in plasma Ca levels may have resulted from decreased bone formation and/or increased bone resorption or from respective changes of intestinal Ca absorption and renal Ca secretion (Baylink *et al.* 1971; Breves & Schröder, 1991). In both Ca-depleted groups increased plasma calcitriol levels occurred which could not be detected when the animals were only P-depleted. The lack of calcitriol stimulation by P-depletion is in contrast to single-stomached species (Baxter & DeLuca, 1976; Fox & Care, 1976; Gray *et al.* 1983; Breves *et al.* 1985; Maunder *et al.* 1986; Shirazi-Beechey *et al.* 1991). Possibly in P-depleted ruminants the vitamin D-hormone system is regulated at the cytosolic calcitriol receptor level of enterocytes since studies in P-depleted lactating goats have shown increased binding affinities of the receptor (Schröder *et al.* 1990). This, however, still needs further investigation since in growing goats diet-induced changes of calcitriol receptor properties were not identical to those in lactating goats (unpublished results). Increased plasma PTH concentrations could only be detected in Ca-depleted goats with an adequate P supply, which is similar to single-stomached animals (Chu *et al.* 1973; Fox, 1992). Possibly the lack of increased PTH levels in the Ca- and P-depleted group was due to hypercalcaemia in those animals (Table 4).

Dietary P depletion resulted in an increased duodenal and jejunal J_{ms} with no significant effect of level of Ca intake (Tables 5 and 6). Since J_{sm} values were not affected, enhanced J_{ms} resulted in increased J_{net} , which clearly indicated the stimulation of active P_i transport in response to dietary P depletion. Similar adaptational processes have been described for the duodenum of chicks and rabbits (Quamme, 1985; Danisi *et al.* 1990) and the jejunum of rats (Caverzasio *et al.* 1987). In these studies it was demonstrated that increases in P_i uptake were due to increases in V_{max} of the Na^+/P_i -cotransport system. Similar findings were obtained by Shirazi-Beechey *et al.* (1991) in sheep concerning the proposed H^+/P_i -cotransport mechanism. In single-stomached species these adaptational processes are controlled by plasma calcitriol (Fuchs & Peterlik, 1980; Matsumoto *et al.* 1980; Hildmann *et al.* 1982; Brandis *et al.* 1987). From the present data it cannot be decided whether the increases in active P_i absorption were due to stimulations of Na^+ -dependent and/or Na^+ -independent P_i transport. For both processes it has to be concluded that in goats stimulation of intestinal P_i absorption was not mediated by increased plasma calcitriol or PTH levels since they were not different from control animals. Evidence for calcitriol-independent stimulation of active P_i -transport has also been found by Brommage *et al.* (1990) in rats. Whether other hormones such as insulin (Peterlik *et al.* 1981), triiodothyronine and thyroxine (Cross *et al.* 1986), glucocorticoids (Cross *et al.* 1990) or somatomedins (Gray, 1987) are involved is not clear.

From the present study it may be concluded that in ruminants, as in single-stomached species, P_i can be absorbed actively from the small intestine and that in the mid-jejunum

of sheep a substantial portion of the active transport consists of a Na^+ -dependent P_i -transport mechanism. Further details on potential differences between intestinal segments and ruminant species have not yet been clarified. At present it is not clear by which mechanism the remaining 30% of J_{net} of P_i was mediated, and it still has to be ascertained whether the increased transport rates in response to dietary P depletion were due to changes of the Na^+ -dependent and/or Na^+ -independent mechanisms and how these changes were induced at the cellular level.

Financial support of the Deutsche Forschungsgemeinschaft and H. Wilhelm Schaumann Stiftung is gratefully acknowledged.

REFERENCES

- Baxter, L. A. & DeLuca, H. F. (1976). Stimulation of 25-hydroxyvitamin D₃-1 α -hydroxylase by phosphate depletion. *Journal of Biological Chemistry* **251**, 3158–3161.
- Baylink, D., Wergedal, J. & Stauffer, M. (1971). Formation, mineralization and resorption of bone in hypophosphatemic rats. *Journal of Clinical Investigation* **50**, 2519–2530.
- Ben-Ghedalia, D., Tagari, H., Zamwel, S. & Bondi, A. (1975). Solubility and net exchange of calcium, magnesium and phosphorus in digesta flowing along the gut of the sheep. *British Journal of Nutrition* **33**, 87–94.
- Berner, W., Kinne, R. & Murer, H. (1976). Phosphate transport into brush-border membrane vesicles isolated from rat small intestine. *Biochemical Journal* **160**, 467–474.
- Blaustein, M. P. (1993). Physiological effects of endogenous ouabain: control of intracellular Ca^{2+} stores and cell responsiveness. *American Journal of Physiology* **264**, C1367–C1387.
- Brandis, M., Harmeyer, J., Kaune, R., Mohrmann, M., Murer, H. & Zimolo, Z. (1987). Phosphate transport in brush-border membranes from control and rachitic pig kidney and small intestine. *Journal of Physiology* **384**, 479–490.
- Breves, G., Ross, R. & Höller, H. (1985). Dietary phosphorus depletion in sheep: effects on plasma inorganic phosphorus, calcium, 1,25(OH)₂-Vit-D₃ and alkaline phosphatase and on gastrointestinal P and Ca balances. *Journal of Agricultural Science, Cambridge* **105**, 623–629.
- Breves, G. & Schröder, B. (1991). Comparative aspects of gastrointestinal phosphorus metabolism. *Nutrition Research Reviews* **4**, 125–140.
- Brommage, R., Baxter, D. C. & Gierke, L. W. (1990). Vitamin D-independent intestinal calcium and phosphorus absorption during reproduction. *American Journal of Physiology* **259**, G631–G638.
- Bruce, J., Goodall, E., Kay, R., Phillipson, A. & Vowles, L. (1966/67). The flow of organic and inorganic materials through the alimentary tract of the sheep. *Proceedings of the Royal Society* **166B**, 46–62.
- Care, A., Barlet, J. & Abdel-Hafeez, H. (1980). Calcium and phosphate homeostasis in ruminants and its relationship to the aetiology and prevention of parturient paresis. In *Digestive Physiology and Metabolism in Ruminants*, pp. 429–446 [Y. Ruckebusch and P. Thivend, editors]. Lancaster: MTP Press Ltd.
- Caverzasio, J., Danisi, G., Murer, H. & Bonjour, J. (1987). Adaptation of phosphate transport to low phosphate diet in renal and intestinal brush border membrane vesicles: influence of sodium and pH. *European Journal of Physiology – Pflügers Archiv* **409**, 333–336.
- Chu, L. L. H., McGregor, R. R., Anast, C. S., Hamilton, J. W. & Cohn, D. V. (1973). Studies on the biosynthesis of rat parathyroid hormone and parathyroid hormone: adaptation of the parathyroid gland to dietary restriction of calcium. *Endocrinology* **93**, 915–924.
- Cross, H., Debiec, H. & Peterlik, M. (1990). Mechanism and regulation of intestinal phosphate absorption. *Mineral and Electrolyte Metabolism* **16**, 115–124.
- Cross, H. S., Pölzleitner, D. & Peterlik, M. (1986). Intestinal phosphate and calcium absorption: joint regulation by thyroid hormones and 1,25-dihydroxyvitamin D₃. *Acta Endocrinologica* **113**, 96–103.
- Daniel, H., Fett, C., Kratz, A. & Rehner, G. (1989). Demonstration and modification of intervillous pH profiles in the rat jejunum *in vitro*. *American Journal of Physiology* **257**, G489–G495.
- Danisi, G., Caverzasio, J., Trechsel, U., Bonjour, J. & Straub, R. (1990). Phosphate transport adaptation in rat jejunum and plasma level of 1,25-dihydroxyvitamin-D₃. *Scandinavian Journal of Gastroenterology* **25**, 210–215.
- Danisi, G., Caverzasio, J., Trechsel, U., Straub, R. & Bonjour, J. (1988). Phosphate transport adaptation in intestinal brush border membrane vesicles (BBMV) and plasma levels of 1,25-dihydroxycholecalciferol. In *Cellular Calcium and Phosphate Transport in Health and Disease*, pp. 65–66 [F. Bronner and M. Peterlik, editors]. New York: Alan R. Liss, Inc.
- Danisi, G. & Murer, H. (1991). Inorganic phosphate absorption in small intestine. In *Handbook of Physiology. The Gastrointestinal System IV*, pp. 323–336 [S. G. Schultz, editor]. New York: Oxford University Press.
- Danisi, G., van Os, C. & Straub, R. (1984). Phosphate transport across brush border and basolateral membrane vesicles of small intestine. In *Epithelial Calcium and Phosphate Transport: Molecular and Cellular Aspects*, pp. 229–234 [F. Bronner and M. Peterlik, editors]. New York: Alan R. Liss, Inc.

- Dixon, W. J. (1987). *BMDP Statistical Software Manual*. Los Angeles: University of California Press.
- Ferrante, P. L., Freeman, D. E., Whitlock, R. H. & Kronfeld, D. S. (1988). Effect of D-glucose on in vitro short-circuit current in neonatal calf jejunum and rabbit ileum. *American Journal of Veterinary Research* **49**, 715–719.
- Fox, J. (1992). Hypocalcemia, but not PTH or hypophosphatemia, induces a rapid increase in 1,25(OH)₂D₃ levels in rats. *American Journal of Physiology* **262**, E211–E215.
- Fox, J. & Care, A. D. (1976). The effects of hydroxylated derivatives of vitamin D₃ and of extracts of *Solanum malacoxylon* on the absorption of calcium, phosphate and water from the jejunum of pigs. *Calcified Tissue Research* **21**, Suppl., 147–152.
- Fox, J. & Care, A. D. (1978). Effect of low calcium and low phosphorus diets on the intestinal absorption of phosphate in intact and parathyroidectomized pigs. *Journal of Endocrinology* **77**, 225–231.
- Fuchs, R. & Peterlik, M. (1980). Vitamin D-induced phosphate transport in intestinal brush border membrane vesicles. *Biochemical and Biophysical Research Communications* **93**, 87–92.
- Grace, N. D., Ulyatt, M. J. & Macrae, J. C. (1974). Quantitative digestion of fresh herbage by sheep. III. The movement of Mg, Ca, P, K and Na in the digestive tract. *Journal of Agricultural Science, Cambridge* **82**, 321–330.
- Gray, R. (1987). Evidence that somatomedins mediate the effect of hypophosphatemia to increase serum 1,25-dihydroxyvitamin D₃ levels in rats. *Endocrinology* **121**, 504–512.
- Gray, R. W., Garthwaite, T. L. & Phillips, L. S. (1983). Growth hormone and triiodothyronine permit an increase in plasma 1,25(OH)₂D concentrations in response to dietary phosphate deprivation in hypophysectomized rats. *Calcified Tissue International* **35**, 100–106.
- Hildmann, B., Storelli, C., Danisi, G. & Murer, H. (1982). Regulation of Na⁺-P_i cotransport by 1,25-dihydroxyvitamin D₃ in rabbit duodenal brush-border membrane. *American Journal of Physiology* **242**, G533–G539.
- Hoffmann, N., Thees, M. & Kinne, R. (1976). Phosphate transport by isolated renal brush border vesicles. *European Journal of Physiology – Pflügers Archiv* **362**, 147–156.
- Hollis, B. (1986). Assay of circulating 1,25-dihydroxyvitamin D involving a novel single-cartridge extraction and purification procedure. *Clinical Chemistry* **32**, 2060–2062.
- Jungbluth, H. & Binswanger, U. (1989). Unidirectional duodenal and jejunal calcium and phosphorus transport in the rat: effects of dietary phosphorus depletion, ethane-1-hydroxy-1,1-diphosphonate and 1,25-dihydroxycholecalciferol. *Research in Experimental Medicine* **189**, 439–449.
- McEwan, G. T. A., Schousboe, B. & Skadhauge, E. (1990). Direct measurement of surface pH of pig jejunum *in vivo*. *Journal of Veterinary Medicine A* **37**, 439–444.
- Matsumoto, T., Fontaine, O. & Rasmussen, H. (1980). Effect of 1,25-dihydroxyvitamin D₃ on phosphate uptake into chick intestinal brush border membrane vesicles. *Biochimica et Biophysica Acta* **599**, 13–23.
- Maunder, E., Pillay, A. & Care, A. (1986). Hypophosphatemia and vitamin D metabolism in sheep. *Quarterly Journal of Experimental Physiology* **71**, 391–399.
- Murer, H. & Hildmann, B. (1981). Transcellular transport of calcium and inorganic phosphate in the small intestinal epithelium. *American Journal of Physiology* **240**, G409–G416.
- Peterlik, M. & Wasserman, R. H. (1980). Regulation by vitamin D of intestinal phosphate absorption. *Hormone and Metabolic Research* **12**, 216–219.
- Peterlik, M., Fuchs, R. & Cross, H. S. (1981). Phosphate transport in the intestine: cellular pathways and hormonal regulation. In *Calcium and Phosphate Transport Across Biomembranes*, pp. 173–179 [F. Bronner and M. Peterlik, editors]. New York: Academic Press, Inc.
- Pfeffer, E., Thompson, A. & Armstrong, D. G. (1970). Studies on intestinal digestion in the sheep. 3. Net movement of certain inorganic elements in the digestive tract on rations containing different proportions of hay and rolled barley. *British Journal of Nutrition* **24**, 197–204.
- Quamme, G. A. (1985). Phosphate transport in intestinal brush-border membrane vesicles: effect of pH and dietary phosphate. *American Journal of Physiology* **249**, G168–G176.
- Reinhardt, T. A. & Horst, R. L. (1990). Competitive protein-binding assay of calcitriol with an advanced preparation of bovine calf thymus cytosol. In *Calcium Regulating Hormones, Vitamin D Metabolites, and Cyclic AMP. Assays and their Clinical Application*, pp. 334–345 [H. Schmidt-Gayk, F. P. Armbruster and R. Bouillon, editors]. Berlin, Heidelberg: Springer Verlag.
- Sachs, L. (1992). *Angewandte Statistik*. Berlin, Heidelberg, New York: Springer Verlag.
- Scharrer, E. (1985). Phosphate absorption at different sites in the developing lamb. *Quarterly Journal of Experimental Physiology* **70**, 615–621.
- Schneider, K. M., Straub, R. W. & Danisi, G. (1985a). Phosphate transport in the gut of the ruminant. *Experientia* **41**, 834–835.
- Schneider, K. M., Straub, R. W. & Danisi, G. (1985b). Sodium-phosphate cotransport in ruminant intestine. *Mineral and Electrolyte Metabolism* **11**, 339.
- Schröder, B., Breves, G. & Pfeffer, E. (1990). Binding properties of duodenal 1,25-dihydroxyvitamin D₃ receptors as affected by phosphorus depletion in lactating goats. *Comparative Biochemistry and Physiology* **96A**, 495–498.
- Schröder, B., Kaune, R. & Harmeyer, J. (1991). Effects of calcitriol on stimulation of ion transport in pig jejunal mucosa. *Journal of Physiology* **433**, 451–465.

- Schröder, B., Kaune, R., Schlumbohm, C., Breves, G. & Harmeyer, J. (1993a). Evidence for vitamin-D-independent active calcium absorption in newborn piglets. *Calcified Tissue International* **52**, 305–309.
- Schröder, B., Rübelt, A. & Breves, G. (1993b). Sodium coupled phosphate transport across jejunal brush border membrane from small ruminants. *Biological Chemistry Hoppe-Seyler* **374**, 157.
- Schultz, S. & Zalusky, R. (1964). Ion transport in isolated rabbit ileum. I. Short-circuit current and Na fluxes. *Journal of General Physiology* **47**, 567–584.
- Scott, D., McLean, A. F. & Buchan, W. (1984). The effects of intravenous phosphate loading on salivary phosphorus secretion, net intestinal phosphorus absorption and fecal endogenous phosphorus excretion in sheep. *Quarterly Journal of Experimental Physiology* **69**, 453–461.
- Shirazi-Beechey, S., Gorvel, J. & Beechey, R. (1988). Phosphate transport in intestinal brush-border membrane. *Journal of Bioenergetics and Biomembranes* **20**, 273–288.
- Shirazi-Beechey, S. P., Beechey, R. B., Penny, J., Vayro, S., Buchan, W. & Scott, D. (1991). Mechanisms of phosphate transport in sheep intestine and parotid gland – response to variation in dietary phosphate supply. *Experimental Physiology* **76**, 231–241.
- Shirazi-Beechey, S. P., Kemp, R. B., Dyer, J. & Beechey, R. B. (1989). Changes in the function of the intestinal brush border membrane during the development of the ruminant habit in lambs. *Comparative Biochemistry and Physiology B* **94**, 801–806.
- Szczepanska-Konkel, M., Yusufi, A. N. K., VanScoy, M., Webster, S. K. & Dousa, T. P. (1986). Phosphono-carboxylic acids as specific inhibitors of Na⁺-dependent transport of phosphate across renal brush border membrane. *Journal of Biological Chemistry* **261**, 6375–6383.
- Wehrle, J. P. & Pedersen, P. L. (1989). Phosphate transport processes in eukaryotic cells. *Journal of Membrane Biology* **111**, 199–213.