


Radial solutions of initial boundary value problems of nonlinear Schrödinger equations in \mathbb{R}^n

Shu-Ming Sun

Department of Mathematics, Virginia Polytechnic Institute and State University, Blacksburg, 24061 VA, USA (sun@math.vt.edu)

Fangchi Yan 

Department of Mathematics, Virginia Polytechnic Institute and State University, Blacksburg, 24061 VA, USA (fyan1@alumni.nd.edu)
(corresponding author)

(Received 19 December 2023; revised 7 September 2024; accepted 1 October 2024)

The article studies an initial boundary value problem (ibvp) for the radial solutions of the nonlinear Schrödinger (NLS) equation in a radially symmetric region $\Omega \in \mathbb{R}^n$ with boundaries. All such regions can be classified into three types: a ball Ω_0 centred at origin, a region Ω_1 outside a ball, and an n -dimensional annulus Ω_2 . To study the well-posedness of those ibvps, the function spaces for the boundary data must be specified in terms of the solutions in appropriate Sobolev spaces. It is shown that when $\Omega = \Omega_1$, the ibvp for the NLS equation is locally well-posed in $C([0, T^*]; H^s(\Omega_1))$ if the initial data is in $H^s(\Omega_1)$ and boundary data is in $H^{\frac{2s+1}{4}}(0, T)$ with $s \geq 0$. This is the optimal regularity for the boundary data and cannot be improved. When $\Omega = \Omega_2$, the ibvp is locally well-posed in $C([0, T^*]; H^s(\Omega_2))$ if the initial data is in $H^s(\Omega_2)$ and boundary data is in $H^{\frac{s+1}{2}}(0, T)$ with $s \geq 0$. In this case, the boundary data requires $1/4$ more derivative compared to the case when $\Omega = \Omega_1$. When $\Omega = \Omega_0$ with $n = 2$ (the case with $n > 2$ can be discussed similarly), the ibvp is locally well-posed in $C([0, T^*]; H^s(\Omega_0))$ if the initial data is in $H^s(\Omega_0)$ and boundary data is in $H^{\frac{s+1}{2}}(0, T)$ with $s > 1$ (or $s > n/2$). Due to the lack of Strichartz estimates for the corresponding boundary integral operator with $0 \leq s \leq 1$, the local well-posedness can only be achieved for $s > 1$. It is noted that the well-posedness results on Ω_0 and Ω_2 are the first ones for the ibvp of NLS equations in bounded regions of higher dimension.

Keywords: nonlinear Schrödinger equations; initial boundary value problems; well-posedness; radial solutions; radially symmetric regions

2020 Mathematics Subject Classification: Primary: 35Q55; 35G16; 35G31

1. Introduction

The article studies the well-posedness of the initial boundary value problem (ibvp) for the general nonlinear Schrödinger (NLS) equation in \mathbb{R}^n , which is given by:

$$iu_t + \Delta u + \lambda|u|^{p-2}u = 0, \quad (x_1, \dots, x_n) \in \Omega, \quad t \in (0, T), \quad T < \frac{1}{2}, \quad (1.1a)$$

$$u(x_1, \dots, x_n, 0) = u_0(x_1, \dots, x_n), \quad (x_1, \dots, x_n) \in \Omega, \quad (1.1b)$$

$$u(x_1, \dots, x_n, t) = g(t), \quad (x_1, \dots, x_n) \in \partial\Omega, \quad t \in (0, T), \quad (1.1c)$$

where $p \geq 3$, $\Delta u = \partial_{x_1}^2 u + \dots + \partial_{x_n}^2 u$ and Ω is a radially symmetric region in \mathbb{R}^n . The radially symmetric regions can be a ball centred at the origin, the outside of this ball, and an annulus between two spheres. In dimension 2, we have sketched the graph of these three regions in figures 1–3. Of course, the most general radially symmetric regions can be a combination of the regions mentioned above. Since we can decompose the general radially symmetric regions into the above three regions and this decomposition allows us to analyse each region independently, in this work, we consider three regions:

$$\begin{aligned} \Omega_0 &\doteq \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1^2 + \dots + x_n^2 < 1\}, \\ \Omega_1 &\doteq \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1^2 + \dots + x_n^2 > 1\}, \\ \Omega_2 &\doteq \{(x_1, \dots, x_n) \in \mathbb{R}^n : \pi^2 < x_1^2 + \dots + x_n^2 < (2\pi)^2\}. \end{aligned}$$

Figures 1–3 depict the regions Ω_0 – Ω_2 , respectively, for the case of $n = 2$.

Equation (1.1a) can be classified as either focusing (indicated by the ‘ $\lambda > 0$ ’) or defocusing (indicated by the ‘ $\lambda < 0$ ’). When $p = 4$, it becomes the well-known cubic NLS equation ($\lambda = \pm 1$)

$$iu_t + u_{xx} \pm |u|^2u = 0, \quad (1.2)$$

which is a ubiquitous model in various areas of mathematical physics, including water waves, plasmas, optics, and Bose–Einstein condensates. The cubic NLS equation has been rigorously derived for water waves of small amplitude over infinite or finite depth and in the context of nonlinear optics [2, 15, 33]. It has also been proposed as a model for rogue waves [14, 31]. In addition to its physical significance, the cubic NLS equation exhibits a complex mathematical framework as the archetypal illustration of a fully integrable system in one dimension. It features an unbounded hierarchy of symmetries and laws of conservation. In the one-dimensional case, the equation’s integrability is particularly noteworthy, as characterized by the existence of a Lax pair of the form:

$$\Psi_x = \begin{pmatrix} -ik & u \\ \mp \frac{1}{2}\bar{u} & ik \end{pmatrix} \Psi, \quad \Psi_t = \begin{pmatrix} -2ik^2 \pm \frac{i}{2}|u|^2 & 2ku + iu_x \\ \mp k\bar{u} \pm \frac{i}{2}\bar{u}_x & 2ik^2 \mp \frac{i}{2}|u|^2 \end{pmatrix} \Psi, \quad k \in \mathbb{C}. \quad (1.3)$$




Figure 1. Region Ω_0 .




Figure 2. Region Ω_1 .




Figure 3. Region Ω_2 .

The Lax pair formulation (1.3) allows the study of the initial value problem (ivp) for the cubic NLS equation under the assumption of initial data with sufficient smoothness and decay at infinity using the inverse scattering transform method [36]. Well-posedness of the ivp for the NLS equation on the circle in Sobolev spaces H^s with $s \geq 0$ has been proven by Bourgain [8] using modern harmonic analysis techniques. Earlier results include the works of Cazenave and Weissler [13], Ginibre and Velo [22], Kenig, Ponce and Vega [29], and Tsutsumi [34]. In addition, the sharp well-posedness result on \mathbb{R} was recently established in [24].

Boundary value problems are particularly significant in real-world applications. For example, in the work [28], the authors study the dynamics of Bose–Einstein condensates confined in cigar-shaped traps, where the trap’s edges impose the boundary conditions. Although ibvps for Eq. (1.1a) are more relevant to real-world applications, they have received relatively little attention due in part to the lack of a Fourier transform in the case of bounded or semi-bounded spatial domains, which poses a significant obstacle to their analysis for dispersive equations like NLS and Korteweg-de Vries (KdV) equations. Nonetheless, researchers have explored various approaches to studying the ibvps, such as using the Riemann–Liouville fractional integration operator [18, 25], the Laplace transform [5–7], and the Fokas unified transform method [20, 21]. Additionally, the ibvp of NLS in two dimensions has been studied in [27, 32]. The regularity properties for the cubic NLS equations on the half-line are discussed in [19] and the global well-posedness in one-dimensional spaces is addressed in [7]. However, the corresponding ibvps for Eq. (1.1a) in bounded regions of higher dimension have not been explored and the goal of this article is to delve into this uncharted territory and establish a foundation for understanding the solution behaviour in more complex and realistic settings.

Here, we investigate the ibvp in $\Omega_j, j = 0, 1, 2$, which are radially symmetric regions of \mathbb{R}^n . We assume that the initial condition u_0 satisfies the following condition of radial symmetry:

$$u_0(x_1, \dots, x_n) = u_0(r), \quad \text{where } r = \sqrt{x_1^2 + \dots + x_n^2}. \tag{1.4}$$

We seek to find radially symmetric solutions of the ibvp. We equip the ibvp with different boundary conditions in three different domains. In Ω_0 , we use the boundary condition:

$$u(x_1, \dots, x_n, t) = g_2(t), \quad x_1^2 + \dots + x_n^2 = 1, \quad t \in (0, T). \tag{1.5}$$

In Ω_1 , we use the boundary condition:

$$u(x_1, \dots, x_n, t) = g(t), \quad x_1^2 + \dots + x_n^2 = 1, \quad t \in (0, T). \tag{1.6}$$

In Ω_2 , we use the boundary condition:

$$\begin{cases} u(x_1, \dots, x_n, t) = g_1(t), & x_1^2 + \dots + x_n^2 = \pi^2, \quad t \in (0, T), \\ u(x_1, \dots, x_n, t) = g_2(t), & x_1^2 + \dots + x_n^2 = (2\pi)^2, \quad t \in (0, T). \end{cases} \tag{1.7}$$

In addition to these boundary conditions, we have established compatibility conditions for each of the problems:

$$\text{The ibvp in } \Omega_0 : \quad u_0(1) = g_2(0), \quad 1 < s < 2, \tag{1.8a}$$

$$\text{The ibvp in } \Omega_1 : \quad u_0(1) = g(0), \quad \frac{1}{2} < s < \frac{3}{2}, \tag{1.8b}$$

$$\text{The ibvp in } \Omega_2 : \quad u_0(\pi) = g_1(0) \quad \text{and} \quad u_0(2\pi) = g_2(0), \quad \frac{1}{2} < s < 2. \tag{1.8c}$$

Furthermore, it is important to highlight that in this article, the lifespan T^* satisfies $0 < T^* \leq T < \frac{1}{2}$, and it depends on both the norm of initial data and the norm of boundary data. In addition, in the theorems that follow, the solutions are radially symmetric. With these clarifications, we can now present the primary outcomes and conclusions of our study.

In the following, $H^s(\Omega)$ is denoted as the classical L^2 -based Sobolev space in Ω with Sobolev index s and $H_0^s(\Omega)$ is the subspace of $H^2(\Omega)$ which is the closure of functions in $H^s(\Omega)$ with compact supports in Ω (formal definitions of those spaces will be given in §2).

THEOREM 1.1 *Let $n=2$. Suppose $u_0 \in H_0^s(\Omega_0)$, satisfying condition (1.4), and $g_2 \in H_0^{\frac{s+1}{2}}(0, T)$. If $1 < s < 2$, $p \geq 3$ and p is an even integer, then the ibvp within domain Ω_0 , subject to the compatibility condition (1.8a), is locally well-posed in $C([0, T^*]; H^s(\Omega_0))$, where the lifespan T^* depends on $\|u_0\|_{H^s(\Omega_0)}$, $\|g_2\|_{H^{\frac{s+1}{2}}(0, T)}$ and p .*

Here, the use of spaces $H_0^s(\Omega_0)$ and $H_0^{\frac{s+1}{2}}(0, T)$ for initial and boundary data merely makes the proof of theorem 1.1 slightly more straightforward since the compatibility conditions for the initial and boundary data are automatically satisfied.

THEOREM 1.2 *Let $n \geq 2$, $u_0 \in H^s(\Omega_1)$, which satisfies the condition (1.4), and $g \in H^{\frac{2s+1}{4}}(0, T)$.*

- If $0 \leq s < \frac{1}{2}$ and $3 \leq p < \frac{6-4s}{1-2s}$, then the ibvp in domain Ω_1 is locally well-posed in $C([0, T^*]; H^s(\Omega_1))$.
- If $\frac{1}{2} < s < \frac{3}{2}$ and $p \geq 3$, then the ibvp in domain Ω_1 with compatibility condition (1.8b) is locally well-posed in $C([0, T^*]; H^s(\Omega_1))$.

In both cases, the lifespan T^ depends on $\|u_0\|_{H^s(\Omega_1)}$, $\|g\|_{H^{\frac{s+1}{4}}(0, T)}$ and p .*

THEOREM 1.3 *Let $n \geq 2$, $u_0 \in H^s(\Omega_2)$, which satisfies the condition (1.4), $g_1 \in H^{\frac{s+1}{2}}(0, T)$ and $g_2 \in H^{\frac{s+1}{2}}(0, T)$.*

- If $0 \leq s < \frac{1}{2}$ and $3 \leq p \leq 4$, then the ibvp in domain Ω_1 is locally well-posed in $C([0, T^*]; H^s(\Omega_2))$.
- If $\frac{1}{2} < s < 2$, $s \neq \frac{3}{2}$ and $p \geq 3$, then the ibvp in domain Ω_2 with compatibility condition (1.8c) is locally well-posed in $C([0, T^*]; H^s(\Omega_2))$.

In both cases, the lifespan T^ depends on $\|u_0\|_{H^s(\Omega_2)}$, $\|g_1\|_{H^{\frac{s+1}{2}}(0, T)}$, $\|g_2\|_{H^{\frac{s+1}{2}}(0, T)}$ and p .*

We remark that the above theorems address a notable gap in current research, which is significantly important in the study of non-homogeneous boundary value

problems for the NLS equations. In particular, [theorem 1.2](#) gives an optimal regularity of the boundary data for the ibvp of NLS equations in Ω_1 , while [theorems 1.1](#) and [1.3](#) provide the first account on the well-posedness issue for the ibvp of NLS equations in bounded regions of higher dimensions. Thus, the results in the article contribute to the limited body of knowledge on the ibvp of NLS equations and add valuable insights to the field, shedding light on a previously under-studied aspect of NLS equations, which we believe will be instrumental to the future research. Moreover, we note that the global well-posedness of ibvps in one-dimensional spaces has been addressed in [\[7\]](#), though it imposes some restrictions on the nonlinearity. For the problems studied in this article, the global well-posedness can be analysed similarly with restrictions on the nonlinearity, utilizing energy conservation and boundary data estimates.

For $s < 0$, [\[17\]](#) showed that the ivp for the cubic NLS equation is ill-posed because the mapping from initial data to solutions fails to be uniformly continuous. For the ibvp in the regions Ω_1 and Ω_2 , we reduce this problem to the one-dimensional ibvp and prove well-posedness holds for $s \geq 0$. Therefore, in terms of the uniform continuity of the data-to-solution map, [theorems 1.2](#) and [1.3](#) are sharp. However, in the recent work [\[24\]](#), the ivp for the cubic NLS equation was shown to be well-posed for $s > -\frac{1}{2}$, which is sharp. We have not yet achieved this sharp result for the ibvp, but we aim to explore this in future work.

Regarding the ibvp in the region Ω_0 , the non-zero boundary conditions complicate the derivation of Strichartz estimates. Thus, in [theorem 1.1](#), we only establish local well-posedness for $s > 1$. The absence of these Strichartz estimates limits the sharpness of our result.

For the unforced case with zero boundary conditions, global well-posedness results have been achieved. In [\[9, 10\]](#), the authors proved global well-posedness for the ibvp of the NLS equation on the two-dimensional and three-dimensional unit balls for $0 < s < \frac{1}{2}$, utilizing and thoroughly discussing Strichartz estimates.

Additionally, in [\[4\]](#), Strichartz estimates were established for the Schrödinger equation on Riemannian manifolds (Ω, g) with zero boundary conditions. This applies both to compact cases and when Ω is the exterior of a smooth, non-trapping obstacle in Euclidean space. Using these estimates, the Schrödinger equation was shown to be well-posed in $H^1(\Omega)$ for three-dimensional space (see [theorems 5.1](#) and [6.3](#) in [\[4\]](#)).

Here, we note that although the works [\[4, 9, 10\]](#) indeed derive Strichartz estimates for problems with zero boundary conditions, our study focuses on the ibvps with non-zero boundary conditions. The Strichartz estimates available in those literatures do not directly apply to such ibvps, which involve boundary integral operators, and as a result, there is a lack of established estimates for the cases we consider. This distinction is crucial to the novelty and challenges for the problem considered here.

In this article, we have not been able to derive the necessary Strichartz estimates for the boundary integral operator $W_{ball}h$ defined by [\(5.27\)](#), which are required to prove the local well-posedness of the ibvp in the region Ω_0 . Hence, the presence of non-zero boundary conditions complicates the derivation of Strichartz estimates, and local well-posedness can only be achieved for $s > 1$. However, we have gained

insights from the works mentioned above and plan to explore this direction further in future research.

Next, we provide an overview of the proof of the main results, which is comprised of four essential steps:

- **Step 1.** We begin by reducing the ibvp (1.1) to the NLS equation in one dimension when $\Omega = \Omega_1$ or $\Omega = \Omega_2$. For $\Omega = \Omega_0$, due to the singularity at $r = 0$ if changing the equation in one dimension, we still use Ω_0 in \mathbb{R}^n .
- **Step 2.** We derive a solution formula for the corresponding linear forced ibvp, which will be crucial in obtaining estimates for the linear problem.
- **Step 3.** Using classical analysis, we obtain linear estimates for the data and forcing in suitable function spaces, which we refer to as ‘good’ solution spaces.
- **Step 4.** We prove that the iteration map defined by the solution formula, with the forcing replaced by the nonlinearity, is a contraction in ‘good’ solution spaces. This will allow us to use the contraction mapping principle to establish the existence of a unique solution to the nonlinear problem.

Here, we remark that, since the solutions and the domains are radially symmetric, we may change the ibvps for $\Omega = \Omega_1$ or Ω_2 to 1D NLS equations on a half line or a finite interval with one or two boundary points. These 1D problems have been studied in [7, 21]. If we require certain linear estimates from [7, 21] in our proof, we may either cite them or provide shorter or more elegant proof.

Paper organization: In §2, we introduce crucial preliminary results that are foundational for our subsequent proofs. Section 3 is dedicated to establishing the well-posedness result for regions outside a ball, with a specific focus on proving theorem 1.2. In §4, we delve into the well-posedness result within an annulus and provide the proof for theorem 1.3. Section 5 is dedicated to demonstrating the well-posedness result within a ball centred at the origin, presenting the proof for theorem 1.1. Additionally, we include an Appendix section where we provide proofs that may have been omitted in earlier sections.

2. Preliminary

In this section, since we only consider the solutions of (1.1) with radial form, we rewrite (1.1) as

$$iu_t + u_{rr} + \frac{n-1}{r}u_r + \lambda|u|^{p-2}u = 0, \quad r \in (r_1, r_2), \quad t \in (0, T), \quad (2.1a)$$

$$u(r, 0) = u_0(r), \quad r \in (r_1, r_2), \quad (2.1b)$$

$$u(r_1, t) = g_1(t), \quad u(r_2, t) = g_2(t), \quad t \in (0, T), \quad (2.1c)$$

where r_1, r_2 are chosen appropriately for $\Omega_j, j = 0, 1, 2$ and $r = (x_1^2 + \dots + x_n^2)^{\frac{1}{2}}$. Here,

$$\Delta u = \partial_{x_1}^2 u + \dots + \partial_{x_n}^2 u = u''(r) + \frac{n-1}{r} u'(r) \tag{2.2}$$

has been used.

The corresponding linear ibvp is

$$i u_t + u_{rr} + \frac{n-1}{r} u_r = f(r, t), \quad r \in (r_1, r_2), \quad t \in (0, T), \tag{2.3a}$$

$$u(r, 0) = u_0(r), \quad r \in (r_1, r_2), \tag{2.3b}$$

$$u(r_1, t) = g_1(t), \quad u(r_2, t) = g_2(t), \quad t \in (0, T). \tag{2.3c}$$

If $r_1 \neq 0$, we can use a change of dependent variable $u(r, t) = r^{-\frac{n-1}{2}} \cdot v(r, t)$ to derive the equations for v , that is

$$i v_t + v_{rr} = r^{\frac{n-1}{2}} f(r, t) + \frac{n^2 - 4n + 3}{4} r^{-2} \cdot v, \quad r \in (r_1, r_2), \quad t \in (0, T), \tag{2.4a}$$

$$v(r, 0) = r^{\frac{n-1}{2}} u_0(r) = v_0(r), \quad r \in (r_1, r_2), \tag{2.4b}$$

$$v(r_1, t) = r_1^{\frac{n-1}{2}} g_1(t), \quad v(r_2, t) = r_2^{\frac{n-1}{2}} g_2(t), \quad t \in (0, T). \tag{2.4c}$$

From the theory of the ivp (1.1) in \mathbb{R}^n , it is known that if the initial data u_0 of radial form is in $W^{s,2}(\Omega_j), j = 0, 1, 2$, then $u_0(r)$ is in $W_{r, n-1}^{s,2}(r_1, r_2)$ for $r_1 \neq 0$, which implies that $v_0(r) \in W^{s,2}(r_1, r_2)$. Here, the weighted Sobolev space $W_\omega^{m,p}$ over the open region Ω is given by (see [35]),

$$\|u\|_{W_\omega^{m,p}(\Omega)} = \left(\sum_{|\alpha| \leq m} \int_\Omega |D^\alpha u|^p \omega dx \right)^{1/p}. \tag{2.5}$$

Hence, we only need to discuss the solutions of (2.4) in L^2 -based Sobolev spaces if $r_1 \neq 0$.

If $r_1 = 0$, then the above change of dependent variables introduces a singularity at $r=0$ and cannot be used, which implies that the weighted Sobolev spaces are necessary. For the case that $r_2 = \infty$, the problem (2.1) is the NLS equation posed in \mathbb{R}^n with radial symmetric initial data. If we consider the following ivp of linear Schrödinger equation

$$iU_t + \Delta U = F, \quad (x_1, \dots, x_n) \in \mathbb{R}^n, \quad t \in (0, T), \tag{2.6a}$$

$$U(x_1, \dots, x_n, 0) = U_0(x_1, \dots, x_n), \quad (x_1, \dots, x_n) \in \mathbb{R}^n, \tag{2.6b}$$

then using Fourier transform, we have

$$U = S_n[U_0; F] \doteq \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\xi \cdot x - i|\xi|^2 t} \widehat{U}_0(\xi_1, \dots, \xi_n) d\xi_1 \cdots d\xi_n \tag{2.7a}$$

$$- \frac{i}{(2\pi)^n} \int_0^t \int_{\mathbb{R}^n} e^{i\xi \cdot x - i|\xi|^2(t-t')} \widehat{F}^x(\xi_1, \dots, \xi_n, t') d\xi_1 \cdots d\xi_n dt', \tag{2.7b}$$

where $\xi \cdot x = \xi_1 x_1 + \cdots + \xi_n x_n$ and $|\xi|^2 = \xi_1^2 + \cdots + \xi_n^2$. Also, we have following claim.

Claim. If U_0 and F are radially symmetric, that is, for $r \geq 0$ we have

$$U_0(x_1, \dots, x_n) = U_0(r), \quad F(x_1, \dots, x_n, t) = F(r, t), \quad x_1^2 + \cdots + x_n^2 = r^2,$$

then the solution of the above ivp (2.6) is also radially symmetric.

Proof. The above claim follows from the next result.

LEMMA 2.1. *If f is radially symmetric, then \widehat{f} is also radially symmetric. Conversely, if \widehat{f} is radially symmetric, then f is radially symmetric.*

Now, since the term (2.7a) is the inverse Fourier transform of $e^{-i|\xi|^2 t} \widehat{U}_0(\xi_1, \dots, \xi_n)$, by lemma 2.1, the term (2.7a) is radially symmetric. Similarly, since term (2.7b) is the integral of the inverse Fourier transform $e^{-i|\xi|^2(t-t')} \widehat{F}(\xi_1, \dots, \xi_n, t')$, lemma 2.1 implies that this term is also radially symmetric. This completes the claim. \square

Proof of lemma 2.1. Here, we only prove that the Fourier transform of a radially symmetric function is radially symmetric. The Fourier transform of function $f(x)$ in \mathbb{R}^n is defined as:

$$\widehat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-i\xi \cdot x} dx, \tag{2.8}$$

where $\xi \cdot x = \xi_1 x_1 + \cdots + \xi_n x_n$ denotes the dot product of the vectors ξ and x . To show that $\widehat{f}(\xi)$ is also radially symmetric, we need to show that $\widehat{f}(\xi)$ is invariant under rotations, i.e., if we rotate the vector ξ in \mathbb{R}^n by an angle θ , the Fourier transform $\widehat{f}(\xi)$ remains the same.

Let R be a rotation matrix in \mathbb{R}^n , i.e., R is an $n \times n$ orthogonal matrix with determinant 1. Then, we have:

$$\begin{aligned} \widehat{f}(R\xi) &= \int_{\mathbb{R}^n} f(r) e^{-i(R\xi) \cdot x} dx = \int_{\mathbb{R}^n} f(r) e^{-i\xi \cdot R^{-1}x} dx \stackrel{y=R^{-1}x}{=} \int_{\mathbb{R}^n} f(r) e^{-i\xi \cdot y} dy \\ &= \widehat{f}(\xi), \end{aligned}$$

where we have used the fact that $R^{-1} = R^T$ for an orthogonal matrix R .

Thus, we have shown that the Fourier transform of a radially symmetric function in \mathbb{R}^n is also radially symmetric, i.e., $\widehat{f}(\xi) = \widehat{f}(|\xi|)$. This completes the proof of lemma 2.1. \square

Therefore, by above discussion, we can use the well-posedness theory of the NLS equations in \mathbb{R}^n to establish the well-posedness of (2.1) with $r_1 = 0$ and $r_2 = \infty$ under the assumption that $u_0(r) \in W_{r^{n-1}}^{s,2}(\mathbb{R})$ with $s \geq 0$. We note that the boundary condition for (2.1) at $r = 0$ must be $v_r(0) = 0$ and the solution space is $W_{r^{n-1}}^{s,2}(\mathbb{R})$ with $s \geq 0$. Hence, in the following, we will only consider the cases with $0 < r_1 < r_2 = \infty$, $0 < r_1 < r_2 < \infty$, and $0 = r_1 < r_2 < \infty$.

Here, we recall the linear estimate for the solution $S_n[U_0; F]$.

PROPOSITION 2.2. *[Strichartz estimates for linear Schrödinger equation]*

For $s \geq 0$, if (q, γ) and (q_1, γ_1) are admissible, which are given below in definition 2.3. Then the solution $U = S_n[U_0; F]$ of ivp (2.6) satisfies

$$\|S_n[U_0; F]\|_{L^q(0,T;W^{s,\gamma}(\mathbb{R}^n))} \lesssim \|U_0\|_{H^s(\mathbb{R}^n)} + \|F\|_{L^{q_1}(0,T;W^{s,\gamma_1}(\mathbb{R}^n))}. \tag{2.9}$$

The proof of proposition 2.2 can be found in [11] (see theorem 2.3.3).

Throughout this work, we shall use the familiar **time localizer** $\psi(t)$, which is defined as follows:

$$\psi \in C_0^\infty(-1, 1), \quad 0 \leq \psi \leq 1 \text{ and } \psi(t) = 1 \text{ for } |t| \leq \frac{1}{2}. \tag{2.10}$$

Moreover, we introduce the notion of admissible pair

DEFINITION 2.3. *We say that a pair (q, γ) is admissible (in n dimension), if*

$$\frac{2}{q} + \frac{n}{\gamma} = \frac{n}{2}, \tag{2.11}$$

and

$$2 \leq \gamma \leq \frac{2n}{n-2}, \quad (2 \leq \gamma \leq \infty \text{ if } n = 1, \quad 2 \leq \gamma < \infty \text{ if } n = 2). \tag{2.12}$$

Additionally, the Sobolev space $H^s(\mathbb{R}^n)$ consists of all temperate distributions F with the norm

$$\|F\|_{H^s(\mathbb{R}^n)} \doteq \left(\int_{\mathbb{R}^n} (1 + |\xi|)^{2s} |\widehat{F}(\xi)|^2 d\xi \right)^{1/2}, \tag{2.13}$$

where $\widehat{F}(\xi)$ is the Fourier transform defined by

$$\widehat{F}(\xi) \doteq \int_{\mathbb{R}^n} e^{-i\xi \cdot x} F(x) dx.$$

For an open set $\Omega \subset \mathbb{R}^n$, the space $H^s(\Omega)$ is defined by

$$H^s(\Omega) \doteq \{f : f = F|_\Omega \text{ where } F \in H^s(\mathbb{R}^n) \text{ and } \|f\|_{H^s(\Omega)} \doteq \inf_{F \in H^s(\mathbb{R}^n)} \|F\|_{H^s(\mathbb{R}^n)} < \infty\}. \tag{2.14}$$

Here, we remind the reader that the space $H_0^s(\Omega)$ is the subspace which is the closure of the class of functions in $H^s(\mathbb{R}^n)$ whose support lies in Ω .

Finally, we define the Sobolev space on a torus, which will be used to study the problem on annulus. For $s \geq 0$, the Sobolev space $H^s(\mathbb{T})$ is defined by

$$H^s(\mathbb{T}) \doteq \left\{ f \in L^2(\mathbb{T}) : \|f\|_s \doteq \left(\sum_{n \in \mathbb{Z}} (1 + |n|)^{2s} |\hat{f}(n)|^2 \right)^{1/2} < \infty \right\}. \tag{2.15}$$

Also, recall the Fourier transform

$$\hat{f}(n) = \int_{-\pi}^{\pi} e^{-inx} f(x) dx, \quad n \in \mathbb{Z}, \tag{2.16}$$

and the inverse Fourier transform

$$f(x) = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{inx}. \tag{2.17}$$

Equations (2.16) and (2.17) present identities that hold in the sense of distributions. Specifically, for functions f such that $f \in L^1$, the Fourier transform \hat{f} is in ℓ^1 , ensuring the validity of the identities as stated.

3. NLS equations on a half-line (i.e., outside of a ball)

In this section, we study (2.1) on a half-line with $r \in (1, \infty)$ (i.e., outside of a ball) and a boundary condition $u(1, t) = g(t)$, where $r_1 = 1$ is chosen for the sake of convenience. We first discuss the corresponding linear problem and then obtain the well-posedness of the nonlinear problem.

3.1. Solutions of linear problems with estimates in Sobolev Spaces

If $f_1(r, t) \doteq r^{\frac{n-1}{2}} f(r, t) + \frac{n^2-4n+3}{4} r^{-2} \cdot v$ and $v_0(r) = r^{\frac{n-1}{2}} u_0(r)$, then (2.4) becomes

$$iv_t + v_{rr} = f_1(r, t), \quad r > 1, \quad t \in (0, T), \tag{3.1a}$$

$$v(r, 0) = v_0(r), \quad r > 1, \tag{3.1b}$$

$$v(1, t) = g(t), \quad t \in (0, T). \tag{3.1c}$$

Also, by using the compatibility (1.8b), the above ibvp is equipped with the following compatibility condition

$$g(0) = v_0(1), \quad \frac{1}{2} < s < \frac{3}{2}. \tag{3.2}$$

Next, we solve the ibvp (3.1) and begin with decomposing the above ibvp into simpler problems. In fact, using superposition principle, the linear ibvp (3.1) can be expressed as the homogeneous ibvp

$$iv_t + v_{rr} = 0, \quad r > 1, \quad t \in (0, T), \quad (3.3a)$$

$$v(r, 0) = v_0(r) \in H^s(1, \infty), \quad r > 1, \quad (3.3b)$$

$$v(1, t) = g(t) \in H^{\frac{2s+1}{4}}(0, T), \quad t \in (0, T), \quad (3.3c)$$

and the forced linear ibvp with zero initial and boundary data

$$iv_t + v_{rr} = f_1(r, t), \quad r > 1, \quad t \in (0, T), \quad (3.4a)$$

$$v(r, 0) = 0, \quad r > 1, \quad (3.4b)$$

$$v(1, t) = 0, \quad t \in (0, T). \quad (3.4c)$$

Also, we can do further decomposition. In fact, the homogeneous ibvp (3.3) can be expressed as the homogeneous ivp and the pure ibvp. The homogeneous ivp is given by:

$$iV_t + V_{rr} = 0, \quad t \in (0, T), \quad (3.5a)$$

$$V(r, 0) = V_0(r) \in H^s(\mathbb{R}), \quad (3.5b)$$

where V_0 is the extension of v_0 from $(1, \infty)$ to \mathbb{R} such that

$$\|V_0\|_{H^s(\mathbb{R})} \leq 2\|v_0\|_{H^s(1, \infty)}. \quad (3.6)$$

The pure ibvp is given by:

$$iv_t + v_{rr} = 0, \quad r > 1, \quad t \in (0, T), \quad (3.7a)$$

$$v(r, 0) = 0, \quad r > 1, \quad (3.7b)$$

$$v(1, t) = g(t) - V(1, t) \in H^{\frac{2s+1}{4}}(0, T), \quad t \in (0, T). \quad (3.7c)$$

For the inhomogeneous ibvp (3.4), it can be decomposed as a forced ivp and a pure ibvp:

$$iW_t + W_{rr} = F(r, t), \quad r > 1, \quad t \in (0, T), \quad (3.8a)$$

$$W(r, 0) = 0, \quad (3.8b)$$

where F is the extension of f_1 from $(1, \infty)$ to \mathbb{R} such that

$$\|F\|_{L_t^{q'}(0, T; W^{s, \gamma'}(\mathbb{R}))} \leq 2\|f_1\|_{L_t^{q'}(0, T; W^{s, \gamma'}(1, \infty))}. \quad (3.9)$$

The pure ibvp is given by:

$$iv_t + v_{rr} = 0, \quad r > 1, t \in (0, T), \tag{3.10a}$$

$$v(r, 0) = 0, \quad r > 1, \tag{3.10b}$$

$$v(1, t) = -W(1, t) \quad t \in (0, T). \tag{3.10c}$$

Linear estimate for homogeneous ivp (3.5). The solution to this problem is obtained by Fourier transform

$$V(r, t) = S[V_0; 0](r, t) \doteq \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\xi r - i\xi^2 t} \widehat{V}_0(\xi) d\xi, \tag{3.11}$$

where \widehat{V}_0 is the Fourier transform of V_0 , that is

$$\widehat{V}_0 \doteq \int_{\mathbb{R}} e^{-i\xi r} V_0(r) dr.$$

We have the following estimates for $S[V_0; 0]$, whose proof can be found in [12, 25].

PROPOSITION 3.1. Homogeneous ivp estimates *The solution $V = S[V_0; 0]$ of the homogeneous linear Schrödinger ivp (3.5) given by formula (3.11) satisfies the space estimate:*

$$\sup_{t \in [0, T]} \|S[V_0; 0](t)\|_{H^s(\mathbb{R})} = \|V_0\|_{H^s(\mathbb{R})}, \quad s \in \mathbb{R}. \tag{3.12}$$

Also, $S[V_0; 0]$ satisfies the time estimate

$$\sup_{r \in \mathbb{R}} \|S[V_0; 0](r)\|_{H^{\frac{2s+1}{4}}(0, T)} \lesssim \|V_0\|_{H^s(\mathbb{R})}, \quad s \in \mathbb{R}. \tag{3.13}$$

In addition, if $\frac{2}{q} + \frac{1}{\gamma} = \frac{1}{2}$ and $\gamma \geq 2$, then $S[V_0; 0](r, t)$ satisfies the following Strichartz estimate

$$\|S[V_0; 0]\|_{L_t^q(\mathbb{R}; W^{s, \gamma}(\mathbb{R}))} \lesssim \|V_0\|_{H^s(\mathbb{R})}, \quad s \geq 0. \tag{3.14}$$

Linear estimate for forced ivp (3.8). The solution to this problem is given by

$$W = S[0; F](r, t) \doteq -\frac{i}{2\pi} \int_{t'=0}^t \int_{\mathbb{R}} e^{i\xi r - i\xi^2(t-t')} \widehat{F}^r(\xi, t') d\xi dt' \tag{3.15a}$$

$$= -i \int_{t'=0}^t S[F(\cdot, t'); 0](r, t - t') dt', \tag{3.15b}$$

where \widehat{F}^r is the Fourier transform of F and $S[F(\cdot, t'); 0]$ denotes the solution of homogeneous ivp (3.5) with initial datum $F(r, t')$. We have the following estimates for $S[0; F](r, t)$, whose proof also can be found in [12, 25]

PROPOSITION 3.2. Forced ivp estimates *The solution $W = S[0; F]$ of the forced ivp (3.8) given by formula (3.15) satisfies the space estimate:*

$$\sup_{t \in [0, T]} \|S[0; F](t)\|_{H^s(\mathbb{R})} \leq T \sup_{t \in [0, T]} \|F(t)\|_{H^s(\mathbb{R})}, \quad s \in \mathbb{R}. \tag{3.16}$$

Also, $S[0; F]$ satisfies the time estimate

$$\sup_{r \in \mathbb{R}} \|S[0; F](r)\|_{H_t^{\frac{2s+1}{4}}(0, T)} \lesssim (1 + T)^{\frac{1}{4}} \|F\|_{L_t^{q'}(0, T; W^{s, \gamma'}(\mathbb{R}))}, \quad -\frac{1}{2} < s < \frac{1}{2}, \tag{3.17}$$

$$\sup_{r \in \mathbb{R}} \|S[0; F](r)\|_{H_t^{\frac{2s+1}{4}}(0, T)} \lesssim \|F\|_{L^1(0, T; H^s(\mathbb{R}))}, \quad \frac{1}{2} < s < \frac{3}{2}. \tag{3.18}$$

In addition, if $\frac{2}{q} + \frac{1}{\gamma} = \frac{1}{2}$, and $\gamma \geq 2$, then $S[0; F](r, t)$ satisfies the following Strichartz estimate

$$\|S[0; F]\|_{L_t^q(0, T; W^{s, \gamma}(\mathbb{R}))} \lesssim \|F\|_{L_t^{q'}(0, T; W^{s, \gamma'}(\mathbb{R}))}, \quad s \geq 0. \tag{3.19}$$

Next, we study the following ibvp

$$iv_t + v_{rr} = 0, \quad r > 1, \quad t \in (0, T), \tag{3.20a}$$

$$v(r, 0) = 0, \quad r > 1, \tag{3.20b}$$

$$v(1, t) = g_1(t) \in H^{\frac{2s+1}{4}}(0, T), \quad t \in (0, T). \tag{3.20c}$$

Also, we extend the boundary data $g_1(t)$ from $(0, T)$ to \mathbb{R} by the following result whose proof can be found in [21, 30].

LEMMA 3.3. For a general function $h^*(t) \in H_t^m(0, 2)$ with $m \geq 0$, let the extension

$$\tilde{h}^*(t) \doteq \begin{cases} h^*(t), & t \in (0, 2), \\ 0, & \text{elsewhere.} \end{cases}$$

If $0 \leq m < \frac{1}{2}$, then the extension $\tilde{h}^* \in H^m(\mathbb{R})$ and for some $c_m > 0$ we have

$$\|\tilde{h}^*\|_{H_t^m(\mathbb{R})} \leq c_m \|h^*\|_{H_t^m(0, 2)}. \tag{3.21}$$

If $\frac{1}{2} < m < \frac{3}{2}$, then for estimate (3.21) to hold we must have the condition

$$h^*(0) = h^*(2) = 0. \tag{3.22}$$

In fact, for $-\frac{1}{2} < s < \frac{1}{2}$ or $0 \leq \frac{2s+1}{4} < \frac{1}{2}$, we define

$$h(t) \doteq \begin{cases} g_1(t), & t \in (0, T), \\ 0, & t \notin (0, T). \end{cases}$$

Then, using lemma 3.3, it is obtained that $h(t)$ is compactly supported in $(0, 2)$ and

$$\|h\|_{\frac{2s+1}{4}(\mathbb{R})} \lesssim \|g_1\|_{\frac{2s+1}{4}(0,T)}.$$

For $\frac{1}{2} < s < \frac{3}{2}$ or $\frac{1}{2} < \frac{2s+1}{4} < 1$, we first extend g_1 from $(0, 2)$ to \mathbb{R} such that $\|g_2\|_{\frac{2s+1}{4}(\mathbb{R})} \leq 2\|g_1\|_{\frac{2s+1}{4}(0,T)}$. Next, we define

$$h(t) \doteq \begin{cases} g_2(t), & t \in (0, 2), \\ 0, & t \notin (0, 2). \end{cases}$$

Again, by lemma 3.3, $h(t)$ is compactly supported in $(0, 2)$ and $\|h\|_{\frac{2s+1}{4}(\mathbb{R})} \lesssim \|g_1\|_{\frac{2s+1}{4}(0,T)}$. Therefore, the ibvp (3.20) becomes

$$iv_t + v_{rr} = 0, \quad r > 1, \quad t \in (0, T), \quad (3.23a)$$

$$v(r, 0) = 0, \quad r > 1, \quad (3.23b)$$

$$v(1, t) = h(t) \in H^{\frac{2s+1}{4}}(0, 2), \quad t \in (0, 2), \quad (3.23c)$$

where $h(t)$ is compactly supported in $(0, 2)$. Using Laplace transform (see [7]) or the Fokas method (see [21]), we derive the solution for the reduced pure ibvp (3.23)

$$\begin{aligned} v &= S_b[0, h; 0] \doteq \frac{1}{2\pi} \int_{-\infty}^0 e^{i\beta t} e^{i\sqrt{-\beta}(r-1)} \tilde{h}(i\beta) d\beta + \frac{1}{2\pi} \int_0^{\infty} e^{i\beta t} e^{-\sqrt{-\beta}(r-1)} \tilde{h}(i\beta) d\beta \\ &= I_1 + I_2, \end{aligned} \quad (3.24)$$

where the integrals I_1 and I_2 are defined by

$$I_1(r, t) \doteq \frac{1}{2\pi} \int_{-\infty}^0 e^{i\beta t} e^{i\sqrt{-\beta}(r-1)} \tilde{h}(i\beta) d\beta = \frac{1}{\pi} \int_0^{\infty} e^{-i\beta^2 t} e^{i\beta(r-1)} \beta \tilde{h}(-i\beta^2) d\beta, \quad (3.25)$$

$$r > 1, \quad (3.25)$$

$$I_2(r, t) \doteq \frac{1}{2\pi} \int_0^{\infty} e^{i\beta t} e^{-\sqrt{-\beta}(r-1)} \tilde{h}(i\beta) d\beta = \frac{1}{\pi} \int_0^{\infty} e^{i\beta^2 t} e^{-\beta(r-1)} \beta \tilde{h}(i\beta^2) d\beta, \quad r > 1. \quad (3.26)$$

Since $h(t)$ is compactly supported in $(0, 2)$, we have

$$\tilde{h}(-i\beta^2) = \int_0^{\infty} e^{i\beta^2 t} h(t) dt = \int_{\mathbb{R}} e^{i\beta^2 t} h(t) dt = \hat{h}(-\beta^2) \quad \text{and} \quad \tilde{h}(i\beta^2) = \hat{h}(\beta^2). \quad (3.27)$$

PROPOSITION 3.4. *The solution $v = S_b[0, h; 0]$ of the ivvp (3.20) given by formula (3.24) satisfies the space estimate:*

$$\sup_{t \in [0, T]} \|S_b[0, h; 0](t)\|_{H^{s(1, \infty)}} \lesssim \|h\|_{H^{\frac{2s+1}{4}}(\mathbb{R})}, \quad s \geq 0. \tag{3.28}$$

Also, if $\frac{2}{q} + \frac{1}{\gamma} = \frac{1}{2}$ then $S_b[0, h; 0](r, t)$ satisfies the following Strichartz estimate

$$\|S_b[0, h; 0]\|_{L_t^q(0, T; W^{s, \gamma}(1, \infty))} \lesssim \|h\|_{H^{\frac{2s+1}{4}}(\mathbb{R})}, \quad s \geq 0. \tag{3.29}$$

Proof of proposition 3.4. For the proof of estimate (3.28), we refer to [7, 21]. Here, we only provide the proof of Strichartz estimate (3.29), which was also discussed in [7]. In this exposition, we offer an alternative proof.

Proof of Strichartz estimate (3.29). The proof for I_1 is similar to that of estimate (3.14) and here we omit it. Next, we prove estimate (3.29) for I_2 . Making the change of variables $\tau = \beta^2$, we get

$$I_2(r, t) \simeq \int_0^\infty e^{i\tau t} e^{-\sqrt{\tau}(r-1)} \widehat{h}(\tau) d\tau = \int_0^\infty K_t(r, \tau) \cdot (1 + |\tau|)^{\frac{1}{4}} \widehat{h}(\tau) d\tau, \quad r > 1, \tag{3.30}$$

where the kernel $K_t(r, \tau)$ is defined as follows

$$K_t(r, \tau) \doteq e^{i\tau t} e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}}, \quad r > 1. \tag{3.31}$$

Also, we see that estimate (3.29) follows from the following result

$$\left\| \int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau \right\|_{L_t^q(0, T; L^\gamma(1, \infty))} \lesssim \|f\|_{L^2}, \quad \frac{2}{q} + \frac{1}{\gamma} = \frac{1}{2} \text{ and } 2 \leq \gamma \leq \infty. \tag{3.32}$$

The proof of estimate (3.32) is provided in Appendix. Now, using (3.32), we show the estimate (3.29). To do this, we will consider the following two cases.

Case 1: $s \in \mathbb{N}$. Taking partial derivative ∂_r^s , we have

$$\partial_r^s I_2(r, t) \simeq \int_0^\infty e^{i\tau t} e^{-\sqrt{\tau}(r-1)} (-\sqrt{\tau})^s \widehat{h}(\tau) d\tau \simeq \int_0^\infty K_t(r, \tau) \cdot |\tau|^{\frac{s}{2}} (1 + |\tau|)^{\frac{1}{4}} \widehat{h}(\tau) d\tau. \tag{3.33}$$

Next, apply (3.32) with $\widehat{f}(\tau) = |\tau|^{\frac{s}{2}} (1 + |\tau|)^{\frac{1}{4}} \widehat{h}(\tau)$ to obtain

$$\begin{aligned} \|I_2\|_{L_t^q(0, T; W^{s, \gamma}(1, \infty))} &= \|\partial_r^s I_2\|_{L_t^q(0, T; L^\gamma(1, \infty))} \lesssim \left(\int_{\mathbb{R}} |\tau|^s (1 + |\tau|)^{\frac{1}{2}} |\widehat{h}(\tau)|^2 d\tau \right)^{1/2} \\ &\lesssim \|h\|_{H^{\frac{2s+1}{4}}}, \end{aligned}$$

which is the desired estimate (3.29).

Case 2: $s \geq 0$ and $s \notin \mathbb{N}$. We prove this by interpolation. In fact, any $s \geq 0$ can be written as $s = (1 - \theta)[s] + \theta([s] + 1)$. Furthermore, in Case 1, we proved that

$$\begin{aligned} \|I_2\|_{L_t^q(0,T;W^{[s],\gamma}(1,\infty))} &\lesssim \|h\|_{H^{\frac{2[s]+1}{4}}} \quad \text{and} \quad \|I_2\|_{L_t^q(0,T;W^{[s]+1,\gamma}(1,\infty))} \\ &\lesssim \|h\|_{H^{\frac{2([s]+1)+1}{4}}}, \end{aligned}$$

which implies that I_2 is a continuous linear operator from $H^{\frac{2[s]+1}{4}}$ to $L_t^q(0, T; W^{[s],\gamma}(1, \infty))$ as well as from $H^{\frac{2([s]+1)+1}{4}}$ to $L_t^q(0, T; W^{[s]+1,\gamma}(1, \infty))$. Thus, according to Theorem 5.1 of [30] (see also [3]), we see that I_2 is a continuous linear operator from $H^{\frac{2s+1}{4}}$ to $L_t^q(0, T; W^{s,\gamma}(1, \infty))$. This completes the proof of Case 2 and estimate (3.29). \square

Now, we can derive the linear estimate for the solution of ibvp (3.1). In fact, this solution is given by

$$v(r, t) = \Lambda[v_0, g; f_1] \doteq S[V_0; 0] + S[0; F] + S_b[0, g - S[V_0; F](1, t); 0], \quad r > 1, \tag{3.34}$$

$$t \in (0, T), \tag{3.34}$$

where V_0 is an extension of v_0 satisfying inequality (3.6) and F is an extension of f_1 satisfying inequality (3.9). Combining propositions 3.1–3.4 and using inequalities (3.6) and (3.9), we obtain the following linear estimate.

THEOREM 3.5 *The following estimates hold.*

- (1) *Suppose that $0 \leq s < \frac{1}{2}$. If $v_0 \in H^s(1, \infty)$, $g \in H^{\frac{2s+1}{4}}(0, T)$ and $f_1 \in L_t^{q'}(0, T; W^{s,\gamma'}(1, \infty))$, where (q, γ) is admissible with $n = 1$, then $\Lambda[v_0, g; f_1]$ defines a solution to the linear ibvp (3.1), which satisfies*

$$\begin{aligned} \sup_{t \in [0, T]} \|\Lambda[v_0, g; f_1](t)\|_{H^s(1, \infty)} + \|\Lambda[v_0, g; f_1]\|_{L_t^q(0, T; W^{s,\gamma}(1, \infty))} \\ \lesssim \|v_0\|_{H^s(1, \infty)} + \|g\|_{H^{\frac{2s+1}{4}}(0, T)} + \|f_1\|_{L_t^{q'}(0, T; W^{s,\gamma'}(1, \infty))}. \end{aligned} \tag{3.35}$$

- (2) *Suppose that $\frac{1}{2} < s < \frac{3}{2}$. If $v_0 \in H^s(1, \infty)$, $g \in H^{\frac{2s+1}{4}}(0, T)$ and $f_1 \in L^1(0, T; H^s(1, \infty))$ then $\Lambda[v_0, g; f_1]$ defines a solution to the linear ibvp (3.1) with compatibility condition (3.2), which satisfies*

$$\begin{aligned} \sup_{t \in [0, T]} \|\Lambda[v_0, g; f_1](t)\|_{H^s(1, \infty)} &\lesssim \|v_0\|_{H^s(1, \infty)} + \|g\|_{H^{\frac{2s+1}{4}}(0, T)} \\ &\quad + \|f_1\|_{L^1(0, T; H^s(1, \infty))}. \end{aligned} \tag{3.36}$$

3.2. Proof of well-posedness for ibvp in domain Ω_1 , i.e., theorem 1.2

Existence of solutions for nonlinear problems on half line. Since the ibvp in Ω_1 is reduced to the ibvp (2.1) for $r \in (1, \infty)$ with $u(1, t) = g(t)$. Now, it suffices to prove the existence of solutions of ibvp (3.1) with forcing f_1 giving by

$$f_1(r, t) = -\lambda r^{\frac{n-1}{2}} |u|^{p-2} u + \frac{n^2 - 4n + 3}{4} r^{-2} \cdot v = -\lambda r^{-\frac{(n-1)(p-2)}{2}} |v|^{p-2} v + \frac{n^2 - 4n + 3}{4} r^{-2} \cdot v. \tag{3.37}$$

The case. $0 \leq s < \frac{1}{2}$. In the solution formula (3.34), replacing f_1 by the nonlinearity above, we obtain the iteration map

$$v = \Lambda[v_0, g; f_1] = \Lambda[v_0, g; -\lambda r^{-\frac{(n-1)(p-2)}{2}} |v|^{p-2} v + \frac{n^2 - 4n + 3}{4} r^{-2} \cdot v]. \tag{3.38}$$

Next, we will show that the iteration map (3.38) is a contraction in the following solution space

$$Z = C([0, T^*]; H^s(1, \infty)) \cap L_t^q(0, T^*; W^{s, \gamma}(1, \infty)), \tag{3.39}$$

where (q, γ) is an admissible pair (with $n = 1$), which are defined as

$$q \doteq \frac{4p}{(p-2)(1-2s)} \quad \text{and} \quad \gamma \doteq \frac{p}{1+(p-2)s}. \tag{3.40}$$

We notice that q and γ satisfy $\gamma \geq 2$ and $q \geq 2(\frac{2}{1-2s} + 1)$. The linear estimate (3.35) implies

$$\begin{aligned} & \sup_{t \in [0, T^*]} \|\Lambda[v_0, g; f_1](t)\|_{H^s(1, \infty)} + \|\Lambda[v_0, g; f_1]\|_{L_t^q(0, T^*; W^{s, \gamma}(1, \infty))} \lesssim \|v_0\|_{H^s(1, \infty)} \\ & + \|g\|_{H^{\frac{2s+1}{4}}(0, T)} + |\lambda| \|r^{-\frac{(n-1)(p-2)}{2}} |v|^{p-2} v\|_{L_t^{q'}(0, T^*; W^{s, \gamma'}(1, \infty))} \\ & + \|r^{-2} \cdot v\|_{L_t^{q'}(0, T^*; W^{s, \gamma'}(1, \infty))}. \end{aligned} \tag{3.41}$$

Now, we need to bound the nonlinear terms in the above inequality.

Estimate for. $\|r^{-\frac{(n-1)(p-2)}{2}} |v|^{p-2} v\|_{L_t^{q'}(0, T^*; W^{s, \gamma'}(1, \infty))}$. We extend $r^{-\frac{(n-1)(p-2)}{2}}$ from $(1, \infty)$ to \mathbb{R} such that the extension $k(r) \in C^\infty(\mathbb{R})$ and it is described as follows

$$\begin{cases} k(r) = |r|^{-\frac{(n-1)(p-2)}{2}}, & |r| > 1, \\ k(r) \leq 2, & |r| \leq 1. \end{cases}$$

Also, we extend v from $(1, \infty) \times (0, T^*)$ to $\mathbb{R} \times (0, T^*)$ such that the extension V satisfies

$$\|V\|_{L_t^q(0, T^*; W^{s, \gamma}(\mathbb{R}))} \leq 2\|v\|_{L_t^q(0, T^*; W^{s, \gamma}(1, \infty))}. \tag{3.42}$$

Now, we have

$$\|r^{-\frac{(n-1)(p-2)}{2}}|v|^{p-2}v\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))} \leq \|k(\cdot)|V|^{p-2}V\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(\mathbb{R}))}.$$

Furthermore, using the chain rule (see lemma 5 in [25]) and using $\|k\|_{L^\infty} \leq 2$, we obtain

$$\begin{aligned} \|k(\cdot)|V|^{p-2}V(t)\|_{W^{s,\gamma'}(\mathbb{R})} &\lesssim \|k(\cdot)|V|^{p-2}(t)\|_{L^{\gamma''}} \|\mathcal{D}^s V(t)\|_{L^\gamma} \\ &\leq \|k\|_{L^\infty} \| |V|^{p-2}(t) \|_{L^{\gamma''}} \|\mathcal{D}^s V(t)\|_{L^\gamma} \\ &\lesssim \| |V|^{p-2}(t) \|_{L^{\gamma''}} \|\mathcal{D}^s V(t)\|_{L^\gamma}, \end{aligned}$$

where $\frac{1}{\gamma''} = \frac{1}{\gamma'} - \frac{1}{\gamma} = 1 - \frac{2}{\gamma}$. Moreover, applying Sobolev–Gagliardo–Nirenberg inequality (we refer to theorem 1.3.4 in [12] and corollary 1.5 in [23]) with $\frac{1}{(p-2)\gamma''} = \frac{1}{\gamma} - s$, it is obtained that $\| |V|^{p-2}(t) \|_{L^{\gamma''}} \lesssim \|\mathcal{D}^s V(t)\|_{L^\gamma}^{p-2}$. Finally, combining above inequalities with Hölder’s inequality and using inequality (3.42) give

$$\|r^{-\frac{(n-1)(p-2)}{2}}|v|^{p-2}v\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))} \lesssim T^{*\sigma} \|V\|_{L_t^q(0,T^*;W^{s,\gamma}(\mathbb{R}))}^{p-1} \tag{3.43}$$

$$\lesssim T^{*\sigma} \|v\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}^{p-1}, \tag{3.43}$$

for some $\sigma > 0$. Working similarly to inequality (3.43) (see also [25, 27]), we have

$$\begin{aligned} &\|r^{-\frac{(n-1)(p-2)}{2}}(|v_1|^{p-2}v_1 - |v_2|^{p-2}v_2)\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))} \\ &\lesssim T^{*\sigma} (\|v_1\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}^{p-2} + \|v_2\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}^{p-2}) \|v_1 - v_2\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}. \end{aligned} \tag{3.44}$$

Estimate for $\|r^{-2}v\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))}$. Since $r^{-2} \in C^\infty(1,\infty)$ and it is bounded by 1, we derive

$$\|r^{-2}v\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))} \lesssim T^{*\sigma} \|v\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}, \tag{3.45}$$

$$\|r^{-2}(v_1 - v_2)\|_{L_t^{q'}(0,T^*;W^{s,\gamma'}(1,\infty))} \lesssim T^{*\sigma} \|v_1 - v_2\|_{L_t^q(0,T^*;W^{s,\gamma}(1,\infty))}. \tag{3.46}$$

Combining linear estimate (3.41) with inequalities (3.44) and (3.45), we can show that the iteration map (3.38) is contraction in solution space Z for small T^* . Since the argument is standard, we omit it here. This completes the proof for the case $0 \leq s < \frac{1}{2}$.

The case $\frac{1}{2} < s < \frac{3}{2}$. In this case, we choose $q = \infty$ and $\gamma = 2$. Since the argument is similar to the case $0 \leq s < \frac{1}{2}$, the proof is omitted here.

Uniqueness and Lipschitz continuity of data-to-solution map. The argument for uniqueness and Lipschitz continuity of data-to-solution map is similar to that in [25] and hence is omitted here. We complete the proof of theorem 1.2.

4. ibvp on annulus

In this section, we study the ibvp on annulus, that is, Eq. (2.1) with $0 < r_1 < r_2 < +\infty$. The boundary conditions are

$$v(r_1, t) = \tilde{g}_1(t) = r_1^{\frac{n-1}{2}} g_1(t), \quad v(r_2, t) = \tilde{g}_2(t) = r_2^{\frac{n-1}{2}} g_2(t).$$

For the sake of convenience, we choose $r_1 = \pi$ and $r_2 = 2\pi$.

4.1. Linear problem on the annulus

We first consider the linear Schrödinger equation on the interval, i.e.,

$$iv_t + v_{rr} = f_1(r, t), \quad \pi < r < 2\pi, \quad t \in (0, T), \tag{4.1a}$$

$$v(r, 0) = v_0(r), \quad \pi < r < 2\pi, \tag{4.1b}$$

$$v(\pi, t) = \tilde{g}_1(t), \quad v(2\pi, t) = \tilde{g}_2(t), \quad t \in (0, T), \tag{4.1c}$$

where $f_1(r, t) \doteq r^{\frac{n-1}{2}} f(r, t) + \frac{n^2-4n+3}{4} r^{-2} \cdot v$. Also, by letting $w(r, t) = v(r + \pi, t)$ and $f_2(r, t) = f_1(r + \pi, t)$, we change the problem (4.1) to the interval $(0, \pi)$. In fact, the ibvp for w is

$$iw_t + w_{rr} = f_2(r, t), \quad 0 < r < \pi, \quad t \in (0, T), \tag{4.2a}$$

$$w(r, 0) = w_0(r) \in H^s(0, \pi), \quad 0 < r < \pi, \tag{4.2b}$$

$$w(0, t) = \tilde{g}_1(t) \in H^{\frac{s+1}{2}}(0, T), \quad w(\pi, t) = \tilde{g}_2(t) \in H^{\frac{s+1}{2}}(0, T), \quad t \in (0, T). \tag{4.2c}$$

Also, from the compatibility condition (1.8c), we have the following compatibility conditions

$$\tilde{g}_1(0) = w_0(0) \quad \text{and} \quad \tilde{g}_2(0) = w_0(\pi), \quad \frac{1}{2} < s < 2. \tag{4.3}$$

Using this, for $\frac{1}{2} < s < 2$, we can assume that $w_0(0) = w_0(\pi) = \tilde{g}_1(0) = \tilde{g}_2(0) = 0$.

Next, we solve the ibvp (4.2) by decomposing it into simpler problems. In fact, using superposition principle, the linear ibvp (4.2) can be expressed as the homogeneous ibvp

$$iw_t + w_{rr} = 0, \quad 0 < r < \pi, \quad t \in (0, T), \tag{4.4a}$$

$$w(r, 0) = w_0(r), \quad 0 < r < \pi, \tag{4.4b}$$

$$w(0, t) = \tilde{g}_1(t), \quad w(\pi, t) = \tilde{g}_2(t), \quad t \in (0, T), \tag{4.4c}$$

and the forced linear ibvp with zero initial and boundary data

$$iw_t + w_{rr} = f_2(r, t), \quad 0 < r < \pi, \quad t \in (0, T), \quad (4.5a)$$

$$w(r, 0) = 0, \quad 0 < r < \pi, \quad (4.5b)$$

$$w(0, t) = 0, \quad w(\pi, t) = 0, \quad t \in (0, T). \quad (4.5c)$$

In addition, we decompose the ibvp (4.4) as an ibvp with homogeneous boundary data

$$iv_t + v_{rr} = 0, \quad 0 < r < \pi, \quad t \in (0, T), \quad (4.6a)$$

$$v(r, 0) = \omega_0(r), \quad 0 < r < \pi, \quad (4.6b)$$

$$v(0, t) = 0, \quad v(\pi, t) = 0, \quad t \in (0, T), \quad (4.6c)$$

and an ibvp with zero initial data:

$$iw_t + w_{rr} = 0, \quad 0 < r < \pi, \quad t \in (0, T), \quad (4.7a)$$

$$w(r, 0) = 0, \quad 0 < r < \pi, \quad (4.7b)$$

$$w(0, t) = h_1(t), \quad w(\pi, t) = h_2(t), \quad t \in (0, T), \quad (4.7c)$$

where $h_1 = \tilde{g}_1(t)$ and $h_2 = \tilde{g}_2(t)$.

Solving the ibvp (4.6). We will solve this problem by reflection. In fact, in order to solve this problem, it suffices to solve the following periodic problem

$$iV_t + V_{rr} = 0, \quad -\pi < r < \pi, \quad t \in (0, T), \quad (4.8a)$$

$$V(r, 0) = V_0(r), \quad -\pi < r < \pi, \quad (4.8b)$$

$$V(-\pi, t) = V(\pi, t), \quad V_r(-\pi, t) = V_r(\pi, t), \quad t \in (0, T), \quad (4.8c)$$

where $V_0(r)$ is the odd extension of $w_0(r)$, i.e.,

$$V_0(r) = \begin{cases} w_0(r), & 0 < r < \pi, \\ 0, & r = 0, \\ -w_0(-r), & -\pi < r < 0. \end{cases} \quad (4.9)$$

By using the standard separation of variables, the solution to the periodic ivp (4.8) is

$$V = W_0(t)w_0 \doteq \sum_{n=1}^{\infty} B_n \sin(nr)e^{-in^2t}, \quad (4.10)$$

where

$$B_n = \frac{2}{\pi} \int_0^\pi w_0(r) \sin(nr) dr, \quad n \in \mathbb{Z}. \tag{4.11}$$

The solution formula $W_0(t)w_0$ can also be written in the complex form

$$W_0(t)w_0 = \frac{1}{2i} \sum_{n=-\infty}^\infty \tilde{B}_n e^{inr} e^{-in^2t}, \quad \text{with} \quad \tilde{B}_n = \begin{cases} B_n, & \text{if } n \geq 1, \\ 0, & \text{if } n = 0, \\ -B_{-n} & \text{if } n \leq -1. \end{cases} \tag{4.12}$$

Furthermore, for w_0 , we define $H^s(0, \pi)$ -norm by its Fourier coefficients as follows

$$\|w_0\|_{H^s(0,\pi)}^2 \doteq \sum_{n=1}^\infty (1+n)^{2s} b_n^2, \tag{4.13}$$

where the Fourier coefficient $b_n = B_n$ is defined as follows

$$b_n \doteq \frac{2}{\pi} \int_0^\pi w_0(r) \sin(nr) dr = B_n, \quad n \geq 1. \tag{4.14}$$

Finally, we state the following result for the solution $W_0(t)w_0$ given by (4.10).

PROPOSITION 4.1. *Let $w_0 \in H^s(0, \pi)$, where $0 \leq s \leq 2$ and $s \neq \frac{1}{2}, \frac{3}{2}$. Then, the solution defined by (4.10) satisfies the following estimates*

$$\sup_{t \in [0, T]} \|W_0(t)w_0\|_{H^s(0,\pi)} \leq C_{T,s} \|w_0\|_{H^s(0,\pi)}. \tag{4.15}$$

In addition, for $s \in \mathbb{N}$, we have

$$\|\psi \partial_r^s W_0(t)w_0\|_{L^4((0,\pi) \times \mathbb{R})} \leq C_T \|w_0\|_{H^s(0,\pi)}. \tag{4.16}$$

Proof of proposition 4.1. By the solution formula (4.10), we see that the Fourier series for $V(x, t)$ is $B_n e^{-in^2t}$, where B_n is given by (4.11). Thus, by the definition (4.13), the space estimate (4.15) is obtained. Next, we prove estimate (4.16). As the proof for $s = 1, 2, 3, \dots$ resembles that of $s = 0$, we focus solely on establishing estimate (4.16) for the case of $s = 0$. It follows from the next result. \square

LEMMA 4.2. *Let $(x, t) \in \mathbb{T} \times \mathbb{R}$ and let $(n, \lambda) \in \mathbb{Z} \times \mathbb{R}$ be the dual variables. Then there is a constant $c > 0$ such that*

$$\|f\|_{L^4(\mathbb{T} \times \mathbb{R})} \leq c \|(1 + |\lambda + n^2|)^{\frac{3}{8}} \widehat{f}\|_{L^2(\mathbb{Z} \times \mathbb{R})}, \tag{4.17}$$

for any test function f on $\mathbb{T} \times \mathbb{R}$.

The above result was proved in [8]. Now, we use it to complete the proof of estimate (4.16). Noticing that the solution formula (4.10) defines an odd function

of r over $(-\pi, \pi)$, we keep the same notation and we shall prove that $\|\psi V\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|w_0\|_{H^s(0, \pi)}$. To do this, letting $f = \psi V$, we have

$$\|f\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} \widehat{f}\|_{L^2(\mathbb{Z} \times \mathbb{R})}.$$

Furthermore, using complex formula (4.12) and taking Fourier transform, it is deduced that $\widehat{f}(n, \lambda) \simeq \widehat{\psi}(\lambda + n^2) \widetilde{B}_n$, which implies that

$$\begin{aligned} \|(1 + |\lambda + n^2|)^{\frac{3}{8}} \widehat{f}\|_{L^2(\mathbb{Z} \times \mathbb{R})}^2 &\lesssim \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{\frac{3}{4}} |\widehat{\psi}(\lambda + n^2)|^2 d\lambda \cdot \widetilde{B}_n^2 \\ &\lesssim \sum_{n=1}^{\infty} B_n^2 \lesssim \|w_0\|_{L^2(0, \pi)}^2. \end{aligned}$$

Combining the above estimates yield the estimate (4.16) for $s = 0$. This completes the proof of proposition 4.1.

Solving the ibvp (4.5). For this problem, we extend f_2 oddly and use the fact that the solution to this ibvp is given by

$$w(r, t) = -i \int_0^t W_0(t - \tau) f_2(\cdot, \tau) d\tau = -i \sum_{n=1}^{\infty} \sin(nr) \int_0^t e^{-in^2(t-t')} \widehat{f}_2^r(n, t') dt', \tag{4.18}$$

where W_0 is given by (4.10) and $\widehat{f}_2^r(n, t')$ is defined as

$$\widehat{f}_2^r(n, t') = \frac{2}{\pi} \int_0^\pi f_2(r, t') \sin(nr) dr, \quad n \in \mathbb{Z}.$$

We define the odd extension of $f_2(r, t)$ with respect to r

$$F_2(r, t) \doteq \begin{cases} f_2(r, t), & 0 < r < \pi, \\ 0, & r = 0, \\ -f_2(-r, t), & -\pi < r < 0. \end{cases} \tag{4.19}$$

Then, we have $\widehat{F}_2^r(n, t') = \int_{-\pi}^\pi F_2(r, t) e^{inr} dr = -\int_{-\pi}^0 f_2(-r, t) e^{inr} dr + \int_0^\pi f_2(r, t) e^{inr} dr = i\pi \widehat{f}_2^r(n, t')$. Using the fact that F_2 is an odd function, we get $\widehat{F}_2^r(n, t') = -\widehat{F}_2^r(-n, t')$. Thus, working similarly as in the formula (4.12), we have

$$w(r, t) \simeq \int_0^t W_0(t - \tau) f_2(\cdot, \tau) d\tau \simeq \sum_{n \in \mathbb{Z}} e^{inr} \int_0^t e^{-in^2(t-t')} \widehat{F}_2^r(n, t') dt'. \tag{4.20}$$

Also, it is derived that

$$\|f_2\|_{L^{4/3}((0, \pi) \times (0, T))} \simeq \|F_2\|_{L^{4/3}(\mathbb{T} \times (0, T))}.$$

Furthermore, for $t \notin (0, T)$ we set $F_2 = 0$ (keeping the same notation) and we obtain

$$\|f_2\|_{L^{4/3}((0,\pi) \times (0,T))} \simeq \|F_2\|_{L^{4/3}(\mathbb{T} \times \mathbb{R})}. \tag{4.21}$$

For the solution formula (4.18), we have the following result.

PROPOSITION 4.3. *Let $f_2 \in L^1(0, T; H^s(0, \pi))$, where $0 \leq s \leq 2$ and $s \neq \frac{1}{2}, \frac{3}{2}$. Then, considering the solution defined by (4.18), the following estimates hold*

$$\sup_{t \in [0, T]} \left\| \int_0^t W_0(t-t') f_2(\cdot, t') dt' \right\|_{H^s(0, \pi)} \leq C_{T,s} \|f_2\|_{L^1(0, T; H^s(0, \pi))}. \tag{4.22}$$

In addition, for $s \in \mathbb{N}$, if $\partial_r^s f_2 \in L^{4/3}((0, \pi) \times (0, T))$, the following estimate holds:

$$\left\| \psi \partial_r^s \int_0^t W_0(t-t') f_2(\cdot, t') dt' \right\|_{L^4((0, \pi) \times \mathbb{R})} \leq C_T \|\partial_r^s f_2\|_{L^{4/3}((0, \pi) \times (0, T))}. \tag{4.23}$$

Proof of proposition 4.3. Using the space estimate (4.15) for the homogeneous IVP (4.8), i.e.,

$$\sup_{t \in [0, T]} \|W_0(t) w_0\|_{H^s(0, \pi)} \leq C_{T,s} \|w_0\|_{H^s(0, \pi)},$$

we have

$$\begin{aligned} \left\| \int_0^t W_0(t-t') f_2(\cdot, t') dt' \right\|_{H^s(0, \pi)} &\leq \int_{t'=0}^t \left\| W_0(t-t') f_2(\cdot, t') \right\|_{H^s(0, \pi)} dt' \\ &\lesssim \int_{t'=0}^t \|f_2(t')\|_{H^s(0, \pi)} dt' \leq \int_{t'=0}^T \|f_2(t')\|_{H^s(0, \pi)} dt', \end{aligned}$$

which is the desired space estimate (4.22). □

Proof of estimate (4.23) with $s = 0$. For this estimate, similar to the proof of estimate (4.16), we will use the formula (4.20). Also, we notice that it defines an odd function of r over $(-\pi, \pi)$. Hence, we prove the following estimate,

$$\left\| \psi \int_0^t W_0(t-t') f_2(\cdot, t') dt' \right\|_{L^4(\mathbb{T} \times \mathbb{R})} \leq C_T \|f_2\|_{L^{4/3}((0, \pi) \times (0, T))}. \tag{4.25}$$

To do this, using space-time Fourier transform, we express F_2 in the phase space (n, λ) . More precisely, substituting in (4.20)

$$\widehat{F}_2^r(n, t') = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\lambda t'} \widehat{F}_2(n, \lambda) d\lambda,$$

we obtain

$$\psi(t) w(r, t) \simeq \psi(t) \sum_{n \in \mathbb{Z}} e^{inr} \int_0^t e^{-in^2(t-t')} \int_{\lambda \in \mathbb{R}} e^{i\lambda t'} \widehat{F}_2(n, \lambda) d\lambda dt'.$$

Performing the t' integration first, it is

$$\int_0^t e^{i(\lambda+n^2)t'} dt' = -i \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2}.$$

Then, the solution $\psi(t)w(r, t)$ becomes

$$\psi(t)w(r, t) \simeq \psi(t) \sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} e^{inr} e^{-in^2t} \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda. \tag{4.27}$$

Finally, adding and subtracting $\psi(\lambda + n^2)$ inside the integral (localizing near the singularity $\lambda = -n^2$) gives the following decomposition of $\psi(t)w(r, t)$

$$\psi(t)w(r, t) \simeq -\psi(t) \sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} e^{inr} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda \tag{4.28}$$

$$+ \psi(t) \sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} e^{inr} e^{-in^2t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda \tag{4.29}$$

$$- \psi(t) \sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} e^{inr} e^{-in^2t} \frac{\psi(\lambda + n^2)(e^{i(\lambda+n^2)t} - 1)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda. \tag{4.30}$$

Estimate for (4.28). Let

$$f(r, t) = \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} e^{inr} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda.$$

Since $\psi \in C_0^\infty(\mathbb{R})$, we have

$$\|(4.28)\|_{L^4(\mathbb{T} \times \mathbb{R})} \simeq \|\psi(\cdot)f(\cdot, \cdot)\|_{L^4(\mathbb{T} \times \mathbb{R})} \leq \|\psi\|_{L_t^\infty} \|f\|_{L^4(\mathbb{T} \times \mathbb{R})}.$$

Also, computing the Fourier transform of f with respect to r gives

$$\widehat{f}^r(n, t) \simeq \int_{\mathbb{R}} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda.$$

Therefore,

$$\widehat{f}(n, \lambda) = \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda).$$

Now, we need the following result, which is provided in [8].

PROPOSITION 4.4. *If the multiplier $M = M(n, \lambda)$ satisfies*

$$|M(n, \lambda)| \lesssim (1 + |\lambda + n^2|)^{-\frac{3}{4}}, \quad \text{for all } n \in \mathbb{Z} \quad \text{and } \lambda \in \mathbb{R}, \tag{4.31}$$

then M acts boundedly from $L^{4/3}(\mathbb{Z} \times \mathbb{R})$ to $L^4(\mathbb{Z} \times \mathbb{R})$, that is

$$\left\| \sum_{n=1}^{\infty} e^{inr} \int_{\mathbb{R}} M(n, \lambda) \widehat{f}(n, \lambda) e^{i\lambda t} d\lambda \right\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|f\|_{L^{4/3}(\mathbb{T} \times \mathbb{R})}. \tag{4.32}$$

Proposition 4.4 is due to lemma 4.2 and duality (see corollary 4.5). Now, from this, we finish the proof of estimate (4.25) for (4.28). In fact, by the definition of $\psi(t)$, the integrand is non-zero only for $|\lambda + n^2| \geq \frac{1}{2}$. Therefore, $1 + |\lambda + n^2| \leq 2|\lambda + n^2| + |\lambda + n^2| = 3|\lambda + n^2|$ and so

$$\|f\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|F_2\|_{L^{4/3}(\mathbb{T} \times \mathbb{R})},$$

which, together with the inequality (4.21), gives us the desired estimate (4.25) for term (4.28).

Estimate for (4.29). To estimate $\|(4.29)\|_{L^4(\mathbb{T} \times \mathbb{R})}$, we first find the Fourier transform of the function

$$f(r, t) = (4.29) = \psi(t) \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} e^{i(nr+n^2t)} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda.$$

It is clearly seen that $\widehat{f}^r(n, t) \simeq \psi(t) e^{in^2t} C_n$, where

$$C_n = \int_{\mathbb{R}} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda.$$

Then, computing the Fourier transform of \widehat{f}^r with respect to t gives

$$\widehat{f}(n, \lambda) = C_n \widehat{\psi}(\lambda + n^2).$$

Applying estimate (4.17), it is obtained that

$$\begin{aligned} \|(4.29)\|_{L^4(\mathbb{T} \times \mathbb{R})} &\lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} C_n \widehat{\psi}(\lambda + n^2)\|_{L^2(\mathbb{Z} \times \mathbb{R})} \\ &= \left(\sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} (1 + |\lambda + n^2|)^{3/4} |C_n \widehat{\psi}(\lambda + n^2)|^2 d\lambda \right)^{\frac{1}{2}} \\ &\lesssim \left(\sum_{n \in \mathbb{Z}} |C_n|^2 \right)^{\frac{1}{2}} = \left(\sum_{n \in \mathbb{Z}} \left| \int_{\mathbb{R}} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda \right|^2 \right)^{\frac{1}{2}}. \end{aligned} \tag{4.33}$$

Furthermore, for the $d\lambda$ -integral in the above estimate, applying the Cauchy–Schwarz inequality gives

$$\begin{aligned} \left| \int_{\mathbb{R}} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \widehat{F}_2(n, \lambda) d\lambda \right|^2 &\leq \int_{\mathbb{R}} \left| \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \right|^{5/4} d\lambda \\ &\quad \cdot \int_{\mathbb{R}} \left| \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \right|^{3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda \\ &\lesssim \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{-3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda. \end{aligned} \tag{4.34}$$

Now, we need the following dual estimate of lemma 4.2 (see [8]).

COROLLARY 4.5. *For any test function f , we have*

$$\|(1 + |\lambda + n^2|)^{-\frac{3}{8}} \widehat{f}\|_{L^2(\mathbb{Z} \times \mathbb{R})} \leq c \|f\|_{L^{4/3}(\mathbb{T} \times \mathbb{R})}. \tag{4.35}$$

Finally, combining estimates (4.33) and (4.34) with corollary 4.5 and using estimate (4.21), it is deduced that

$$\begin{aligned} \|(4.29)\|_{L^4(\mathbb{T} \times \mathbb{R})} &\lesssim \left(\sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{-3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda \right)^{\frac{1}{2}} \\ &= \|(1 + |\lambda + n^2|)^{-\frac{3}{8}} \widehat{F}_2\|_{L^2(\mathbb{Z} \times \mathbb{R})} \\ &\lesssim \|F_2\|_{L^{4/3}(\mathbb{T} \times \mathbb{R})} \simeq \|f_2\|_{L^{4/3}(\mathbb{T} \times (0, T))}. \end{aligned}$$

This completes the estimate for (4.29).

Estimate for (4.30). From Taylor’s series at $\lambda + n^2 = 0$, we can expand

$$e^{i(\lambda+n^2)t} - 1 = \sum_{k=1}^{\infty} \frac{t^k (\lambda + n^2)^k}{k!}.$$

Thus, (4.30) $\simeq \sum_{k=1}^{\infty} \frac{i^{k+1}}{k!} f_k$, where

$$f_k(r, t) \doteq t^k \psi(t) \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} e^{inr} e^{-in^2t} \psi(\lambda + n^2) (\lambda + n^2)^{k-1} \widehat{F}_2(n, \lambda) d\lambda.$$

The Fourier transform of the function f_k yields

$$\widehat{f}_k(n, t) = C_k(n) t^k \psi(t) e^{-in^2t},$$

where

$$C_k(n) = \int_{\mathbb{R}} \psi(\lambda + n^2) (\lambda + n^2)^{k-1} \widehat{F}_2(n, \lambda) d\lambda.$$

Therefore,

$$\widehat{f}_k(n, \lambda) = C_k(n)\widehat{t^k\psi}(\lambda + n^2).$$

Let us now estimate (4.30) using the expression of $\widehat{f}_k(n, \lambda)$.

$$\|(4.30)\|_{L^4(\mathbb{T}\times\mathbb{R})} \lesssim \left\| \sum_{k=1}^{\infty} \frac{i^{k+1}}{k!} f_k \right\|_{L^4(\mathbb{T}\times\mathbb{R})} \lesssim \sum_{k=1}^{\infty} \frac{1}{k!} \|f_k\|_{L^4(\mathbb{T}\times\mathbb{R})}. \tag{4.36}$$

Applying estimate (4.17), we obtain

$$\begin{aligned} \|f_k\|_{L^4(\mathbb{T}\times\mathbb{R})} &\lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} C_k(n)\widehat{t^k\psi}(\lambda + n^2)\|_{L^2(\mathbb{Z}\times\mathbb{R})} \\ &= \left(\sum_{n\in\mathbb{Z}} \int_{\lambda\in\mathbb{R}} (1 + |\lambda + n^2|)^{3/4} |C_k(n)\widehat{t^k\psi}(\lambda + n^2)|^2 d\lambda \right)^{\frac{1}{2}} \\ &\lesssim \left(\sum_{n\in\mathbb{Z}} |C_k(n)|^2 \right)^{\frac{1}{2}} = \left(\sum_{n\in\mathbb{Z}} \left| \int_{\mathbb{R}} \psi(\lambda + n^2)(\lambda + n^2)^{k-1} \widehat{F}_2(n, \lambda) d\lambda \right|^2 \right)^{\frac{1}{2}}. \end{aligned} \tag{4.37}$$

Furthermore, for the $d\lambda$ -integral in the above estimate, the Cauchy–Schwartz inequality gives

$$\begin{aligned} &\left| \int_{\mathbb{R}} \psi(\lambda + n^2)(\lambda + n^2)^{k-1} \widehat{F}_2(n, \lambda) d\lambda \right|^2 \\ &\leq \int_{\mathbb{R}} |\psi(\lambda + n^2)(\lambda + n^2)^{k-1}|^2 (1 + |\lambda + n^2|)^{3/4} d\lambda \cdot \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{-3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda \\ &\lesssim \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{-3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda. \end{aligned} \tag{4.38}$$

Finally, combining estimates (4.36), (4.37), and (4.38) with corollary 4.5 and using estimate (4.21), it is obtained that

$$\begin{aligned} \|(4.30)\|_{L^4(\mathbb{T}\times\mathbb{R})} &\lesssim \sum_{k=1}^{\infty} \frac{1}{k!} \left(\sum_{n\in\mathbb{Z}} \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{-3/4} |\widehat{F}_2(n, \lambda)|^2 d\lambda \right)^{\frac{1}{2}} \\ &\lesssim \|(1 + |\lambda + n^2|)^{-\frac{3}{8}} \widehat{F}_2\|_{L^2(\mathbb{Z}\times\mathbb{R})} \\ &\lesssim \|F_2\|_{L^{4/3}(\mathbb{T}\times\mathbb{R})} \simeq \|f_2\|_{L^{4/3}(\mathbb{T}\times(0,T))}. \end{aligned}$$

We complete the proof of estimate (4.23) with $s = 0$.

Proof of estimate (4.23) with $s \in \mathbb{N}^+$. Consider the periodic function $F_2(r, t)$. Employing integration by parts, we can express the Fourier transform of the spatial derivative $\partial_r^s F_2$ as

$$\widehat{\partial_r^s F_2}^r(n, t') \simeq n^s \widehat{F_2}^r(n, t').$$

Utilizing this result and the complex formula (4.20), we derive

$$\begin{aligned} \partial_r^s \int_0^t W_0(t - \tau) f_2(\cdot, \tau) d\tau &\simeq \sum_{n \in \mathbb{Z}} e^{inr} n^s \int_0^t e^{-in^2(t-t')} \widehat{F_2}^r(n, t') dt' \\ &\simeq \sum_{n \in \mathbb{Z}} e^{inr} \int_0^t e^{-in^2(t-t')} \widehat{\partial_r^s F_2}^r(n, t') dt'. \end{aligned}$$

Consequently, by applying estimate (4.23) with $s = 0$ for $\partial_r^s F_2$, we obtain the desired estimate (4.23) with $s \in \mathbb{N}^+$. The proof of proposition 4.3 is completed. \square

Solving the ibvp (4.7). For this problem, the solution is given by the following formula, which is provided in [7]

$$u(r, t) = 2\pi i \sum_{n=1}^{\infty} n \sin(nr) \int_0^t e^{-in^2(t-t')} [h_1(t') - (-1)^n h_2(t')] dt' \tag{4.39}$$

$$= \int_0^t W_0(t - t') q(\cdot, t') dt', \tag{4.39}$$

where

$$q(x, t) \doteq 2\pi i \left[\sum_{n=1}^{\infty} n \sin(nx) h_1(t) - \sum_{n=1}^{\infty} (-1)^n n \sin(nx) h_2(t) \right]. \tag{4.40}$$

Next, we define the following boundary operator

$$W_b h \doteq 2i \sum_{n=1}^{\infty} n \sin(nr) \int_0^t e^{-in^2(t-t')} h(t') dt', \tag{4.41}$$

which has the form

$$W_b h = \sum_{n \in \mathbb{Z}} n e^{inr} \int_0^t e^{-in^2(t-t')} h(t') dt', \quad r \in \mathbb{T}, t \in \mathbb{R}. \tag{4.42}$$

The following result can be proved.

PROPOSITION 4.6. *Let $h \in H_{00}^{1/2}(0, T)$. Then, the solution defined by (4.41) satisfies the following estimates*

$$\sup_{t \in [0, T]} \left\| \psi W_b h \right\|_{L^2(\mathbb{T})} \lesssim \|h\|_{H^{1/2}(0, T)}, \tag{4.43}$$

$$\left\| \psi W_b h \right\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|h\|_{H^{1/2}(0,T)}. \tag{4.44}$$

In addition, for $0 \leq s \leq 2$, if $h \in H_0^{\frac{s+1}{2}}(0,T)$ (for s an even integer $h \in H_0^{\frac{s+1}{2}}(0,T)$), then we have

$$\sup_{t \in [0,T]} \left\| \psi \partial_r^s W_b h \right\|_{L^2(\mathbb{T})} + \left\| \psi \partial_r^s W_b h \right\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|h\|_{H^{(s+1)/2}(0,T)}. \tag{4.45}$$

Proof of proposition 4.6. The proof of this proposition is similar to the proof of proposition 4.3 and was also given in [7] (see propositions 4.7 and 4.8). Here, we only provide the proof for estimates (4.43) and (4.44), and we give a different proof by modifying and simplifying the proof in [7]. We start with writing $h(t') = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\lambda t'} \widehat{h}(\lambda) d\lambda$, which, together with formula (4.42), implies

$$\psi \cdot W_b h = \frac{\psi(t)}{2\pi} \sum_{n=-\infty}^{\infty} n e^{inr} \int_0^t e^{-in^2(t-t')} \int_{\mathbb{R}} e^{i\lambda t'} \widehat{h}(\lambda) d\lambda dt'. \tag{4.46}$$

Performing the t' integration first gives

$$\int_0^t e^{i(\lambda+n^2)t'} dt' = -i \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2}.$$

Then, the above solution $\psi(t)w(r,t)$ becomes

$$\psi(t)W_b h(r,t) = -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} e^{inr} e^{-in^2 t} \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda = I^+ + I^-, \tag{4.47}$$

where I^+ is the integral over $(0, \infty)$ and I^- is the integral over $(-\infty, 0)$. More precisely, we have

$$I^+ \doteq -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_0^{\infty} e^{inr} e^{-in^2 t} \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda, \tag{4.48}$$

$$I^- \doteq -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{-in^2 t} \frac{e^{i(\lambda+n^2)t} - 1}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda. \tag{4.49}$$

To estimate I^+ , we split it as $I^+ = I_1^+ - I_2^+$, where

$$I_1^+ \doteq -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_0^{\infty} e^{inr} \frac{e^{i\lambda t}}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda \quad \text{and}$$

$$I_2^+ \doteq -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_0^{\infty} e^{inr} \frac{e^{in^2 t}}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda.$$

Estimate for I_1^+ . Let

$$f(r, t) = \sum_{n \in \mathbb{Z}} e^{inr} \int_0^\infty e^{i\lambda t} \frac{1}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda.$$

First, we prove L^2 estimate (4.43) for I_1^+ . Since ψ is compactly supported in $(0, 1)$, we have

$$\sup_{t \in [0, T]} \|I_1^+\|_{L^2(\mathbb{T})} \leq \sup_{t \in [0, 1]} \|f\|_{L^2(\mathbb{T})}.$$

Taking the Fourier transform with respect to r yields

$$\widehat{f}^r(n, t) \simeq \int_0^\infty e^{i\lambda t} \frac{1}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda, \tag{4.50}$$

which, by Plancherel theorem, implies that

$$\|f\|_{L^2(\mathbb{T})}^2 \lesssim \sum_{n \in \mathbb{Z}} \left| \int_0^\infty e^{i\lambda t} \frac{1}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda \right|^2 \leq \sum_{n \in \mathbb{Z}} \left[\int_0^\infty \left| \frac{1}{\lambda + n^2} \cdot n\widehat{h}(\lambda) \right| d\lambda \right]^2. \tag{4.51}$$

Now, applying Cauchy–Schwarz inequality in $d\lambda$, for $\varepsilon > 0$ and small, it is derived that

$$\begin{aligned} \left[\int_0^\infty \left| \frac{1}{\lambda + n^2} \cdot n\widehat{h}(\lambda) \right| d\lambda \right]^2 &\leq \int_0^\infty \frac{n^2(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda \cdot \int_0^\infty \frac{1}{(1 + \lambda)^{1+2\varepsilon}} d\lambda \\ &\lesssim \int_0^\infty \frac{n^2(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Thus, combining above estimates, we get

$$\begin{aligned} \|f\|_{L^2(\mathbb{T})}^2 &\lesssim \sum_{n \in \mathbb{Z}} \int_0^\infty \frac{n^2(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda \\ &\leq \int_0^\infty \left[\sum_{n \in \mathbb{Z}} \frac{n^2(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^2} \right] (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Furthermore, using $\lambda + n^2 \geq \max\{\lambda, n^2\}$, for all $\lambda \geq 0$ and for $2 - 2\varepsilon > \frac{3}{2}$ or $\varepsilon < \frac{1}{4}$, we obtain

$$\sum_{n \in \mathbb{Z}} \frac{n^2(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^2} \leq \sum_{n \in \mathbb{Z}} \frac{n^2}{(\lambda + n^2)^{2-2\varepsilon}} \cdot \frac{(1 + \lambda)^{2\varepsilon}}{(\lambda + n^2)^{2\varepsilon}} \leq \sum_{n \in \mathbb{Z}} \frac{n^2}{(\lambda + n^2)^{\frac{3}{2} +}} \lesssim \sum_{n=1}^\infty \frac{1}{n^{1+}} \lesssim 1.$$

Therefore, the desired L^2 estimate (4.43) for I_1^+ is reached.

L^4 estimate (4.44) for I_1^+ . Fubini's theorem implies

$$\begin{aligned} I_1^+(r, t) &= -\frac{i\psi(t)}{2\pi} \int_0^\infty \sum_{n \in \mathbb{Z}} \frac{ne^{inr}}{\lambda + n^2} \cdot e^{i\lambda t} \widehat{h}(\lambda) d\lambda = -\frac{i\psi(t)}{2\pi} \cdot 2i \int_0^\infty \sum_{n=1}^\infty \frac{n \sin(nr)}{\lambda + n^2} \\ &\quad \cdot e^{i\lambda t} \widehat{h}(\lambda) d\lambda \\ &= \frac{\psi(t)}{\pi} \int_0^\infty \sum_{n=1}^\infty \frac{n \sin(nr)}{\lambda + n^2} \cdot e^{i\lambda t} \widehat{h}(\lambda) d\lambda. \end{aligned}$$

Since $\lambda \geq 0$, we see that $\lambda + n^2$ is away from 0 and $\lambda + n^2 \geq \max\{\lambda, n^2\}$. Thus, by Cauchy–Schwarz inequality, we have

$$|I_1^+(r, t)|^2 \lesssim \psi(t) \sum_{n \in \mathbb{Z}} \int_0^\infty \frac{n^2}{|\lambda + n^2|^2(1 + \lambda)} d\lambda \cdot \int_0^\infty (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda.$$

Then, for any $0 < \varepsilon < \frac{1}{2}$,

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \int_0^\infty \frac{n^2}{|\lambda + n^2|^2(1 + \lambda)} d\lambda &\lesssim \sum_{n \in \mathbb{Z}} \int_0^\infty \frac{n^2}{|\lambda + n^2|^{2-\varepsilon} |n^2 + \lambda|^\varepsilon (1 + \lambda)} d\lambda \lesssim \sum_{n=1}^\infty \frac{1}{|n^2|^{1-\varepsilon}} \\ &\quad \cdot \int_0^\infty \frac{1}{(1 + \lambda)^{1+\varepsilon}} d\lambda \lesssim 1. \end{aligned}$$

Hence, the above estimate yields

$$|I_1^+(r, t)| \leq \psi(t) \|h\|_{H^{1/2}}, \quad -\pi < r < \pi, \quad t \in \mathbb{R},$$

which gives

$$\|I_1^+\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|h\|_{H^{1/2}}.$$

Estimate for I_2^+ . The L^2 estimate(4.43) for I_2^+ is similar to the L^2 estimate for I_1^+ and hence we omit it here. For the L^4 estimate (4.44), taking Fourier transform with respect to r , we have

$$\widehat{I_2^+}^r(n, t) \simeq \psi(t) e^{in^2 t} C_n,$$

where

$$C_n = \int_0^\infty \frac{n}{\lambda + n^2} \widehat{h}(\lambda) d\lambda.$$

Then, the Fourier transform of \widehat{f}^r with respect to t gives

$$\widehat{f}(n, \lambda) = C_n \widehat{\psi}(\lambda + n^2).$$

Applying (4.17) yields

$$\begin{aligned} \|I_2^+\|_{L^4(\mathbb{T} \times \mathbb{R})} &\lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} C_n \widehat{\psi}(\lambda + n^2)\|_{L^2(\mathbb{Z} \times \mathbb{R})} \\ &= \left(\sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} (1 + |\lambda + n^2|)^{3/4} |C_n \widehat{\psi}(\lambda + n^2)|^2 d\lambda \right)^{\frac{1}{2}} \\ &\lesssim \left(\sum_{n \in \mathbb{Z}} |C_n|^2 \right)^{\frac{1}{2}} = \left(\sum_{n \in \mathbb{Z}} \left| \int_0^\infty \frac{n}{\lambda + n^2} \widehat{h}(\lambda) d\lambda \right|^2 \right)^{\frac{1}{2}}, \end{aligned} \tag{4.52}$$

which reduces to the estimate (4.51). We complete the estimates for I_2^+ .

Estimate for I^- . Adding and subtracting $\psi(\lambda + n^2)$ inside the integral (localizing near the singularity $\lambda = -n^2$) gives the following decomposition of I^- ,

$$I^- = -\frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda \tag{4.53}$$

$$+ \frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{-in^2 t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda \tag{4.54}$$

$$- \frac{i\psi(t)}{2\pi} \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{-in^2 t} \frac{\psi(\lambda + n^2)(e^{i(\lambda+n^2)t} - 1)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda. \tag{4.55}$$

Estimates for (4.53). First, we prove (4.43) for this term. Let

$$f(r, t) = \sum_{n \in \mathbb{Z}} e^{inr} \int_{-\infty}^0 e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda.$$

Since ψ is compactly supported in $(0, 1)$, we have $\sup_{t \in [0, T]} \|(4.53)\|_{L^2(\mathbb{T})} \leq \sup_{t \in [0, 1]} \|f\|_{L^2(\mathbb{T})}$. Taking the Fourier transform of f with respect to r gives

$$\widehat{f}^r(n, t) \simeq \int_{-\infty}^0 e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda, \tag{4.56}$$

which implies that

$$\|f\|_{L^2(\mathbb{T})}^2 \lesssim \sum_{n \in \mathbb{Z}} \left| \int_{-\infty}^0 e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) d\lambda \right|^2 \tag{4.57}$$

$$\leq \sum_{n \in \mathbb{Z}} \left[\int_{-\infty}^0 \left| \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n\widehat{h}(\lambda) \right| d\lambda \right]^2. \tag{4.57}$$

Also, making the change of variables $\lambda = -\mu^2$ and using $\frac{n}{n^2-\mu^2} = \frac{1}{2} \left[\frac{1}{n-\mu} + \frac{1}{n+\mu} \right]$, it is deduced that

$$\|f\|_{L^2(\mathbb{T})}^2 \lesssim \sum_{n \in \mathbb{Z}} \left[\int_0^\infty |\mu \widehat{h}(-\mu^2)| \left| \frac{1}{n-\mu} + \frac{1}{n+\mu} \right| [1 - \psi(n^2 - \mu^2)] d\lambda \right]^2.$$

Now we need the following estimate, whose proof is provided in [7],

$$\sum_{n \in \mathbb{Z}} \left| \int_0^\infty \widehat{F}(\mu) \frac{1}{n-\mu} (1 - \psi(n^2 - \mu^2)) d\mu \right|^2 \lesssim \int_0^\infty (1 + \mu) |\widehat{F}(\mu)|^2 d\mu.$$

In fact, applying the above estimate twice with $\widehat{F}(\mu) = |\mu \widehat{h}(-\mu^2)|$ implies

$$\|f\|_{L^2(\mathbb{T})}^2 \lesssim \int_0^\infty (1 + \mu) |\mu \widehat{h}(\mu^2)|^2 d\mu \lesssim \int_{\mathbb{R}} (1 + |\lambda|^{1/2}) |\lambda|^{1/2} |\widehat{h}(\lambda)|^2 d\lambda \lesssim \|h\|_{H^{1/2}}^2.$$

L^4 estimate (4.44) for (4.53). To show this estimate, we will split the $d\lambda$ -integral at $\lambda = -\frac{n^2}{2}$. More precisely, we have

$$(4.53) = -\frac{i\psi(t)}{2\pi} f_1 - \frac{i\psi(t)}{2\pi} f_2,$$

where

$$\begin{aligned} f_1 &= \sum_{n \in \mathbb{Z}} \int_{-\infty}^{-\frac{n^2}{2}} e^{inr} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda, \\ f_2 &= \sum_{n \in \mathbb{Z}} \int_{-\frac{n^2}{2}}^0 e^{inr} e^{i\lambda t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda. \end{aligned}$$

For $\frac{i\psi(t)}{2\pi} f_1$, since ψ is compactly supported in $(0, 1)$, we have $\|\frac{i\psi(t)}{2\pi} f_1\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|f_1\|_{L^4(\mathbb{T} \times \mathbb{R})}$. Also, we have $|n|^2 \lesssim |\lambda|$. Thus, applying estimate (4.17) gives

$$\begin{aligned} \|f_1\|_{L^4(\mathbb{T} \times \mathbb{R})}^2 &\lesssim \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} (1 + |\lambda + n^2|)^{\frac{3}{4}} \chi_{\lambda < -\frac{n^2}{2}}(\lambda) \left| \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) \right|^2 d\lambda \\ &\lesssim \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} \frac{1}{(1 + |\lambda + n^2|)^{5/4}} \cdot |\lambda| |\widehat{h}(\lambda)|^2 d\lambda \\ &\lesssim \int_{\mathbb{R}} \sum_{n \in \mathbb{Z}} \frac{1}{(1 + |\lambda + n^2|)^{5/4}} \cdot |\lambda| |\widehat{h}(\lambda)|^2 d\lambda \\ &\lesssim \int_{\mathbb{R}} |\lambda| |\widehat{h}(\lambda)|^2 d\lambda \lesssim \|h\|_{H^{1/2}}. \end{aligned}$$

For $\frac{i\psi(t)}{2\pi} f_2$, again since $\psi(t)$ is compactly supported in $(0, 1)$, it suffices to show that

$$\|f_2\|_{L^\infty_{\mathbb{T}} \times \mathbb{R}} \lesssim \|h\|_{H^{1/2}}. \tag{4.58}$$

By Cauchy–Schwarz inequality,

$$|f_2| \leq \sum_{n \in \mathbb{Z}} \int_{-\frac{n^2}{2}}^0 n^2 \frac{1 - \psi(\lambda + n^2)}{|\lambda + n^2|^2(1 + |\lambda|)} d\lambda \cdot \int_{-\frac{n^2}{2}}^0 (1 + |\lambda|) |\widehat{h}(\lambda)|^2 d\lambda.$$

For the first integral in the above estimate, choosing $0 < \varepsilon < \frac{1}{2}$, we have

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \int_{-\frac{n^2}{2}}^0 n^2 \frac{1 - \psi(\lambda + n^2)}{|\lambda + n^2|^2(1 + |\lambda|)} d\lambda &\lesssim \sum_{n \in \mathbb{Z}} \int_{-\frac{n^2}{2}}^0 \frac{n^2}{(1 + |\lambda + n^2|)^{2-\varepsilon}} \\ &\quad \cdot \frac{1}{(1 + |\lambda|)(1 + |\lambda + n^2|)^\varepsilon} d\lambda \\ &\lesssim \sum_{n \in \mathbb{Z}} \int_{-\frac{n^2}{2}}^0 \frac{n^2}{(1 + n^2)^{2-\varepsilon}} \cdot \frac{1}{(1 + |\lambda|)^{1+\varepsilon}} d\lambda \\ &\leq \sum_{n \in \mathbb{Z}} \frac{n^2}{(1 + n^2)^{2-\varepsilon}} \cdot \int_{\mathbb{R}} \frac{1}{(1 + |\lambda|)^{1+\varepsilon}} d\lambda \lesssim 1. \end{aligned}$$

Combining the above estimates yields the desired estimate (4.58).

Estimates for (4.54). We let

$$f(r, t) \doteq \psi(t) \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{-in^2t} \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda.$$

Taking Fourier transform with respect to r , it is obtained that

$$\widehat{f}^r(n, t) \simeq \psi(t) e^{-in^2t} C_n,$$

where

$$C_n = \int_{-\infty}^0 \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda.$$

Hence, by Plancherel’s theorem, we have

$$\begin{aligned} \|(4.54)\|_{L^2(\mathbb{T})} &\simeq \|f\|_{L^2(\mathbb{T})} \simeq \left(\sum_{n \in \mathbb{Z}} |C_n|^2 \right)^{\frac{1}{2}} \\ &= \left(\sum_{n \in \mathbb{Z}} \left| \int_{-\infty}^0 \frac{1 - \psi(\lambda + n^2)}{\lambda + n^2} \cdot n \widehat{h}(\lambda) d\lambda \right|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

which is reduced to the estimate (4.57). Concerning L^4 estimate of (4.54), computing the Fourier transform of \widehat{f}^r with respect to t gives

$$\widehat{f}(n, \lambda) = C_n \widehat{\psi}(\lambda + n^2).$$

Estimate (4.17) gives $\|I_2^+\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} C_n \widehat{\psi}(\lambda + n^2)\|_{L^2(\mathbb{Z} \times \mathbb{R})}$. This implies

$$\|I_2^+\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \left(\sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} (1 + |\lambda + n^2|)^{3/4} |C_n \widehat{\psi}(\lambda + n^2)|^2 d\lambda \right)^{\frac{1}{2}} \lesssim \left(\sum_{n \in \mathbb{Z}} |C_n|^2 \right)^{\frac{1}{2}}.$$

Again, we arrive at estimate (4.57). This completes L^4 estimate of (4.54).

Estimates for (4.55). Using Taylor’s series at $\lambda + n^2 = 0$, we obtain

$$e^{i(\lambda+n^2)t} - 1 = \sum_{k=1}^{\infty} \frac{(it)^k (\lambda + n^2)^k}{k!}.$$

Thus, (4.55) $\simeq \sum_{k=1}^{\infty} \frac{i^{k+1}}{k!} f_k$, where

$$f_k(r, t) \doteq t^k \psi(t) \sum_{n \in \mathbb{Z}} \int_{-\infty}^0 e^{inr} e^{-in^2 t} \psi(\lambda + n^2) (\lambda + n^2)^{k-1} n \widehat{h}(\lambda) d\lambda.$$

For the L^2 estimate of (4.55), we have

$$\|(4.55)\|_{L^2(\mathbb{T})} \lesssim \left\| \sum_{k=1}^{\infty} \frac{i^{k+1}}{k!} f_k \right\|_{L^2(\mathbb{T})} \lesssim \sum_{k=1}^{\infty} \frac{1}{k!} \|f_k\|_{L^2(\mathbb{T})}.$$

Taking the Fourier transform of the function f_k , it is deduced that

$$\widehat{f}_k^r(n, t) = C_k(n) t^k \psi(t) e^{-in^2 t},$$

where

$$C_k(n) = \int_{-\infty}^0 \psi(\lambda + n^2) (\lambda + n^2)^{k-1} n \widehat{h}(\lambda) d\lambda.$$

Since ψ is compactly supported in $(0, 1)$, by the Plancherel theorem and Cauchy–Schwarz inequality, it is obtained that

$$\begin{aligned} \|f_k(t)\|_{L^2(\mathbb{T})}^2 &\lesssim \sum_{n \in \mathbb{Z}} |C_k(n)|^2 = \left| \int_{-\infty}^0 \psi(\lambda + n^2) (\lambda + n^2)^{k-1} n \widehat{h}(\lambda) d\lambda \right|^2 \\ &\lesssim \int_{-\infty}^0 \psi(\lambda + n^2) (\lambda + n^2)^{2(k-1)} d\lambda \cdot \int_{-\infty}^0 \psi(\lambda + n^2) n^2 |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Since $\psi \in C_0^\infty(0, 1)$, the first integral is bounded. Also, for the second integral, $|\lambda + n^2| < 1$, which implies that $n^2 \lesssim (1 + |\lambda|)$. Hence,

$$\|f_k(t)\|_{L^2(\mathbb{T})}^2 \lesssim \int_{-\infty}^0 (1 + |\lambda|) |\widehat{h}(\lambda)|^2 d\lambda \lesssim \|h\|_{H^{1/2}}^2.$$

For the L^4 estimate of (4.55), we have

$$\|(4.55)\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \left\| \sum_{k=1}^{\infty} \frac{i^{k+1}}{k!} f_k \right\|_{L^4(\mathbb{T} \times \mathbb{R})} \lesssim \sum_{k=1}^{\infty} \frac{1}{k!} \|f_k\|_{L^4(\mathbb{T} \times \mathbb{R})}.$$

To bound the L^4 norm of f_k , taking the full Fourier transform, we obtain

$$\widehat{f}_k(n, \lambda) = C_k(n) \widehat{t^k \psi}(\lambda + n^2).$$

Applying estimate (4.17) gives

$$\begin{aligned} \|f_k\|_{L^4(\mathbb{T} \times \mathbb{R})} &\lesssim \|(1 + |\lambda + n^2|)^{\frac{3}{8}} C_k(n) \widehat{t^k \psi}(\lambda + n^2)\|_{L^2(\mathbb{Z} \times \mathbb{R})} \\ &= \left(\sum_{n \in \mathbb{Z}} \int_{\lambda \in \mathbb{R}} (1 + |\lambda + n^2|)^{3/4} |C_k(n) \widehat{t^k \psi}(\lambda + n^2)|^2 d\lambda \right)^{\frac{1}{2}} \\ &\lesssim \left(\sum_{n \in \mathbb{Z}} |C_k(n)|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

which is the estimate needed. This completes the proof of proposition 4.6. □

Now, we can derive linear estimate for the solution to the linear ibvp (4.2). In fact, for $0 < r < \pi$ and $0 < t < T$, this solution is given by

$$w(r, t) = K[w_0, \tilde{g}_1, \tilde{g}_2; f_2] \doteq W_0(t)w_0 - i \int_0^t W_0(t - \tau) f_2(\cdot, \tau) d\tau \tag{4.59}$$

$$+ \int_0^t W_0(t - t') q(\cdot, t') dt', \tag{4.59}$$

where W_0 is defined by (4.12) and q is given by (4.40) with $h_1(t) = \tilde{g}_1(t)$, $h_2(t) = \tilde{g}_2(t)$. Combining propositions 4.1, 4.3, and 4.6, the following linear estimate holds for solution formula $K[w_0, \tilde{g}_1, \tilde{g}_2; f_2]$.

THEOREM 4.7 (1) *Let $s \in \mathbb{N}$. If $w_0 \in H^s(0, \pi)$, $\tilde{g}_1, \tilde{g}_2 \in H_0^{\frac{s+1}{2}}(0, T)$ (for even s , $\tilde{g}_1, \tilde{g}_2 \in H_0^{\frac{s+1}{2}}(0, T)$), and $\partial_r^s f_2 \in L^{4/3}((0, \pi) \times (0, T))$, then $K[w_0, \tilde{g}_1, \tilde{g}_2; f_2]$ defines a solution to the ibvp (4.2), which satisfies*

$$\begin{aligned} &\sup_{t \in [0, T]} \|K[w_0, \tilde{g}_1, \tilde{g}_2; f_2](t)\|_{H^s(0, \pi)} + \|\partial_r^s K[w_0, \tilde{g}_1, \tilde{g}_2; f_2]\|_{L^4((0, \pi) \times (0, T))} \\ &\lesssim \|w_0\|_{H^s(0, \pi)} + \|\tilde{g}_1\|_{H^{\frac{s+1}{2}}(0, T)} + \|\tilde{g}_2\|_{H^{\frac{s+1}{2}}(0, T)} + \|\partial_r^s f_2\|_{L^{4/3}((0, \pi) \times (0, T))}. \end{aligned} \tag{4.60}$$

(2) Suppose $\frac{1}{2} < s < 2$ and $s \neq \frac{3}{2}$. If $w_0 \in H^s(0, \pi)$, $\tilde{g}_1, \tilde{g}_2 \in H^{\frac{s+1}{2}}(0, T)$ and $f_2 \in L^1(0, T; H^s(0, \pi))$ then $K[w_0, \tilde{g}_1, \tilde{g}_2; f_2]$ defines a solution to the ibvp (4.2) with compatibility condition (4.3), satisfying

$$\sup_{t \in [0, T]} \|K[w_0, \tilde{g}_1, \tilde{g}_2; f_2](t)\|_{H^s(0, \pi)} \lesssim \|\omega_0\|_{H^s(1, \infty)} + \|\tilde{g}_1\|_{H^{\frac{s+1}{2}}(0, T)} + \|\tilde{g}_2\|_{H^{\frac{s+1}{2}}(0, T)} + \|f_2\|_{L^1(0, T; H^s(0, \pi))}.$$

4.2. Proof of well-posedness for ibvp in domain Ω_2 , i.e., theorem 1.3

Since the ibvp in Ω_2 is reduced to the ibvp (2.1) for $r \in (\pi, 2\pi)$ with $u(\pi, t) = g_1(t)$ and $u(2\pi, t) = g_2(t)$, it suffices to prove the well-posedness of ibvp (4.2) with forcing f_2 giving by

$$f_2(r, t) = f_1(r + \pi, t) = -\lambda(r + \pi)^{-\frac{(n-1)(p-2)}{2}} |w|^{p-2} w + \frac{n^2 - 4n + 3}{4} (r + \pi)^{-2} \cdot w. \tag{4.61}$$

Furthermore, since both multipliers $-(r + \pi)^{-\frac{(n-1)(p-2)}{2}}$ and $(r + \pi)^{-2}$ are smooth and bounded for $0 < r < \pi$, similar to the proof of theorem 1.2, we can bound them by using the L^∞ -norm. Therefore, the well-posedness proof is similar to the proof of theorem 4.10 and proposition 4.11 in [7]. Hence, the details are omitted here.

5. ibvp in a ball centred at origin

In this section, we study the ibvp of (2.1) for $r \in (0, 1)$. For the sake of convenience and simplicity, we only discuss the case with $n = 2$. The cases for $n \geq 3$ can be studied similarly.

5.1. Linear problem

The forced linear ibvp within the region Ω_0 can be described by the following equations

$$iu_t + \Delta u = f(x_1, x_2, t), \quad (x_1, x_2, t) \in \Omega_0, \quad t \in (0, T), \tag{5.1a}$$

$$u(x_1, x_2, 0) = u_0(x_1, x_2), \quad (x_1, x_2) \in \Omega_0, \tag{5.1b}$$

$$u(x_1, x_2, t) = g(t), \quad x_1^2 + x_2^2 = 1, \quad t \in (0, T). \tag{5.1c}$$

We also decompose the ibvp (5.1) into the following two separate problems.

Problem 1. ibvp with homogeneous boundary condition

$$iu_t + \Delta u = f(x_1, x_2, t), \quad (x_1, x_2, t) \in \Omega_0, \quad t \in (0, T), \tag{5.2a}$$

$$u(x_1, x_2, 0) = u_0(x_1, x_2), \quad (x_1, x_2) \in \Omega_0, \tag{5.2b}$$

$$u(x_1, x_2, t) = 0, \quad x_1^2 + x_2^2 = 1, \quad t \in (0, T). \tag{5.2c}$$

Problem 2. ibvp with non-homogeneous boundary condition

$$iu_t + \Delta u = 0, \quad (x_1, x_2, t) \in \Omega_0, \quad t \in (0, T), \tag{5.3a}$$

$$u(x_1, x_2, 0) = 0, \quad (x_1, x_2) \in \Omega_0, \tag{5.3b}$$

$$u(x_1, x_2, t) = g(t), \quad x_1^2 + x_2^2 = 1, \quad t \in (0, T). \tag{5.3c}$$

In the subsequent sections, we will investigate these two problems separately.

Estimate for Problem 1. In order to analyse this problem, we will employ the semigroup method. Following the methodology outlined in [11] (refer to section 2.1), we introduce the operator A acting on functions in $L^2(\Omega_0)$, defined as follows:

$$\begin{cases} D(A) = \{u \in H_0^1(\Omega_0), \Delta u \in L^2(\Omega_0)\}, \\ A(u) = \Delta u \quad \text{for } u \in D(A). \end{cases} \tag{5.4}$$

It is worth noting that $D(A) = H^2(\Omega_0) \cap H_0^1(\Omega_0)$. Additionally, we observe that A is a self-adjoint operator and $A \leq 0$, as indicated by the following

$$\langle Au, v \rangle = \langle \Delta u, v \rangle = - \langle \nabla u, \nabla v \rangle = \langle u, \Delta v \rangle = \langle u, A^*v \rangle \implies A = A^*,$$

and

$$\langle Au, u \rangle = - \langle \nabla u, \nabla u \rangle \leq 0 \implies A \leq 0.$$

Moreover, let $(\mathcal{J}(t))_{t \in \mathbb{R}}$ represent the group of isometries generated by iA within any of the following spaces: $D(A)$, $H_0^1(\Omega_0)$, $L^2(\Omega_0)$, $H^{-1}(\Omega_0)$, or $(D(A))^*$. By utilizing the property that iA is skew-adjoint, i.e., $(iA)^* = -iA$, we deduce the following relation

$$\mathcal{J}(t)^* = \mathcal{J}(-t). \tag{5.5}$$

Furthermore, the solution for the forced linear ibvp (5.2) is defined by

$$u(x, t) = S_J[u_0; f] \doteq \mathcal{J}(t)u_0(x) + i \int_0^t \mathcal{J}(t-s)f(x, s)ds, \quad x \in \Omega_0, \quad t \in [0, T]. \tag{5.6}$$

With these foundations in place, we can now proceed to establish the following results.

PROPOSITION 5.1. If $u_0 \in L^2(\Omega_0)$ and $f \in L^1_t(0, T; L^2_x(\Omega_0))$, then we have $S_J[u_0; f] \in C([0, T]; L^2_x(\Omega_0))$ and it satisfies the following estimate

$$\sup_{t \in [0, T]} \|S_J[u_0; f](\cdot, t)\|_{L^2_x(\Omega_0)} \lesssim \|u_0\|_{L^2(\Omega_0)} + \|f(\cdot, t)\|_{L^1_t(0, T; L^2_x(\Omega_0))}. \tag{5.7}$$

In addition, if $u_0 \in H^2_0(\Omega_0)$ and $f \in L^1_t(0, T; H^2_0(\Omega_0))$, then we have $S_J[u_0; f] \in C([0, T]; H^2_0(\Omega_0))$ and it satisfies the following estimate

$$\sup_{t \in [0, T]} \|S_J[u_0; f](\cdot, t)\|_{H^2(\Omega_0)} \lesssim \|u_0\|_{H^2(\Omega_0)} + \|f(\cdot, t)\|_{L^1_t(0, T; H^2(\Omega_0))}. \tag{5.8}$$

Finally, for $1 < s < 2$, if $u_0 \in H^s_0(\Omega_0)$ and $f \in L^1_t(0, T; H^s_0(\Omega_0))$, then we have $S_J[u_0; f] \in C([0, T]; H^s_0(\Omega_0))$ and it satisfies the following estimate

$$\sup_{t \in [0, T]} \|S_J[u_0; f](\cdot, t)\|_{H^s(\Omega_0)} \lesssim \|u_0\|_{H^s(\Omega_0)} + \|f(\cdot, t)\|_{L^1_t(0, T; H^s(\Omega_0))}. \tag{5.9}$$

Proof 5.1. Proof of proposition 5.1. The proof of estimate (5.9) can be deduced through interpolation and the estimates (5.7) and (5.8). Additionally, the proof of estimate (5.8) closely parallels the proof of estimate (5.7). Therefore, in this exposition, we shall exclusively present the proof for estimate (5.7). Using the relation (5.5), we obtain

$$\begin{aligned} \langle \mathcal{J}(t)u_0(x), \mathcal{J}(t)u_0(x) \rangle_{L^2(\Omega_0)} &= \langle u_0(x), \mathcal{J}(t)^* \mathcal{J}(t)u_0(x) \rangle_{L^2(\Omega_0)} \\ &= \langle u_0(x), \mathcal{J}(-t)\mathcal{J}(t)u_0(x) \rangle_{L^2(\Omega_0)} \\ &= \langle u_0(x), u_0(x) \rangle_{L^2(\Omega_0)}. \end{aligned}$$

This gives us that

$$\sup_{t \in [0, T]} \|\mathcal{J}(t)u_0(\cdot)\|_{L^2(\Omega_0)} = \|u_0\|_{L^2(\Omega_0)}. \tag{5.10}$$

Using the identity (5.10), we derive

$$\begin{aligned} \sup_{t \in [0, T]} \left\| \int_0^t \mathcal{J}(t-s)f(x, s)ds \right\| &\leq \sup_{t \in [0, T]} \int_0^t \|\mathcal{J}(t-s)f(\cdot, s)\|_{L^2(\Omega_0)} ds \\ &\leq \sup_{t \in [0, T]} \int_0^t \|f(\cdot, s)\|_{L^2(\Omega_0)} ds \leq \|f(\cdot, t)\|_{L^1_t(0, T; L^2(\Omega_0))}. \end{aligned} \tag{5.11}$$

Combining estimates (5.10) with (5.11), we establish the inequality (5.7). This concludes the proof of proposition 5.1 \square

Estimate for Problem 2. We initiate our analysis by finding the solution to the ibvp (5.3). Utilizing the expression $\Delta u = u''(r) + \frac{1}{r}u'(r)$, we reformulate this ibvp as follows

$$iu_t + u_{rr} + \frac{1}{r}u_r = 0, \quad 0 < r < 1, \quad t \in (0, T), \tag{5.12a}$$

$$u(r, 0) = 0, \quad 0 < r < 1, \tag{5.12b}$$

$$u(1, t) = h(t), \quad t \in \mathbb{R}. \tag{5.12c}$$

Here, $h(t)$ extends the boundary data $g(t)$ from the interval $(0, T)$ to the entire real line \mathbb{R} . By making use of lemma 3.3, we assume that h is compactly supported in the interval $(0, 2)$. Additionally, at $r = 0$, the equation in (5.12) implies the following boundary condition

$$u_r(0, t) = 0. \tag{5.13}$$

We will solve the **reduced pure ibvp** (5.12) and commence with solving the following ibvp

$$iU_t + U_{rr} + \frac{1}{r}U_r = 0, \quad 0 < r < 1, \quad t \in (0, T), \tag{5.14a}$$

$$U(r, 0) = U_0(r), \quad 0 < r < 1, \tag{5.14b}$$

$$U_r(0, t) = U(1, t) = 0, \quad t \in (0, T). \tag{5.14c}$$

By employing the standard method of separation of variables, we can find the general solution for (5.14) as

$$U(r, t) = S_{ball}(t)U_0(r) \doteq \sum_{n=0}^{\infty} \beta_n J_0(\lambda_n r) e^{-i\lambda_n^2 t}, \tag{5.15}$$

where $J_0(z)$ represents the Bessel function of order 0, and $\lambda_n, n = 0, 1, 2, \dots$, are the positive zeros of the Bessel function J_0 , meaning $J_0(\lambda_n) = 0$. The coefficients $\beta_n, n = 0, 1, \dots$, are determined by the initial condition $U_0(r)$ and can be calculated as follows

$$\sum_{n=0}^{\infty} \beta_n J_0(\lambda_n r) = U_0(r),$$

which gives us that

$$\beta_n = \frac{2}{J_1^2(\lambda_n)} \int_0^1 r U_0(r) J_0(\lambda_n r) dr. \tag{5.16}$$

REMARK 5.2. The Fourier–Bessel series for a function $f(x)$ on the interval $(0, 1)$ with respect to the Bessel function J_0 can be expressed as:

$$f(x) = \sum_{n=0}^{\infty} a_n J_0(\lambda_n x),$$

where the coefficients a_n are given by

$$a_n = \frac{2}{J_1^2(\lambda_n)} \int_0^1 x f(x) J_0(\lambda_n x) dx, \quad (5.17)$$

and $\lambda_n, n = 0, 1, 2, \dots$, are the positive zeros of the Bessel function J_0 , i.e., $J_0(\lambda_n) = 0$.

REMARK 5.3. The Bessel functions $J_0(\lambda_n x), n = 0, 1, 2, \dots$, are orthogonal over the interval $(0, 1)$ with respect to the weight function x , meaning:

$$\int_0^1 x J_0(\lambda_m x) J_0(\lambda_n x) dx = \frac{1}{2} \delta_{mn} [J_1(\lambda_n)]^2, \quad (5.18)$$

where δ_{mn} is the Kronecker delta, defined as:

$$\delta_{mn} = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{otherwise.} \end{cases}$$

This allows us to express the coefficient a_n in terms of inner products of $f(x)$ with the Bessel function $J_0(\lambda_n x)$.

Solving forced ibvp. Now, let us address the following forced problem

$$iV_t + V_{rr} + \frac{1}{r}V_r = F(r, t), \quad 0 < r < 1, \quad t \in (0, T), \quad (5.19a)$$

$$V(r, 0) = 0, \quad 0 < r < 1, \quad (5.19b)$$

$$V_r(0, t) = V(1, t) = 0, \quad t \in (0, T). \quad (5.19c)$$

Utilizing Duhamel's principle, the solution to the forced ibvp (5.19) is given by

$$V(r, t) = -i \int_0^t S_{ball}(t-t') F(r, t') dt' \doteq -i \sum_{n=0}^{\infty} J_0(\lambda_n r) \int_0^t e^{-i\lambda_n^2(t-t')} B_n(t') dt', \quad (5.20)$$

where

$$B_n(t') = \frac{2}{J_1^2(\lambda_n)} \int_0^1 r F(r, t') J_0(\lambda_n r) dr. \quad (5.21)$$

Solving ibvp (5.12). Defining $v(r, t) = u(r, t) - h(t)$, we obtain the ibvp for v as follows

$$iv_t + v_{rr} + \frac{1}{r}v_r = F_h(r, t), \quad 0 < r < 1, \quad t \in (0, T), \tag{5.22a}$$

$$v(r, 0) = 0, \quad 0 < r < 1, \tag{5.22b}$$

$$v_r(0, t) = v(1, t) = 0, \quad t \in (0, T), \tag{5.22c}$$

where $F_h = -ih'(t)$. Employing the solution formula (5.20) with $F = F_h$, we have

$$v = - \sum_{n=0}^{\infty} J_0(\lambda_n r) b_n \int_0^t h'(t') e^{-i\lambda_n^2(t-t')} dt', \tag{5.23}$$

where b_n represents the Fourier–Bessel coefficient of 1, and can be computed as

$$b_n = \frac{2}{J_1^2(\lambda_n)} \int_0^1 r \cdot J_0(\lambda_n r) dr = \frac{2}{J_1^2(\lambda_n)} \cdot \frac{J_1(\lambda_n)}{\lambda_n} = \frac{2}{J_1(\lambda_n)\lambda_n}. \tag{5.24}$$

Further, by performing integration by parts and utilizing the initial condition $h(0) = 0$, we obtain

$$v = -h(t) - i \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{2\lambda_n}{J_1(\lambda_n)} \int_0^t h(t') e^{-i\lambda_n^2(t-t')} dt'. \tag{5.25}$$

Therefore, the solution to the ibvp (5.12), denoted as $u = v + h$, is expressed as

$$u = W_{ball}h \doteq -2i \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{\lambda_n}{J_1(\lambda_n)} \int_0^t h(t') e^{-i\lambda_n^2(t-t')} dt'. \tag{5.26}$$

In addition, if we switch back to the variables (x_1, x_2) , then we have

$$u(x_1, x_2, t) = W_{ball}h \doteq -2i \sum_{n=0}^{\infty} J_0\left(\lambda_n \sqrt{x_1^2 + x_2^2}\right) \frac{\lambda_n}{J_1(\lambda_n)} \int_0^t h(t') e^{-i\lambda_n^2(t-t')} dt'. \tag{5.27}$$

Next, we will prove the following result.

PROPOSITION 5.4. *Let $h \in H_{00}^{1/2}(0, T)$. Then, the solution defined by (5.26) satisfies the following estimate*

$$\sup_{t \in [0, T]} \left\| r^{1/2} \cdot W_{ball}h \right\|_{L^2(0,1)} \lesssim \|h\|_{H^{1/2}(0, T)}. \tag{5.28}$$

Furthermore, if $h \in H_0^{(s+1)/2}(0, T)$ for $0 \leq s \leq 2$ (for $s = 0, 2$, $h \in H_{00}^{\frac{s+1}{2}}(0, T)$), then the solution defined by (5.27) satisfies the following estimates

$$\sup_{t \in [0, T]} \left\| W_{ball}h \right\|_{H^s(\Omega_0)} \lesssim \|h\|_{H^{(s+1)/2}(0, T)}. \tag{5.29}$$

To show the above result, we need **Parseval’s identity for Fourier–Bessel series**, which is stated as follows.

LEMMA 5.5. *Let $x^{1/2}f(x) \in L^2(0, 1)$. If the Fourier–Bessel series of $f(x)$ is given by*

$$f(x) = \sum_{n=0}^{\infty} c_n J_0(\lambda_n x),$$

where $c_n = \frac{2}{J_1^2(\lambda_n)} \int_0^1 x f(x) J_0(\lambda_n x) dx$, then we have

$$\int_0^1 x |f(x)|^2 dx = \frac{1}{2} \sum_{n=0}^{\infty} |c_n|^2 [J_1(\lambda_n)]^2. \tag{5.30}$$

Proof 5.1. Proof of lemma 5.5. By straightforward computation, we get

$$\int_0^1 x |f(x)|^2 dx = \int_0^1 x f(x) \overline{f(x)} dx = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} c_n c_m \int_0^1 x J_0(\lambda_n x) J_0(\lambda_m x) dx.$$

Now, applying the orthogonal identity (5.18), i.e.,

$$\int_0^1 x J_0(\lambda_m x) J_0(\lambda_n x) dx = \frac{1}{2} \delta_{mn} [J_1(\lambda_n)]^2, \tag{5.31}$$

we obtain the desired identity (5.30). This completes the proof of lemma 5.5. \square

Proof 5.1. Proof of proposition 5.4. We will begin by providing the proof of estimate (5.28), which is similar to the proof of estimate (4.43). We start by expressing $h(t')$ as follows

$$h(t') = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i\lambda t'} \widehat{h}(\lambda) d\lambda.$$

Using this expression and formula (5.26), we can write

$$\psi \cdot W_{ball} h = \frac{-i\psi(t)}{\pi} \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{\lambda_n}{J_1(\lambda_n)} \int_0^t e^{-i\lambda_n^2(t-t')} \int_{\mathbb{R}} e^{i\lambda t'} \widehat{h}(\lambda) d\lambda dt'. \tag{5.32}$$

Now, let us perform the integration with respect to t' using the computation

$$\int_0^t e^{i(\lambda + \lambda_n^2)t'} dt' = -i \frac{e^{i(\lambda + \lambda_n^2)t} - 1}{\lambda + \lambda_n^2}.$$

With this, the expression for $\psi(t)W_{\text{ball}}h$ becomes

$$\psi(t)W_{\text{ball}}h = \frac{i\psi(t)}{\pi} \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{\lambda \in \mathbb{R}} e^{-i\lambda_n^2 t} \frac{e^{i(\lambda+\lambda_n^2)t} - 1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda \tag{5.33}$$

$$\simeq I^+ + I^-, \tag{5.33}$$

where I^+ is the integral over $(0, \infty)$ and I^- is the integral over $(-\infty, 0)$. More precisely, we have

$$I^+ \doteq \psi(t) \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_0^{\infty} e^{-i\lambda_n^2 t} \frac{e^{i(\lambda+\lambda_n^2)t} - 1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda, \tag{5.34}$$

$$I^- \doteq \psi(t) \sum_0^{\infty} J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 e^{-i\lambda_n^2 t} \frac{e^{i(\lambda+\lambda_n^2)t} - 1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda. \tag{5.35}$$

□

Estimate of weighted L^2 -norm for I^+ . Let

$$f(r, t) = r^{1/2} \cdot \sum_{n=0}^{\infty} J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_0^{\infty} e^{-i\lambda_n^2 t} \frac{e^{i(\lambda+\lambda_n^2)t} - 1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda.$$

First, we prove L^2 estimate (5.28) for I_1^+ . Since ψ is compactly supported in $(0, 1)$, we have $\sup_{t \in [0, T]} \|r^{1/2} \cdot I_1^+\|_{L^2(0,1)} \leq \sup_{t \in [0, 1]} \|f\|_{L^2(0,1)}$. Using Parseval’s identity for Fourier–Bessel series (identity (5.30)), we get

$$\|f\|_{L^2(0,1)}^2 \lesssim \sum_{n \in \mathbb{Z}} \left| \int_0^{\infty} e^{-i\lambda_n^2 t} \frac{e^{i(\lambda+\lambda_n^2)t} - 1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda \right|^2 \tag{5.36}$$

$$\leq \sum_{n \in \mathbb{Z}} \left[\int_0^{\infty} \left| \frac{1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) \right| d\lambda \right]^2. \tag{5.36}$$

Now, applying Cauchy–Schwarz inequality in $d\lambda$, for $\varepsilon > 0$ (small), we obtain

$$\begin{aligned} \left[\int_0^{\infty} \left| \frac{1}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) \right| d\lambda \right]^2 &\leq \int_0^{\infty} \frac{\lambda_n^2 (1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda \cdot \int_0^{\infty} \frac{1}{(1 + \lambda)^{1+2\varepsilon}} d\lambda \\ &\lesssim \int_0^{\infty} \frac{\lambda_n^2 (1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Thus, combining above estimates, it is deduced that

$$\begin{aligned} \|f\|_{L^2(0,1)}^2 &\lesssim \sum_{n \in \mathbb{Z}} \int_0^{\infty} \frac{\lambda_n^2 (1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^2} (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda \\ &\leq \int_0^{\infty} \left[\sum_{n \in \mathbb{Z}} \frac{\lambda_n^2 (1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^2} \right] (1 + \lambda) |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Furthermore, using $\lambda + \lambda_n^2 \geq \max\{\lambda, \lambda_n^2\}$ and $\lambda_n \approx (n - \frac{1}{4})\pi \simeq n$, for all $\lambda \geq 0$ and $2 - 2\varepsilon > \frac{3}{2}$ or $\varepsilon < \frac{1}{4}$, we get

$$\sum_{n \in \mathbb{Z}} \frac{\lambda_n^2 (1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^2} \leq \sum_{n \in \mathbb{Z}} \frac{\lambda_n^2}{(\lambda + \lambda_n^2)^{2-2\varepsilon}} \cdot \frac{(1 + \lambda)^{2\varepsilon}}{(\lambda + \lambda_n^2)^{2\varepsilon}} \lesssim \sum_{n \in \mathbb{Z}} \frac{n^2}{(\lambda + n^2)^{\frac{3}{2} +}} \lesssim \sum_{n=1}^{\infty} \frac{1}{n^{1+}} \lesssim 1.$$

Therefore, the desired estimate (5.28) for I^+ is obtained.

Estimate of weighted L^2 -norm for I^- . Adding and subtracting $\psi(\lambda + \lambda_n^2)$ inside the integral (localizing near the singularity $\lambda = -\lambda_n^2$) gives the following decomposition of I^-

$$I^- = \psi(t) \sum_0^\infty J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 [e^{-i\lambda_n^2 t} (e^{i(\lambda + \lambda_n^2)t} - 1)] \frac{1 - \psi(\lambda + \lambda_n^2)}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda \tag{5.37}$$

$$+ \psi(t) \sum_0^\infty J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 e^{-i\lambda_n^2 t} \frac{(e^{i(\lambda + \lambda_n^2)t} - 1)\psi(\lambda + \lambda_n^2)}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda. \tag{5.38}$$

Estimate of weighted L^2 -norm for (5.37). Let us start with the first part of I^- , as given by (5.37). We define a function $f(r, t)$ as follows

$$f(r, t) = r^{1/2} \cdot \sum_0^\infty J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 [e^{-i\lambda_n^2 t} (e^{i(\lambda + \lambda_n^2)t} - 1)] \frac{1 - \psi(\lambda + \lambda_n^2)}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda.$$

Since ψ is compactly supported in $(0, 1)$, we have $\sup_{t \in [0, T]} \|r^{1/2} \cdot (5.37)\|_{L^2(0,1)} \leq \sup_{t \in [0,1]} \|f\|_{L^2(0,1)}$. Using Parseval's identity (5.30), we deduce that

$$\begin{aligned} \|f\|_{L^2(0,1)}^2 &\lesssim \sum_{n \in \mathbb{Z}} \left| \int_{-\infty}^0 [e^{-i\lambda_n^2 t} (e^{i(\lambda + \lambda_n^2)t} - 1)] \frac{1 - \psi(\lambda + \lambda_n^2)}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) d\lambda \right|^2 \\ &\lesssim \sum_{n \in \mathbb{Z}} \left[\int_{-\infty}^0 \left| \frac{1 - \psi(\lambda + \lambda_n^2)}{\lambda + \lambda_n^2} \cdot \lambda_n \widehat{h}(\lambda) \right| d\lambda \right]^2. \end{aligned} \tag{5.39}$$

Making the change of variables $\lambda = -\mu^2$ and using the identity $\frac{\lambda_n}{\lambda_n^2 - \mu^2} = \frac{1}{2} \left(\frac{1}{\lambda_n - \mu} + \frac{1}{\lambda_n + \mu} \right)$, we get

$$\|f\|_{L^2(0,1)}^2 \lesssim \sum_{n \in \mathbb{Z}} \left[\int_0^\infty |\mu \widehat{h}(-\mu^2)| \left| \frac{1}{\lambda_n - \mu} + \frac{1}{\lambda_n + \mu} \right| [1 - \psi(\lambda_n^2 - \mu^2)] d\lambda \right]^2.$$

Now we need the following estimate (similar to the proof of Lemma A-1 in [7])

$$\sum_{n \in \mathbb{Z}} \left| \int_0^\infty \widehat{F}(\mu) \frac{1}{\lambda_n - \mu} (1 - \psi(\lambda_n^2 - \mu^2)) d\mu \right|^2 \lesssim \int_0^\infty (1 + \mu) |\widehat{F}(\mu)|^2 d\mu.$$

Applying this estimate twice with $\widehat{F}(\mu) = |\mu\widehat{h}(-\mu^2)|$, we obtain

$$\|f\|_{L^2(\mathbb{T})}^2 \lesssim \int_0^\infty (1 + \mu)|\mu\widehat{h}(\mu^2)|^2 d\mu \lesssim \int_{\mathbb{R}} (1 + |\lambda|^{1/2})|\lambda|^{1/2}|\widehat{h}(\lambda)|^2 d\lambda \lesssim \|h\|_{H^{1/2}}^2.$$

This completes the proof of desired estimate (5.28) for (5.37).

Estimate of weighted L^2 -norm for (5.38). Now, we move on to the second part of I^- , as given by (5.38). We will first consider a decomposition of the term. Using Taylor’s series at $\lambda + \lambda_n^2 = 0$, we have

$$e^{i(\lambda + \lambda_n^2)t} - 1 = \sum_{k=1}^\infty \frac{(it)^k (\lambda + \lambda_n^2)^k}{k!}.$$

Thus, we can write (5.38) $\simeq \sum_{k=1}^\infty \frac{i^k}{k!} f_k$, where

$$f_k(r, t) \doteq t^k \psi(t) \sum_0^\infty J_0(\lambda_n r) \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 e^{-i\lambda_n^2 t} \psi(\lambda + \lambda_n^2) (\lambda + \lambda_n^2)^{k-1} \lambda_n \widehat{h}(\lambda) d\lambda.$$

We can now estimate the weighted L^2 -norm for this term, starting with the following

$$\|r^{1/2} \cdot (5.38)\|_{L^2(0,1)} \lesssim \|r^{1/2} \cdot \sum_{k=1}^\infty \frac{i^k}{k!} f_k\|_{L^2(0,1)} \lesssim \sum_{k=1}^\infty \frac{1}{k!} \|r^{1/2} \cdot f_k\|_{L^2(0,1)}.$$

The Fourier–Bessel series coefficients for f_k are given by $t^k \psi(t) e^{-i\lambda_n^2 t} C_k(n)$, where

$$C_k(n) = \frac{1}{J_1(\lambda_n)} \int_{-\infty}^0 \psi(\lambda + \lambda_n^2) (\lambda + \lambda_n^2)^{k-1} \lambda_n \widehat{h}(\lambda) d\lambda.$$

Since ψ is compactly supported in $(0, 1)$, we can use Parseval’s identity (5.30) and Cauchy–Schwarz inequality to obtain

$$\begin{aligned} \|r^{1/2} \cdot f_k(t)\|_{L^2(\mathbb{T})}^2 &\lesssim \sum_{n \in \mathbb{Z}} |C_k(n)|^2 \left| \frac{1}{J_1(\lambda_n)} \right|^2 = \left| \int_{-\infty}^0 \psi(\lambda + \lambda_n^2) (\lambda + \lambda_n^2)^{k-1} \lambda_n \widehat{h}(\lambda) d\lambda \right|^2 \\ &\lesssim \int_{-\infty}^0 \psi(\lambda + \lambda_n^2) (\lambda + \lambda_n^2)^{2(k-1)} d\lambda \cdot \int_{-\infty}^0 \psi(\lambda + \lambda_n^2) \lambda_n^2 |\widehat{h}(\lambda)|^2 d\lambda. \end{aligned}$$

Using the fact that $\psi \in C_0^\infty(0, 1)$, we can show that the first integral is bounded. For the second integral, note that $|\lambda + \lambda_n^2| < 1$, which implies that $\lambda_n^2 \lesssim (1 + |\lambda|)$. Therefore, we have

$$\|r^{1/2} \cdot f_k(t)\|_{L^2(\mathbb{T})}^2 \lesssim \int_{-\infty}^0 (1 + |\lambda|) |\widehat{h}(\lambda)|^2 d\lambda \lesssim \|h\|_{H^{1/2}}^2.$$

This completes the desired estimate (5.28) for (5.38).

Proof of estimate (5.29). Using estimate (5.28), we prove this estimate for $s = 0$. Then, we will extend the proof to $s = 2$. We note that $W_{ball}h$ defined by (5.27) satisfies the ibvp (5.12), which can be expressed as

$$\begin{aligned} iu_t + u_{x_1x_1} + u_{x_2x_2} &= 0, & x_1^2 + x_2^2 &< 1, \quad t \in (0, T), \\ u(x_1, x_2, 0) &= 0, & x_1^2 + x_2^2 &< 1, \\ u(x_1, x_2, t) &= h(t), & x_1^2 + x_2^2 &= 1, \quad t \in (0, T). \end{aligned}$$

If we let $u_t = v$, then $v(x_1, x_2, 0) = 0$ and v satisfies the same ibvp with h replaced by h_t , and we have

$$\sup_{t \in [0, T]} \|r^{1/2}v\|_{L^2(0,1)} = \sup_{t \in [0, T]} \|(W_{ball}h)_t\|_{L^2(\Omega_0)} \leq C_T \|h_t\|_{H^{1/2}(0, T)}.$$

Hence, for a fixed value of t , the function $u = W_{ball}h$ satisfies

$$\begin{aligned} u_{x_1x_1} + u_{x_2x_2} &= -i(W_{ball}h)_t, & x_1^2 + x_2^2 &< 1, \\ u(x_1, x_2, t) &= h(t), & x_1^2 + x_2^2 &= 1, \end{aligned}$$

which is an elliptic problem on Ω_0 . Since $(W_{ball}h)_t \in L^2(\Omega_0)$ and $h(t)$ is a constant for a fixed t , the theory of elliptic equations implies that $u \in H^2(\Omega_0)$ and

$$\begin{aligned} \|W_{ball}h\|_{H^2(\Omega_0)} &\leq C \left(\|(W_{ball}h)_t\|_{L^2(\Omega_0)} + |h(t)| \right) \leq C \left(\|h_t\|_{H^{1/2}(0, T)} + \|h\|_{H^1(0, T)} \right) \\ &\leq C \|h\|_{H^{3/2}(0, T)}, \end{aligned}$$

where C is a constant dependent only on the domain Ω_0 . This completes the proof of estimate (5.29) for $s = 2$. The result for $0 < s < 2$ follows from interpolation. Therefore, we complete the proof of proposition 5.4.

Now, we are able to state the linear estimate for the solution of ibvp (5.1). To do this, we first express the solution for this ibvp

$$u = B[u_0, g; f] \doteq S_J[u_0; f] + W_{ball}(g), \quad (x_1, x_2) \in \Omega_0, \quad t \in (0, T). \tag{5.41}$$

Furthermore, using propositions 5.1 and 5.4, we obtain the linear estimate in the following solution spaces.

THEOREM 5.6 *Suppose that $0 \leq s \leq 2$. If $u_0 \in H_0^s(\Omega_0)$, $g \in H_0^{\frac{s+1}{2}}(0, T)$ (for $s = 0, 2$, $g \in H_0^{\frac{s+1}{2}}(0, T)$) and $f \in L^1(0, T; H_0^s(\Omega_0))$, then $B[u_0, g_2; f]$ defines a solution to the linear ibvp (5.1) with compatibility condition (3.2), which satisfies*

$$\sup_{t \in [0, T]} \|B[u_0, g; f](t)\|_{H^s(\Omega_0)} \lesssim \|u_0\|_{H^s(\Omega_0)} + \|g\|_{H^{\frac{s+1}{2}}(0, T)} + \|f\|_{L^1(0, T; H^s(\Omega_0))}. \tag{5.42}$$

5.2. Nonlinear problem

We will now investigate the well-posedness of the nonlinear problem (1.1) for $(x_1, x_2) \in \Omega_0$ with the boundary condition $u(x_1, x_2, t) = g(t)$ at $x_1^2 + x_2^2 = 1$. In the solution formula (5.41), by replacing the forcing term f with $-\lambda|u|^{p-2}u$, we obtain the following iteration map

$$u = B[u_0, g; f] = B[u_0, g; -\lambda|u|^{p-2}u]. \tag{5.43}$$

REMARK. In (5.43), it is worth noting that for $s > 1$, we need the condition that $f = -\lambda|u|^{p-2}u \in L^1(0, T; H_0^s(\Omega_0))$. Thus, we can express (5.43) differently. Let

$$w(t) = -\lambda|u|^{p-2}u|_{x_1^2+x_2^2=1} = -\lambda|g|^{p-2}g,$$

and $v(t) = -i \int_0^t w(s) ds$. Then, (5.43) can be transformed into the form

$$u = v(t) + B[u_0, g - v; -\lambda|u|^{p-2}u - w],$$

where $-\lambda|u|^{p-2}u - w = 0$ at $x_1^2 + x_2^2 = 1$, which is the desired boundary condition. Here, $v(t)$ is one order smoother than g if $s > 1$, which does not introduce any difficulties in deriving the relevant estimates. For the sake of simplicity, we will only consider (5.43) in the following.

Next, we will demonstrate that the iteration map defined by (5.43) is a contraction in the solution space $C([0, T^*]; H^s(\Omega_0))$, for $1 < s \leq 2$. To do this, we can use the linear estimate (5.42) to obtain

$$\begin{aligned} \sup_{t \in [0, T^*]} \|B[u_0, g; f](t)\|_{H^s(\Omega_0)} &\lesssim \|u_0\|_{H^s(\Omega_0)} + \|g\|_{H^{\frac{s+1}{2}}(0, T)} \\ &\quad + |\lambda| \| |u|^{p-2}u \|_{L^1(0, T^*; H^s(\Omega_0))}. \end{aligned}$$

To estimate the nonlinear term $|u|^{p-2}u$, we extend u from $\Omega_0 \times (0, T^*)$ to $\mathbb{R}^2 \times (0, T^*)$, such that the extension U satisfies

$$\|U\|_{L^1(0, T^*; H^s(\mathbb{R}^2))} \leq 2\|u\|_{L^1(0, T^*; H^s(\Omega_0))}.$$

Hence, by applying Sobolev–Gagliardo–Nirenberg inequality (see [1]) and Sobolev embedding theorem in \mathbb{R}^2 , for $s > \frac{2}{2} = 1$ (if Ω_0 is in \mathbb{R}^n , then $s > \frac{n}{2}$), we obtain

$$\| |u|^{p-2}u(t) \|_{H^s(\Omega_0)} \lesssim \| |U|^{p-2}U(t) \|_{H^s(\mathbb{R}^2)} \stackrel{s > 1}{\leq} \|U(t)\|_{H^s(\mathbb{R}^2)}^{p-1} \lesssim \|u(t)\|_{H^s(\Omega_0)}^{p-1}.$$

Working similarly, for $s > 1$, we get

$$\| |u|^{p-2}u(t) - |v|^{p-2}v(t) \|_{H^s(\Omega_0)} \lesssim (\|u(t)\|_{H^s(\Omega_0)}^{p-2} + \|v(t)\|_{H^s(\Omega_0)}^{p-2}) \|(u - v)(t)\|_{H^s(\Omega_0)}.$$

Finally, using Hölder's inequality in t integral, we arrive at

$$\begin{aligned} \| |u|^{p-2}u \|_{L^1(0,T^*;H^s(\Omega_0))} &\lesssim T^* \sup_{t \in [0, T^*]} \|u(t)\|_{H^s(\Omega_0)}^{p-1}, \\ \| |u|^{p-2}u(t) - |v|^{p-2}v(t) \|_{L^1(0,T^*;H^s(\Omega_0))} \\ &\lesssim T^* \sup_{t \in [0, T^*]} (\|u(t)\|_{H^s(\Omega_0)}^{p-2} + \|v(t)\|_{H^s(\Omega_0)}^{p-2}) \|(u-v)(t)\|_{H^s(\Omega_0)}. \end{aligned}$$

Using the above estimates, we find that the iteration map (5.43) is a contraction in $C([0, T^*]; H^s(\Omega_0))$ for $s > 1$, as long as $T^* > 0$ is sufficiently small. Thus, we obtain a unique solution $u \in C([0, T^*]; H^s(\Omega_0))$. Given the radial symmetry of the initial and boundary conditions, and due to the uniqueness and rotational invariance of this problem, the solution is also radially symmetric. Consequently, u is a function of $r = \sqrt{x_1^2 + x_2^2}$ in terms of spatial variables.

Funding

The first author was partially supported by a grant from the Simons Foundation (712822, SMS).

References

- [1] R. A. Adams and J. J. F. Fournier. *Sobolev spaces*, Pure and Applied Mathematics, Vol. 140, 2nd ed., p. xiv+305 (Elsevier/Academic Press, Amsterdam, 2003).
- [2] D. Benney and A. Newell. Propagation of nonlinear wave envelopes. *Stud. Appl. Math.* **46** (1967), 133–139.
- [3] J. Bergh and J. Löfström. *Interpolation spaces*, An Introduction, Grundlehren der Mathematischen Wissenschaften, Vol. 223, p. x+207 (Springer-Verlag, Berlin-New York, 1976).
- [4] M. D. Blair, H. F. Smith and C. D. Sogge. Strichartz estimates and the nonlinear Schrödinger equation on manifolds with boundary. *Math. Ann.* **354** (2012), 1397–1430.
- [5] J. L. Bona, S. M. Sun and B.-Y. Zhang. A non-homogeneous boundary-value problem for the Korteweg-de Vries equation in a quarter plane. *Trans. Amer. Math. Soc.* **354** (2002), 427–490.
- [6] J. L. Bona, S. M. Sun and B.-Y. Zhang. A nonhomogeneous boundary-value problem for the Korteweg-de Vries equation posed on a finite domain. *Comm. Partial Differential Equations* **28** (2003), 1391–1436.
- [7] J. L. Bona, S. M. Sun and B.-Y. Zhang. Nonhomogeneous boundary value problems of one-dimensional nonlinear Schrödinger equations. *J. Math. Pures Appl.* **109** (2018), 1–66.
- [8] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. *Geom. Funct. Anal.* **3** (1993), 157–178.
- [9] J. Bourgain and A. Bulut. Almost sure global well-posedness for the radial nonlinear Schrödinger equation on the unit ball II: the 3d case. *J. Eur. Math. Soc. (JEMS)* **16** (2014), 1289–1325.
- [10] J. Bourgain and A. Bulut. Almost sure global well posedness for the radial nonlinear Schrödinger equation on the unit ball I: the 2D case. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **31** (2014), 1267–1288.
- [11] T. Cazenave. *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics, 10. New York University, Courant Institute of Mathematical Sciences, New York. p. xiv+323 (American Mathematical Society, Providence, RI, 2003).
- [12] T. Cazenave and A. Haraux. *An introduction to semilinear evolution equations* (Clarendon, Oxford, 1998).
- [13] T. Cazenave and F. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in H^s . *Nonlin. Anal.* **14** (1990), 807–836.

- [14] J. Chabchoub, N. Hoffmann and N. Akhmediev. Rogue wave observation in a water wave tank. *Phys. Rev. Lett.* **106** (2011), 204502.
- [15] R. Chiao, E. Garmire and C. Townes. Self-trapping of optical beams. *Phys. Rev. Lett.* **13** (1964), 479.
- [16] R. Chiao, E. Garmire and C. Townes. *Phys. Rev. Lett.* **14** (1965), 1056 (erratum).
- [17] M. Christ, J. Colliander and T. Tao. Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math.* **125** (2003), 1235–1293.
- [18] J. E. Colliander and C. E. Kenig. The generalized Korteweg-de Vries equation on the half-line. *Comm. Partial Differential Equations* **27** (2002), 2187–2266.
- [19] M. B. Erdoğan and N. Tzirakis. Regularity properties of the cubic nonlinear Schrödinger equation on the half line. *J. Funct. Anal.* **271** (2016), 2539–2568.
- [20] A. S. Fokas, A. A. Himonas and D. Mantzavinos. The Korteweg-de Vries equation on the half-line. *Nonlinearity* **29** (2016), 489–527.
- [21] A. S. Fokas, A. A. Himonas and D. Mantzavinos. The nonlinear Schrödinger equation on the half-line. *Trans. Amer. Math. Soc.* **369** (2017), 681–709.
- [22] J. Ginibre and G. Velo. On a class of nonlinear Schrödinger equations. II. Scattering theory, general case. *J. Funct. Anal.* **32** (1979), 33–71.
- [23] H. Hajaiej, L. Molinet, T. Ozawa and B. Wang. *Sufficient and necessary conditions for the fractional Gagliardo-Nirenberg inequalities and applications to Navier-Stokes and generalized Bose equations*, RIMS Kokyuroku Bessatsu B26: Harmonic Analysis and Nonlinear Partial Differential Equations (ed. T. Ozawa M. Sugimoto), pp. 159–75 (Kyoto University, Kyoto, 2011).
- [24] B. Harrop-Griffiths, R. Killip and M. Visan. Sharp well-posedness for the cubic NLS and mKdV in $H^s(\mathbb{R})$. *Forum Math. Pi* **12** (2024), 86.
- [25] J. Holmer. The initial-boundary-value problem for the 1D nonlinear Schrödinger equation on the half-line. *Diff. Int. Eq.* **18** (2005), 647–668.
- [26] G. H. Hardy, J. E. Littlewood and G. Pólya. *Inequalities*, 2nd ed., p. xii+324 (University Press, Cambridge, 1952).
- [27] A. Himonas and D. Mantzavinos. Well-posedness of the nonlinear Schrödinger equation on the half-plane. *Nonlinearity* **33** (2020), 5567–5609.
- [28] A. M. Kamchatnov and V. S. Shchesnovich. Dynamics of Bose-Einstein condensates in cigar-shaped traps. *Phys. Rev. A* **70** (2004), 023604.
- [29] C. Kenig, G. Ponce and L. Vega. Oscillatory integrals and regularity of dispersive equations. *Indiana Univ. Math. J.* **40** (1991), 33–69.
- [30] J.-L. Lions and E. Magenes. *Non-homogeneous boundary value problems and applications. Vol. I*, Translated from the French by P. Kenneth, Die Grundlehren der Mathematischen Wissenschaften, Band 182 (Springer-Verlag, New York-Heidelberg, 1972).
- [31] D. Peregrine. Water waves, nonlinear Schrödinger equations and their solutions. *J. Austral. Math. Soc. Ser. B* **25** (1983), 16–43.
- [32] Y. Ran, S. M. Sun and B.-Y. Zhang. Nonhomogeneous boundary value problems of nonlinear Schrödinger equations in a half plane. *SIAM J. Math. Anal.* **50** (2018), 2773–2806.
- [33] V. Talanov. Self-focusing of electromagnetic waves in nonlinear media. *Radiophysics* **8** (1964), 254–257.
- [34] Y. Tsutsumi. L^2 -solutions for nonlinear Schrödinger equations and nonlinear groups. *Funkcial. Ekvac.* **30** (1987), 115–125.
- [35] B. O. Turesson. *Nonlinear potential theory and weighted Sobolev spaces*, Lecture Notes in Mathematics, Vol. 1736, p. xiv+173 (Springer-Verlag, Berlin, 2000).
- [36] V. Zakharov and A. Shabat. Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media. *Sov. Phys. JETP* **34** (1972), 63–69.

Appendix A

Proof of estimate (3.32). We define the operator $J : L^2 \rightarrow L_t^q(\mathbb{R}; L^\gamma(1, \infty))$ as follows

$$J[f](r, t) \doteq \int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau. \tag{A.1}$$

To establish estimate (3.32), it suffices to demonstrate that J is bounded. To do this, we will show that the adjoint operator of J is bounded. Utilizing duality, we have

$$\begin{aligned} & \left\| \int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau \right\|_{L_t^q(\mathbb{R}; L^\gamma(1, \infty))} \\ &= \sup_{\substack{\|\psi\|_{L^{q'}(\mathbb{R}; L^{\gamma'})} = 1 \\ \psi \in C_c([0, T]; \mathcal{D}(1, \infty))}} \int_{\mathbb{R}} \int_{r=1}^\infty \left[\int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau \right] \cdot \psi(r, t) dr dt. \end{aligned}$$

Now, utilizing (3.31), we obtain the t -Fourier transform of $\int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau$, which is

$$\mathcal{F} \left[\int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau \right] \simeq \begin{cases} e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{f}(\tau), & \tau > 0, \\ 0, & \tau < 0. \end{cases}$$

Combined with the Plancherel theorem, we obtain

$$\begin{aligned} & \int_{\mathbb{R}} \int_{r=1}^\infty \left[\int_0^\infty K_t(r, \tau) \widehat{f}(\tau) d\tau \right] \cdot \psi(r, t) dr dt = \int_{r=1}^\infty \int_{\tau=0}^\infty \\ & \left[e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{f}(\tau) \right] \cdot \overline{\widehat{\psi}(r, \tau)} d\tau dr \\ &= \int_{\tau=0}^\infty \widehat{f}(\tau) \overline{\int_{r=1}^\infty \left[e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \cdot \widehat{\psi}(r, \tau) \right] dr d\tau} \\ &\simeq \int_{t \in \mathbb{R}} f(t) \overline{\left[\int_{\tau=0}^\infty \int_{r=1}^\infty e^{i\tau t} e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \cdot \widehat{\psi}(r, \tau) dr d\tau \right]} dt \\ &= \int_{t \in \mathbb{R}} f(t) \overline{\left[\int_{r=1}^\infty J[\psi](r, t) dr \right]} dt. \end{aligned}$$

Thus, the adjoint operator of J is given by

$$J^*[\psi](t) \doteq \overline{\int_{r=1}^\infty J[\psi](r, t) dr}. \tag{A.2}$$

Furthermore, using the Hölder inequality, to establish that J is bounded, it suffices to show

$$\boxed{\left\| \int_{r=1}^\infty J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})} \leq C \|\psi\|_{L^{q'}(0, T; L^{\gamma'})}, \quad \psi \in C_c([0, T]; \mathcal{D}(\mathbb{R})).} \tag{A.3}$$

We have

$$\begin{aligned} \left\| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})}^2 &= \int_{t \in \mathbb{R}} \left| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right|^2 dt \\ &= \int_{t \in \mathbb{R}} \left| \int_{r=1}^{\infty} \int_0^{\infty} K_t(r, \tau) \widehat{f}(\tau) d\tau dr \right|^2 dt \\ &= \int_{t \in \mathbb{R}} \left| \int_0^{\infty} e^{i\tau t} \left(\int_{r=1}^{\infty} e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{\psi}(r, \tau) dr \right) d\tau \right|^2 dt. \end{aligned}$$

Applying Plancherel theorem to dt , we have

$$\begin{aligned} \left\| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})}^2 &= \int_{\tau=0}^{\infty} \left| \int_{r=1}^{\infty} e^{-\sqrt{\tau}(r-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{\psi}(r, \tau) dr \right|^2 d\tau \\ &= \int_{\tau=0}^{\infty} \left(\int_{r_1=1}^{\infty} e^{-\sqrt{\tau}(r_1-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{\psi}(r_1, \tau) dr_1 \right) \\ &\quad \left(\overline{\int_{r_2=1}^{\infty} e^{-\sqrt{\tau}(r_2-1)} (1 + |\tau|)^{-\frac{1}{4}} \widehat{\psi}(r_2, \tau) dr_2} \right) d\tau. \end{aligned}$$

Using Fubini's theorem, we can simplify the expression to

$$\begin{aligned} \left\| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})}^2 &= \int_{r_1=1}^{\infty} \int_{\tau=0}^{\infty} \widehat{\psi}(r_1, \tau) \\ &\quad \left(\overline{\int_{r_2=1}^{\infty} e^{-\sqrt{\tau}(r_1+r_2-2)} (1 + |\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2} \right) d\tau dr_1. \end{aligned}$$

Furthermore, applying Plancherel's theorem to $d\tau$, we have

$$\begin{aligned} \left\| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})}^2 &\simeq \int_{r_1=1}^{\infty} \int_{t \in \mathbb{R}} \psi(r_1, t) \left(\overline{\int_{\tau=0}^{\infty} \int_{r_2=1}^{\infty} e^{i\tau t} e^{-\sqrt{\tau}(r_1+r_2-2)} (1 + |\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2 d\tau} \right) \\ &\quad dt dr_1. \end{aligned}$$

Moreover, applying Holder's inequality to dr_1 and dt , we get

$$\begin{aligned} &\left\| \int_{r=1}^{\infty} J[\psi](r, \cdot) dr \right\|_{L_t^2(\mathbb{R})}^2 \\ &\leq \|\psi\|_{L_t^{q'}[\mathbb{R}; L_{r_1}^{\gamma'}(1, \infty)]} \left\| \int_{\tau=0}^{\infty} \int_{r_2=1}^{\infty} e^{i\tau t} e^{-\sqrt{\tau}(r_1+r_2-2)} (1 + |\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2 d\tau \right\|_{L_t^q[\mathbb{R}; L_{r_1}^{\gamma}(1, \infty)]} \\ &= \|\psi\|_{L_t^{q'}[\mathbb{R}; L_{r_1}^{\gamma'}(1, \infty)]} \left\| \int_{\tau=0}^{\infty} \int_{r_2=1}^{\infty} e^{i\tau t} e^{-\sqrt{\tau}(r_1+r_2-2)} (1 + |\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2 d\tau \right\|_{L_t^q[\mathbb{R}; L_{r_1}^{\gamma}(1, \infty)]}. \end{aligned}$$

Now, to prove the inequality (A.3), it is enough to show that

$$\left\| \int_{\tau=0}^{\infty} \int_{r_2=1}^{\infty} e^{i\tau t} e^{-\sqrt{\tau}(r_1+r_2-2)} (1+|\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2 d\tau \right\|_{L_t^q[\mathbb{R}; L_{r_1}^\gamma(1, \infty)]} \\ \lesssim \|\psi\|_{L_t^{q'}[\mathbb{R}; L^{\gamma'}(1, \infty)]}. \quad (\text{A.6})$$

To proceed, we will use the convolution property: $\mathcal{F}^{-1}(f \cdot g) = \mathcal{F}^{-1}(g) * \mathcal{F}^{-1}(f)$. This allows us to rewrite the integral as

$$\int_{\tau=0}^{\infty} \int_{r_2=1}^{\infty} e^{i\tau t} e^{-\sqrt{\tau}(r_1+r_2-2)} (1+|\tau|)^{-\frac{1}{2}} \widehat{\psi}(r_2, \tau) dr_2 d\tau \\ \simeq \int_{t_1 \in \mathbb{R}} \left[\int_{r_2=1}^{\infty} \left(\int_{\tau=0}^{\infty} e^{i\tau(t-t_1)} e^{-\sqrt{\tau}(r_1+r_2-2)} (1+|\tau|)^{-\frac{1}{2}} d\tau \right) \psi(r_2, t_1) dr_2 \right] dt_1.$$

Additionally, following a similar approach to the proof in reference [7], we obtain

$$\left\| \int_{r_2=1}^{\infty} \left(\int_{\tau=0}^{\infty} e^{i\tau(t-t_1)} e^{-\sqrt{\tau}(r_1+r_2-2)} (1+|\tau|)^{-\frac{1}{2}} d\tau \right) \psi(r_2, t_1) dr_2 \right\|_{L_{r_1}^\gamma(1, \infty)} \\ \lesssim |t-t_1|^{\frac{1}{\gamma}-\frac{1}{2}} \|\psi(\cdot, t_1)\|_{L_{r_2}^{\gamma'}(1, \infty)}, \quad 2 \leq \gamma \leq \infty. \quad (\text{A.7})$$

Finally, we combine the above estimate with the following result to obtain the desired estimate (A.6).

LEMMA A.1. Hardy–Littlewood–Polya [26] section 10.17, theorem 382 *If $F, G \geq 0$ then*

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{F(x)G(y)}{|x-y|^\lambda} dx dy \leq C \|F\|_{L_x^P(\mathbb{R})} \|G\|_{L_x^Q(\mathbb{R})}, \quad C > 0, \quad (\text{A.8})$$

with $P > 1, Q > 1, \frac{1}{P} + \frac{1}{Q} > 1, \lambda = 2 - \frac{1}{P} - \frac{1}{Q}$.

It can be observed that

$$\begin{aligned} A &\doteq \left(\int_{t=-\infty}^{\infty} \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right|^q dt \right)^{1/q} \\ &= \left(\int_{t=-\infty}^{\infty} \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right| \right. \\ &\quad \cdot \left. \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right|^{q-1} dt \right)^{1/q} \\ &= \left(\int_{t=-\infty}^{\infty} \int_{t_1=-\infty}^{\infty} |t - t_1|^{-\lambda} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} G(t) dt_1 dt \right)^{1/q} \\ &\lesssim \left(\|\psi\|_{L^{\gamma'} L^q_{t_1}} \|G\|_{L^q_t} \right)^{1/q}, \end{aligned}$$

where we have taken $\lambda = \frac{1}{2} - \frac{1}{\gamma}$, $G(t) = \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right|^{q-1}$, and set $P = Q = q'$. It is easy to verify that the admissible condition $\frac{2}{q} + \frac{1}{\gamma} = \frac{1}{2}$ implies that $\lambda = 2 - \frac{1}{P} - \frac{1}{Q}$. In fact, we have

$$\lambda = \frac{1}{2} - \frac{1}{\gamma} = \frac{1}{2} - \left(\frac{1}{2} - \frac{2}{q} \right) = \frac{2}{q} = 2 - \frac{2}{q'} = 2 - \frac{1}{P} - \frac{1}{Q}.$$

Additionally, we have

$$\|G\|_{L^q_t(\mathbb{R})} = \left(\int_{t \in \mathbb{R}} \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right|^{(q-1)q'} dt \right)^{1/q'}.$$

Combining the above identity with $\frac{1}{q'} = 1 - \frac{1}{q}$, or $q' = \frac{q}{q-1}$, we get

$$\|G\|_{L^q_t(\mathbb{R})} = \left(\int_{t \in \mathbb{R}} \left| \int_{t_1=-\infty}^{\infty} |t - t_1|^{\frac{1}{\gamma} - \frac{1}{2}} \|\psi(\cdot, t_1)\|_{L^{\gamma'}} dt_1 \right|^q dt \right)^{1/q} = A^{q/q'}.$$

Thus, we find

$$A \lesssim \left(\|\psi\|_{L^{\gamma'} L^q_{t_1}} \|G\|_{L^q_t(\mathbb{R})} \right)^{1/q} = \left(\|\psi\|_{L^{\gamma'} L^q_{t_1}} A^{q/q'} \right)^{1/q} = \|\psi\|_{L^{\gamma'} L^q_{t_1}}^{1/q} A^{1/q'}.$$

Therefore, we have $A^{1-1/q'} = A^{1/q} \lesssim \|\psi\|_{L^{\gamma'} L^q_{t_1}}^{1/q}$ or $A \lesssim \|\psi\|_{L^{\gamma'}_{[\mathbb{R}; L^q_{t_1}(1, \infty)]}}$, which combined with inequality (A.7) gives the desired estimate (A.6). This completes the proof of estimate (3.32). \square