


# COMPOSITIO MATHEMATICA

## Cubical setting for discrete homotopy theory, revisited

D. Carranza and K. Kapulkin 

Compositio Math. **160** (2024), 2856–2903.

[doi:10.1112/S0010437X24007486](https://doi.org/10.1112/S0010437X24007486)



FOUNDATION  
COMPOSITIO  
MATHEMATICA



LONDON  
MATHEMATICAL  
SOCIETY  
EST. 1865





# Cubical setting for discrete homotopy theory, revisited

D. Carranza and K. Kapulkin 

## ABSTRACT

We construct a functor associating a cubical set to a (simple) graph. We show that cubical sets arising in this way are Kan complexes, and that the A-groups of a graph coincide with the homotopy groups of the associated Kan complex. We use this to prove a conjecture of Babson, Barcelo, de Longueville, and Laubenbacher from 2006, and a strong version of the Hurewicz theorem in discrete homotopy theory.

## Contents

<b>Introduction</b>	<b>2857</b>
Organization of the paper	2859
<b>1 Discrete homotopy theory</b>	<b>2859</b>
The category of graphs	2859
Examples of graphs	2861
Monoidal structure on the category of graphs	2862
Homotopy theory of graphs	2863
Path and loop graphs	2865
<b>2 Cubical sets and their homotopy theory</b>	<b>2866</b>
Cubical sets	2866
Kan complexes	2868
Anodyne maps	2870
Homotopies and homotopy groups	2871
<b>3 Cubical nerve of a graph</b>	<b>2873</b>
<b>4 Main result</b>	<b>2879</b>
Statement	2879
Proof of part (i)	2879
Proof of part (ii)	2884
<b>5 Consequences</b>	<b>2890</b>
Proof of the conjecture of Babson, Barcelo, de Longueville, and Laubenbacher	2890
Discrete homology of graphs	2890
Fibration category of graphs	2892

Received 17 March 2023, accepted in final form 16 May 2024.

*2020 Mathematics Subject Classification* 05C25 (primary), 55U35, 18N40, 18N45 (secondary).

*Keywords:* discrete homotopy theory, cubical set, reflexive graph, Kan complex.

© The Author(s), 2025. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited. *Compositio Mathematica* is

© Foundation Compositio Mathematica.

Cubical enrichment of the category of graphs .....	2899
<b>Acknowledgements</b>	<b>2901</b>
<b>References</b>	<b>2901</b>

### Introduction

Discrete homotopy theory, introduced in [BKLW01, BBdLL06], is a homotopy theory in the category of simple graphs. It builds on the earlier work of Atkin [Atk74, Atk76], made precise in [KL98], on the homotopy theory of simplicial complexes, and can also be generalized to the homotopy theory of finite metric spaces [BCW14]. It has found numerous applications [BL05, § 5–6], including in matroid theory, hyperplane arrangements, combinatorial time series analysis, and, more recently, topological data analysis [MZ19].

The key invariants associated to graphs in discrete homotopy theory are the A-groups  $A_n(G, v)$ , named after Atkin, which are the discrete analogue of homotopy groups  $\pi_n(X, x)$ , studied in the homotopy theory of topological spaces. In [BBdLL06], Babson, Barcelo, de Longueville, and Laubenbacher construct an assignment  $G \mapsto X_G$ , taking a graph  $G$  to a topological space, constructed as a certain cubical complex, and conjecture that the A-groups of  $G$  coincide with the homotopy groups of  $X_G$ . They further prove [BBdLL06, Theorem 5.2] their conjecture under an assumption of a cubical approximation property [BBdLL06, Proposition 5.1], a cubical analogue of the simplicial approximation theorem, which remains open.

The assignment  $G \mapsto X_G$  arises as a composite

$$\text{Graph} \xrightarrow{N_1} \text{cSet} \xrightarrow{|\cdot|} \text{Top}.$$

Here **cSet** denotes a particular category of cubical sets, which are well-studied combinatorial models for the homotopy theory of spaces [Cis06]. (Specifically, cubical sets used in this paper are cubical sets with positive and negative connections; cf. [BH81, BM17, DKLS24].) Informally, a cubical set consists of a family of sets  $\{X_n\}_{n \in \mathbb{N}}$  of  $n$ -cubes together with a family of structure maps indicating how different cubes ‘fit together’, e.g. that a certain  $(n - 1)$ -cube is a face of another  $n$ -cube. More formally, it is a presheaf on the category  $\square$  of combinatorial cubes. The functor  $N_1: \text{Graph} \rightarrow \text{cSet}$  is obtained by taking the  $n$ -cubes of  $N_1G$  to be maps  $I_1^{\otimes n} \rightarrow G$ , where  $I_1^{\otimes n}$  denotes the  $n$ -dimensional hypercube graph. The functor  $|\cdot|: \text{cSet} \rightarrow \text{Top}$ , called the geometric realization, assigns the topological  $n$ -cube  $[0, 1]^n$  to each (formal)  $n$ -cube of  $X$  and then glues these cubes together according to structure maps. (We will of course give precise definitions of all these notions later in the paper.)

A reader familiar with [BBdLL06] will recognize a change in notation: the functor  $N_1$  above was in [BBdLL06, §4] denoted by  $M_*$ . This change is intentional, as we wish to emphasize that this is only the first in a sequence of functors  $\text{Graph} \rightarrow \text{cSet}$  and natural transformations:

$$N_1 \longrightarrow N_2 \longrightarrow N_3 \longrightarrow \dots \tag{1}$$

It is this sequence and its colimit

$$N := \text{colim}(N_1 \rightarrow N_2 \rightarrow N_3 \rightarrow \dots)$$

that we investigate. Our main technical results can be summarized by the following theorem.


**THEOREM** (cf. Theorems 4.1 and 4.6).

- (i) For a graph  $G$ , the cubical set  $NG$  is a Kan complex.
- (ii) The natural transformations in (1) are natural weak equivalences.

(iii) For a based graph  $(G, v)$ , there is a natural group isomorphism  $A_n(G, v) \cong \pi_n(NG, v)$ .

A few words of explanation are in order. A Kan complex is a cubical set satisfying a certain lifting property making it particularly convenient for the development of homotopy theory. In particular, in a companion paper [CK23], we develop the theory of homotopy groups of Kan complexes and show that these agree with their topological analogues under the geometric realization functor.

This establishes the key ingredient required for the proof of the conjecture of Babson, Barcelo, de Longueville, and Laubenbacher, which asks for the commutativity of the outer square in the following diagram.



By the theorem above, the two triangles on the left commute, with the upper one commuting up to a natural weak equivalence (which is sent by the composite functor  $\pi_n \circ |\cdot|: \text{cSet}_* \rightarrow \text{Grp}$  to a natural isomorphism). The upper right triangle commutes on the nose and the lower right triangle commutes since it expresses the compatibility between homotopy groups of cubical sets and those of topological spaces, established in [CK23]. Thus, the  $A$ -groups of  $(G, v)$  agree with those of  $(NG, v)$ , which in turn agree with those of  $(|N_1G|, v)$ , since the map  $N_1G \rightarrow NG$  is a weak equivalence.

**THEOREM** (Conjectured in [BBdLL06]; cf. Theorem 5.1). *There is a natural group isomorphism  $A_n(G, v) \cong \pi_n(|N_1G|, v)$ .*

The insight of [BBdLL06] cannot be overstated. It is *a priori* not clear, or at the very least it was not clear to us, that the  $A$ -groups of a graph should correspond to the homotopy groups of any space. The fact that the space can be obtained from a graph in such a canonical and simple way is what drove us to the problem and to the field of discrete homotopy theory. The title of our paper, ‘Cubical setting for discrete homotopy theory, revisited’, pays tribute to this insight by alluding to the title ‘A cubical set setting for the  $A$ -theory of graphs’ of [BBdLL06, § 3].

It is also worth noting that the case of  $n = 1$  of Theorem 5.1 was previously proven in [BKLW01, Proposition 5.12] and perhaps helped inspire the statement of the conjecture in the general case.

Our main theorem allows us to derive a few more consequences of interest in discrete homotopy theory. The first of those is a strong form of the Hurewicz theorem for graphs. The Hurewicz theorem relates the first non-trivial homotopy group of a sufficiently connected space to its homology. In discrete homotopy theory, it relates the first non-trivial  $A$ -group of a graph to its reduced discrete homology, introduced in [BCW14].

**THEOREM** (Discrete Hurewicz theorem; cf. Theorem 5.8). *Let  $n \geq 2$  and  $(G, v)$  be a connected pointed graph. Suppose  $A_i(G, v) = 0$  for all  $i \in \{1, \dots, n - 1\}$ . Then the Hurewicz map  $A_n(G, v) \rightarrow \widetilde{DH}_n(G, v)$  from the  $n$ th  $A$ -group to the  $n$ th reduced discrete homology group is an isomorphism.*

This generalizes the results of Lutz [Lut21, Theorem 5.10], who proves surjectivity of the Hurewicz map for a more restrictive class of graphs, and complements the result of Barcelo,

Capraro, and White [BCW14, Theorem 4.1], who prove the one-dimensional analogue of the Hurewicz theorem, namely that the Hurewicz map  $A_1(G, v) \rightarrow \widetilde{DH}_1(G, v)$  is surjective with kernel given by the commutator  $[A_1(G, v), A_1(G, v)]$  subgroup.

Finally, our main theorem allows us equip the category of graphs with additional structure making it amenable to techniques of abstract homotopy theory and higher category theory.

By our theorem, the functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  takes values in the full subcategory  $\mathbf{Kan}$  of  $\mathbf{cSet}$  spanned by Kan complexes. This subcategory is known to carry the structure of a fibration category in the sense of Brown [Bro73]. In brief, a fibration category is a category equipped with two classes of maps: fibrations and weak equivalences, subject to some axioms. Fibration categories are one of the main frameworks used in abstract homotopy theory (see, for instance, [Szu16]). We declare a map  $f$  of graphs to be a fibration/weak equivalence if  $Nf$  is one in the fibration category of Kan complexes. Our theorem guarantees that this gives a well-defined fibration category structure on  $\mathbf{Graph}$  (Theorem 5.9), hence allowing for the use of techniques from abstract homotopy theory in discrete homotopy theory. Furthermore, it follows that the weak equivalences of this fibration category are precisely graph maps inducing isomorphisms on all  $A$ -groups for all choices of the basepoint.

This also allows us to put the results of [BBdLL06, § 6] in the context of abstract homotopy theory by proving that the loop graph functor constructed there is an exact functor in the sense of fibration category theory (Theorem 5.27).

In a different direction, we observe that the functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  is lax monoidal (Lemma 5.28). The category of graphs is enriched over itself, meaning that the collection of graph maps between two graphs forms not merely a set, but a graph, and that the composition of graph maps defines a graph homomorphism. Enrichment can be transferred along lax monoidal functors, which means that the category of graphs is, via  $N$ , canonically enriched over cubical sets, and, more precisely, over Kan complexes (Theorem 5.36). This establishes a presentation of the  $(\infty, 1)$ -category of graphs, thus allowing for the use of techniques of higher category theory, developed extensively by Joyal and Lurie [Lur09], in discrete homotopy theory, via the cubical homotopy coherent nerve construction of [KV20, § 2].

**Organization of the paper**

This paper is organized as follows. In §§ 1, 2, we review the background on discrete homotopy theory and cubical sets, respectively. In § 3, we explain the link between graphs and cubical sets by defining the functors  $N_m$  and  $N$ , and proving their basic properties. The technical heart of the paper is contained in § 4, where we prove our main results. In § 5, we proceed to deduce the consequences of our main theorem, as described above.

**1. Discrete homotopy theory**

**The category of graphs**

We define the category of simple undirected graphs without loops as a reflective subcategory of a presheaf category.

Let  $\mathbf{G}$  be the category generated by the diagram

$$V \begin{array}{c} \xrightarrow{s} \\ \xleftarrow{r} \\ \xrightarrow{t} \end{array} E \curvearrowright \sigma$$




FIGURE 1. An example depiction of a graph.

subject to the identities

$$\begin{aligned} rs = rt = \text{id}_V, \quad \sigma^2 = \text{id}_E, \\ \sigma s = t, \quad \sigma t = s, \\ r\sigma = r. \end{aligned}$$

We write  $\widehat{\mathbf{G}}$  for the functor category  $\mathbf{Set}^{\mathbf{G}^{\text{op}}}$ .

For  $G \in \widehat{\mathbf{G}}$ , we write  $G_V$  and  $G_E$  for the sets  $G(V)$  and  $G(E)$ , respectively. Explicitly, such a functor consists of sets  $G_V$  and  $G_E$  together with the following functions between them:

$$G_V \begin{array}{c} \xleftarrow{Gs} \\ \xleftarrow{Gr} \rightarrow \\ \xleftarrow{Gt} \end{array} G_E \begin{array}{c} \xrightarrow{G\sigma} \\ \xrightarrow{G\sigma} \end{array} G_E$$

subject to the dual versions of identities in  $\mathbf{G}$ .

DEFINITION 1.1. A *graph* is a functor  $G \in \widehat{\mathbf{G}}$  such that the map  $(Gs, Gt): G_E \rightarrow G_V \times G_V$  is a monomorphism (Figure 1).

Let  $\mathbf{Graph}$  denote the full subcategory of  $\widehat{\mathbf{G}}$  spanned by graphs.

In more concrete terms, a graph  $G$  consists of a set  $G_V$  of vertices and a set  $G_E$  of ‘half-edges’. A half-edge  $e \in G_E$  has source and target, and these are given by the maps  $Gs$  and  $Gt$ , respectively. Each half-edge is paired with its other half via the map  $G\sigma: G_E \rightarrow G_E$ . Note that the edges paired by  $G\sigma$  have swapped source and target, making the pair (i.e. whole edge) undirected. The map  $Gr: G_V \rightarrow G_E$  takes a vertex to an edge whose source and target is that vertex (i.e. a loop). Finally, the condition that  $(Gs, Gt)$  is a monomorphism ensures that there is at most one (whole) edge between any two vertices. This is equivalent to specifying a binary ‘incidence’ relation on  $G_V$  which is reflexive and symmetric.

A map of graphs  $f: G \rightarrow H$  is a natural transformation between such functors. However, since  $(Hs, Ht)$  is a monomorphism, such a map is completely determined by a function  $f_V: G_V \rightarrow H_V$  that preserves the incidence relation.

We may therefore assume that our graphs have no loops, but the maps between them, rather than merely preserving edges, are allowed to contract them to a single vertex. That is, a graph map  $f: G \rightarrow H$  is a function  $f: G_V \rightarrow H_V$  such that if  $v, w \in G_V$  are connected by an edge, then either  $f(v)$  and  $f(w)$  are connected by an edge or  $f(v) = f(w)$ .

PROPOSITION 1.2.

- (i) *The inclusion  $\mathbf{Graph} \hookrightarrow \widehat{\mathbf{G}}$  admits a left adjoint.*
- (ii) *The category  $\mathbf{Graph}$  is (co)complete.*
- (iii) *The functor  $\mathbf{Graph} \rightarrow \mathbf{Set}$  mapping a graph  $G$  to its set of vertices  $G_V$  admits both adjoints.*

*Proof.* (i) The left adjoint is  $\text{Im}: \widehat{\mathbf{G}} \rightarrow \mathbf{Graph}$  given by

$$(\text{Im } G)_V = G_V, \quad (\text{Im } G)_E = (Gs, Gt)(G_E),$$

where  $(Gs, Gt)(G_E)$  is the image of  $G_E$  under the map  $(Gs, Gt): G_E \rightarrow G_V \times G_V$ .

(ii) The category  $\widehat{\mathbf{G}}$  is (co)complete as a presheaf category. The conclusion follows from (i) as  $\mathbf{Graph}$  is a reflective subcategory of a (co)complete category.

(iii) The left adjoint  $\mathbf{Set} \rightarrow \mathbf{Graph}$  takes a set  $A$  to the discrete graph with vertex set  $A$ . The right adjoint takes a set  $A$  to the complete graph with vertex set  $A$ .  $\square$

*Remark 1.3.* Proposition 1.2 gives a procedure for constructing limits and colimits in  $\mathbf{Graph}$ . Given a diagram  $F: J \rightarrow \mathbf{Graph}$ , the set of vertices of  $\lim F$  is the limit  $\lim UF$  of the diagram  $UF: J \rightarrow \mathbf{Set}$ . The set of edges is the largest set such that the limit projections  $\lim F \rightarrow F_j$  are graph maps. The set of vertices of  $\operatorname{colim} F$  is the colimit  $\operatorname{colim} UF$  of the diagram  $UF: J \rightarrow \mathbf{Set}$  and the set of edges is the smallest set such that the colimit inclusions  $F_j \rightarrow \operatorname{colim} F$  are graph maps.

**Examples of graphs**

DEFINITION 1.4. For  $m \geq 0$ ,

- (i) the  $m$ -interval  $I_m$  is the graph which has
  - as vertices, integers  $0 \leq i \leq m$ ;
  - an edge between  $i$  and  $i + 1$  (see Figure 2);
- (ii) the  $m$ -cycle  $C_m$  is the graph which has
  - as vertices, integers  $0 \leq i \leq m - 1$ ;
  - an edge between  $i$  and  $i + 1$  and an edge between  $m - 1$  and  $0$  (see Figure 2);
- (iii) the infinite interval  $I_\infty$  is the graph which has
  - as vertices, integers  $i \in \mathbb{Z}$ ;
  - an edge between  $i$  and  $i + 1$  for all  $i \in \mathbb{Z}$ .

*Remark 1.5.* The graphs  $I_0$  and  $I_1$  are representable when regarded as functors  $\mathbf{G}^{\text{op}} \rightarrow \mathbf{Set}$ , represented by  $V$  and  $E$ , respectively.

For  $m \geq 0$ , we have a map  $l: I_{m+1} \rightarrow I_m$  defined by

$$l(v) = \begin{cases} 0 & \text{if } v = 0 \\ v - 1 & \text{otherwise.} \end{cases}$$

As well, we have a map  $r: I_{m+1} \rightarrow I_m$  defined by

$$r(v) = \begin{cases} m & \text{if } v = m + 1 \\ v & \text{otherwise.} \end{cases}$$

We write  $c: I_{m+2} \rightarrow I_m$  for the composite  $lr = rl$ . Explicitly, this map is defined by

$$c(v) = \begin{cases} 0 & \text{if } v = 0 \\ m & \text{if } v = m + 2 \\ v - 1 & \text{otherwise.} \end{cases}$$

We show the inclusion  $\mathbf{Graph} \hookrightarrow \widehat{\mathbf{G}}$  preserves filtered colimits and use this to show all finite graphs are compact, i.e. if  $G$  is a finite graph then the functor  $\mathbf{Graph}(G, -): \mathbf{Graph} \rightarrow \mathbf{Set}$  preserves filtered colimits.

PROPOSITION 1.6. *The inclusion  $\mathbf{Graph} \hookrightarrow \widehat{\mathbf{G}}$  preserves filtered colimits.*

*Proof.* Fix a filtered category  $J$  and a diagram  $D: J \rightarrow \mathbf{Graph}$ . Let  $i$  denote the inclusion  $\mathbf{Graph} \hookrightarrow \widehat{\mathbf{G}}$ . Recall that  $\operatorname{colim} D$  is computed by  $\operatorname{lm}(\operatorname{colim}(iD))$ . It suffices to show  $\operatorname{colim}(iD) \in \widehat{\mathbf{G}}$  is a graph,




FIGURE 2. The graphs  $I_3$  and  $C_3$ , respectively.

since then the unit map  $\text{colim}(iD) \rightarrow \text{ilm}(\text{colim}(iD))$  is an isomorphism  $\text{colim}(iD) \cong i(\text{colim } D)$  natural in  $D$ .

Let  $\lambda: iD \rightarrow \text{colim}(iD)$  denote the colimit cone. Suppose two edges  $e, e' \in \text{colim}(iD)_E$  have the same source and target. Regarding  $e, e'$  as maps  $I_1 \rightarrow \text{colim}(iD)$ , these maps factor as

$$\begin{array}{ccc}
 I_1 & \xrightarrow{e, e'} & \text{colim}(iD) \\
 \searrow_{\bar{e}, \bar{e}'} & & \nearrow_{\lambda_x} \\
 & iDx &
 \end{array}$$

for some  $x \in J$  since  $I_1 \in \widehat{\mathbf{G}}$  is representable. Let  $s, s' \in (iDx)_V$  denote the sources of  $\bar{e}, \bar{e}'$  and let  $t, t' \in (iDx)_V$  denote the targets, respectively. Using an explicit description of the colimit (Remark 1.3), since  $\lambda_x(s) = \lambda_x(s')$  and  $\lambda_x(t) = \lambda_x(t')$ , there exist arrows  $f, g: y \rightarrow x$  and  $h, k: z \rightarrow x$  with vertices  $v \in iDy$  and  $w \in iDz$  such that

$$iDf(v) = s, \quad iDg(v) = s', \quad iDh(w) = t, \quad iDk(w) = t'.$$

As  $J$  is filtered, there exists an arrow  $l: x \rightarrow w$  in  $J$  such that  $lf = lg$  and  $lh = lk$ . This implies the edges  $iDl(\bar{e}), iDl(\bar{e}') \in (iDw)_E$  have the same source and target. As  $iDw$  is a graph, it follows that  $\bar{e} = \bar{e}'$ , thus  $e = e'$ . □

**COROLLARY 1.7.** *For a finite graph  $G$ , the functor  $\text{Graph}(G, -): \mathbf{Graph} \rightarrow \mathbf{Set}$  preserves filtered colimits.*

*Proof.* Given a filtered category  $J$  and a diagram  $D: J \rightarrow \mathbf{Graph}$ , we have a natural isomorphism

$$\text{Graph}(G, \text{colim } D) \cong \widehat{\mathbf{G}}(G, \text{colim } D)$$

by Proposition 1.6 since  $\mathbf{Graph}$  is a full subcategory. This then follows since  $G \in \widehat{\mathbf{G}}$  is a finite colimit of representable presheaves. □

**Monoidal structure on the category of graphs**

Define a functor  $\otimes: \mathbf{G} \times \mathbf{G} \rightarrow \widehat{\mathbf{G}}$  by

$$\begin{aligned}
 (V, V) &\mapsto I_0, & (V, E) &\mapsto I_1, \\
 (E, V) &\mapsto I_1, & (E, E) &\mapsto C_4.
 \end{aligned}$$

Left Kan extension along the Yoneda embedding yields a monoidal product  $\otimes: \widehat{\mathbf{G}} \times \widehat{\mathbf{G}} \rightarrow \widehat{\mathbf{G}}$ :

$$\begin{array}{ccc}
 \mathbf{G} \times \mathbf{G} & \longrightarrow & \widehat{\mathbf{G}} \\
 \downarrow & \nearrow_{\otimes} & \\
 \widehat{\mathbf{G}} \times \widehat{\mathbf{G}} & &
 \end{array}$$

Note that  $\mathbf{Graph}$  is closed with respect to this product, i.e. if  $G, H \in \mathbf{Graph}$  then  $G \otimes H \in \widehat{\mathbf{G}}$  is a graph. Thus, this product descends to a monoidal product  $\otimes: \mathbf{Graph} \times \mathbf{Graph} \rightarrow \mathbf{Graph}$ , called the *cartesian product*.



Explicitly, the graph  $G \otimes H$  has:

- as vertices, pairs  $(v, w)$  where  $v \in G$  is a vertex of  $G$  and  $w \in H$  is a vertex of  $H$ ;
- an edge from  $(v, w)$  to  $(v', w')$  if either  $v = v'$  and  $w$  is connected to  $w'$  in  $H$  or  $w = w'$  and  $v$  is connected to  $v'$  in  $G$ .

DEFINITION 1.8. Let  $G$  and  $H$  be graphs. The graph  $\text{hom}^\otimes(G, H)$  has:

- as vertices, morphisms  $G \rightarrow H$  in  $\mathbf{Graph}$ ;
- an edge from  $f$  to  $g$  if there exists  $H: G \otimes I_1 \rightarrow H$  such that  $H|_{G \otimes \{0\}} = f$  and  $H|_{G \otimes \{1\}} = g$ .

This structure makes the category of graphs into a closed symmetric monoidal category.

PROPOSITION 1.9.  $(\mathbf{Graph}, \otimes, I_0, \text{hom}^\otimes(-, -))$  is a closed symmetric monoidal category.

### Homotopy theory of graphs

We now review the basics of discrete homotopy theory. Our treatment is brief and more categorically oriented, but the reader can find full details in any of the references [Mal83, BKLW01, BL05, BBdLL06].

DEFINITION 1.10. Let  $f, g: G \rightarrow H$  be graph maps. An  $A$ -homotopy (or just a homotopy) from  $f$  to  $g$  is a map  $\eta: G \otimes I_m \rightarrow H$  for some  $m \geq 0$  such that  $\eta|_{G \otimes \{0\}} = f$  and  $\eta|_{G \otimes \{m\}} = g$ .

Note that one needs to allow the parameter  $m$  appearing above to vary, as otherwise the notion of homotopy is not transitive. This is reminiscent of the notion of Moore path in topological spaces, which is a path parameterized by an interval of arbitrary length [May75, vdBG12, BR13].

PROPOSITION 1.11 [Mal83, Proposition 2.1]. For graphs  $G$  and  $H$ , homotopy is an equivalence relation on the set of graphs maps  $G \rightarrow H$ .

We also define the  $A$ -homotopy groups of a graph. As in topological spaces, this requires the definition of based graph maps and based homotopies between them.

DEFINITION 1.12. Let  $A \hookrightarrow G$  and  $B \hookrightarrow H$  be graph monomorphisms. A relative graph map, denoted  $(G, A) \rightarrow (H, B)$ , is a morphism from  $A \hookrightarrow G$  to  $B \hookrightarrow H$  in the arrow  $\mathbf{Graph}^{[1]}$ , where  $[1]$  denotes the poset  $\{0 \leq 1\}$  viewed as a category.

Explicitly, this data consists of maps  $G \rightarrow H$  and  $A \rightarrow B$  such that the following square commutes:

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ G & \longrightarrow & H \end{array}$$

That is,  $A$  is a subgraph of  $G$  and the map  $A \rightarrow B$  is the restriction of the map  $G \rightarrow H$  to  $A$ , whose image is contained in the subgraph  $B$ .

In the context of relative graph maps, we denote a monomorphism  $A \hookrightarrow G$  by  $(G, A)$ , suppressing the data of the map itself. An exception to this is for monomorphisms of the form  $I_0 \rightarrow G$ . This is exactly the data of a pointed graph, which we denote by  $(G, v)$ , where  $v$  is the unique vertex in the image of the map  $I_0 \rightarrow G$ .

For a relative graph map  $(f, g): (G, A) \rightarrow (H, B)$ , the map  $g$  is uniquely determined by  $f$ . If a map  $h: A \rightarrow B$  also forms a commutative square with  $f$  then  $g = h$  as the map  $B \hookrightarrow H$  is monic. Thus, we denote a relative graph map by  $f: (G, A) \rightarrow (H, B)$ . We additionally write  $f: G \rightarrow H$  for the bottom map in the square and  $f|_A: A \rightarrow B$  for the top map.

We define relative homotopies between relative graph maps as follows.

DEFINITION 1.13. Let  $f, g: (G, A) \rightarrow (H, B)$  be relative graph maps. A *relative homotopy* from  $f$  to  $g$  is a relative map  $(G \otimes I_m, A \otimes I_m) \rightarrow (H, B)$  for some  $m \geq 0$  such that

- the map  $A \otimes I_m \rightarrow B$  is a homotopy from  $f|_A$  to  $g|_A$ ;
- the map  $G \otimes I_m \rightarrow H$  is a homotopy from  $f$  to  $g$ .

Explicitly, a relative homotopy from  $f$  to  $g$  consists of

- a homotopy  $\eta: G \otimes I_m \rightarrow H$  from  $f$  to  $g$ ,
- a homotopy  $\eta|_A: A \otimes I_m \rightarrow B$  from  $f|_A$  to  $g|_A$ ,

such that the following square commutes:

$$\begin{array}{ccc} A \otimes I_m & \xrightarrow{\eta|_A} & B \\ \downarrow & & \downarrow \\ G \otimes I_m & \xrightarrow{\eta} & H \end{array}$$

Remark 1.14. For graph maps  $f, g: G \rightarrow H$ , a path of length  $m$  in  $\text{hom}^\otimes(G, H)$  from the vertex  $f$  to the vertex  $g$  is exactly a homotopy  $G \otimes I_m \rightarrow H$  from  $f$  to  $g$ . For relative graph maps  $f, g: (G, A) \rightarrow (H, B)$ , a path of length  $m$  from  $f$  to  $g$  in the pullback graph

$$\begin{array}{ccc} \bullet & \longrightarrow & \text{hom}^\otimes(A, B) \\ \downarrow & \lrcorner & \downarrow \\ \text{hom}^\otimes(G, H) & \longrightarrow & \text{hom}^\otimes(A, H) \end{array}$$

is exactly a relative homotopy from  $f$  to  $g$ .

It follows that relative homotopy is an equivalence relation on relative graph maps.

PROPOSITION 1.15. *Relative homotopy is an equivalence relation on the set of relative graph maps  $(G, A) \rightarrow (H, B)$ .*

Given a relative graph map  $f: (G, A) \rightarrow (H, B)$ , if the subgraph  $B$  consists of a single vertex  $v$  then we refer to  $f$  as a graph map *based at  $v$*  or a *based graph map*. We refer to a homotopy between two such maps as a *based homotopy*.

DEFINITION 1.16. Let  $G$  be a graph and  $n \geq 1$ . For  $i = 0, \dots, n$  and  $\varepsilon = 0, 1$ , a map  $f: I_\infty^{\otimes n} \rightarrow G$  is *stable in direction  $(i, \varepsilon)$*  if there exists  $M \geq 0$  so that for  $v_i > M$  we have

$$f(v_1, \dots, v_{i-1}, (2\varepsilon - 1)v_i, v_{i+1}, \dots, v_n) = f(v_1, \dots, v_{i-1}, (2\varepsilon - 1)M, v_{i+1}, \dots, v_n).$$

In words, a map  $f: I_\infty^{\otimes n} \rightarrow G$  is stable in direction  $(i, 0)$  if it becomes constant (with respect to change in the  $i$ th coordinate) once the  $i$ th coordinate is sufficiently large in the negative direction. It is stable in direction  $(i, 1)$  if it becomes constant (again, with respect to change in the  $i$ th coordinate) once the  $i$ th coordinate is sufficiently large in the positive direction.

For  $n, M \geq 0$ , let  $I_{\geq M}^{\otimes n}$  denote the subgraph of  $I_\infty^{\otimes n}$  consisting of vertices  $(v_1, \dots, v_n)$  such that  $|v_i| \geq M$  for some  $i = 1, \dots, n$ . Given a based graph map  $f: (I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v)$  we may also regard  $f$  as a based graph map  $(I_\infty^{\otimes n}, I_{\geq K}^{\otimes n}) \rightarrow (G, v)$  for any  $K \geq M$ . This gives a notion of based homotopy between maps  $(I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v)$  which, for some  $M \geq 0$ , are based at  $v$ .

PROPOSITION 1.17 [Mal83, Proposition 3.2]. *Based homotopy is an equivalence relation on the set of based maps*

$$\{(I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v) \mid M \geq 0\}.$$

DEFINITION 1.18. Let  $n \geq 0$  and  $v \in G$  be a vertex of a graph  $G$ . The  $n$ th  $A$ -homotopy group of  $G$  at  $v$  is the set of based homotopy classes of maps  $(I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v)$  based at  $v$  for some  $M \geq 0$ .

Let  $n \geq 1$  and  $i = 1, \dots, n$ . Given  $f: (I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v)$  and  $g: (I_\infty^{\otimes n}, I_{\geq M'}^{\otimes n}) \rightarrow (G, v)$ , we define a binary operation  $f \cdot_i g: (I_\infty^{\otimes n}, I_{\geq M+M'}^{\otimes n}) \rightarrow (G, v)$  by

$$(f \cdot_i g)(v_1, \dots, v_n) := \begin{cases} f(v_1, \dots, v_n) & v_i \leq M \\ g(v_1, \dots, v_i - M - M', \dots, v_n) & v_i > M. \end{cases}$$

This induces a group operation on homotopy groups  $\cdot_i: A_n(G, v) \times A_n(G, v) \rightarrow A_n(G, v)$ . For  $n \geq 2$  and  $1 \leq i < j \leq n$ , it is straightforward to construct a homotopy witnessing that

$$[[f] \cdot_i [g]] \cdot_j [[f] \cdot_i [g]] = [[f] \cdot_j [g]] \cdot_i [[f] \cdot_j [g]]$$

for any  $[f], [g] \in A_n(G, v)$ . The Eckmann–Hilton argument gives  $[f] \cdot_i [g] = [f] \cdot_j [g]$  and that this operation is abelian.

**Path and loop graphs**

DEFINITION 1.19. For a graph  $G$ , we define the *path graph*  $PG$  to be the induced subgraph of  $\text{hom}^\otimes(I_\infty, G)$  consisting of maps which stabilize in the  $(1, 0)$  and  $(1, 1)$  directions.

As vertices of  $PG$  are paths that stabilize, we have graph maps  $\partial_{1,0}, \partial_{1,1}: PG \rightarrow G$  which send a vertex  $v: I_\infty \rightarrow G$  to its left and right endpoints, respectively.

PROPOSITION 1.20. *For a graph  $G$ , we have an isomorphism*

$$PG \cong \text{colim} \left( \text{hom}^\otimes(I_1, G) \xrightarrow{l^*} \text{hom}^\otimes(I_2, G) \xrightarrow{r^*} \text{hom}^\otimes(I_3, G) \xrightarrow{l^*} \dots \right)$$

*natural in  $G$ .*

*Proof.* For each  $m \geq 1$ , we write  $m = 2k + b$ , where  $k \geq 0$  and  $b \in \{0, 1\}$ , and define a graph map  $I_\infty \rightarrow I_m$  by

$$v \mapsto \begin{cases} 0 & v \leq -k \\ k + v & -k \leq v \leq k + b \\ m & v \geq k + b. \end{cases}$$

Geometrically, this function maps the subinterval  $[-k, k + b]$  surjectively onto  $I_m$ , and collapses all other vertices to the endpoints. By pre-composition, this induces a cone

$$\begin{array}{ccccccc} \text{hom}^\otimes(I_1, G) & \xrightarrow{l^*} & \text{hom}^\otimes(I_2, G) & \xrightarrow{r^*} & \text{hom}^\otimes(I_3, G) & \xrightarrow{l^*} & \dots \\ & \searrow & \downarrow & \swarrow & & & \\ & & PG & & & & \end{array}$$

which one verifies is a colimit cone. □

DEFINITION 1.21. For a pointed graph  $(G, v)$ , the *loop graph*  $\Omega(G, v)$  is the subgraph of  $PG$  of paths  $I_\infty \rightarrow G$  whose left and right endpoints are  $v$ .

From the definition, the following proposition is immediate.

PROPOSITION 1.22. For a pointed graph  $(G, v)$ , the square

$$\begin{array}{ccc} \Omega(G, v) & \longrightarrow & PG \\ \downarrow & \lrcorner & \downarrow (\partial_{1,0}, \partial_{1,1}) \\ I_0 & \xrightarrow{(v,v)} & G \times G \end{array}$$

is a pullback.

The loop graph of a pointed graph  $(G, v)$  has a distinguished vertex which is the constant path at  $v$ . This gives an endofunctor  $\Omega: \mathbf{Graph}_* \rightarrow \mathbf{Graph}_*$ . From this, we define the notion of  $n$ th loop graphs.

DEFINITION 1.23. For  $n \geq 0$ , we define the  $n$ th loop graph to be

$$\Omega^n(X, x) := \begin{cases} (X, x) & n = 0 \\ \Omega(\Omega^{n-1}(X, x)) & \text{otherwise.} \end{cases}$$

In [BBdLL06], it is shown that the  $n$ th homotopy groups of a graph correspond to the connected components of  $\Omega^n(G, v)$ .

PROPOSITION 1.24 [BBdLL06, Proposition 7.4]. For  $n \geq 0$ , we have an isomorphism

$$A_n(G, v) \cong A_0(\Omega^n(G, v)).$$

## 2. Cubical sets and their homotopy theory

### Cubical sets

We begin by defining the box category  $\square$ . As used in this paper, the box category will include both positive and negative connections. This variant of the box category was introduced in [BH81] and was later used in [DKLS24] to model  $(\infty, 1)$ -categories. The objects of  $\square$  are posets of the form  $[1]^n = \{0 \leq 1\}^n$  and the maps are generated (inside the category of posets) under composition by the following four special classes:

- faces  $\partial_{i,\varepsilon}^n: [1]^{n-1} \rightarrow [1]^n$  for  $i = 1, \dots, n$  and  $\varepsilon = 0, 1$  given by

$$\partial_{i,\varepsilon}^n(x_1, x_2, \dots, x_{n-1}) = (x_1, x_2, \dots, x_{i-1}, \varepsilon, x_i, \dots, x_{n-1});$$

- degeneracies  $\sigma_i^n: [1]^n \rightarrow [1]^{n-1}$  for  $i = 1, 2, \dots, n$  given by

$$\sigma_i^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n);$$

- negative connections  $\gamma_{i,0}^n: [1]^n \rightarrow [1]^{n-1}$  for  $i = 1, 2, \dots, n - 1$  given by

$$\gamma_{i,0}^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, \max\{x_i, x_{i+1}\}, x_{i+2}, \dots, x_n).$$

- positive connections  $\gamma_{i,1}^n: [1]^n \rightarrow [1]^{n-1}$  for  $i = 1, 2, \dots, n - 1$  given by

$$\gamma_{i,1}^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, \min\{x_i, x_{i+1}\}, x_{i+2}, \dots, x_n).$$

These maps obey the following *cubical identities*:

$$\begin{aligned} \partial_{j,\varepsilon'}\partial_{i,\varepsilon} &= \partial_{i+1,\varepsilon}\partial_{j,\varepsilon'} \quad \text{for } j \leq i; & \sigma_j\partial_{i,\varepsilon} &= \begin{cases} \partial_{i-1,\varepsilon}\sigma_j & \text{for } j < i \\ \text{id} & \text{for } j = i \\ \partial_{i,\varepsilon}\sigma_{j-1} & \text{for } j > i; \end{cases} \\ \sigma_i\sigma_j &= \sigma_j\sigma_{i+1} \quad \text{for } j \leq i; & \gamma_{j,\varepsilon'}\gamma_{i,\varepsilon} &= \begin{cases} \gamma_{i,\varepsilon}\gamma_{j+1,\varepsilon'} & \text{for } j > i \\ \gamma_{i,\varepsilon}\gamma_{i+1,\varepsilon} & \text{for } j = i, \varepsilon' = \varepsilon; \end{cases} \\ \gamma_{j,\varepsilon'}\partial_{i,\varepsilon} &= \begin{cases} \partial_{i-1,\varepsilon}\gamma_{j,\varepsilon'} & \text{for } j < i-1 \\ \text{id} & \text{for } j = i-1, i, \varepsilon = \varepsilon' \\ \partial_{j,\varepsilon}\sigma_j & \text{for } j = i-1, i, \varepsilon = 1-\varepsilon' \\ \partial_{i,\varepsilon}\gamma_{j-1,\varepsilon'} & \text{for } j > i; \end{cases} & \sigma_j\gamma_{i,\varepsilon} &= \begin{cases} \gamma_{i-1,\varepsilon}\sigma_j & \text{for } j < i \\ \sigma_i\sigma_i & \text{for } j = i \\ \gamma_{i,\varepsilon}\sigma_{j+1} & \text{for } j > i. \end{cases} \end{aligned}$$

This category enjoys many good properties, making it suitable for modeling homotopy theory. Formally speaking, these properties can be encapsulated in saying that  $\square$  is an Eilenberg–Zilber category, which in particular implies that the object  $[1]^n$  has no non-identity automorphisms. However, we will not explicitly rely on the notion of Eilenberg–Zilber categories.

A *cubical set* is a presheaf  $X: \square^{\text{op}} \rightarrow \mathbf{Set}$ . A *cubical map* is a natural transformation of such presheaves. We write  $\mathbf{cSet}$  for the category of cubical sets and cubical maps.

Given a cubical set  $X$ , we write  $X_n$  for the value of  $X$  at  $[1]^n$  and refer to the elements of  $X_n$  as *n-cubes* of  $X$ . We write cubical operators on the right, e.g. given an  $n$ -cube  $x \in X_n$  of  $X$ , we write  $x\partial_{1,0}$  for the  $\partial_{1,0}$ -face of  $x$ .

By a *degenerate cube*, we always mean a cube that is in the image of a degeneracy or a connection map. This nomenclature is borrowed from the theory of Reedy categories, where one can speak abstractly of degenerate elements in a presheaf ([RV14], [Hov99, § 5.2], [Hir03, Ch. 15]).

DEFINITION 2.1. Let  $n \geq 0$ .

- The *combinatorial n-cube*  $\square^n$  is the representable functor  $\square(-, [1]^n): \square^{\text{op}} \rightarrow \mathbf{Set}$ ,
- The *boundary of the n-cube*  $\partial\square^n$  is the subobject of  $\square^n$  defined by

$$\partial\square^n := \bigcup_{\substack{j=0,\dots,n \\ \eta=0,1}} \text{im } \partial_{j,\eta}.$$

- When  $n \geq 1$ , given  $i = 0, \dots, n$  and  $\varepsilon = 0, 1$ , the  $(i, \varepsilon)$ -open box  $\square_{i,\varepsilon}^n$  is the subobject of  $\partial\square^n$  defined by

$$\square_{i,\varepsilon}^n := \bigcup_{(j,\eta) \neq (i,\varepsilon)} \text{im } \partial_{j,\eta}.$$

Observe  $\square^0 \in \mathbf{cSet}$  is the terminal object in  $\mathbf{cSet}$ .

Example 2.2. Define a functor  $\square \rightarrow \mathbf{Top}$  from the box category to the category of topological spaces which sends  $[1]^n$  to  $[0, 1]^n$ , where  $[0, 1]$  is the unit interval. The face and degeneracy maps are sent to the face inclusion and product projection maps, respectively. The negative connection  $\gamma_{i,0}$  is sent to the map  $[0, 1]^n \rightarrow [0, 1]^{n-1}$  defined by

$$(x_1, \dots, x_n) \mapsto (x_1, \dots, x_{i-1}, \max(x_i, x_{i+1}), x_{i+2}, \dots, x_n),$$

and the image of the positive connection  $\gamma_{i,1}$  is defined analogously.

Left Kan extension along the Yoneda embedding gives the *geometric realization* functor  $|-|: \mathbf{cSet} \rightarrow \mathbf{Top}$ :

$$\begin{array}{ccc} \square & \xrightarrow{[1]^n \mapsto [0,1]^n} & \mathbf{Top} \\ \downarrow & \nearrow \text{\scriptsize } |-| & \\ \mathbf{cSet} & & \end{array}$$

This functor is left adjoint to the *singular cubical complex* functor  $\mathbf{Sing}: \mathbf{Top} \rightarrow \mathbf{cSet}$  defined by

$$(\mathbf{Sing} S)_n := \mathbf{Top}([0, 1]^n, S).$$

Define a functor  $\otimes: \square \times \square \rightarrow \square$  on the cube category which sends  $([1]^m, [1]^n)$  to  $[1]^{m+n}$ . Post-composing with the Yoneda embedding and left Kan extending gives a monoidal product on cubical sets:

$$\begin{array}{ccc} \square \times \square & \longrightarrow & \square & \longrightarrow & \mathbf{cSet} \\ \downarrow & & & \nearrow \otimes & \\ \mathbf{cSet} \times \mathbf{cSet} & & & & \end{array}$$

This is the *geometric product* of cubical sets. Although, for  $[1]^m, [1]^n \in \square$ , there is an isomorphism  $[1]^m \otimes [1]^n \cong [1]^n \otimes [1]^m$ , this isomorphism is not natural. As a result, the geometric product of cubical sets is not symmetric, i.e.  $X \otimes Y$  is not in general isomorphic to  $Y \otimes X$ .

This product is however biclosed. For a cubical set  $X$ , we write  $\mathbf{hom}_L(X, -): \mathbf{cSet} \rightarrow \mathbf{cSet}$  and  $\mathbf{hom}_R(X, -): \mathbf{cSet} \rightarrow \mathbf{cSet}$  for the right adjoints to the functors  $- \otimes X$  and  $X \otimes -$ , respectively. As the geometric product is not symmetric, the functors  $\mathbf{hom}_L(X, -)$  and  $\mathbf{hom}_R(X, -)$  are not naturally isomorphic.

### Kan complexes

DEFINITION 2.3.

- (i) A cubical map  $X \rightarrow Y$  is a *Kan fibration* if it has the right lifting property with respect to open box inclusions. That is, if for any commutative square,

$$\begin{array}{ccc} \square^n_{i,\varepsilon} & \longrightarrow & X \\ \downarrow & & \downarrow f \\ \square^n & \longrightarrow & Y \end{array}$$

there exists a map  $\square^n \rightarrow X$  so that the triangles

$$\begin{array}{ccc} \square^n_{i,\varepsilon} & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \square^n & \longrightarrow & Y \end{array}$$

commute.

- (ii) A cubical set  $X$  is a *Kan complex* if the unique map  $X \rightarrow \square^0$  is a Kan fibration.

We write **Kan** for the full subcategory of **cSet** consisting of Kan complexes.

*Example 2.4.* For any  $S \in \mathbf{Top}$ , the cubical set  $\mathbf{Sing} S$  is a Kan complex. A map  $\square^n_{i,\varepsilon} \rightarrow \mathbf{Sing} S$  is, by adjointness, a map  $|\square^n_{i,\varepsilon}| \rightarrow S$ . The inclusion  $|\square^n_{i,\varepsilon}| \hookrightarrow |\square^n|$  has a retract in **Top**. Pre-composing

with this retract gives a map  $|\square^n| \rightarrow S$  which restricts to the open box map  $|\square^n_{i,\varepsilon}| \rightarrow S$ :

$$\begin{array}{ccc} |\square^n_{i,\varepsilon}| & \longrightarrow & S \\ \downarrow \wr & & \\ |\square^n| & & \end{array}$$

By adjointness, this gives a suitable map  $\square^n \rightarrow \text{Sing } S$ .

**DEFINITION 2.5.** A map  $f: X \rightarrow Y$  is a *weak equivalence* if the map  $|f|: |X| \rightarrow |Y|$  is a weak homotopy equivalence, i.e. for any  $n \geq 0$  and  $x \in |X|$ , the map  $\pi_n|f|: \pi_n(|X|, x) \rightarrow \pi_n(|Y|, |f|(x))$  is an isomorphism.

We move towards describing the fibration category of Kan complexes.

**DEFINITION 2.6** [Bro73, Definition 1.1]. A *fibration category* is a category  $\mathcal{C}$  with two subcategories of *fibrations* and *weak equivalences* such that (in what follows, an *acyclic fibration* is a map that is both a fibration and a weak equivalence) the following statements hold:

- (i) weak equivalences satisfy the two-out-of-three property, i.e. given two composable morphisms

$$X \xrightarrow{f} Y \xrightarrow{g} Z.$$

if two of  $f, g, gf$  are weak equivalences then all three are;

- (ii) all isomorphisms are acyclic fibrations;
- (iii) pullbacks along fibrations exist; fibrations and acyclic fibrations are stable under pullback;
- (iv)  $\mathcal{C}$  has a terminal object  $1$ ; the canonical map  $X \rightarrow 1$  is a fibration for any object  $X \in \mathcal{C}$  (i.e. all objects are *fibrant*);
- (v) every map can be factored as a weak equivalence followed by a fibration.

*Example 2.7* [Hov99, Theorem 2.4.19]. The category  $\text{Top}$  of topological spaces is a fibration where:

- fibrations are Serre fibrations;
- weak equivalences are weak homotopy equivalences, i.e. maps  $f: S \rightarrow S'$  such that, for all  $s \in S$  and  $n \geq 0$ , the map  $\pi_n f: \pi_n(S, s) \rightarrow \pi_n(S', f(s))$  is an isomorphism.

**DEFINITION 2.8.** A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between fibration categories is *exact* if it preserves fibrations, acyclic fibrations, pullbacks along fibrations, and the terminal object.

Given a fibration category  $\mathcal{C}$  with finite coproducts and a terminal object, the category of pointed objects  $1 \downarrow \mathcal{C}$  is a fibration category as well.

**PROPOSITION 2.9.** *Let  $\mathcal{C}$  be a fibration category with finite coproducts and a terminal object  $1 \in \mathcal{C}$ .*

- (i) *The slice category  $1 \downarrow \mathcal{C}$  under  $1$  is a fibration category where a map is a fibration/weak equivalence if the underlying map in  $\mathcal{C}$  is.*
- (ii) *The projection functor  $1 \downarrow \mathcal{C} \rightarrow \mathcal{C}$  is exact.*

*Proof.* Statement (i) follows from [Hov99, Proposition 1.1.8].

For statement (ii), the projection functor preserves fibrations/weak equivalences by definition. It is a right adjoint to the functor  $-\sqcup 1: \mathcal{C} \rightarrow 1 \downarrow \mathcal{C}$  which adds a disjoint basepoint, hence preserves finite limits. □

THEOREM 2.10.

- (i) The category  $\mathbf{Kan}$  of Kan complexes is a fibration category where fibrations are Kan fibrations and weak equivalences are as defined above.
- (ii)  $\text{Sing} : \mathbf{Top} \rightarrow \mathbf{Kan}$  is an exact functor
- (iii) The category  $\mathbf{Kan}_*$  of pointed Kan complexes is a fibration category where a map is a fibration/weak equivalence if the underlying map in  $\mathbf{Kan}$  is.

*Proof.* (i) This is shown in [CK23, Theorem 2.17].

(ii) This is [CK23, Corollary 2.25].

(iii) This follows from Proposition 2.9. □

**Anodyne maps**

We move towards defining anodyne maps of cubical sets, i.e. maps which are both monomorphisms and weak equivalences.

DEFINITION 2.11. A class of morphisms  $S$  in a cocomplete category  $\mathcal{C}$  is *saturated* if it is closed under

- pushouts: if  $s: A \rightarrow B$  is in  $S$  and  $f: A \rightarrow C$  is any map then the pushout  $C \rightarrow B \cup C$  of  $s$  along  $f$  is in  $S$ ;

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ s \downarrow & & \downarrow \in S \\ B & \longrightarrow & B \cup C \end{array}$$

- retracts: if  $s: A \rightarrow B$  is in  $S$  and  $r: C \rightarrow D$  is a retract of  $s$  in  $\mathcal{C}^{[1]}$  then  $r$  is in  $S$ ;

$$\begin{array}{ccccc} & & \text{id}_A & & \\ & \curvearrowright & & \curvearrowleft & \\ A & \longrightarrow & B & \longrightarrow & A \\ r \downarrow & & \downarrow s & & \downarrow r \\ C & \longrightarrow & D & \longrightarrow & C \\ & \curvearrowleft & \text{id}_C & \curvearrowright & \end{array}$$

- transfinite composition: given a limit ordinal  $\lambda$  and a diagram  $\lambda \rightarrow \mathcal{C}$  whose morphisms lie in  $S$ ,

$$A_1 \xrightarrow{s_1} A_2 \xrightarrow{s_2} A_3 \xrightarrow{s_3} \dots$$

writing  $A$  for the colimit of this diagram, the components of the colimit cone  $\lambda_i: A_i \rightarrow A$  are in  $S$ .

$$\begin{array}{ccccccc} A_1 & \longrightarrow & A_2 & \longrightarrow & A_3 & \longrightarrow & \dots \\ & \searrow & \downarrow \lambda_2 & \swarrow & & & \\ & \lambda_1 & A & \lambda_3 & & & \end{array}$$

DEFINITION 2.12. For a set of morphisms  $S$  in a cocomplete category  $\mathcal{C}$ , the *saturation*  $\text{Sat } S$  of  $S$  is the smallest saturated class containing  $S$ .

For a saturated class of maps, closure under pushouts and transfinite composition gives the following proposition.



PROPOSITION 2.13 [Hov99, Lemma 2.1.13]. *Saturated classes of maps are closed under coproduct. That is, given a collection  $\{s_i: A_i \rightarrow B_i \mid i \in I\}$  of morphisms in a saturated class  $S$ , the coproduct*

$$\coprod_{i \in I} s_i: \coprod_{i \in I} A_i \rightarrow \coprod_{i \in I} B_i$$

is in  $S$ .

DEFINITION 2.14. A map of cubical sets is *anodyne* if it is in the saturation of open box inclusions

$$\text{Sat}\{\square_{i,\varepsilon}^n \hookrightarrow \square^n \mid n \geq 1, i = 0, \dots, n, \varepsilon = 0, 1\}.$$

We use the following property of anodyne maps, which follows since saturations are closed under the left lifting property.

THEOREM 2.15 [DKLS24, Theorem 1.34]. *Let  $g: A \rightarrow B$  be an anodyne map and  $f: X \rightarrow Y$  be a Kan fibration. Given a commutative square,*

$$\begin{array}{ccc} A & \longrightarrow & X \\ g \downarrow & & \downarrow f \\ B & \longrightarrow & Y \end{array}$$

there exists a map  $B \rightarrow X$  so that the triangles

$$\begin{array}{ccc} A & \longrightarrow & X \\ g \downarrow & \nearrow & \downarrow f \\ B & \longrightarrow & Y \end{array}$$

commute.

THEOREM 2.16. *A cubical map is anodyne if and only if it is a monomorphism and a weak equivalence.*

*Proof.* This follows from [DKLS24, Theorem 1.34], as the saturation of open box inclusions is exactly the class of maps which have the left lifting property with respect to fibrations.  $\square$

In particular, we use that anodyne maps are sent to weak homotopy equivalences under geometric realization.

COROLLARY 2.17. *If  $f: X \rightarrow Y$  is anodyne then, for all  $n \geq 0$  and  $x \in |X|$ , the map  $\pi_n|f|: \pi_n(|X|, x) \rightarrow \pi_n(|Y|, |f|(x))$  is an isomorphism.*

**Homotopies and homotopy groups**

Using the geometric product, we may define a notion of homotopy between cubical maps.

DEFINITION 2.18. Given cubical maps  $f, g: X \rightarrow Y$ , a *homotopy* from  $f$  to  $g$  is a map  $H: X \otimes \square^1 \rightarrow Y$  such that the diagram

$$\begin{array}{ccc} X \otimes \square^0 & & \\ \partial_{1,0} \downarrow & \searrow f & \\ X \otimes \square^1 & \xrightarrow{H} & Y \\ \partial_{1,1} \uparrow & \nearrow g & \\ X \otimes \square^0 & & \end{array}$$

commutes.

Let  $\mathbf{cSet}_2$  denote the full subcategory of  $\mathbf{cSet}^{[1]}$  spanned by monomorphisms. Explicitly, its objects are monic cubical maps  $A \hookrightarrow X$ . A morphism from  $A \hookrightarrow X$  to  $B \hookrightarrow Y$  is a pair of maps  $(f, g)$  which form a commutative square of the following form:

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

We refer to the objects and morphisms of  $\mathbf{cSet}_2$  as *relative cubical sets* and *relative cubical maps*, respectively. Following our convention for relative graph maps, we denote a relative cubical set  $A \hookrightarrow X$  by  $(X, A)$ , suppressing the data of the map itself. If the domain of the map  $A \hookrightarrow X$  is the 0-cube  $A = \square^0$ , we denote it by  $(X, x)$ , where  $x$  is the unique 0-cube in the image of the map  $\square^0 \hookrightarrow X$ .

For a relative cubical map  $(f, g): (X, A) \rightarrow (Y, B)$ , the map  $g$  is uniquely determined by  $f$  since  $B \hookrightarrow Y$  is monic. As a result, we denote a relative cubical map by  $f: (X, A) \rightarrow (Y, B)$ . Mirroring our convention for relative graph maps, we write  $f$  for the map  $X \rightarrow Y$  and  $f|_A$  for the map  $A \rightarrow B$ .

We have a corresponding notion of homotopy between relative cubical maps.

**DEFINITION 2.19.** Let  $f, g: (X, A) \rightarrow (Y, B)$  be relative cubical maps. A *relative homotopy* from  $f$  to  $g$  is a morphism  $(X \otimes \square^1, A \otimes \square^1) \rightarrow (Y, B)$  in  $\mathbf{cSet}_2$  such that:

- the map  $A \otimes \square^1 \rightarrow B$  is a homotopy from  $f|_A$  to  $g|_A$ ;
- the map  $X \otimes \square^1 \rightarrow Y$  is a homotopy from  $f$  to  $g$ .

**PROPOSITION 2.20** [CK23, Proposition 2.30]. *If  $B, Y$  are Kan complexes then relative homotopy is an equivalence relation on relative cubical maps  $(X, A) \rightarrow (B, Y)$ .*

It is essential in Proposition 2.20 that  $B$  and  $Y$  are Kan complexes. If not, the relation of relative homotopy is neither symmetric nor transitive (and in this case, one considers the symmetric transitive closure of relative homotopy).

We write  $[(X, A), (Y, B)]$  for the set of relative homotopy classes of relative maps  $(X, A) \rightarrow (Y, B)$ . With this, we define the homotopy groups of a Kan complex.

**DEFINITION 2.21** [CK23, Corollary 3.16]. Let  $(X, x)$  be a pointed Kan complex. We define the  $n$ th homotopy group  $\pi_n(X, x)$  of  $(X, x)$  as the relative homotopy classes of relative maps  $(\square^n, \partial \square^n) \rightarrow (X, x)$ .

We give an explicit description of multiplication in the first homotopy group  $\pi_1(X, x)$ .

**DEFINITION 2.22.** Given two 1-cubes  $u, v: \square^1 \rightarrow X$  in a cubical set  $X$ ,

- (i) a *concatenation square* for  $u$  and  $v$  is a map  $\eta: \square^2 \rightarrow X$  such that
  - $\eta \partial_{1,0} = u$ ,
  - $\eta \partial_{1,1} = v \partial_{1,1} \sigma_1$ ,
  - $\eta \partial_{2,1} = v$ .
- (ii) a *concatenation* of  $u$  and  $v$  is a 1-cube  $w: \square^1 \rightarrow X$  which is the  $\partial_{2,0}$ -face of some concatenation square for  $u$  and  $v$ .

**PROPOSITION 2.23** [CK23, Theorem 3.11]. *If  $(X, x)$  is a pointed Kan complex then composition induces a well-defined binary operation*

$$[(\square^1, \partial \square^1), (X, x)] \times [(\square^1, \partial \square^1), (X, x)] \rightarrow [(\square^1, \partial \square^1), (X, x)]$$

on relative homotopy classes of relative maps which gives a group structure on  $[(\square^1, \partial \square^1), (X, x)]$ .

As with spaces, the homotopy groups of a Kan complex are the connected components of its loop space, which we define.

DEFINITION 2.24. For a pointed Kan complex  $(X, x)$ ,

- the loop space  $\Omega(X, x)$  of  $X$  is the pullback

$$\begin{array}{ccc} \Omega(X, x) & \longrightarrow & \text{hom}(\square^1, X) \\ \downarrow & \lrcorner & \downarrow (\partial_{1,0}^*, \partial_{1,1}^*) \\ \square^0 & \xrightarrow{x} & X \times X \end{array}$$

with a distinguished 0-cube  $x\sigma_1: \square^0 \rightarrow \Omega(X, x)$ .

- for  $n \geq 0$ , the  $n$ th loop space  $\Omega^n(X, x)$  of  $X$  is defined to be

$$\Omega^n(X, x) := \begin{cases} (X, x) & n = 0 \\ \Omega(\Omega^{n-1}(X, x)) & n > 0. \end{cases}$$

PROPOSITION 2.25 [CK23, Corollary 3.16]. For a pointed Kan complex  $(X, x)$  and  $0 \leq k \leq n$ , we have an isomorphism

$$\pi_n(X, x) \cong \pi_{n-k}(\Omega^k(X, x))$$

natural in  $X$ .

Using Proposition 2.25, the group structure on higher homotopy groups is induced by the bijection  $\pi_n(X, x) \cong \pi_1(\Omega^{n-1}(X, x))$ .

This definition of homotopy groups agrees with the homotopy groups of its geometric realization.

THEOREM 2.26 [CK23, Theorem 3.25]. There is an isomorphism

$$\pi_n(X, x) \cong \pi_n(|X|, x)$$

natural in  $X$ .

We know that the loop space functor is exact.

THEOREM 2.27 [CK23, Theorem 3.6]. The loop space functor  $\Omega: \text{Kan}_* \rightarrow \text{Kan}_*$  is exact.

### 3. Cubical nerve of a graph

Let  $m \geq 1$ . Geometrically, we view the graph  $I_m^{\otimes n}$  as an  $n$ -dimensional cube. Making this intuition formal, we have face, degeneracy, and connection maps defined as follows.

- the face map  $\partial_{i,\varepsilon}^n: I_m^{\otimes n-1} \rightarrow I_m^{\otimes n}$  for  $1 \leq i \leq n$  and  $\varepsilon = 0$  or  $1$  is given by

$$\partial_{i,\varepsilon}^n(v_1, \dots, v_n) = (v_1, \dots, v_{i-1}, \varepsilon m, v_i, \dots, v_n);$$

- the degeneracy map  $\sigma_i^n: I_m^{\otimes n+1} \rightarrow I_m^{\otimes n}$  for  $1 \leq i \leq n$  is given by

$$\sigma_i^n(v_1, \dots, v_n) = (v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n);$$


- the negative connection map  $\gamma_{i,0}^n: I_m^{\otimes n} \rightarrow I_m^{\otimes n-1}$  for  $1 \leq i \leq n - 1$  is given by

$$\gamma_{i,0}^n(v_1, \dots, v_n) = (v_1, \dots, v_{i-1}, \max(v_i, v_{i+1}), v_{i+2}, \dots, v_n);$$

– the positive connection map  $\gamma_{i,1}^n: I_m^{\otimes n} \rightarrow I_m^{\otimes n-1}$  for  $1 \leq i \leq n - 1$  is given by

$$\gamma_{i,1}^n(v_1, \dots, v_n) = (v_1, \dots, v_{i-1}, \min(v_i, v_{i+1}), v_{i+2}, \dots, v_n).$$

It is straightforward to verify that these maps satisfy cubical identities. This defines a functor  $\square \rightarrow \mathbf{Graph}$  which sends  $[1]^n$  to  $I_m^{\otimes n}$ . Left Kan extension along the Yoneda embedding gives an adjunction  $\mathbf{cSet} \rightleftarrows \mathbf{Graph}$ :



DEFINITION 3.1. For  $m \geq 1$ ,

- (i) the  $m$ -realization functor  $|-|_m: \mathbf{cSet} \rightarrow \mathbf{Graph}$  is the left Kan extension of the functor  $\square \rightarrow \mathbf{Graph}$  which sends  $[1]^n$  to  $I_m^{\otimes n}$ .
- (ii) the  $m$ -nerve functor  $N_m: \mathbf{Graph} \rightarrow \mathbf{cSet}$  is the right adjoint of the  $m$ -realization functor defined by

$$(N_m G)_n := \mathbf{Graph}(I_m^{\otimes n}, G).$$

Remark 3.2. The 1-nerve of a graph  $G$  is constructed in [BBdLL06] as the *cubical set associated to  $G$* , denoted  $M_*(G)$ . The geometric realization of this cubical set is referred to as the *cell complex associated to  $G$* , denoted  $X_G$ .

For a cubical set  $X \in \mathbf{cSet}$ , the graph  $|X|_m$  may be explicitly described as the colimit

$$|X|_m := \text{colim}(G_{m,0} \hookrightarrow G_{m,1} \hookrightarrow G_{m,2} \hookrightarrow \dots)$$

where:

- $G_{m,0}$  is the discrete graph whose vertices are  $X_0$ ;
- $G_{m,n+1}$  is obtained from  $G_{m,n}$  via the following pushout (where  $(X_n)_{\text{nd}}$  is the subset of  $X_n$  consisting of non-degenerate cubes)

$$\begin{array}{ccc} \coprod_{x \in (X_n)_{\text{nd}}} \partial I_m^{\otimes n} & \longrightarrow & G_{m,n} \\ \downarrow & & \downarrow \\ \coprod_{x \in (X_n)_{\text{nd}}} I_m^{\otimes n} & \longrightarrow & G_{m,n+1} \end{array} \quad \lrcorner$$

in which  $\partial I_m^{\otimes n}$  is the subgraph of  $I_m^{\otimes n}$  defined by

$$\partial I_m^{\otimes n} := \{(v_1, \dots, v_n) \in I_m^{\otimes n} \mid v_i = 0 \text{ or } m \text{ for some } i = 0, \dots, n\}$$

where the edge set is discrete if  $m = n = 1$  and full otherwise.

Example 3.3.

- (i) We describe the graph  $|\square_{2,1}^2|_1$ . The cubical set  $\square_{2,1}^2$  has four 0-cubes; thus, the graph  $G_{1,0}$  is the discrete graph with four vertices. The open box  $\square_{2,1}^2$  has three non-degenerate 1-cubes. The pushout constructed to obtain  $G_{1,1}$  glues three copies of  $I_1$  to  $G_{1,0}$  (Figure 3). As  $\square_{2,1}^2$  contains only degenerate cubes above dimension 1, constructing  $G_{1,1}$  completes the construction of the graph  $|\square_{2,1}^2|_1$ .
- (ii) To construct  $|\square_{2,1}^2|_3$ , we instead glue three copies of  $I_3$  (Figure 4). Observe that this process adds new vertices to the graph.




FIGURE 3. The embedding of  $G_{1,0} \hookrightarrow G_{1,1}$  for  $|\square_{2,1}^2|_1$ .




FIGURE 4. The graph  $|\square_{2,1}^2|_3$ . The image of the embedding  $G_{3,0} \hookrightarrow |\square_{2,1}^2|_3$  is shaded.

For a graph  $G$ , let  $l^*, r^*: N_m G \rightarrow N_{m+1} G$  denote the cubical maps obtained by precomposition with the surjections  $l^{\otimes n}, r^{\otimes n}: I_{m+1}^{\otimes n} \rightarrow I_m^{\otimes n}$  (Figure 5). We think of these maps as inclusions of  $n$ -cubes of size  $m$  into  $n$ -cubes of size  $m + 1$  (Figure 6). We write  $c^*: N_m G \rightarrow N_{m+2} G$  for the composite  $l^* r^* = r^* l^*$ .




FIGURE 5 (colour online). For a 1-cube  $f: I_1 \rightarrow G$  of  $N_1 G$ , the map  $l^*: N_1 G \rightarrow N_2 G$  sends  $f$  to the 1-cube  $fl: I_2 \rightarrow G$  of  $N_2 G$ , whereas  $r^*: N_1 G \rightarrow N_2 G$  sends  $f$  to the 1-cube  $fr: I_2 \rightarrow G$  of  $N_2 G$ .




FIGURE 6 (colour online). For a 2-cube  $g: I_1^{\otimes 2} \rightarrow G$  of  $N_1 G$ , the map  $l^*: N_1 G \rightarrow N_2 G$  sends  $g$  to the 2-cube  $gl^{\otimes 2}: I_2^{\otimes 2} \rightarrow G$  of  $N_2 G$ , whereas  $r^*: N_1 G \rightarrow N_2 G$  sends  $g$  to the 2-cube  $gr^{\otimes 2}: I_2^{\otimes 2} \rightarrow G$  of  $N_2 G$ .

*Remark 3.4.* While the maps  $l^{\otimes n}, r^{\otimes n}$  have sections  $I_m^{\otimes n} \rightarrow I_{m+1}^{\otimes n}$ , these maps do not commute with face maps, hence do not give retractions  $N_{m+1} G \rightarrow N_m G$ . To demonstrate this, we show the map  $l^*: N_1 I_2 \rightarrow N_2 I_2$  does not have a retraction. The identity map  $\text{id}_{I_2}: I_2 \rightarrow I_2$  gives a 1-cube of  $N_2 I_2$  whose faces are the 0-cubes 0 and 2. Observe that a retraction of  $l^*$  must send the 0-cubes 0 and 2 to 0 and 2, respectively. There is no map  $f: I_1 \rightarrow I_2$  such that  $f\partial_{1,0} = 0$  and  $f\partial_{1,1} = 2$ . That is, there is no 1-cube of  $N_1 I_2$  which  $\text{id}_{I_2} \in (N_2 I_2)_1$  can be mapped to. Thus, the map  $l^*$  does not have a retraction.

For a cubical set  $X$ , we analogously have maps  $l_*, r_*: |X|_{m+1} \rightarrow |X|_m$ . We write  $c_*: |X|_{m+2} \rightarrow |X|_m$  for the composite  $l_* r_* = r_* l_*$ .

We define the *nerve* functor  $N: \text{Graph} \rightarrow \text{cSet}$  by

$$(NG)_n := \{f: I_\infty^{\otimes n} \rightarrow G \mid f \text{ is stable in all directions } (i, \varepsilon)\}.$$

Cubical operators of  $NG$  are given as follows.

- The map  $\partial_{i,\varepsilon}^n: (NG)_n \rightarrow (NG)_{n-1}$  for  $i = 1, \dots, n$  and  $\varepsilon = 0, 1$  is given by  $f\partial_{i,\varepsilon}^n: I_\infty^{\otimes n-1} \rightarrow G$  defined by

$$f\partial_{i,\varepsilon}^n(v_1, \dots, v_{n-1}) = f(v_1, \dots, v_{i-1}, (2\varepsilon - 1)M, v_i, \dots, v_{n-1})$$

where  $M$  is such that  $f$  is stable in direction  $(i, \varepsilon)$ .

- The map  $\sigma_i^n: (NG)_n \rightarrow (NG)_{n+1}$  for  $i = 1, \dots, n$  is given by  $f\sigma_i^n: I_\infty^{\otimes n+1} \rightarrow G$  defined by

$$f\sigma_i^n(v_1, \dots, v_{n+1}) = f(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_{n+1}).$$

- The map  $\gamma_{i,0}^n: (NG)_n \rightarrow (NG)_{n+1}$  for  $i = 1, \dots, n - 1$  is given by  $f\gamma_{i,0}^n: I_\infty^{\otimes n+1} \rightarrow G$  defined by

$$f\gamma_{i,0}^n(v_1, \dots, v_{n+1}) = f(v_1, \dots, v_{i-1}, \max(v_i, v_{i+1}), v_{i+1}, \dots, v_{n+1}).$$

- The map  $\gamma_{i,1}^n: (NG)_n \rightarrow (NG)_{n+1}$  for  $i = 1, \dots, n - 1$  is given by  $f\gamma_{i,1}^n: I_\infty^{\otimes n+1} \rightarrow G$  defined by

$$f\gamma_{i,1}^n(v_1, \dots, v_{n+1}) = f(v_1, \dots, v_{i-1}, \min(v_i, v_{i+1}), v_{i+1}, \dots, v_{n+1}).$$

One verifies these maps satisfy cubical identities, thus  $NG$  is a cubical set. A straightforward computation gives the following statement.

PROPOSITION 3.5. *We have an isomorphism*

$$NG \cong \text{colim}(N_1G \xrightarrow{l^*} N_2G \xrightarrow{r^*} N_3G \xrightarrow{l^*} N_4G \xrightarrow{r^*} \dots)$$

natural in  $G$ .

The nerve and realization functors satisfy the following categorical properties.

PROPOSITION 3.6. *For cubical sets  $X, Y$  and  $m \geq 1$ , we have an isomorphism*

$$|X \otimes Y|_m \cong |X|_m \otimes |Y|_m$$

natural in  $X$  and  $Y$ .

*Proof.* The composite functors

$$|- \otimes -|_m, |-|_m \otimes |-|_m: \text{cSet} \times \text{cSet} \rightarrow \text{Graph}$$

preserve all colimits. As  $\text{cSet}$  is a presheaf category, every cubical set is a colimit of representable presheaves. Thus, it suffices to show these composites are naturally isomorphic on pairs  $(\square^a, \square^b)$  for  $a, b \geq 0$ . We compute

$$\begin{aligned} |\square^a \otimes \square^b|_m &\cong |\square^{a+b}|_m \\ &\cong I_m^{\otimes a+b} \\ &\cong I_m^{\otimes a} \otimes I_m^{\otimes b} \\ &\cong |\square^a|_m \otimes |\square^b|_m. \end{aligned} \quad \square$$

COROLLARY 3.7. *Let  $X$  be a cubical set and  $G$  be a graph. For  $m \geq 1$ , we have isomorphisms*

$$\text{hom}_L(X, N_mG) \cong N_m(\text{hom}^\otimes(|X|_m, G)) \cong \text{hom}_R(X, N_mG)$$

natural in  $X$  and  $G$ .

*Proof.* The square

$$\begin{array}{ccc} \mathbf{cSet} & \xrightarrow{-\otimes X} & \mathbf{cSet} \\ \downarrow |-\!|_m & & \downarrow |-\!|_m \\ \mathbf{Graph} & \xrightarrow{-\otimes |X|_m} & \mathbf{Graph} \end{array}$$

commutes up to natural isomorphism by Proposition 3.6, thus the corresponding square of right adjoints

$$\begin{array}{ccc} \mathbf{Graph} & \xrightarrow{\mathrm{hom}^\otimes(|X|_m, -)} & \mathbf{Graph} \\ \downarrow N_m & & \downarrow N_m \\ \mathbf{cSet} & \xrightarrow{\mathrm{hom}_L(X, -)} & \mathbf{cSet} \end{array}$$

commutes up to natural isomorphism. For naturality in  $X$ , the required square commutes by faithfulness of the Yoneda embedding  $\mathbf{cSet} \rightarrow \mathbf{Set}^{\mathbf{cSet}^{\mathrm{op}}}$ . A similar argument involving  $X \otimes -$  constructs the isomorphism involving  $\mathrm{hom}_R(X, -)$ .  $\square$

PROPOSITION 3.8. *The nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  preserves finite limits.*

*Proof.* By Proposition 3.5, the nerve of  $G$  is a filtered colimit. This then follows as filtered colimits commute with finite limits and  $N_m: \mathbf{Graph} \rightarrow \mathbf{cSet}$  is a right adjoint for all  $m \geq 1$ .  $\square$

We prove that the nerve functors preserve filtered colimits, which we use to give an analogue of Corollary 3.7 for the nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$ .

PROPOSITION 3.9. *For  $m \geq 0$ , the functors  $N_m, N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  preserve filtered colimits.*

*Proof.* For  $N_m$ , it suffices to show filtered colimits are preserved componentwise, i.e. that

$$\mathbf{Graph}(I_m^{\otimes n}, -): \mathbf{Graph} \rightarrow \mathbf{Set}$$

preserves filtered colimits. This follows from Corollary 1.7.

For  $N$ , this follows since  $N_m$  preserves filtered colimits and colimits commute with colimits.  $\square$

For a cubical set  $X$ , define a functor  $P_X: \mathbf{Graph} \rightarrow \mathbf{Graph}$  by

$$P_X G := \mathrm{colim} \left( \mathrm{hom}^\otimes(|X|_1, G) \xrightarrow{(l_*)^*} \mathrm{hom}^\otimes(|X|_2, G) \xrightarrow{(r_*)^*} \mathrm{hom}^\otimes(|X|_3, G) \xrightarrow{(l_*)^*} \dots \right).$$

As an example, the path graph  $PG$  of a graph is exactly  $P_{\square_1} G$ .

PROPOSITION 3.10. *Let  $X$  be a cubical set with finitely many non-degenerate cubes. We have isomorphisms*

$$N(P_X G) \cong \mathrm{hom}_L(X, NG) \cong \mathrm{hom}_R(X, NG)$$

*natural in  $X$  and  $G$ .*

*Proof.* By Proposition 3.5, the left term  $N(P_X G)$  is the colimit

$$N(P_X G) \cong \operatorname{colim} \left( \begin{array}{ccc} N_1 \operatorname{hom}^\otimes(|X|_1, G) \xleftarrow{l^*} N_2 \operatorname{hom}^\otimes(|X|_1, G) \xleftarrow{r^*} \dots & & \\ \downarrow (l_*)^* & & \downarrow (l_*)^* \\ N_1 \operatorname{hom}^\otimes(|X|_2, G) \xleftarrow{l^*} N_2 \operatorname{hom}^\otimes(|X|_2, G) \xleftarrow{r^*} \dots & & \\ \downarrow (r_*)^* & & \downarrow (r_*)^* \\ \dots & & \dots \end{array} \right),$$

where the vertical maps are monomorphisms since  $l_*, r_* : |X|_{m+1} \rightarrow |X|_m$  are epimorphisms for all  $m \geq 1$ . Computing this colimit componentwise in **Set**, this colimit is naturally isomorphic to the colimit

$$N(P_X G) \cong \operatorname{colim}(N_1 \operatorname{hom}^\otimes(|X|_1, G) \rightarrow N_2 \operatorname{hom}^\otimes(|X|_2, G) \rightarrow \dots)$$

along the diagonal. Applying Corollary 3.7, we may write

$$N(P_X G) \cong \operatorname{colim}(\operatorname{hom}_L(X, N_1 G) \rightarrow \operatorname{hom}_L(X, N_2 G) \rightarrow \dots).$$

As  $X$  has finitely many non-degenerate cubes, the functor  $\operatorname{hom}_L(X, -)$  preserves filtered colimits. Thus,

$$N(P_X G) \cong \operatorname{hom}_L(X, NG).$$

An analogous proof applies in the case of  $\operatorname{hom}_R(X, NG)$ . □

We show that the nerve functors ‘detect’ concatenation of paths. That is, they contain all possible composition squares.

**PROPOSITION 3.11.** *Let  $f : (I_\infty, I_{\geq M}) \rightarrow (G, v)$  and  $g : (I_\infty, I_{\geq N}) \rightarrow (G, v)$  be based graph maps for some  $M, N \geq 0$  such that  $f\partial_{1,1} = g\partial_{1,0}$ . The concatenation  $f \cdot g$  of  $f$  followed by  $g$  is a concatenation of  $f$  and  $g$  in the  $(M + N)$ -nerve  $N_{(M+N)}G$ .*

*Proof.* The horizontal concatenation of  $f\gamma_{1,0} : I_M^{\otimes 2} \rightarrow G$  on the left and  $g\sigma_2 : I_N^{\otimes 2} \rightarrow G$  on the right gives a square  $\eta : I_{M+N}^{\otimes 2} \rightarrow G$  such that

$$\begin{aligned} \eta\partial_{1,0} &= f, & \eta\partial_{1,1} &= g\partial_{1,1}\sigma_1, \\ \eta\partial_{2,0} &= f \cdot g, & \eta\partial_{2,1} &= g. \end{aligned}$$

This is exactly a composition square witnessing  $f \cdot g$  as a composition of  $f$  and  $g$  in  $N_{(M+N)}G$ . □

As well, the nerve functors reflect isomorphisms.

**PROPOSITION 3.12.** *The nerve functors reflect isomorphisms. That is, given a graph map  $f : G \rightarrow H$ , if either*

- (i)  $N_m f : N_m G \rightarrow N_m H$  is an isomorphism for some  $m \geq 1$ , or
- (ii)  $Nf : NG \rightarrow NH$  is an isomorphism,

then  $f$  is.

*Proof.* We prove that  $N : \mathbf{Graph} \rightarrow \mathbf{cSet}$  reflects isomorphisms as the case for  $N_m : \mathbf{Graph} \rightarrow \mathbf{cSet}$  is analogous.

The inverse of  $Nf$  is, in particular, an inverse on 0-cubes  $(NH)_0 \rightarrow (NG)_0$ , i.e. an inverse of  $f$  on vertices  $H_V \rightarrow G_V$ . It suffices to show this map  $g : H_V \rightarrow G_V$  is a graph map.



An edge in  $H$  gives a map  $e: I_1 \rightarrow H$ . This gives a 1-cube  $\bar{e}: \square^1 \rightarrow NG$  by the inclusion  $N_1H \hookrightarrow NH$ . As  $Nf: NG \rightarrow NH$  is an isomorphism, there is a unique 1-cube  $\bar{p}: \square^1 \rightarrow NG$  such that  $Nf(\bar{p}) = \bar{e}$ . This corresponds to a map  $p: I_\infty \rightarrow G$  such that  $p$  stabilizes in both directions and  $fp = e$ .

As  $f$  is injective on vertices and  $e$  is a path of length 1 (i.e. it consists of two vertices and one edge), we deduce that  $p$  is a path of length 1. That is,  $p$  is an edge, hence  $g$  is a graph map.  $\square$

Recall that a functor which reflects isomorphisms also reflects any (co)limits which it preserves. Thus, we have the following corollary.

COROLLARY 3.13.

- (i) For  $m \geq 1$ , the  $m$ -nerve  $N_m: \mathbf{Graph} \rightarrow \mathbf{cSet}$  reflects all limits.
- (ii) The nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  reflects finite limits.

#### 4. Main result

##### Statement

Our main theorem is the following result.

THEOREM 4.1. For any graph  $G$ ,

- (i) the nerve  $NG$  of  $G$  is a Kan complex;
- (ii) the natural inclusion  $N_1G \hookrightarrow NG$  is anodyne.

Before proving Theorem 4.1, we explain the proof strategy and establish some auxiliary lemmas.

To show that the nerve of any graph is a Kan complex, we construct a map  $\Phi: I_{3m}^{\otimes n} \rightarrow |\Pi_{i,\varepsilon}^n|_m$  so that the triangle

$$\begin{array}{ccc} |\Pi_{i,\varepsilon}^n|_{3m} & \xrightarrow{c_*^m} & |\Pi_{i,\varepsilon}^n|_m \\ \downarrow & \nearrow \Phi & \\ I_{3m}^{\otimes n} & & \end{array}$$

commutes. We show that every map  $\Pi_{i,\varepsilon}^n \rightarrow NG$  must factor through some  $m$ -nerve  $\Pi_{i,\varepsilon}^n \rightarrow N_mG \rightarrow NG$ . The map  $\Phi$  gives a filler for the composite  $\Pi_{i,\varepsilon}^n \rightarrow N_mG \rightarrow N_{3m}G$ , thus a filler in  $NG$ .

To show that the map  $N_1G \hookrightarrow NG$  is anodyne, we show the maps  $l^*, r^*: N_mG \hookrightarrow N_{m+1}G$  are anodyne. This is done by an explicit construction establishing  $l^*$  and  $r^*$  as a transfinite composition of pushouts along coproducts of open box inclusions. This implies  $N_1G \hookrightarrow NG$  is anodyne as the nerve of  $G$  is a transfinite composition of  $l^*$  and  $r^*$ .

##### Proof of part (i)

CONSTRUCTION 4.2. Fix  $m, n \geq 1$ ,  $i \in \{1, \dots, n\}$ , and  $\varepsilon \in \{0, 1\}$ . We construct the map  $\Phi: I_{3m}^{\otimes n} \rightarrow |\Pi_{i,\varepsilon}^n|_m$ . For  $n = 1$ , we compute  $|\Pi_{i,\varepsilon}^1|_m \cong I_0$ . In this case, the map  $\Phi$  is immediate. Thus, we assume  $n \geq 2$ .

Define a map  $\varphi: I_{3m} \rightarrow I_{3m}$  by

$$\varphi v := \begin{cases} m & v \leq m \\ i & m \leq v \leq 2m \\ 2m & v \geq 2m. \end{cases}$$

It is straightforward to verify this definition gives a graph map. From this, we have a map  $\varphi^{\otimes n-1}: I_{3m}^{\otimes n-1} \rightarrow I_{3m}^{\otimes n-1}$  which sends  $(v_1, \dots, v_{n-1})$  to  $(\varphi v_1, \dots, \varphi v_{n-1})$ .

For  $v = (v_1, \dots, v_{n-1}) \in I_{3m}^{\otimes n-1}$ , let  $d(v)$  denote the node distance between  $v$  and  $\varphi^{\otimes n-1}v$ . That is,

$$d(v) := \sum_{i=1}^{n-1} |v_i - \varphi v_i|.$$

Observe this gives a graph map  $d: I_{3m}^{\otimes n-1} \rightarrow I_\infty$  (Figure 7).




FIGURE 7. Vertices of  $I_6^{\otimes 2}$  labeled by their image under  $d: I_6^{\otimes 2} \rightarrow I_\infty$ .

For  $t \geq 0$ , we have a graph map  $\beta[t]: I_\infty \rightarrow I_t$  defined by

$$\beta[t](v) := \begin{cases} 0 & v \leq 0 \\ v & 0 \leq v \leq t \\ t & v \geq t. \end{cases}$$

One thinks of  $\beta[t]$  as bounding the graph  $I_\infty$  between 0 and  $t$ . Recall that the map  $c^m: I_{3m} \rightarrow I_m$  is given by

$$c^m v := \begin{cases} 0 & v \leq m \\ v - m & m \leq v \leq 2m \\ m & v \geq 2m. \end{cases}$$

For a vertex  $(v_1, \dots, v_n) \in I_{3m}^{\otimes n}$ , we write  $\sigma_i v$  for the vertex  $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n) \in I_{3m}^{\otimes n-1}$ . As well, let  $\alpha_\varepsilon^{v_i}$  denote the value

$$\alpha_\varepsilon^{v_i} := \varepsilon m + (1 - 2\varepsilon)(c^m v_i).$$

We may also write this as

$$\alpha_\varepsilon^{v_i} = \begin{cases} c^m v_i & \varepsilon = 0 \\ m - c^m v_i & \varepsilon = 1. \end{cases}$$

We define  $\Phi: I_{3m}^{\otimes n} \rightarrow |\Gamma_{i,\varepsilon}^n|_m$  by

$$\begin{aligned} &\Phi(v_1, \dots, v_n) \\ &:= (c^m v_1, \dots, c^m v_{i-1}, (1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}])(d(\sigma_i v) - \alpha_\varepsilon^{v_i}), c^m v_{i+1}, \dots, c^m v_n). \end{aligned}$$

That is,

$$\begin{aligned} &\Phi(v_1, \dots, v_n) \\ &= \begin{cases} (c^m v_1, \dots, c^m v_{i-1}, m - \beta[m - c^m v_i](d(\sigma_i v) - c^m v_i), c^m v_{i+1}, \dots, c^m v_n) & \text{if } \varepsilon = 0 \\ (c^m v_1, \dots, c^m v_{i-1}, \beta[c^m v_i](d(\sigma_i v) + c^m v_i - m), c^m v_{i+1}, \dots, c^m v_n) & \text{if } \varepsilon = 1. \end{cases} \end{aligned}$$

We first show this formula is well defined, i.e. that this tuple lies in the subgraph  $|\Gamma_{i,\varepsilon}^n|_m$  of  $I_m^{\otimes n}$ . Observe that if  $c^m v_k = 0$  or  $m$  for some  $k \neq i$  in  $\{1, \dots, n\}$  then this tuple indeed lies in  $|\Gamma_{i,\varepsilon}^n|_m$ . For  $k < i$ , this tuple lies on the  $(k, 0)$ - or  $(k, 1)$ -face. For  $k > i$ , this tuple lies on the  $(k + 1, 0)$ - or  $(k + 1, 1)$ -face. Otherwise, if  $0 < c^m v_k < m$  for all  $k \neq i$  then  $d(\sigma_i v) = 0$  since  $v_k = \varphi v_k$  for all  $k \neq i$ . From this, it follows that

$$\beta[\alpha_{1-\varepsilon}^{v_i}](d(\sigma_i v) - \alpha_\varepsilon^{v_i}) = \beta[\alpha_{1-\varepsilon}^{v_i}](-\alpha_\varepsilon^{v_i}) = 0,$$

thus  $\Phi(v_1, \dots, v_n)$  lies on the  $(i, 1 - \varepsilon)$ -face, i.e. the face opposite the missing face.

To see this formula gives a graph map, suppose  $(v_1, \dots, v_n)$  and  $(w_1, \dots, w_n)$  are connected vertices in  $I_{3m}^{\otimes n}$ . By definition, there exists  $k = 1, \dots, n$  so that  $v_k$  and  $w_k$  are connected in  $I_{3m}$  and  $v_j = w_j$  for all  $j \neq k$ . We first consider the case where  $k \neq i$ . Observe that if  $c^m v_k = c^m w_k$  then this is immediate. If  $c^m v_k \neq c^m w_k$  then  $\sigma_i v = \varphi(\sigma_i w)$ . This gives that  $d(\sigma_i v) = d(\sigma_i w)$ , hence  $\Phi(v_1, \dots, v_n)$  and  $\Phi(w_1, \dots, w_n)$  are equal on all components except the  $k$ th component. That is, they are connected. In the case where  $k = i$ , we have that  $dv$  and  $dw$  differ by at most 1. This implies  $(1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](dv - \alpha_\varepsilon^{v_i}))$  and  $(1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{w_i}](dw - \alpha_\varepsilon^{w_i}))$  differ by at most 1, thus  $\Phi(v_1, \dots, v_n)$  and  $\Phi(w_1, \dots, w_n)$  are connected.

*Example 4.3.* We look at the map  $\Phi: I_{3m}^{\otimes n} \rightarrow |\Gamma_{i,\varepsilon}^n|_m$  in the case of  $n = 3, m = 2$ , and  $(i, \varepsilon) = (3, 1)$  (Figure 8).

We may write the map  $\Phi: I_6^{\otimes 3} \rightarrow |\Gamma_{3,1}^3|_2$  as

$$\Phi(v_1, v_2, v_3) = \begin{cases} (c^2 v_1, c^2 v_2, 0) & \text{if } v_3 \leq 2 \\ (c^2 v_1, c^2 v_2, \beta[1](d(v_1, v_2) - 1)) & \text{if } v_3 = 3 \\ (c^2 v_1, c^2 v_2, \beta[2](d(v_1, v_2))) & \text{if } v_3 \geq 4. \end{cases}$$

If  $v_3 \leq 2$  then  $(v_1, v_2, v_3)$  is contained in the  $\partial_{3,0}$ -face of  $|\Gamma_{3,1}^3|_2$ , which is the face opposite the missing face. For the cross-section where  $v_3 = 3$ , if  $d(v_1, v_2) \geq 2$  then  $(v_1, v_2, v_3)$  is sent to a vertex in the  $v_3 = 1$  cross-section of  $|\Gamma_{3,1}^3|_2$ . For  $v_3 \geq 3$ , if  $d(v_1, v_2) = 1$  then  $(v_1, v_2, v_3)$  is sent to a vertex in the  $v_3 = 1$  cross-section. If  $d(v_1, v_2) \geq 2$  then  $(v_1, v_2, v_3)$  is sent to a vertex in the  $v_3 = 2$  cross-section. In all three cross-sections, if  $d(v_1, v_2) = 0$  then  $(v_1, v_2, v_3)$  is contained in the face opposite the missing face of  $|\Gamma_{3,1}^3|_2$  (Figure 9).




FIGURE 8 (colour online). The graph  $|\square_{3,1}^3|_2$  as a net. Vertices connected by a dotted line are identical.




FIGURE 9 (colour online). The map  $\Phi: I_6^{\otimes 3} \rightarrow |\square_{3,1}^3|_2$  split into cross-sections.

LEMMA 4.4. *The diagram*

$$\begin{array}{ccc}
 |\square_{i,\varepsilon}^n|_{3m} & \xrightarrow{(c_*)^m} & |\square_{i,\varepsilon}^n|_m \\
 \downarrow & \nearrow \Phi & \\
 I_{3m}^{\otimes n} & & 
 \end{array}$$

commutes.

*Proof.* For  $n = 1$ , we have  $|\square_{i,\varepsilon}^1|_m \cong I_0$ . The diagram then commutes as  $I_0$  is terminal in Graph.

For  $n \geq 2$ , fix  $(v_1, \dots, v_n) \in |\square_{i,\varepsilon}^n|_{3m}$ . It suffices to show

$$(1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](d(\sigma_i v) - \alpha_\varepsilon^{v_i})) = c^m v_i.$$

If  $v_i = (1 - \varepsilon)(3m)$  then  $d(\sigma_i v) = 0$ . Thus,

$$\begin{aligned}
 (1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](d(\sigma_i v) - \alpha_\varepsilon^{v_i})) &= (1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](-\alpha_\varepsilon^{v_i})) \\
 &= (1 - \varepsilon)m + (2\varepsilon - 1)(0) \\
 &= (1 - \varepsilon)m \\
 &= c^m((1 - \varepsilon)(3m)) \\
 &= c^m v_i.
 \end{aligned}$$

Otherwise, if  $v_i \neq (1 - \varepsilon)(3m)$  then  $d(\sigma_i v) \geq m$  (since  $v_k = 0, 3m$  for some  $k \neq i$ ). For  $\varepsilon = 0$ , we compute

$$\begin{aligned} (1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](d(\sigma_i v) - \alpha_\varepsilon^{v_i})) &= m - \beta[\alpha_1^{v_i}](d(\sigma_i v) - \alpha_0^{v_i}) \\ &= m - \beta[m - c^m v_i](d(\sigma_i v) - c^m v_i) \\ &= m - (m - c^m v_i) \\ &= c^m v_i. \end{aligned}$$

For  $\varepsilon = 1$ , we compute

$$\begin{aligned} (1 - \varepsilon)m + (2\varepsilon - 1)(\beta[\alpha_{1-\varepsilon}^{v_i}](d(\sigma_i v) - \alpha_\varepsilon^{v_i})) &= \beta[\alpha_0^{v_i}](d(\sigma_i v) - \alpha_1^{v_i}) \\ &= \beta[c^m v_i](d(\sigma_i v) + c^m v_i - m) \\ &= c^m v_i. \end{aligned} \quad \square$$

**THEOREM 4.5.** *For any graph  $G$ , the nerve  $NG$  of  $G$  is a Kan complex.*

*Proof.* Fix a map  $f: \square_{i,\varepsilon}^n \rightarrow NG$ . We know that  $NG \cong \text{colim}(N_1 G \hookrightarrow N_3 G \hookrightarrow N_5 G \hookrightarrow \dots)$  by Proposition 3.5. Recall that in a presheaf category, any map from a representable presheaf to a colimit must factor through some component of the colimit cone by the Yoneda lemma. Thus, for any  $k \geq 0$  and  $x: \square^k \rightarrow \square_{i,\varepsilon}^n$ , the map  $fx: \square^k \rightarrow NG$  factors through an inclusion  $N_m G \hookrightarrow NG$  for some  $m \geq 1$ . As  $\square_{i,\varepsilon}^n$  has only finitely many non-degenerate cubes,  $f$  factors through the natural inclusion  $N_m G \hookrightarrow NG$  as a map  $g: \square_{i,\varepsilon}^n \rightarrow N_m G$  for some  $m \geq 0$ .

By adjointness,  $g$  corresponds to a map  $\bar{g}: |\square_{i,\varepsilon}^n|_m \rightarrow G$ . Lemma 4.4 shows that  $\bar{g}\Phi: I_{3m}^{\otimes n} \rightarrow G$  is a lift of the composite map  $\bar{g}(c^*)^m: |\square_{i,\varepsilon}^n|_{3m} \rightarrow G$ :

$$\begin{array}{ccc} |\square_{i,\varepsilon}^n|_{3m} & \xrightarrow{(c^*)^m} & |\square_{i,\varepsilon}^n|_m \xrightarrow{\bar{g}} G \\ \downarrow & \nearrow \Phi & \\ |\square^n|_{3m} & & \end{array}$$

By adjointness, this gives a filler  $\square^n \rightarrow N_{3m} G$  for the map  $(c^*)^m g: \square_{i,\varepsilon}^n \rightarrow N_{3m} G$ . Post-composing with the natural inclusion  $N_{3m} G \hookrightarrow NG$ , this gives a filler  $\square^n \rightarrow NG$  of  $f$ .  $\square$

Via Theorem 4.5, we may speak of the homotopy groups of  $NG$ . We prove that the  $A$ -homotopy groups of  $G$  are exactly the cubical homotopy groups of  $NG$ .

**THEOREM 4.6.** *We have an isomorphism  $A_n(G, v) \cong \pi_n(NG, v)$  natural in  $(G, v)$ .*

*Proof.* It is straightforward to verify that a relative graph map  $(I_\infty^{\otimes n}, I_{\geq M}^{\otimes n}) \rightarrow (G, v)$  is exactly a relative cubical map  $(\square^n, \partial \square^n) \rightarrow (NG, v)$  and that a relative homotopy between two such relative graph maps is exactly a relative homotopy between two such relative cubical maps. This gives a set bijection  $A_n(G, v) \rightarrow \pi_n(NG, v)$ . Proposition 3.11 shows this map is a group homomorphism.  $\square$

*Remark 4.7.* Theorem 4.6 shows why the functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  fails to preserve arbitrary limits, even though, by Proposition 3.8, it preserves finite ones. To see that, we first note that

$$A_1 \left( \prod_{k \geq 5} (C_k, 0) \right) \cong \bigoplus_{k \geq 5} \mathbb{Z},$$

since a map  $I_\infty \rightarrow \prod_{k \geq 5} C_k$  that stabilizes outside of a finite interval must necessarily be null-homotopic in all but finitely many  $C_k$ . That is,  $A_1: \mathbf{Graph}_* \rightarrow \mathbf{Grp}$  does not preserve infinite

products. Theorem 4.6 shows that  $A_1 \cong \pi_1 \circ N$ , and  $\pi_1 : \mathbf{cSet}_* \rightarrow \mathbf{Grp}$  preserves infinite products by [CK23, Proposition 4.1]. Thus, we conclude that

$$N\left(\prod_{k \geq 5} (C_k, 0)\right) \not\cong \prod_{k \geq 5} (NC_k, 0).$$

**Proof of part (ii)**

Now we show that the inclusions  $l^*, r^* : N_m G \hookrightarrow N_{m+1} G$  are anodyne. We first explain the intuition for why this statement holds.

The 1-nerve  $N_1 G$  of a graph  $G$  contains, as 1-cubes, all paths of length 1 in  $G$  (i.e. paths with two vertices and one edge). Consider the image of the embedding  $l^* : N_1 G \rightarrow N_2 G$  as a cubical subset  $X \subseteq N_2 G$  of the 2-nerve of  $G$ . A 1-cube of  $N_2 G$  is exactly a path of length 2 in  $G$ ; a 1-cube of  $X$  is a path of length 1 regarded as a path of length 2 whose first two vertices are the same. Given a path  $f : I_2 \rightarrow G$  of length 2 in  $G$ , we may define a  $3 \times 3$  square  $g : I_2^{\otimes 2} \rightarrow G$  in  $G$  by

$$g(v_1, v_2) := \begin{cases} f(l(v_1)) & v_2 < 2 \\ f(v_1) & v_2 = 2. \end{cases}$$

Observe that the  $\partial_{1,0^-}$ ,  $\partial_{1,1^-}$ , and  $\partial_{2,0}$ -faces of  $g$  are 1-cubes of  $X$ , i.e. they are paths of length 2 whose first two vertices are the same, whereas the  $\partial_{2,1}$ -face of  $g$  is  $f$ . That is, we have shown the restriction  $g|_{|\square_{2,1}^2|_2} : |\square_{2,1}^2|_2 \rightarrow G$  of  $g$  to the open box corresponds to a map  $|\square_{2,1}^2|_2 \rightarrow N_2 G$  whose image is contained in  $X$  (Figure 10).




FIGURE 10 (colour online). The square  $g : I_2^{\otimes 2} \rightarrow G$  constructed from the path  $f : I_2 \rightarrow G$ .

Let  $Y \subseteq N_2 G$  denote the cubical subset generated by  $X$  and  $g$ , i.e. the cubical subset containing  $X$ , the 2-cube  $g \in (N_2 G)_2$ , and all faces, degeneracies, and connections of  $g$ . The square

$$\begin{array}{ccc} |\square_{2,1}^2|_2 & \xrightarrow{g|_{|\square_{2,1}^2|_2}} & X \\ \downarrow & \lrcorner & \downarrow \\ \square^2 & \xrightarrow{g} & Y \end{array}$$

is a pushout by definition of  $Y$ . This gives exactly that the inclusion  $X \hookrightarrow Y$  is anodyne. Following this approach, one may construct an anodyne inclusion  $X \hookrightarrow X_n$  such that  $X_n$  contains all  $n$ -cubes of  $N_2 G$ . With this, the colimit of the sequence  $X \hookrightarrow X_1 \hookrightarrow X_2 \hookrightarrow \dots$  is exactly  $N_2 G$  and the inclusion  $X \hookrightarrow N_2 G$  is anodyne by closure under transfinite composition.

For  $n > 0$  and  $j \in \{0, \dots, n\}$ , the maps  $l, r : I_{m+1} \rightarrow I_m$  yield maps

$$(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j}), (\text{id}_{I_{m+1}}^{\otimes j} \otimes r^{\otimes n-j}) : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j}.$$

For  $j \in \{0, \dots, n-1\}$ , we define maps  $\lambda_{m,j}^n, \rho_{m,j}^n : I_{m+1}^{\otimes n} \otimes I_1 \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  by

$$\lambda_{m,j}^n(v_1, \dots, v_n, v_{n+1}) = \begin{cases} (v_1, \dots, v_j, lv_{j+1}, lv_{j+2}, \dots, lv_n) & v_{n+1} = 0 \\ (v_1, \dots, v_{j+1}, lv_{j+2}, \dots, lv_n) & v_{n+1} = 1, \end{cases}$$

$$\rho_{m,j}^n(v_1, \dots, v_n, v_{n+1}) = \begin{cases} (v_1, \dots, v_j, rv_{j+1}, rv_{j+2}, \dots, rv_n) & v_{n+1} = 0 \\ (v_1, \dots, v_{j+1}, rv_{j+2}, \dots, rv_n) & v_{n+1} = 1. \end{cases}$$

That is, the restriction of  $\lambda_{m,j}^n$  to  $I_{m+1}^{\otimes n} \otimes \{0\}$  is the map  $\text{id}_{I_{m+1}^{\otimes j}} \otimes l^{\otimes n-j}$  and its restriction to  $I_{m+1}^{\otimes n} \otimes \{1\}$  is  $\text{id}_{I_{m+1}^{\otimes j+1}} \otimes l^{\otimes n-j-1}$  (likewise for  $\rho_{m,j}^n$ ).

The maps  $l^m, r^m: I_{m+1} \rightarrow I_1$  denote application of the maps  $l, r$  a total of  $m$  times. That is,

$$l^m(v) = \begin{cases} 0 & v < m + 1 \\ 1 & v = m + 1, \end{cases}$$

$$r^m(v) = \begin{cases} 0 & v = 0 \\ 1 & v > 0. \end{cases}$$

We write  $\bar{\lambda}_{m,j}^n: I_{m+1}^{\otimes n+1} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  for the composition of  $\text{id}_{I_{m+1}^{\otimes n}} \otimes l^m: I_{m+1}^{\otimes n+1} \rightarrow I_{m+1}^{\otimes n} \otimes I_1$  followed by  $\lambda_{m,j}^n: I_{m+1}^{\otimes n} \otimes I_1 \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  (Figure 11), and we write  $\bar{\rho}_{m,j}^n: I_{m+1}^{\otimes n+1} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  for the composition of  $\text{id}_{I_{m+1}^{\otimes n}} \otimes r^n: I_{m+1}^{\otimes n+1} \rightarrow I_{m+1}^{\otimes n} \otimes I_1$  followed  $\rho_{m,j}^n: I_{m+1}^{\otimes n} \otimes I_1 \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$ , respectively (Figure 12).







FIGURE 11. The graph  $I_2 \otimes I_1$  with vertices labeled by their image under  $\lambda_{1,0}^1: I_2 \otimes I_1 \rightarrow I_2$ .




(a) The subgraph  $I_2^{\otimes 2} \otimes \{0\}$  under  $\lambda_{1,0}^2$



(b) The subgraph  $I_2^{\otimes 2} \otimes \{1\}$  under  $\lambda_{1,1}^2$



(c) The subgraph  $I_2^{\otimes 2} \otimes \{0\}$  under  $\lambda_{1,1}^2$



(d) The subgraph  $I_2^{\otimes 2} \otimes \{1\}$  under  $\lambda_{1,1}^2$

FIGURE 12. Cross-sections of the graph  $I_2^{\otimes 2} \otimes I_1$  with vertices labeled by their image under the maps  $\lambda_{1,0}^2: I_2^{\otimes 2} \otimes I_1 \rightarrow I_2 \otimes I_1$  and  $\lambda_{1,1}^2: I_2^{\otimes 2} \otimes I_1 \rightarrow I_2^{\otimes 2}$ .

PROPOSITION 4.8. Let  $m, n > 0$  and  $j \in \{0, \dots, n\}$ . For  $i = 1, \dots, n$  such that  $i \neq j + 1$  and  $\varepsilon = 0, 1$ ,

- (i)  $\bar{\lambda}_{m,j}^n \partial_{i,\varepsilon} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes n}$  factors through  $\bar{\lambda}_{m,j}^{n-1} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes n-1}$ ;
- (ii)  $\bar{\rho}_{m,j}^n \partial_{i,\varepsilon} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes n}$  factors through  $\bar{\rho}_{m,j}^{n-1} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes n-1}$ .

*Proof.* We consider the result for  $\bar{\lambda}_{m,j}^n$ , as the result for  $\bar{\rho}_{m,j}^n$  is analogous.

Fix  $(v_1, \dots, v_n) \in I_{m+1}^{\otimes n}$ . If  $i < j + 1$  then we have

$$\begin{aligned} \bar{\lambda}_{m,j}^n \partial_{i,\varepsilon}(v_1, \dots, v_n) &= \bar{\lambda}_{m,j}^n(v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_n) \\ &= \lambda_{m,j}^n(v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_{n-1}, l^m v_n) \\ &= \begin{cases} (v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_j, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_{j+1}, lv_{j+2}, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= \partial_{i,\varepsilon}^n \lambda_{m,j}^n(v_1, \dots, v_{n-1}, l^m v_n) \\ &= \partial_{i,\varepsilon}^n \bar{\lambda}_{m,j}^n(v_1, \dots, v_n). \end{aligned}$$

Thus,  $\bar{\lambda}_{m,j}^n \partial_{i,\varepsilon} = \partial_{i,\varepsilon} \bar{\lambda}_{m,j}^{n-1}$ .

Otherwise, we have  $i > j + 1$ . Consider the embedding  $\iota : I_{m+1}^{\otimes n-1} \rightarrow I_{m+1}^{\otimes n}$  given by

$$\iota(v_1, \dots, v_{n-1}) = (v_1, \dots, v_{i-1}, \varepsilon m, v_i, \dots, v_{n-1}).$$

With this, we may write

$$\begin{aligned} \bar{\lambda}_{m,j}^n \partial_{i,0}^{n+1}(v_1, \dots, v_n) &= \bar{\lambda}_{m,j}^n(v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_n) \\ &= \lambda_{m,j}^n(v_1, \dots, v_{i-1}, \varepsilon(m+1), v_i, \dots, v_{n-1}, l^m v_n) \\ &= \begin{cases} (v_1, \dots, v_j, lv_{j+1}, \dots, lv_{i-1}, l(\varepsilon(m+1)), lv_i, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_{j+1}, lv_{j+2}, \dots, lv_{i-1}, l(\varepsilon(m+1)), lv_i, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= \begin{cases} (v_1, \dots, v_j, lv_{j+1}, \dots, lv_{i-1}, \varepsilon m, lv_i, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_{j+1}, lv_{j+2}, \dots, lv_{i-1}, \varepsilon m, lv_i, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= \iota \lambda_{m,j}^n(v_1, \dots, v_{n-1}, l^m v_n) \\ &= \iota \bar{\lambda}_{m,j}^n(v_1, \dots, v_n). \end{aligned}$$

Thus,  $\bar{\lambda}_{m,j}^n \partial_{i,\varepsilon} = \iota \bar{\lambda}_{m,j}^{n-1}$ . □

PROPOSITION 4.9. Let  $m, n > 0$  and  $j \in \{0, \dots, n\}$ . Then

- (i)  $\bar{\lambda}_{m,j}^n \partial_{j+1,0}$  and  $\bar{\lambda}_{m,j}^n \partial_{j+1,1}$  factor through  $\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j}$ ;
- (ii)  $\bar{\rho}_{m,j}^n \partial_{j+1,0}$  and  $\bar{\rho}_{m,j}^n \partial_{j+1,1}$  factor through  $\text{id}_{I_{m+1}}^{\otimes j} \otimes r^{\otimes n-j} : I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j}$ .

*Proof.* We show the result for  $\bar{\lambda}_{m,j}^n$  as the result for  $\bar{\rho}_{m,j}^n$  is analogous.



Fix  $(v_1, \dots, v_n) \in I_{m+1}^{\otimes n}$ . We compute

$$\begin{aligned} \bar{\lambda}_{m,j}^n \partial_{j+1,0}(v_1, \dots, v_n) &= \bar{\lambda}_{m,j}^n(v_1, \dots, v_j, 0, v_{j+1}, \dots, v_n) \\ &= \lambda_{m,j}^n(v_1, \dots, v_j, 0, v_{j+1}, \dots, l^m v_n) \\ &= \begin{cases} (v_1, \dots, v_j, l0, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_j, 0, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= (v_1, \dots, v_j, 0, lv_{j+1}, \dots, lv_{n-1}) \\ &= \partial_{j+1,0} \sigma_n(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j})(v_1, \dots, v_n), \end{aligned}$$

thus  $\bar{\lambda}_{m,j}^n \partial_{j+1,0} = \partial_{j+1,0} \sigma_n(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j})$ .

Consider the map  $f: I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  defined by

$$f(a_1, \dots, a_j, b_1, \dots, b_{n-j}) = \begin{cases} (a_1, \dots, a_j, m, b_1, \dots, b_{n-j-1}) & \text{if } l^{m-1} b_{n-j} = 0 \\ (a_1, \dots, a_j, m+1, b_1, \dots, b_{n-j-1}) & \text{if } l^{m-1} b_{n-j} = 1. \end{cases}$$

It is straightforward to verify this is a graph map. With this, we may write

$$\begin{aligned} \bar{\lambda}_{m,j}^n \partial_{j+1,1}(v_1, \dots, v_n) &= \bar{\lambda}_{m,j}^n(v_1, \dots, v_j, m+1, v_{j+1}, \dots, v_n) \\ &= \lambda_{m,j}^n(v_1, \dots, v_j, m+1, v_{j+1}, \dots, l^m v_n) \\ &= \begin{cases} (v_1, \dots, v_j, l(m+1), lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_j, m+1, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= \begin{cases} (v_1, \dots, v_j, m, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 0 \\ (v_1, \dots, v_j, m+1, lv_{j+1}, \dots, lv_{n-1}) & \text{if } l^m v_n = 1 \end{cases} \\ &= f(v_1, \dots, v_j, lv_{j+1}, \dots, lv_n) \\ &= f(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j})(v_1, \dots, v_n). \end{aligned}$$

Thus,  $\bar{\lambda}_{m,j}^n \partial_{j+1,1} = f(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j})$ . □

LEMMA 4.10. Let  $m, n > 0$ ,  $j \in \{0, \dots, n-1\}$ , and  $G$  be a graph. Consider a subobject  $X$  of  $N_{m+1}G$  which contains:

- all  $n$ -cubes of  $N_{m+1}G$  which factor through  $l^{\otimes n}: I_{m+1}^{\otimes n} \rightarrow I_m^{\otimes n}$ ;
- (if  $n > 1$ ) for any  $x: I_{m+1}^{\otimes h} \otimes I_m^{\otimes k-h} \rightarrow G$  where  $k < n$  and  $h \leq k$ , the  $(k+1)$ -cube  $x\bar{\lambda}_{m,h-1}^k: I_{m+1}^{\otimes k+1} \rightarrow G$ ;
- (if  $j > 0$ ) for any  $x: I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j} \rightarrow G$ , the  $(n+1)$ -cube  $x\bar{\lambda}_{m,j-1}^n: I_{m+1}^{\otimes n+1} \rightarrow G$  of  $N_{m+1}G$ .

Then, for any  $n$ -cube of  $N_{m+1}G$  which factors through  $\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}: I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  as some  $x: I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1} \rightarrow G$ , the restriction of the  $(n+1)$ -cube  $x\bar{\lambda}_{m,j}^n: I_{m+1}^{\otimes n+1} \rightarrow G$  to the open box  $x\bar{\lambda}_{m,j}^n|_{\square_{n+1,1}^{n+1}}: \square_{n+1,1}^{n+1} \rightarrow N_{m+1}G$  factors through the inclusion  $X \hookrightarrow N_{m+1}G$ .

*Proof.* We first show that  $X$  contains all  $n$ -cubes which factor through the map  $\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j}: I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j}$ . If  $j = 0$  then this follows by assumption. Otherwise, consider such an  $n$ -cube, which we write as  $y(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j}): I_{m+1}^{\otimes n} \rightarrow G$  for some  $y: I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j} \rightarrow G$ . Recall  $\bar{\lambda}_{m,j-1}^n \partial_{n+1,1} = \text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j}$ . This gives that  $y(\text{id}_{I_{m+1}}^{\otimes j} \otimes l^{\otimes n-j}) =$

$y\bar{\lambda}_{m,j-1}^n \partial_{n+1,1}$ . By assumption,  $X$  contains  $y\bar{\lambda}_{m,j-1}^n$ . Thus, it contains all faces of  $y\bar{\lambda}_{m,j-1}^n$ , including  $y$ .

To see that each face of  $x\bar{\lambda}_{m,j}^n|_{\square_{n+1,1}^{n+1}}$  is contained in  $X$ , fix  $(i, \varepsilon) \neq (n + 1, 1)$ . For  $i = n + 1$  and  $\varepsilon = 0$ , this follows as  $\bar{\lambda}_{m,j}^n \partial_{n+1,0} = \text{id}_{I_{m+1}^{\otimes j}} \otimes l^{\otimes n-j}$ . Otherwise, if  $i \neq j + 1$ , this follows from Proposition 4.8. If  $i = j + 1$  then Proposition 4.9 gives that  $\bar{\lambda}_{m,j}^n$  factors through  $\text{id}_{I_{m+1}^{\otimes j}} \otimes l^{\otimes n-j}$ . □

We have an analogous result for  $\bar{\rho}_{m,j}^n$  as well.

LEMMA 4.11. *Let  $m, n > 0$ ,  $j \in \{0, \dots, n - 1\}$ , and  $G$  be a graph. Consider a subobject  $X$  of  $N_{m+1}G$  which contains:*

- all  $n$ -cubes of  $N_{m+1}G$  which factor through  $r^{\otimes n}: I_{m+1}^{\otimes n} \rightarrow I_m^{\otimes n}$ ;
- (if  $n > 1$ ) for any  $x: I_{m+1}^{\otimes h} \otimes I_m^{\otimes k-h} \rightarrow G$  where  $k < n$  and  $h \leq k$ , the  $(k + 1)$ -cube  $x\bar{\rho}_{m,h-1}^n: I_{m+1}^{\otimes n+1} \rightarrow G$ ;
- (if  $j > 0$ ) for any  $x: I_{m+1}^{\otimes j} \otimes I_m^{\otimes n-j} \rightarrow G$ , the  $(n + 1)$ -cube  $x\bar{\rho}_{m,j-1}^n: I_{m+1}^{\otimes n+1} \rightarrow G$  of  $N_{m+1}G$ .

Then, for any  $n$ -cube of  $N_{m+1}G$  which factors through  $\text{id}_{I_{m+1}^{\otimes j+1}} \otimes r^{\otimes n-j-1}: I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  as some  $x: I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1} \rightarrow G$ , the restriction of the  $(n + 1)$ -cube  $x\bar{\rho}_{m,j}^n: I_{m+1}^{\otimes n+1} \rightarrow G$  to the open box  $x\bar{\rho}_{m,j}^n|_{\square_{n+1,1}^{n+1}}: \square_{n+1,1}^{n+1} \rightarrow N_{m+1}G$  factors through the inclusion  $X \hookrightarrow N_{m+1}G$ .

THEOREM 4.12. *Let  $m > 0$  and  $G$  be a graph. The maps  $l^*, r^*: N_mG \rightarrow N_{m+1}G$  are anodyne.*

*Proof.* We show that  $l^*$  is anodyne as the result for  $r^*$  is analogous. Let  $X_0$  denote the image of  $N_mG$  in  $N_{m+1}G$  under the embedding  $l^*: N_mG \hookrightarrow N_{m+1}G$ . We show  $N_{m+1}G$  can be obtained from  $X_0$  by a transfinite composition of pushouts along coproducts of open box inclusion. For  $n > 0$  and  $j \in \{1, \dots, n\}$ , let  $X_{n,j}$  be the subobject of  $N_{m+1}G$  generated by:

- $X_0$ ;
- (if  $n > 1$ ) for any  $x: I_{m+1}^{\otimes l} \otimes I_m^{\otimes k-l} \rightarrow G$  where  $k < n$  and  $l \leq k$ , the  $(k + 1)$ -cube  $x\bar{\lambda}_{m,l-1}^k: I_{m+1}^{\otimes n+1} \rightarrow G$  of  $N_{m+1}G$ ;
- for any  $x: I_{m+1}^{\otimes i} \otimes I_m^{\otimes n-i} \rightarrow G$  where  $i \leq j$ , the  $(n + 1)$ -cube  $x\bar{\lambda}_{m,i-1}^n: I_{m+1}^{\otimes n+1} \rightarrow G$  of  $N_{m+1}G$ .

By construction, there is a sequence of embeddings

$$X_0 \hookrightarrow X_{1,1} \hookrightarrow X_{2,1} \hookrightarrow X_{2,2} \hookrightarrow X_{3,1} \hookrightarrow \dots$$

Note that the subobject  $X_{n,n}$  contains all  $n$ -cubes of  $N_{m+1}G$ . This is because any  $n$ -cube  $x: I_{m+1}^{\otimes n} \rightarrow G$  is the  $\partial_{n+1,1}$ -face of  $x\bar{\lambda}_{m,n-1}^n: I_{m+1}^{\otimes n+1} \rightarrow G$ . By construction,  $X_{n,n}$  contains  $x\bar{\lambda}_{m,n-1}^n$ . Thus, it contains all faces of  $x\bar{\lambda}_{m,n-1}^n$ , including  $x$ . With this, we have

$$N_{m+1}G \cong \text{colim}(X_0 \hookrightarrow X_{1,1} \hookrightarrow X_{2,1} \hookrightarrow X_{2,2} \hookrightarrow \dots).$$

It remains to show  $X_{n,j+1}$  is a pushout of  $X_{n,j}$  along a coproduct of open box inclusions and  $X_{n+1,1}$  is a pushout of  $X_{n,n}$  along a coproduct of open box inclusions.

Fix  $n > 0$  and  $j \in \{1, \dots, n - 1\}$ . Let  $S_{n,j+1}$  be the set of  $n$ -cubes  $I_{m+1}^{\otimes n} \rightarrow G$  which factor through the map  $\text{id}_{I_{m+1}^{\otimes j+1}} \otimes l^{\otimes n-j-1}: I_{m+1}^{\otimes n} \rightarrow I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1}$  and are not contained in  $X_{n,j}$ . We write an element of  $S_{n,j+1}$  as  $x(\text{id}_{I_{m+1}^{\otimes j+1}} \otimes l^{\otimes n-j-1})$  for some  $x: I_{m+1}^{\otimes j+1} \otimes I_m^{\otimes n-j-1} \rightarrow G$ . By construction,  $X_{n,j+1}$  contains all  $n$ -cubes of  $S_{n,j+1}$  as such an  $n$ -cube  $x(\text{id}_{I_{m+1}^{\otimes j+1}} \otimes l^{\otimes n-j-1})$  is the

$\partial_{n+1,1}$ -face of  $x\bar{\lambda}_{m,j}^n$  (which is contained in  $X_{n,j+1}$  by construction). This gives a map

$$\coprod_{x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}) \in S_{n,j+1}} \square^{n+1} \xrightarrow{x\bar{\lambda}_{m,j}^n} X_{n,j+1}.$$

For each  $x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}) \in S_{n,j+1}$ , the restriction of the map  $x\bar{\lambda}_{m,j}^n$  to the open box  $\square_{n+1,1}^{n+1}$  factors through  $X_{n,j}$  by Lemma 4.10:

$$\coprod_{x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}) \in S_{n,j+1}} \square_{n+1,1}^{n+1} \xrightarrow{x\bar{\lambda}_{m,j}^n|_{\square}} X_{n,j}.$$

As  $x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1})$  is not in  $X_{n,j}$ , the  $(n+1)$ -cube  $x\bar{\lambda}_{m,j}^n$  is also not in  $X_{n,j}$  (as one of its faces is  $x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1})$ ). The generating cubes of  $X_{n,j+1}$  which are not contained in  $X_{n,j}$  are exactly those of the form  $x\bar{\lambda}_{m,j}^n$  for some  $x \in S_{n,j+1}$ . Thus, we may write  $X_{n,j+1}$  as the following pushout.

$$\begin{array}{ccc} \coprod_{x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}) \in S_{n,j+1}} \square_{n+1,1}^{n+1} & \xrightarrow{x\bar{\lambda}_{m,j}^n|_{\square}} & X_{n,j} \\ \downarrow & \lrcorner & \downarrow \\ \coprod_{x(\text{id}_{I_{m+1}}^{\otimes j+1} \otimes l^{\otimes n-j-1}) \in S_{n,j+1}} \square^{n+1} & \xrightarrow{x\bar{\lambda}_{m,j}^n} & X_{n,j+1} \end{array}$$

Thus, the map  $X_{n,j} \rightarrow X_{n,j+1}$  is a pushout along a coproduct of open box inclusions.

We now show the map  $X_{n,n} \rightarrow X_{n+1,1}$  is a pushout along a coproduct of open box inclusions. Let  $S_{n+1,1}$  be the set of  $(n+1)$ -cubes which factor through the map  $\text{id}_{I_{m+1}} \otimes l^{\otimes n}: I_{m+1}^{\otimes n+1} \rightarrow I_{m+1} \otimes I_m^{\otimes n}$  and are not contained in  $X_{n,n}$ . Similar to before, the generating cubes of  $X_{n+1,1}$  which are not contained in  $X_{n,n}$  are exactly those of the form  $x\bar{\lambda}_{m,0}^{n+1}$  for some  $x \in S_{n+1,1}$ . Thus, Lemma 4.10 similarly shows that  $X_{n+1,1}$  may be written as the following pushout.

$$\begin{array}{ccc} \coprod_{x(\text{id}_{I_{m+1}} \otimes l^{\otimes n}) \in S_{n+1,1}} \square_{n+1,0}^{n+1} & \xrightarrow{x\bar{\lambda}_{m,0}^{n+1}|_{\square}} & X_{n,n} \\ \downarrow & \lrcorner & \downarrow \\ \coprod_{x(\text{id}_{I_{m+1}} \otimes l^{\otimes n}) \in S_{n+1,1}} \square^{n+1} & \xrightarrow{x\bar{\lambda}_{m,0}^{n+1}} & X_{n+1,1} \end{array}$$

Thus, the map  $X_{n,n} \rightarrow X_{n+1,1}$  is a pushout along a coproduct of open box inclusions. □

From this, we conclude that the inclusion of the 1-nerve of a graph into its nerve is anodyne.

**COROLLARY 4.13.** *The natural map  $N_1G \rightarrow NG$  is anodyne.*

*Proof.* By Proposition 3.5,  $NG$  is a transfinite composition of the maps  $c^*: N_mG \rightarrow N_{m+2}G$  for  $m > 0$ . By Theorem 4.12, each  $c^*$  is anodyne. Thus, each component of the colimit cone  $N_m \rightarrow NG$  is anodyne. □

This gives us our main theorem.

*Proof of Theorem 4.1.* The first result is proven in Theorem 4.5. The second result is proven in Corollary 4.13. □

5. Consequences

**Proof of the conjecture of Babson, Barcelo, de Longueville, and Laubenbacher**

Using our main result, we obtain a proof of [BBdLL06, Theorem 5.2] which does not rely on the cubical approximation hypothesis [BBdLL06, Proposition 5.1].

THEOREM 5.1. *There is a natural group isomorphism  $A_n(G, v) \cong \pi_n(|N_1G|, v)$ .*

*Proof.* We have

$$\begin{aligned} A_n(G, v) &\cong \pi_n(NG, v) && \text{by Theorem 4.6} \\ &\cong \pi_n(|N|G, v) && \text{by Theorem 2.26} \\ &\cong \pi_n(|N|_1G, v) && \text{by Theorem 4.1, Corollary 2.17.} \end{aligned}$$

□

**Discrete homology of graphs**

In this subsection, we prove the discrete Hurewicz theorem, relating discrete homotopy and homology groups. To do so, we begin with a quick review of cubical homology (cf. [Mas80, BHS11, BGJW21]).

First, we recall the standard definition of homology (with integral coefficients) of a chain complex. A (bounded) *chain complex* (over  $\mathbb{Z}$ ) consists of a collection  $\{C_n \mid n \geq 0\}$  of abelian groups and, for  $n \geq 1$ , a group homomorphism  $\partial_n: C_n \rightarrow C_{n-1}$  such that  $\partial_{n-1}\partial_n = 0$ . A map  $f: C \rightarrow D$  of chain complexes consists of maps  $f_n: C_n \rightarrow D_n$  for  $n \geq 0$  which make the respective squares commute. We write  $\text{Ch}$  for the category of chain complexes. Define a functor  $H_*: \text{Ch} \rightarrow \text{Ab}^{\mathbb{N}}$  by taking a chain complex  $C$  to a sequence of homology groups given by

$$H_n C := \begin{cases} C_0 / \text{im } \partial_0 & n = 0 \\ \ker \partial_n / \text{im } \partial_{n+1} & n > 0. \end{cases}$$

We refer to  $H_n C$  as the *n*th homology of  $C$  with integer coefficients. One verifies that a map of chain complexes  $C \rightarrow D$  induces maps  $H_n C \rightarrow H_n D$  between their *n*th homologies.

Next, we explain how to construct a chain complex out of a cubical set via a construction analogous to the normalized complex in simplicial singular homology. We construct a functor  $N: \text{cSet}_* \rightarrow \text{Ch}$  by

$$(NX)_n := F_* X_n / DX_n$$

where  $F_* X_n$  is the free abelian group on the pointed set  $X_n$  and  $DX_n$  is the subgroup of  $F_* X_n$  generated by degenerate cubes, i.e. those that lie in the image of a degeneracy or a connection  $X_{n-1} \rightarrow X_n$ . The chain differentials  $\partial: (NX)_n \rightarrow (NX)_{n-1}$  are given by

$$\partial(x) := \sum_{\substack{i=1, \dots, n \\ \varepsilon=0,1}} (-1)^{i+\varepsilon} x \partial_{i,\varepsilon}.$$

DEFINITION 5.2. The *reduced cubical homology with integer coefficients* functor is the functor  $\tilde{H}_*: \text{cSet}_* \rightarrow \text{Ab}^{\mathbb{N}}$  given by the composite

$$\text{cSet}_* \xrightarrow{N} \text{Ch} \xrightarrow{H_*} \text{Ab}^{\mathbb{N}}.$$

We contrast the definition of *reduced* homology with that of *unreduced homology*, which is defined on the category of non-pointed cubical sets by using non-pointed free abelian groups.

Unlike in the construction of simplicial homology, we are forced to take the quotient of  $F_* X_\bullet$  by the subcomplex of cubes in the image of a degeneracy map. We do, however, have flexibility

regarding whether or not to quotient by the subcomplex of cubes in the image of a connection map, as the two complexes are quasi-isomorphic [BGJW21, Corollary 3.10].

DEFINITION 5.3 (cf. [BCW14, § 2]). The *reduced discrete homology* functor  $\widetilde{DH}_* : \text{Graph}_* \rightarrow \text{Ab}^{\mathbb{N}}$  is the composite of functors

$$\text{Graph}_* \xrightarrow{N_1} \text{cSet}_* \xrightarrow{\widetilde{H}_*} \text{Ab}^{\mathbb{N}}.$$

By the homotopy invariance of cubical homology [CKT23, Theorem 3.11], we have the following fact.

PROPOSITION 5.4. *Let  $f : (X, x) \rightarrow (Y, y)$  be a pointed cubical map. If  $f$  is a weak equivalence then  $H_*f : H_*(X, x) \rightarrow H_*(Y, y)$  is an isomorphism of graded abelian groups.*

From this, we deduce that the discrete homology of a graph is the same as the cubical homology of its nerve.

COROLLARY 5.5. *For a pointed graph  $(G, v)$ , the natural map  $N_1G \rightarrow NG$  induces an isomorphism*

$$\widetilde{H}_*(N_1G, v) \cong \widetilde{H}_*(NG, v).$$

of graded abelian groups.

*Proof.* This follows from Proposition 5.4 and Theorem 4.1. □

For any pointed Kan complex  $(X, x)$ , using the unit of the adjunction

$$F_* : \text{cSet}_* \rightleftarrows \text{Ab}^{\square^{\text{op}}} : U_*$$

between pointed cubical sets and cubical abelian groups, we may construct a natural map  $\pi_n(X, x) \rightarrow \widetilde{H}_*(X, x)$ , since  $\pi_n(U_*F_*(X, x)) \cong \widetilde{H}_n(X, x)$  by [CKT23, Theorem 4.11]. This is the *Hurewicz homomorphism*; cf. [CKT23, Definition 4.14].

We then have the classical theorem of Hurewicz, phrased in the language of cubical sets.

THEOREM 5.6 [CKT23, Theorem 4.16]. *Let  $n \geq 2$  and  $(X, x)$  be a pointed connected Kan complex. Suppose  $\pi_i(X, x) = 0$  for all  $i \in \{1, \dots, n - 1\}$ , i.e.  $X$  is  $n$ -connected. Then the Hurewicz homomorphism  $\pi_n(X, x) \rightarrow \widetilde{H}_n(X, x)$  is an isomorphism.*

DEFINITION 5.7. For any pointed connected graph  $(G, v)$  and  $n \geq 2$ , we therefore obtain the *discrete Hurewicz homomorphism*  $A_n(G, v) \rightarrow \widetilde{DH}_n(G, v)$  as the composite

$$\begin{aligned} A_n(G, v) &\cong \pi_n(NG, v) && \text{by Theorem 4.6} \\ &\rightarrow \widetilde{H}_n(NG) && \text{the Hurewicz homomorphism} \\ &\cong \widetilde{H}_n(N_1G) && \text{by Corollary 5.5} \\ &= \widetilde{DH}_n(G, v) && \text{by Definition 5.3.} \end{aligned}$$

One may verify that this map recovers the homomorphism defined in [Lut21, § 5.2]. We then have the expected discrete analogue of the Hurewicz theorem.

THEOREM 5.8 (Discrete Hurewicz theorem). *Let  $n \geq 2$  and  $(G, v)$  be a pointed connected graph. Suppose  $A_i(G, v) = 0$  for all  $i \in \{1, \dots, n - 1\}$ . Then the induced Hurewicz map  $A_n(G, v) \rightarrow \widetilde{DH}_n(G, v)$  is an isomorphism.*

*Proof.* This is an immediate consequence of the definition of the discrete Hurewicz homomorphism and Theorem 5.6. □

**Fibration category of graphs**

Via Theorem 4.1, we may view the nerve functor as a functor  $N: \mathbf{Graph} \rightarrow \mathbf{Kan}$  taking values in Kan complexes. From this, we induce a fibration category structure on the category of graphs.

**THEOREM 5.9.** *The category  $\mathbf{Graph}$  of graphs and graph maps carries a fibration category structure where:*

- the weak equivalences are the weak homotopy equivalences, i.e. maps  $f: G \rightarrow H$  such that, for all  $v \in G$  and  $n \geq 0$ , the map  $A_n f: A_n(G, v) \rightarrow A_n(H, f(v))$  is an isomorphism;
- the fibrations are maps  $f: G \rightarrow H$  which are sent to fibrations  $Nf: NG \rightarrow NH$  under the nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$ .

Before proving this, we consider factorization of the diagonal map separately. Recall that, for any graph  $G$ , we have a commutative triangle

$$\begin{array}{ccc}
 & PG & \\
 \swarrow & & \searrow^{(\partial_{1,0}^*, \partial_{1,1}^*)} \\
 G & \xrightarrow{(\text{id}_G, \text{id}_G)} & G \times G
 \end{array}$$

where  $G \hookrightarrow PG$  sends a vertex  $v$  to the constant path on  $v$ .

**LEMMA 5.10.** *For any graph  $G$ , applying  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  to the diagram*

$$\begin{array}{ccc}
 & PG & \\
 \swarrow & & \searrow^{(\partial_{1,0}^*, \partial_{1,1}^*)} \\
 G & \xrightarrow{(\text{id}_G, \text{id}_G)} & G \times G
 \end{array}$$

gives a factorization of the diagonal map  $NG \rightarrow NG \times NG$  as a weak equivalence followed by a fibration.

*Proof.* Applying naturality of the isomorphism in Proposition 3.10 to the maps  $\square^1 \rightarrow \square^0$  and  $\partial \square^1 \rightarrow \square^1$  gives that the diagram

$$\begin{array}{ccccc}
 NG & \longrightarrow & N(PG) & \longrightarrow & N(G \times G) \\
 \cong \downarrow & & \downarrow \cong & & \downarrow \cong \\
 NG & \longrightarrow & \text{hom}_R(\square^1, NG) & \longrightarrow & NG \times NG
 \end{array}$$

commutes. Our earlier paper [CK23, Proposition 2.22] shows that the bottom composite provides a factorization of the diagonal map into a weak equivalence followed by a fibration. Thus, the top left map is a weak equivalence and the top right map is a fibration.  $\square$

*Proof of Theorem 5.9.* Both the two-out-of-six property for weak equivalences and closure of acyclic fibrations under isomorphisms follow from Theorem 2.10. Pullbacks exist as  $\mathbf{Graph}$  is complete; they preserve (acyclic) fibrations by Proposition 3.8. Theorem 4.1 shows that the nerve of every graph is a Kan complex. Factorization of the diagonal map is shown in Lemma 5.10, and this gives the factorization of an arbitrary map via [Bro73, Factorization lemma].  $\square$

The following proposition gives a useful criterion for verifying whether a map is a fibration.

PROPOSITION 5.11. A graph map  $f: G \rightarrow H$  is a fibration if and only if, for any commutative square

$$\begin{array}{ccc} |\square_{i,\varepsilon}^n|_m & \longrightarrow & G \\ \downarrow & & \downarrow f \\ I_m^{\otimes n} & \longrightarrow & H \end{array}$$

there exist  $k \geq 0$  and a lift  $I_{m+2k}^{\otimes n} \rightarrow G$  of the following square.

$$\begin{array}{ccccc} |\square_{i,\varepsilon}^n|_{m+2k} & \xrightarrow{c_*^k} & |\square_{i,\varepsilon}^n|_m & \longrightarrow & G \\ \downarrow & & \downarrow & \nearrow \gamma & \downarrow f \\ I_{m+2k}^{\otimes n} & \xrightarrow{c_*^k} & I_m^{\otimes n} & \longrightarrow & H \end{array}$$

*Proof.* The map  $f$  is a fibration if and only if any commutative square

$$\begin{array}{ccc} \square_{i,\varepsilon}^n & \longrightarrow & NG \\ \downarrow & & \downarrow Nf \\ \square^n & \longrightarrow & NH \end{array}$$

admits a lift. As  $N$  is a sequential colimit, every such square and lift (if it exists) factors as

$$\begin{array}{ccccccc} \square_{i,\varepsilon}^n & \longrightarrow & N_m G & \xrightarrow{(c^*)^k} & N_{m+2k} G & \longrightarrow & NG \\ \downarrow & & \downarrow N_m f & \nearrow \gamma & \downarrow N_{m+2k} f & & \downarrow Nf \\ \square^n & \longrightarrow & N_m H & \xrightarrow{(c^*)^k} & N_{m+2k} H & \longrightarrow & NH \end{array}$$

for some  $m \geq 1$  and  $k \geq 0$ . The left two squares in this diagram correspond to a diagram

$$\begin{array}{ccccc} |\square_{i,\varepsilon}^n|_{m+2k} & \xrightarrow{(c_*)^k} & |\square_{i,\varepsilon}^n|_m & \longrightarrow & G \\ \downarrow & & \downarrow & \nearrow \gamma & \downarrow f \\ I_{m+2k}^{\otimes n} & \xrightarrow{(c_*)^k} & I_m^{\otimes n} & \longrightarrow & H \end{array}$$

in **Graph** by the realization-nerve adjunction. □

By definition, the nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  reflects (and preserves) fibrations. We show that it reflects weak equivalences as well.

PROPOSITION 5.12. The nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  reflects weak equivalences. That is, given a map  $f: G \rightarrow H$ , if  $Nf: NG \rightarrow NH$  is a weak equivalence then  $f$  is a weak equivalence.

*Proof.* This follows from Theorem 4.6 and [CK23, Theorem 4.7]. □

As the nerve functor preserves finite limits, it is straightforward to show it is exact.

PROPOSITION 5.13. The nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  is exact.

*Proof.* The nerve functor preserves fibrations and acyclic fibrations by definition and by Proposition 5.12, respectively. Proposition 3.8 shows that it preserves all finite limits. □

A consequence of Proposition 5.13 is that *fibration sequences* induce a long exact sequence of homotopy groups.

**THEOREM 5.14.** *Let  $f: (G, v) \rightarrow (H, w)$  be a fibration between pointed graphs and  $(F, v)$  be the fiber of  $f$  over  $w$ , i.e. the pullback*

$$\begin{array}{ccc} (F, v) & \xrightarrow{i} & (G, v) \\ \downarrow & \lrcorner & \downarrow f \\ (I_0, 0) & \xrightarrow{w} & (H, w) \end{array}$$

in  $\text{Graph}_*$ . Then, there is a long exact sequence

$$\begin{array}{ccccccc} \dots & \longrightarrow & A_n(F, v) & \xrightarrow{A_n i} & A_n(G, v) & \xrightarrow{A_n f} & A_n(H, w) \\ & & \searrow & & \searrow & & \searrow \\ & & A_{n-1}(F, v) & \xrightarrow{A_{n-1} i} & A_{n-1}(G, v) & \xrightarrow{A_{n-1} f} & A_{n-1}(H, w) \\ & & \searrow & & \searrow & & \searrow \\ & & \dots & \longleftarrow & \dots & & \dots \\ & & \searrow & & \searrow & & \searrow \\ & & A_1(F, v) & \xrightarrow{A_1 i} & A_1(G, v) & \xrightarrow{A_1 f} & A_1(H, w) \\ & & \searrow & & \searrow & & \searrow \\ & & A_0(F, v) & \xrightarrow{A_0 i} & A_0(G, v) & \xrightarrow{A_0 f} & A_0(H, w). \end{array}$$

*Proof.* By Proposition 5.13, the map  $Nf: (NG, v) \rightarrow (NH, w)$  is a fibration and its fiber is naturally isomorphic to  $(NF, v)$ . The result then follows by applying [CK23, Corollary 4.6] and Theorem 4.6. □

There are two large classes of examples of fibrations: fiber bundles and  $m$ -fibrations.

**DEFINITION 5.15** (Babson). For  $m \geq 1$ , a graph map  $f: G \rightarrow H$  is an  $m$ -fiber bundle with fiber  $F$  if, for any  $u: I_1^{\otimes n} \rightarrow H$ , the left map in the pullback diagram

$$\begin{array}{ccc} P & \longrightarrow & G \\ \downarrow & \lrcorner & \downarrow f \\ I_1^{\otimes n} & \xrightarrow{u} & H \end{array}$$

is a product projection  $I_1^{\otimes n} \times F \rightarrow I_1^{\otimes n}$  from the categorical product of  $I_1^{\otimes n}$  and  $F$ .

**DEFINITION 5.16.** A fiber bundle  $G \rightarrow H$  is *trivial* if it is a product projection  $H \times F \rightarrow H$ .

**PROPOSITION 5.17.** *Any pullback of an  $m$ -fiber bundle is an  $m$ -fiber bundle.*

*Proof.* This follows from the two-pullback lemma. □

**PROPOSITION 5.18.** *For  $m \geq 1$ , every  $(m + 1)$ -fiber bundle is an  $m$ -fiber bundle.*

*Proof.* Fix an  $(m + 1)$ -fiber bundle  $f: G \rightarrow H$  and a map  $u: I_m^{\otimes n} \rightarrow H$ . The map  $l^{\otimes n}: I_{m+1}^{\otimes n} \rightarrow I_m^{\otimes n}$  admits a section  $i: I_m^{\otimes n} \rightarrow I_{m+1}^{\otimes n}$ . By assumption, the left map in the pullback diagram

$$\begin{array}{ccccc} P & \longrightarrow & G & & \\ \downarrow & \lrcorner & \downarrow f & & \\ I_{m+1}^{\otimes n} & \xrightarrow{l^{\otimes n}} & I_m^{\otimes n} & \xrightarrow{u} & H \end{array}$$



is a product projection  $I_{m+1}^{\otimes n} \times F \rightarrow I_{m+1}^{\otimes n}$ . As the right map in the pullback square

$$\begin{array}{ccc} I_m^{\otimes n} \times F & \xrightarrow{i \times \text{id}} & I_{m+1}^{\otimes n} \times F \\ \downarrow & \lrcorner & \downarrow \\ I_m^{\otimes n} & \xrightarrow{i} & I_{m+1}^{\otimes n} \end{array}$$

is a product projection, the left map is as well. Thus, the outer square in

$$\begin{array}{ccccc} I_m^{\otimes n} \times F & \xrightarrow{i \times \text{id}} & I_{m+1}^{\otimes n} \times F & \longrightarrow & G \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow f \\ I_m^{\otimes n} & \xrightarrow{i} & I_{m+1}^{\otimes n} & \xrightarrow{uol^{\otimes n}} & H \end{array}$$

is a pullback whose left map is a product projection. □

Recall that a graph  $G$  is a *retract* of a graph  $H$  if there is an inclusion  $i: G \hookrightarrow H$  with a retraction  $r: H \rightarrow G$ , i.e.  $ri = \text{id}$ .

LEMMA 5.19. *If  $H$  is a retract of  $I_1^{\otimes n}$  for some  $n \geq 0$  then any 1-fiber bundle  $G \rightarrow H$  is trivial.*

*Proof.* As  $H$  is a retract of  $I_1^{\otimes n}$ , we fix an inclusion  $i: H \hookrightarrow I_1^{\otimes n}$  and retraction  $r: I_1^{\otimes n} \rightarrow H$ . Given a fiber bundle  $f: G \rightarrow H$ , the left map in the pullback diagram

$$\begin{array}{ccc} F \times I_1^{\otimes n} & \longrightarrow & G \\ \downarrow & \lrcorner & \downarrow f \\ I_1^{\otimes n} & \xrightarrow{r} & H \end{array}$$

is a product projection. By universal property of the pullback, the outer square in

$$\begin{array}{ccc} G & \xrightarrow{\text{id}} & H \\ \downarrow f & \lrcorner & \downarrow f \\ F \times I_1^{\otimes n} & \longrightarrow & H \\ \downarrow & \lrcorner & \downarrow \\ I_1^{\otimes n} & \xrightarrow{r} & H \end{array}$$

induces a map  $G \rightarrow F \times I_1^{\otimes n}$ . By the two-pullback lemma, the left square in

$$\begin{array}{ccccc} G & \cdots \longrightarrow & F \times I_1^{\otimes n} & \longrightarrow & G \\ f \downarrow & & \downarrow & \lrcorner & \downarrow f \\ H & \xleftarrow{i} & I_1^{\otimes n} & \xrightarrow{r} & H \end{array}$$

is a pullback. Therefore, the outer square in

$$\begin{array}{ccccc} G & \cdots \longrightarrow & F \times I_1^{\otimes n} & \longrightarrow & F \\ f \downarrow & & \downarrow & \lrcorner & \downarrow \\ H & \xleftarrow{i} & I_1^{\otimes n} & \longrightarrow & I_0 \end{array}$$

is a pullback, which proves that  $G$  is a product of  $H$  and  $F$ , and  $f$  is the projection onto the first component. □

**THEOREM 5.20.** *For any  $m \geq 1$ , a 1-fiber bundle is an  $m$ -fiber bundle.*

*Proof.* Fix a 1-fiber bundle  $f: G \rightarrow H$  and a map  $u: I_m^{\otimes n} \rightarrow H$ . By Proposition 5.17, the left map in the pullback diagram

$$\begin{array}{ccc} u^*G & \longrightarrow & G \\ \downarrow & \lrcorner & \downarrow f \\ I_m^{\otimes n} & \xrightarrow{u} & H \end{array}$$

is a 1-fiber bundle. By Lemma 5.19, it suffices to show  $I_m^{\otimes n}$  is a retract of  $I_1^{\otimes k}$  for some  $k \geq 0$ .

Define an embedding  $i: I_m \hookrightarrow I_1^{\otimes m}$  by sending a vertex  $k \in I_m$  to the tuple  $(x_1, \dots, x_m)$  where  $x_i = 1$  if  $i \leq k$  and  $x_i = 0$  if  $i > k$ . This map has a retraction  $r: I_1^{\otimes m} \rightarrow I_m$  which sends a tuple  $(x_1, \dots, x_m)$  to its sum  $\sum_{i=1}^m x_i$ . Thus, the map  $i^{\otimes n}: I_m^{\otimes n} \rightarrow I_1^{\otimes mn}$  has a retraction  $r^{\otimes n}: I_1^{\otimes mn} \rightarrow I_m^{\otimes n}$ . □

In particular, if  $f: X \rightarrow Y$  is an  $m$ -fiber bundle for some  $m \geq 1$  then it is a fiber bundle for all  $m \geq 1$ . In the remainder of this section, we refer to such an  $f$  as simply a *fiber bundle*.

**THEOREM 5.21.** *Every fiber bundle is a fibration.*

*Proof.* Fix a fiber bundle  $f: G \rightarrow H$ . We apply Proposition 5.11 and consider a lifting problem

$$\begin{array}{ccc} |\square_{i,\varepsilon}^n|_m & \longrightarrow & G \\ \downarrow & & \downarrow f \\ I_m^{\otimes n} & \longrightarrow & H \end{array}$$

Taking a pullback of the right and bottom map gives a factorization of this square as

$$\begin{array}{ccccc} |\square_{i,\varepsilon}^n|_m & \dashrightarrow & I_1^{\otimes n} \times F & \longrightarrow & G \\ \downarrow & & \downarrow & \lrcorner & \downarrow f \\ I_m^{\otimes n} & \xrightarrow{\text{id}} & I_m^{\otimes n} & \longrightarrow & H \end{array}$$

The middle map is a fibration since it is a product projection. Hence, there exists  $k \geq 0$  such that the outer square in

$$\begin{array}{ccccc} |\square_{i,\varepsilon}^n|_{m+2k} & \xrightarrow{c_*^k} & |\square_{i,\varepsilon}^n|_m & \dashrightarrow & I_1^{\otimes n} \times F \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ I_{m+2k}^{\otimes n} & \xrightarrow{c_*^k} & I_m^{\otimes n} & \xrightarrow{\text{id}} & I_m^{\otimes n} \end{array}$$

admits a lift. Post-composing this lift with the map  $I_1^{\otimes n} \times F \rightarrow G$  gives a lift of the outer square in

$$\begin{array}{ccccc} |\square_{i,\varepsilon}^n|_{m+2k} & \xrightarrow{c_*^k} & |\square_{i,\varepsilon}^n|_m & \longrightarrow & G \\ \downarrow & & \downarrow & \nearrow & \downarrow f \\ I_{m+2k}^{\otimes n} & \xrightarrow{c_*^k} & I_m^{\otimes n} & \longrightarrow & H \end{array} \quad \square$$

COROLLARY 5.22. A pointed fiber bundle  $f: (G, v) \rightarrow (H, w)$  with fiber  $(F, v)$  induces a long exact sequence

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & A_n(F, v) & \xrightarrow{A_n i} & A_n(G, v) & \xrightarrow{A_n f} & A_n(H, w) \\
 & & \searrow & & \searrow & & \searrow \\
 & & A_{n-1}(F, v) & \xrightarrow{A_{n-1} i} & A_{n-1}(G, v) & \xrightarrow{A_{n-1} f} & A_{n-1}(H, w) \\
 & & \searrow & & \searrow & & \searrow \\
 & & \dots & \longleftarrow & \dots & & \dots \\
 & & A_1(F, v) & \xrightarrow{A_1 i} & A_1(G, v) & \xrightarrow{A_1 f} & A_1(H, w) \\
 & & \searrow & & \searrow & & \searrow \\
 & & A_0(F, v) & \xrightarrow{A_0 i} & A_0(G, v) & \xrightarrow{A_0 f} & A_0(H, w).
 \end{array}$$

of  $A$ -homotopy groups.

*Proof.* This follows from Theorems 5.21 and 5.14. □

The second class of maps which are fibrations is the class of  $m$ -fibrations.

DEFINITION 5.23. For  $m \geq 1$ , a graph map  $f: G \rightarrow H$  is an  $m$ -fibration if  $N_m f: N_m G \rightarrow N_m H$  is a Kan fibration.

PROPOSITION 5.24. Let  $m, k \geq 1$ .

- (i) The inclusion  $\{0, km\} \rightarrow I_{km}$  is in the saturation of  $\{\emptyset \rightarrow I_0, \{0, m\} \rightarrow I_m\}$ .
- (ii) For  $\varepsilon = 0, 1$ , the end-point inclusion  $\{\varepsilon km\} \rightarrow I_{km}$  is in the saturation of  $\{\{\varepsilon m\} \rightarrow I_m\}$ .
- (iii) The inclusion  $\partial I_{km}^{\otimes n} \rightarrow I_{km}^{\otimes n}$  is in the saturation of  $\{\partial I_m^{\otimes j} \rightarrow I_m^{\otimes j} \mid j \leq n\}$ .
- (iv) The inclusion  $|\Gamma_{i,\varepsilon}^n|_{km} \rightarrow I_{km}^{\otimes n}$  is in the saturation of  $\{|\Gamma_{i,\varepsilon}^j|_m \rightarrow I_m^{\otimes j} \mid j \leq n\}$ .

*Proof.* (i) By induction, if  $\partial I_{km} \rightarrow I_m$  lies in the saturation then the bottom map in the pushout

$$\begin{array}{ccc}
 \{0, km\} \sqcup \{0, m\} & \longrightarrow & I_{km} \sqcup I_m \\
 \downarrow [f,g] & & \downarrow \\
 \{0, km, (k+1)m\} & \longrightarrow & I_{(k+1)m}
 \end{array}$$

is in the saturation as a pushout of the coproduct of maps in the saturation, where  $[f, g]$  is defined by

$$\begin{aligned}
 f(0) &= 0, & f(km) &= km, \\
 g(0) &= km, & g(m) &= (k+1)m.
 \end{aligned}$$

The inclusion  $\{0, (k+1)m\} \rightarrow \{0, km, (k+1)m\}$  is a pushout along  $\emptyset \rightarrow I_0$ , thus the composite  $\{0, (k+1)m\} \rightarrow I_{(k+1)m}$  lies in the saturation.

(ii) By induction, if  $\{\varepsilon m\} \rightarrow I_m$  lies in the saturation then the bottom map in the pushout

$$\begin{array}{ccc}
 I_0 & \xrightarrow{\varepsilon km} & I_{km} \\
 (1-\varepsilon)m \downarrow & & \downarrow \\
 I_m & \longrightarrow & I_{(k+1)m}
 \end{array}$$

lies in the saturation. Pre-composing with the end-point inclusion  $\{\varepsilon m\} \rightarrow I_m$  gives the endpoint inclusion  $\{\varepsilon(k+1)m\} \rightarrow I_{(k+1)m}$ .

(iii) The inclusion  $\partial I_{km}^{\otimes n} \rightarrow I_{km}^{\otimes n}$  may be written as a pushout product

$$(\partial I_{km}^{\otimes n} \rightarrow I_{km}^{\otimes n}) = (\{0, km\} \rightarrow I_{km})^{\hat{\otimes} n}.$$

Noting the equality

$$\{\emptyset \rightarrow I_0, \{0, m\} \rightarrow I_m\}^{\hat{\otimes} n} = \{\partial I_m^{\otimes j} \rightarrow I_m^{\otimes j} \mid j \leq n\},$$

this follows from (i) by [HSS00, Proposition 5.3.4].

(iv) The inclusion  $|\square_{i,\varepsilon}^n|_{km} \rightarrow I_{km}^{\otimes n}$  may be written as a pushout product

$$(|\square_{i,\varepsilon}^n|_{km} \rightarrow I_{km}^{\otimes n}) = (\{0, km\} \rightarrow I_{km})^{\hat{\otimes}(i-1)} \hat{\otimes} (\{\varepsilon km\} \rightarrow I_{km}) \hat{\otimes} (\{0, km\} \rightarrow I_{km})^{\hat{\otimes}n-i}.$$

Noting the equality

$$\begin{aligned} \{\emptyset \rightarrow I_0, \{0, m\} \rightarrow I_m\}^{\hat{\otimes}i-1} \hat{\otimes} \{\{\varepsilon m\} \rightarrow I_m\} \hat{\otimes} \{\emptyset \rightarrow I_0, \{0, m\} \rightarrow I_m\}^{\hat{\otimes}n-i} \\ = \{|\square_{i,\varepsilon}^j|_m \rightarrow I_m^{\otimes j} \mid j \leq n\}, \end{aligned}$$

this follows from (i) and (ii) by [HSS00, Proposition 5.3.4]. □

**THEOREM 5.25.** *Let  $k, m \geq 1$ . Every  $m$ -fibration is both a  $km$ -fibration and a fibration.*

*Proof.* Let  $f: G \rightarrow H$  be an  $m$ -fibration. We have that  $f$  is a  $km$ -fibration by Proposition 5.24. To see  $f$  is a fibration, we apply Proposition 5.11. For a commutative square

$$\begin{array}{ccc} |\square_{i,\varepsilon}^n|_t & \longrightarrow & G \\ \downarrow & & \downarrow \\ I_t^{\otimes n} & \longrightarrow & H \end{array}$$

let  $k \geq 0$  be the smallest non-negative integer such that  $t + k$  is a multiple of  $m$ . As  $f$  is a  $(t + k)$ -fibration, the required lift exists. □

Proposition 5.13 also shows that the nerve functor preserves loop spaces.

**PROPOSITION 5.26.** *The square*

$$\begin{array}{ccc} \text{Graph}_* & \xrightarrow{N} & \text{Kan}_* \\ \Omega \downarrow & & \downarrow \Omega \\ \text{Graph}_* & \xrightarrow{N} & \text{Kan}_* \end{array}$$

*commutes up to natural isomorphism.*

*Proof.* Fix a pointed graph  $(G, v)$ . By Proposition 1.22, the square

$$\begin{array}{ccc} \Omega(G, v) & \longrightarrow & PG \\ \downarrow & \ulcorner & \downarrow (\partial_{1,0}^*, \partial_{1,1}^*) \\ I_0 & \xrightarrow{(v,v)} & G \times G \end{array}$$

is a pullback. Proposition 3.8 shows that the nerve functor preserves this pullback, thus the square

$$\begin{array}{ccc} N(\Omega(G, v)) & \longrightarrow & N(PG) \\ \downarrow & \ulcorner & \downarrow (\partial_{1,0}^*, \partial_{1,1}^*) \\ NI_0 & \xrightarrow{(v,v)} & N(G \times G) \end{array}$$

is a pullback. Proposition 3.8 also shows the nerve preserves products and the terminal object. Lemma 5.10 implies that  $N(PG) \cong \text{hom}_R(\square^1, NG)$ , hence the square

$$\begin{array}{ccc} N(\Omega(G, v)) & \longrightarrow & \text{hom}_R(\square^1, NG) \\ \downarrow & \lrcorner & \downarrow (\partial_{1,0}^*, \partial_{1,1}^*) \\ \square^0 & \xrightarrow{(v,v)} & NG \times NG \end{array}$$

is a pullback. That is,  $N(\Omega(G, v)) \cong \Omega(NG, v)$ . □

By Proposition 2.9, the category  $\text{Graph}_*$  of pointed graphs has a fibration category structure as well. Thus, the loop graph functor  $\Omega: \text{Graph}_* \rightarrow \text{Graph}_*$  is a functor between fibration categories.

**THEOREM 5.27.** *The loop graph functor  $\Omega: \text{Graph}_* \rightarrow \text{Graph}_*$  is exact.*

*Proof.* By Proposition 5.13, Theorem 2.27, the composite  $\Omega(N-, -): \text{Graph}_* \rightarrow \text{cSet}_*$  is exact. Applying Proposition 5.26, this gives that the composite  $N(\Omega(-, -)): \text{Graph}_* \rightarrow \text{cSet}_*$  is exact. The nerve functor reflects fibrations and acyclic fibrations as, by definition, it creates them. It reflects finite limits by Corollary 3.13. From this, it follows that  $\Omega: \text{Graph}_* \rightarrow \text{Graph}_*$  preserves fibrations, acyclic fibrations, and finite limits. □

**Cubical enrichment of the category of graphs**

Recall (e.g. from [Rie14, Definition 3.4.1]) that a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between monoidal categories  $(\mathcal{C}, \otimes_{\mathcal{C}}, I_{\mathcal{C}})$  and  $(\mathcal{D}, \otimes_{\mathcal{D}}, I_{\mathcal{D}})$  is *lax monoidal* if there exist natural transformations

$$Fc \otimes_{\mathcal{D}} Fc' \rightarrow F(c \otimes_{\mathcal{C}} c') \quad \text{and} \quad I_{\mathcal{D}} \rightarrow FI_{\mathcal{C}}$$

subject to the associativity and unitality conditions, which we omit here.

**LEMMA 5.28.**

- (i) *For  $m \geq 1$ , the functor  $N_m: \text{Graph} \rightarrow \text{cSet}$  is lax monoidal.*
- (ii) *The functor  $N: \text{Graph} \rightarrow \text{cSet}$  is lax monoidal.*

*Proof.* By Proposition 3.6, the functors  $|-|_m: \text{cSet} \rightarrow \text{Graph}$  are strong monoidal, and hence, in particular, oplax monoidal. Thus, their right adjoints  $N_m: \text{Graph} \rightarrow \text{cSet}$  are lax monoidal, proving (i).

Clearly,  $NI_0 \cong \square^0$ . As the geometric product preserves colimits in each variable and colimits commute with colimits, the cubical set  $NG \otimes NH$  (for graphs  $G$  and  $H$ ) is the colimit of the following diagram.

$$NG \otimes NH = \text{colim} \left( \begin{array}{ccccccc} N_1G \otimes N_1H & \longleftarrow & N_2G \otimes N_1H & \longleftarrow & N_3G \otimes N_1H & \longleftarrow & \dots \\ \downarrow & & \downarrow & & \downarrow & & \\ N_1G \otimes N_2H & \longleftarrow & N_2G \otimes N_2H & \longleftarrow & N_3G \otimes N_2H & \longleftarrow & \dots \\ \downarrow & & \downarrow & & \downarrow & & \\ N_1G \otimes N_3H & \longleftarrow & N_2G \otimes N_3H & \longleftarrow & N_3G \otimes N_3H & \longleftarrow & \dots \\ \downarrow & & \downarrow & & \downarrow & & \\ \vdots & & \vdots & & \vdots & & \ddots \end{array} \right)$$

Computing this colimit componentwise in **Set**, one verifies that the colimit of the diagonal

$$\operatorname{colim}(N_1G \otimes N_1H \hookrightarrow N_2G \otimes N_2H \hookrightarrow N_3G \otimes N_3H \hookrightarrow \dots)$$

computes the same cubical set. As  $N(G \otimes H)$  is the colimit

$$N(G \otimes H) = \operatorname{colim}(N_1(G \otimes H) \hookrightarrow N_2(G \otimes H) \hookrightarrow N_3(G \otimes H) \hookrightarrow \dots),$$

the lax monoidal maps  $N_mG \otimes N_mH \rightarrow N_m(G \otimes H)$  induce a map on colimits  $NG \otimes NH \rightarrow N(G \otimes H)$  which satisfies the required associativity and unitality conditions.  $\square$

*Remark 5.29.* An alternative proof of (ii) can be given using [DKLS24, Proposition 1.24], which gives an explicit description of the geometric product of cubical sets. Explicitly, the  $n$ -cubes of  $NG \otimes NH$  are in bijective correspondence with pairs of cubes

$$(\square^k \rightarrow NG, \square^l \rightarrow NH),$$

where  $k + l = n$ . By definition of  $N$ , each such pair corresponds in turn to pair of maps

$$(I_\infty^{\otimes k} \rightarrow G, I_\infty^{\otimes l} \rightarrow H)$$

that stabilize in all directions. Taking products of these maps, we obtain a map  $I_\infty^{\otimes n} \rightarrow G \otimes H$  that stabilizes in all directions.

**COROLLARY 5.30.** *The nerve functor  $N: \mathbf{Graph} \rightarrow \mathbf{cSet}$  preserves homotopy equivalences.*

We describe the notion of enriched categories informally, with a reference to [Rie14, Definition 3.3.1] in lieu of a fully formal statement.

**DEFINITION 5.31.** For a monoidal category  $(\mathcal{V}, \otimes, 1)$ , a  $(\mathcal{V}, \otimes, I)$ -enriched category  $\mathcal{C}$  consists of

- a class of objects  $\operatorname{ob} \mathcal{C}$ ;
- for  $X, Y \in \operatorname{ob} \mathcal{C}$ , a *morphism* object  $\mathcal{C}(X, Y) \in \mathcal{V}$ ;
- for  $X, Y, Z \in \operatorname{ob} \mathcal{C}$ , a *composition* morphism  $\circ: \mathcal{C}(Y, Z) \otimes \mathcal{C}(X, Y) \rightarrow \mathcal{C}(X, Z)$  in  $\mathcal{V}$ ;
- for  $X \in \operatorname{ob} \mathcal{C}$ , an *identity* morphism  $1 \rightarrow \mathcal{C}(X, X)$  in  $\mathcal{V}$ ,

subject to appropriate associativity and unitality axioms (cf. [Rie14, Definition 3.3.1]).

*Example 5.32.* Any locally small category is a  $(\mathbf{Set}, \times, \{*\})$ -enriched category, where the objects, morphism sets, composition function, and identity morphisms are as usually defined.

*Example 5.33.* Any closed monoidal category is enriched over itself. In particular,  $\mathbf{Graph}_{\mathcal{C}}$  is a  $(\mathbf{Graph}, \otimes, I_0)$ -enriched category where:

- $\operatorname{ob} \mathbf{Graph}_{\mathcal{C}}$  is the collection of all graphs;
- for graphs  $G, H$ , the morphism graph is  $\operatorname{hom}^{\otimes}(G, H)$ ;
- for graphs  $G, H, J$ , the composition morphism is the graph map given by composition of graph maps regarded as vertices,

$$\operatorname{hom}^{\otimes}(H, J) \otimes \operatorname{hom}^{\otimes}(G, H) \rightarrow \operatorname{hom}^{\otimes}(G, J);$$

- the identity morphism is the identity map on  $G$  as a vertex  $\operatorname{id}_G: I_0 \rightarrow \operatorname{hom}^{\otimes}(G, G)$ .

*Example 5.34.* Since enrichment can be transferred along lax monoidal functors [Rie14, Lemma 3.4.3], Lemma 5.28 implies there is a  $(\mathbf{cSet}, \otimes, \square^0)$ -enriched category  $\mathbf{Graph}_{\square}$  of graphs where  $N(\operatorname{hom}^{\otimes}(G, H))$  is the morphism cubical set. Composition and identity morphisms are defined analogously.

DEFINITION 5.35. A  $(\mathbf{cSet}, \otimes, \square^0)$ -enriched category  $\mathcal{C}$  is *locally Kan* if, for all  $X, Y \in \text{ob } \mathcal{C}$ , the cubical set  $\mathcal{C}(X, Y)$  is a Kan complex.

THEOREM 5.36. *The  $(\mathbf{cSet}, \otimes, \square^0)$ -enriched category  $\text{Graph}_{\square}$  of graphs is locally Kan.*

*Proof.* This follows from Theorem 4.1. □

By the results of [KV20], we have established a presentation of the  $(\infty, 1)$ -category of graphs (localized at  $A$ -homotopy equivalences). In subsequent work, we use this presentation to show that  $A$ -homotopy equivalences are not part of a model structure on the category of graphs, as well as to study homotopy limits in discrete homotopy theory.

#### ACKNOWLEDGEMENTS

We thank Bob Lutz for comments on an earlier draft of this paper and for the beautiful talks on discrete homotopy theory (at MSRI in Spring 2020 and at Western University in Fall 2020) that got us interested in this area. We are also very grateful to Yeonjoon Choi, Udit Mavinkurve, and Mohabat Tarkeshian, members of an informal seminar on discrete homotopy theory that we organized in Spring 2021. In particular, Choi was the first to observe Lemma 5.28. Finally, we would like to thank Eric Babson for suggesting to us the notion of a fiber bundle (cf. Definition 5.15).

#### CONFLICTS OF INTEREST

None.

#### FINANCIAL SUPPORT

During the work on this paper, the first author was partially supported by an NSERC Undergraduate Student Research Award and the second author was partially supported by an NSERC Discovery Grant. We thank NSERC for its generosity.

#### JOURNAL INFORMATION

*Compositio Mathematica* is owned by the Foundation Compositio Mathematica and published by the London Mathematical Society in partnership with Cambridge University Press. All surplus income from the publication of *Compositio Mathematica* is returned to mathematics and higher education through the charitable activities of the Foundation, the London Mathematical Society and Cambridge University Press.

#### REFERENCES

- Atk74 R. H. Atkin, *An algebra for patterns on a complex. I*, Internat. J. Man-Mach. Stud. **6** (1974), 285–307.
- Atk76 R. H. Atkin, *An algebra for patterns on a complex. II*, Internat. J. Man-Mach. Stud. **8** (1976), 483–498.
- BBdLL06 E. Babson, H. Barcelo, M. de Longueville and R. Laubenbacher, *Homotopy theory of graphs*, J. Algebraic Combin. **24** (2006), 31–44.
- BCW14 H. Barcelo, V. Capraro and J. A. White, *Discrete homology theory for metric spaces*, Bull. Lond. Math. Soc. **46** (2014), 889–905.
- BGJW21 H. Barcelo, C. Greene, A. S. Jarrah and V. Welker, *Homology groups of cubical sets with connections*, Appl. Categ. Structures **29** (2021), 415–429.
- BKLW01 H. Barcelo, X. Kramer, R. Laubenbacher and C. Weaver, *Foundations of a connectivity theory for simplicial complexes*, Adv. Appl. Math. **26** (2001), 97–128.

- BL05 H. Barcelo and R. Laubenbacher, *Perspectives on A-homotopy theory and its applications*, Discrete Math. **298** (2005), 39–61.
- BR13 T. Barthel and E. Riehl, *On the construction of functorial factorizations for model categories*, Algebr. Geom. Topol. **13** (2013), 1089–1124.
- BH81 R. Brown and P. J. Higgins, *Colimit theorems for relative homotopy groups*, J. Pure Appl. Algebra **22** (1981), 11–41.
- BHS11 R. Brown, P. J. Higgins and R. Sivera, *Nonabelian algebraic topology: Filtered spaces, crossed complexes, cubical homotopy groupoids*, EMS Tracts in Mathematics, vol. 15 (European Mathematical Society (EMS), Zürich, 2011); with contributions by Christopher D. Wensley and Sergei V. Soloviev.
- BM17 U. Buchholtz and E. Morehouse, *Varieties of cubical sets*, in *Relational and algebraic methods in computer science*, Lecture Notes in Computer Science, vol. 10226 (Springer, Cham, 2017), 77–92.
- Bro73 K. S. Brown, *Abstract homotopy theory and generalized sheaf cohomology*, Trans. Amer. Math. Soc. **186** (1973), 419–458.
- CK23 D. Carranza and K. Kapulkin, *Homotopy groups of cubical sets*, Expo. Math. **41** (2023), 125518.
- CKT23 D. Carranza, K. Kapulkin and A. Tonks, *The Hurewicz theorem for cubical homology*, Math. Z. **305** (2023), 61.
- Cis06 D.-C. Cisinski, *Les préfaisceaux comme modèles des types d'homotopie*, Astérisque, vol. 308 (2006).
- DKLS24 B. Doherty, K. Kapulkin, Z. Lindsey and C. Sattler, *Cubical models of  $(\infty, 1)$ -categories*, Mem. Amer. Math. Soc. **297** (2024), 1484.
- Hir03 P. S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99 (American Mathematical Society, Providence, RI, 2003).
- Hov99 M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63 (American Mathematical Society, Providence, RI 1999).
- HSS00 M. Hovey, B. Shipley and J. Smith, *Symmetric spectra*, J. Amer. Math. Soc. **13** (2000), 149–208.
- KV20 K. Kapulkin and V. Voevodsky, *A cubical approach to straightening*, J. Topol. **13** (2020), 1682–1700.
- KL98 X. H. Kramer and R. C. Laubenbacher, *Combinatorial homotopy of simplicial complexes and complex information systems*, in *Applications of computational algebraic geometry (San Diego, CA, 1997)*, Proceedings of Symposia in Applied Mathematics, vol. 53 (American Mathematical Society, Providence, RI, 1998), 91–118.
- Lur09 J. Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170 (Princeton University Press, Princeton, NJ, 2009).
- Lut21 B. Lutz, *Higher discrete homotopy groups of graphs*, Algebr. Comb. **4** (2021), 69–88.
- Mal83 G. Malle, *A homotopy theory for graphs*, Glas. Mat. Ser. III **18** (1983), 3–25.
- Mas80 W. S. Massey, *Singular homology theory*, Graduate Texts in Mathematics, vol. 70 (Springer, New York, 1980).
- May75 J. P. May, *Classifying spaces and fibrations*, Mem. Amer. Math. Soc. **1** (1975), 155.
- MZ19 F. Mémoli and L. Zhou, *Persistent homotopy groups of metric spaces*, Preprint (2019), [arxiv:1912.12399](https://arxiv.org/abs/1912.12399).
- Rie14 E. Riehl, *Categorical homotopy theory*, New Mathematical Monographs, vol. 24 (Cambridge University Press, Cambridge, 2014).
- RV14 E. Riehl and D. Verity, *The theory and practice of Reedy categories*, Theory Appl. Categ. **29** (2014), 256–301.



- Szu16 K. Szumilo, *Homotopy theory of cofibration categories*, Homology Homotopy Appl. **18** (2016), 345–357.
- vdBG12 B. van den Berg and R. Garner, *Topological and simplicial models of identity types*, ACM Trans. Comput. Log. **13** (2012), 3.

D. Carranza dcarran3@jh.edu

Department of Mathematics, Krieger Hall 211, Johns Hopkins University,  
3400 N. Charles Street, Baltimore, MD 21218, USA

K. Kapulkin kkapulki@uwo.ca

Department of Mathematics, Middlesex College 255C, The University of Western Ontario,  
1151 Richmond Street, London, ON, Canada N6A 5B7