

⁸Hematology Department, Hedi chaker University hospital, sfax, Tunisia; ⁹Occupational Department, HediChaker Hospital, sfax, Tunisia; ¹⁰Community Health And Epidemiology Department, HediChaker Hospital, sfax, Tunisia; ¹¹Pneumology, HediChaker Hospital, sfax, Tunisia and ¹²Department Of Occupational Medicine, HEDI CHAKER hospital, SFAX, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.753

Introduction: Quarantine for suspected patients of being infected by the COVID-19 can lead to negative consequences for mental health and the appearance of depressive symptoms.

Objectives: To assess the prevalence of depression in quarantined patients, and to analyze the associated factors.

Methods: This was a descriptive and analytical survey, carried out from April 4 to May 30, 2020, with 149 patients consulting the COVID-19 sorting box at the Hedi Chaker CHU in Sfax. Suspected COVID-19 patients were contacted by phone during their quarantine and invited to participate in our study. The Patient Health Questionnaire (PHQ-9) scale was used to assess the severity of depression. Cutoffs of 5, 10, 15, and 20 represent minimal, mild, moderate, moderately severe, and severe levels of depression based on PHQ-9 scores. A cutoff score of 10 determines major depression.

Results: The results showed a prevalence of major depression of 10.7%. Of all patients, 89.3% had minimal to mild depression; 10% had moderate to moderately severe depression and 0.7% had severe depression. The PHQ-9 score was statistically correlated with travel to a suspect area during the 14 days preceding the consultation ($p = 0.008$), contact with a subject confirmed COVID-19 ($p = 0.01$), previous follow-up in psychiatry ($p = 0.047$), the change of residence during quarantine ($p = 0.045$), the fear of transmitting the disease to relatives ($p = 0.00$) and the positive result of the nasopharyngeal swab ($p = 0.00$). **Conclusions:** Psychological distress was felt in our patients. We recommend that necessary measures should be taken to combat depression.

Keywords: quarantine; suspected patients; Depression; COVID-19 pandemic

EPP0371

Beliefs about coronavirus: Relationship with magical thinking and adherence to self-isolation regimen

S. Kumchenko^{1*}, E. Rasskazova¹ and A. Tkhostov²

¹Clinical Psychology, Moscow State University, Moscow, Russian Federation and ²Department Of Neuro- And Pathopsychology, Faculty Of Psychology, Lomonosov Moscow State University, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.754

Introduction: Pandemic is accompanied by “infodemic” that is related to higher anxiety (Moghanibashi-Mansourieh, 2020; Roy et al., 2020; Huang, Zhao, 2020). We suggest that indefinite and stressful situation of pandemic provoke magical thinking leading to lower adherence with recommendations for self-isolation.

Objectives: The aim was to reveal the structure of beliefs about reasons, manifestation and consequences of coronavirus and their relationship with magical thinking, anxiety and COVID-19-related behaviour.

Methods: In April 2020 (2-3 weeks of self-isolation regimen) 402 adults aged 18-64 years old filled checklist including beliefs about pandemic (based on the model of Leventhal et al., 2003), Magical Ideation Scale Eckblad, Chapman, 1983) as well as scales measuring anxiety and protective behaviour in pandemic and monitoring of information about coronavirus (Tkhostov, Rasskazova, 2020).

Results: Factor analysis revealed three groups of radical beliefs about coronavirus (48.6% of variance, Cronbach’s alphas .62-.75). Belief about the particular meaning of coronavirus was associated with the magical thinking ($r=.21$), less anxiety about infection ($r=-.19$) and poorer adherence to self-isolation ($r=-.26$). Belief in the negligence as a cause of coronavirus was more typical for those with better adherence ($r=.18$) while catastrophic beliefs about the consequences of pandemic were related to frequent monitoring of the information about the pandemic ($r=.24$), and anxiety regarding future negative consequences of the pandemic ($r=.46$).

Conclusions: Dysfunctional beliefs about coronavirus could be a factor of poorer adherence related to magical thinking and could be addressed in psychological interventions. Research is supported by the Russian Foundation for Basic Research, project No. 20-04-60072.

Keywords: infodemic; coronavirus; magical thinking

EPP0372

The spectrum and severity of psychopathological symptoms in previously healthy individuals who have had severe COVID-19 pneumonia.

I. Belokrylov^{1*}, N. Kotovskaya², N. Deryugina³, A. Izyumov⁴, K. Eruslanova⁵ and M. Balaeva⁶

¹Department Of Psychiatry And Medical Psychology, Peoples Friendship University of Russia (RUDN University), Moscow, Russian Federation; ²Department Of Psychiatry And Medical Psychology, Peoples Friendship University of Russia (RUDN University), Moscow, Russian Federation; ³Department Of Math, Lomonosov Moscow State University, Moscow, Russian Federation; ⁴Laboratory Of General Geriatrics And Neurogeriatrics, Pirogov RNMU – RCRCG, Moscow, Russian Federation; ⁵Laboratory Of Cardiovascular Aging, Pirogov RNMU – RCRCG, Moscow, Russian Federation and ⁶Department Of Aging Diseases, Pirogov RNMU – RCRCG, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.755

Introduction: The medical novelty of COVID-19 requires a comprehensive study of its impact on various areas of human health, including mental health.

Objectives: To study the spectrum and severity of psychopathological disorders in previously healthy patients of different age groups who have had moderate and severe COVID-19 pneumonia.

Methods: Immediately after stabilization of the physical condition, patients completed the Symptom Checklist-90-R, designed to assess 11 parameters: somatization (SOM), obsessive-compulsive (OS), interpersonal sensitivity (INT), depression (DEP), anxiety (ANX), hostility (HOS), phobic anxiety (PHOB), paranoid ideas (PAR), psychoticism (PSY). Patients with cognitive impairment were excluded.

Results: The study involved 148 patients aged from 26 to 84 years. In the general sample, psychopathological symptoms were detected

mainly on the SOM, DEP, ANX, HOS scales. To a lesser extent - on the INT and PAR scales; were practically not determined on the PSY and PHOB scales. Most of the symptoms are significantly more intense in patients over 46 years old ($n = 129$) compared with the younger population (<46 years old, $n = 19$). Older patients according to SOM revealed 1.23 points (IQR 0.5) versus 0.85 (IQR 0.7) among young people, DEP - 0.88 (IQR 0.44) vs. 0.47 (IQR 0.44), ANX - 0.66 (IQR 0.44) vs. 0.43 (IQR 0.29), OS - 0.55 (IQR 0.5) vs. 0.31 (IQR 0.25) and HOS - 0.46 (IQR 0.34) vs. 0.29 (IQR 0.09).

Conclusions: Patients recovering from severe COVID-19 pneumonia require psychiatric evaluation and subsequent differentiated psychotherapeutic rehabilitation, especially for the age group over 46.

Conflict of interest: No significant relationships.

EPP0373

Anxiety and depression among students in a greek university amidst COVID-19 pandemic

E. Sazakli¹, M. Leotsinidis¹, M. Bakola², K.S. Kitsou^{2*}, K. Argyropoulos¹, A. Konstantopoulou¹, A. Katsifara¹, P. Gourzis³ and E. Jelastopulu¹

¹Department Of Public Health, Medical School, University of Patras, Patras, Greece; ²Postgraduate Program Of Public Health, Medical School, University of Patras, Patras, Greece and ³Department Of Psychiatry, School of Medicine, University of Patras, Patras, Greece

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.756

Introduction: The coronavirus pandemic has challenged the world with an unprecedented situation. Social distancing, self or quarantine isolation, personal hand hygiene, self-protection, and the fear of becoming infected with the virus, come with a psychological fallout. The COVID-19 pandemic has affected students around the world, in terms of their education and lifestyle.

Objectives: To investigate the impact of COVID-19 pandemic on the students' mental health and well-being at the University of Patras, in Western Greece.

Methods: An online questionnaire was prepared to collect responses from students during April 2020. Socio-demographic data, academic status, opinions about distance learning, changes in daily routine during the lockdown and anxiety and depression scores, according to the Greek version of the Hospital Anxiety and Depression Scale (HADS), were gathered.

Results: The total number of responders was 2009, of which 67.3% women. During lockdown, the 68% of the students returned to their family home. Anxiety and depression scores were higher in students with a low income, poor self-rated health, not informed about COVID-19, not satisfied with distance learning and being annoyed at staying home. Prevalence of anxiety and depression was found to be 35.8% and 51.2%, ranging from 26.7% to 48.2% for anxiety and from 36.3% to 60.5% for depression in Health Sciences and Humanities and Social Sciences, respectively.

Conclusions: Depression rates among university students in Greece were alarmingly high, denoting the impact of lockdown and changes in students' life, due to the COVID-19 pandemic.

Keywords: Anxiety; Depression; Students; COVID-19

EPP0377

Hopelessness and externality as predictors of experiencing anger during COVID-19 lockdown in Russia

T. Gordeeva^{1*}, O. Sychev² and M. Dovger¹

¹Psychology, Lomonosov Moscow State University, Moscow, Russian Federation and ²Research Department, Shukshin Altai State University for Humanities and Pedagogy, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.757

Introduction: Following Italy and many other European countries Russia entered a nationwide lockdown in March 2020. Since quarantine had impact on mental health (Gualano et al., 2020, Stanton et al., 2020), this study aimed to study the psychological predictors of low mental health and anger in Russian university students. Previous studies have shown that young people are most vulnerable part of population during Covid-19 pandemic (Pervichko et al., 2020).

Objectives: The purpose of this research was to assess the effects of externality and hopelessness on anger and irritation during COVID-19 lockdown.

Methods: The sample comprised 120 university students (86% women, $M=18.84$, $SD=1.58$) from Moscow. Online survey has been conducted in April 2020. Measures included Russian externality-internality scale based on Rotter's scale and three new scales specific for COVID-19 pandemic developed for this study to assess feeling of hopelessness ($\alpha = 0.72$), anger ($\alpha = 0.70$) and positive reformulation ($\alpha = 0.84$).

Results: Anger shows significant correlations with hopelessness ($r=0.43$; $p<0.001$), externality ($r=0.29$; $p<0.01$) and positive reformulation ($r=-0.41$; $p<0.001$). Structural equation modeling confirms theoretical model according to which the effect of externality on anger is mediated by hopelessness and positive reformulation (negatively) (indirect effects sig. at $p<0.01$, $\chi^2 = 1.32$; $df = 1$; $p = 0.251$; CFI = 0.995; TLI = 0.969; RMSEA = 0.052).

Conclusions: Conclusions. Anger and irritation regarding the necessity to stay at home during COVID-19 lockdown may be caused by external locus of control which effect on anger is mediated by hopelessness and limited capacities for positive reframing.

Keywords: anger; COVID-19 lockdown; Hopelessness; externality

EPP0378

Anxiety on the lockdown resolution

S. Ajmi^{1*}, S. Hentati², R. Masmoudi³, R. Sellami⁴, I. Baati², I. Feki³ and J. Masmoudi³

¹Psychiatrie A, hospital University Hedi Chaker, sfax, Tunisia;

²Psychiatry A, hedi chaker hospital, Sfax, Tunisia; ³Psychiatrie "a" Department, Hedi Chaker Hospital University -Sfax - Tunisia, sfax, Tunisia and ⁴Psychiatry "a" Department, Hedi Chaker Hospital University -Sfax - Tunisia, sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.758

Introduction: Lockdown due to the management of infectious diseases such as corona virus disease affect mental health. We would think that with the end of the lockdown due to the corona virus pandemic, the feeling of regaining freedom and movement would be good for our morale.