

Health, an administrative officer, and mental welfare officers of the local health department. The hospital representatives are the Consultant Psychiatrists, the hospital's Director of Rehabilitation, Psychiatric Social Worker and Senior Ward Nurses.

The meeting serves a number of purposes. Firstly, they bring together members of the local authority and hospital staffs to form a team to discuss patients' progress and placement. Secondly patients who are ready for accommodation outside hospital are brought to the notice of the local authority. Thirdly, there is the opportunity to consider the priorities of patients who may be awaiting admission. Fourthly, the need to renew and reinforce communications between home and patient is often revealed.

Unfortunately, not all the excellent recommendations in the Royal Medico-Psychological Association's Memorandum are likely to be implemented in the near future; but the mental health liaison committee provides one means of overcoming some of the inadequacies and differences which arise in the present arrangement for the care of the mentally handicapped.

D. A. SPENCER.

*Oulton Hall Hospital,
Leeds Road,
Oulton,
Leeds, LS26 8HL.*

MIGRAINE, ANOREXIA NERVOSA AND SCHIZOPHRENIA

DEAR SIR,

Whilst I appreciate Dr. Gosling's interesting points (*Journal*, November, 1970), some doubt has been expressed about the use of oral contraceptives in migrainous subjects in view of the preliminary stage of cerebral arterial constriction, at which time it was thought that thrombus formation might be encouraged. No doubt this risk is less since the 'Scowen Scare', but should still be borne in mind.

T. L. AVERY.

*The Central Hospital,
Near Warwick.*

UNIPOLAR AND BIPOLAR AFFECTIVE PSYCHOSIS. THE PROBLEM OF CLASSIFICATION ILLUSTRATED BY A CASE HISTORY

DEAR SIR,

Leonhard (1) suggested that on purely clinical grounds affective psychosis could be classed as

unipolar (attacks of depression only) or bipolar (attacks of depression and mania). Perris (2) compared the expectation of bipolar and unipolar psychosis in the relatives of a series of 138 patients with bipolar psychosis (patients who had had at least one attack of depression and one of mania) and 139 patients with unipolar psychosis (patients who had had three discrete attacks of depression), and produced evidence that these two classes might be genetically distinct. A difficulty with such studies is that until the patient's death there is no certain way of classifying his psychosis. Perris (3) later showed that attacks of mania were very uncommon after more than four depressive attacks. However, the present case history shows that typical mania may occur after as many as 13 distinct depressive attacks.

Case History

The patient was born in 1907. She was the second in a family of six and has had three daughters. There is no family history of mental illness. She married in 1935, and her husband, a boilerman, died in 1955. For many years she has worked as a laundry presser. She is described by her daughter as being energetic when well, but of a suspicious nature.

Her first depressive illness occurred in 1933 at the age of 26. This lasted a month, and a year later she had another lasting about the same period and requiring hospital admission. She had no further mental illness for over 20 years, but between the ages of 49 and 59 she had eleven hospital admissions for depression. These recurrences tended to develop acutely (in a week or two) and to be characterized by retardation, ideas of unworthiness, insomnia, anorexia, and (in the early attacks only) hostility towards her family. On five occasions she attempted suicide. Bouts of drenching perspiration heralded the onset of each attack, but no significant abnormality in her physical health has been found to account for this. The attacks each lasted 2 to 4 months and were variously treated with drugs and ECT. In between her psychotic episodes she kept at work and lived a normal active life.

At the age of 59, she had her twelfth hospital admission for depression, but after 26 days (and on antidepressant drugs) she became manic, elated, overactive and overtalkative and making amorous advances to male patients. This was an astonishing change in a patient so well known to the hospital staff. After two months she recovered her normal health. During the next 18 months she had two more attacks of depression, but during her fifteenth admission (age 61), again for depression, she had a further manic episode. Since then she has had two further admissions for depression only.

HOWARD ROBERTS.

*The Bethlem Royal and Maudsley Hospitals,
London, S.E.5.*

REFERENCES

1. LEONHARD, K. (1959). *Aufteilung der Endogenen Psychosen*. Berlin.
2. PERRIS, C. (1966). 'A study of bipolar (manic-depressive) and unipolar recurrent depressive psychosis.' *Acta psychiat. Scand.*, 42, Suppl. 194.
3. PERRIS, C. (1968). 'The course of depressive psychosis.' *Acta psychiat. Scand.*, 44, 238-48.

MENTAL HEALTH RESEARCH FUND:
THE SIR GEOFFREY VICKERS LECTURE

DEAR SIR,

The 1971 Sir Geoffrey Vickers Lecture will be given by Dr. Michael Rutter, Reader in Child Psychiatry, Institute of Psychiatry, on Wednesday, February 24, 1970, at 5.30 p.m. at the Edward Lewis Theatre, Middlesex Hospital Medical School, Cleveland Street, London W.1. His title will be 'The Effect of Adverse Family Influences on Children's Psychological Development'. Admission is by ticket which can be obtained from the Fund at 38 Wigmore Street, London W1H 9DF.

J. M. TANNER.

*Mental Health Research Fund,
38 Wigmore Street,
London, W1H 9DF.*

'GROUP AND CREATIVITY'

DEAR SIR,

The Third International Symposium on Analytical Group Therapy, organized by the German Academy for Psychoanalysis and the German Group Thera-

peutic Society, with the above title, will be held from 1st to 7th August, 1971, in Stelzerreut/Kumreut, near Passau in the Bavarian Forest.

The preliminary programme and further details are available from me at the address below.

GISLINDE BASS.

*Deutsche Akademie für Psychoanalyse,
1 Berlin 15,
Wielandstrasse 27-28,
Germany.*

FIRST BRITISH INTERNATIONAL
CONFERENCE ON ALCOHOLISM AND DRUG
DEPENDENCE

DEAR SIR,

The above Conference, the first of its kind to be held in the United Kingdom on the scale planned, will take place at Liverpool University from 28 March to 1 April, 1971. Papers will be presented by psychiatrists, physicians, sociologists and social workers, all either internationally or nationally well known for their work in the field of alcoholism and/or drug dependence.

Early application is advised for either residential or non-residential participation.

Further information can be obtained from the address below.

W. H. KENYON.

*Merseyside Council on Alcoholism,
B15, The Temple,
Dale Street,
Liverpool L2 5RU,
England.*