

Research Article

Cite this article: Chauhan BS and Walsh M (2022) Differential response of winter- and summer-emerging accessions of rigid ryegrass (*Lolium rigidum* Gaudin) to postemergence herbicides in Australia. *Weed Technol.* 36: 663–670. doi: [10.1017/wet.2022.63](https://doi.org/10.1017/wet.2022.63)

Received: 5 May 2022
Revised: 25 July 2022
Accepted: 26 July 2022
First published online: 22 August 2022

Associate Editor:

Rodrigo Werle, University of Wisconsin

Nomenclature:

Butoxydim; clethodim; glyphosate; glufosinate; haloxyfop; imazamox; imazapyr; mesosulfuron; paraquat; amitrole; pinoxaden; rigid ryegrass, *Lolium rigidum* Gaudin; cotton, *Gossypium hirsutum* L.; sorghum, *Sorghum bicolor* (L.) Moench

Keywords:

Cotton; fallows; herbicide resistance; summer crops; temperature

Author for correspondence:

Bhagirath S. Chauhan, Queensland Alliance for Agriculture and Food Innovation, University of Queensland, Gatton, QLD 4343, Australia.
Email: b.chauhan@uq.edu.au

Differential response of winter- and summer-emerging accessions of rigid ryegrass (*Lolium rigidum* Gaudin) to postemergence herbicides in Australia

Bhagirath S. Chauhan¹  and Michael Walsh² 

¹Professor, Queensland Alliance for Agriculture and Food Innovation and Professor, School of Agriculture and Food Sciences, University of Queensland, Gatton, Queensland, Australia and ²Director Weed Research, Sydney Institute of Agriculture, University of Sydney, Brownlow Hill, New South Wales, Australia

Abstract

Rigid ryegrass is a problematic weed species in winter crops and winter fallow; however, recently, this weed species has been observed in summer crops and fallow. These observations warrant the evaluation of different postemergence herbicides for its control. Outdoor pot studies were conducted during the spring and summer of 2021 to 2022 to determine the performance of POST herbicides on two summer-emerging (S3 and S6) and two winter-emerging (W3 and W8) accessions of rigid ryegrass. Across all accessions, butoxydim, clethodim, paraquat, and paraquat + amitrole at the field rate provided complete control of rigid ryegrass. Both summer-emerging accessions were found to be resistant to the field rates of glufosinate (750 g ai ha⁻¹), glyphosate (454 g ae ha⁻¹), haloxyfop (52 g ai ha⁻¹), and pinoxaden (30 g ai ha⁻¹). The S6 accession had the highest dose required for a 50% reduction in biomass for these herbicides (glufosinate 1,120 g ai ha⁻¹, glyphosate 1,210 g ae ha⁻¹, haloxyfop 140 g ai ha⁻¹, and pinoxaden 55 g ai ha⁻¹). This summer-emerging accession (S6) was also resistant to iodoflurofen. All four accessions were found susceptible to imazamox + imazapyr (a commercial mixture) and mesosulfuron. The study provides the first evidence of poor control of summer-emerging accessions of rigid ryegrass with different herbicides. Multiple-herbicide-resistant summer-emerging rigid ryegrass accessions would be a challenge to the production of summer crops (e.g., cotton and sorghum) as well as winter crops that rely on weed-free summer fallows for soil moisture retention; therefore these accessions warrant diversified management strategies.

Introduction

Rigid ryegrass is the most problematic grass weed species in winter grain crops in Australia. This weed currently infests more than 8 million ha of grain cropping areas, resulting in losses of AU\$93 million annually for Australian grain growers (Llewellyn et al. 2016). Uncontrolled rigid ryegrass can cause a significant reduction in the yields of wheat (*Triticum aestivum* L.), barley (*Hordeum vulgare* L.), and canola (*Brassica napus* L.), for example, up to 80% in wheat (Lemerle et al. 1996). It is a prolific seed producer (Chauhan et al. 2007), and the lightweight seeds are easily dispersed by natural (water, wind, animals, etc.) and anthropogenic (farm implements, produce, grain, etc.) means (Beckie and Jasieniuk 2021).

A recent Australian study reported that rigid ryegrass accessions can germinate across a wide range of diurnally fluctuating temperatures, from 15/5 C to 35/25 C (day/night temperatures) (Thompson et al. 2021). More than 90% of seeds germinated at 30/20 C and 35/25 C, suggesting that this weed could germinate in the summer months in Australia provided that soil moisture is sufficient. Although rigid ryegrass is a well-recognized problem in winter crops and winter fallow, there are recent reports of its occurrence in summer crops (e.g., cotton) and summer fallow in the southeastern region of Australia (Thompson and Chauhan 2022; Thompson et al. 2021).

In Australia, herbicides have become the most common method of weed control, and over-reliance on herbicides has resulted in the evolution of resistance in rigid ryegrass (Heap 2022). This weed has developed resistance to several classes of POST herbicides, including acetyl coenzyme-A carboxylase (ACCase) inhibitors, acetolactate synthase (ALS) inhibitors, and the 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitor glyphosate. Some factors contributing to the evolution of resistance in rigid ryegrass are its high seed production and high genetic diversity within accessions (Saini et al. 2015).

Glyphosate is the most commonly used herbicide to control weeds in glyphosate-resistant cotton and summer fallow in Australia. Owing to the continuous use of this herbicide in these

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Table 1. Herbicide treatments, their sites of action, doses, and adjuvants used in Experiment 1.^{a,b}

Herbicide	SOA	Dose	Adjuvant
ae/ai %		—g ai/ae ha ⁻¹ —	
Control	—	—	—
Butoxydim (25)	ACCCase inhibitor	45	1% Supercharge®
Clethodim (24)	ACCCase inhibitor	60	1% Supercharge®
Glufosinate (20)	Glutamine synthetase inhibitor	750	—
Glyphosate (57)	EPSP inhibitor	454	—
Haloxypop (52)	ACCCase inhibitor	52	1% UpTake™
*Imazamox (3.3) + imazapyr (1.5)	ALS inhibitors	36	0.5% HASTEN™
Iodosulfuron-methyl sodium (10)	ALS inhibitor	10	—
Mesosulfuron-methyl (3)	ALS inhibitor	3	1% HASTEN™
Pinoxaden (10)	ACCCase inhibitor	30	0.5% Adigor®
Paraquat (36)	Photosystem I inhibitor	600	1% HASTEN™
*Paraquat (12.5) + amitrole (25)	Photosystem I inhibitor + lycopene cyclase inhibitor	563	—

^aAdigor® contains 440 g L⁻¹ methyl esters of canola oil fatty acids; HASTEN™ contains 704 g L⁻¹ ethyl and methyl esters of vegetable oil with 196 g L⁻¹ nonionic surfactants; Supercharge® is 471 g L⁻¹ paraffin oil; and UpTake™ contains 582 g L⁻¹ paraffinic oil and 240 g L⁻¹ alkoxyated alcohol nonionic surfactants. An asterisk indicates a commercial mixture.

^bAbbreviations: ACCCase, acetyl-coenzyme-A carboxylase; ae, acid equivalent; ai, active ingredient; ALS, acetolactate synthase; EPSP, 5-enolpyruvylshikimate-3-phosphate; SOA, site of action.

situations, several summer annual weed species have evolved resistance (Heap 2022). Rigid ryegrass has also evolved resistance to glyphosate and ACCase inhibitors, such as haloxypop, in winter crop production systems. However, there is no information on the effectiveness of these and other selective and nonselective herbicides on summer-emerging accessions of rigid ryegrass.

Before rigid ryegrass becomes a widespread weed in summer fallow and summer crops (Thompson and Chauhan 2022), there is an urgent need to identify different herbicide options for its control. Therefore an outdoor pot study was conducted with two summer-emerging and two winter-emerging accessions of rigid ryegrass to answer the following questions: (1) what POST herbicide options exist for selective control of summer-emerging rigid ryegrass accessions and (2) if summer-emerging and winter-emerging rigid ryegrass accessions respond differently to various POST herbicide treatments.

Material and Methods

Seed Description

Four accessions of rigid ryegrass were used in this study: W3 and W8 as winter-emerging accessions and S3 and S6 as summer-emerging accessions. Seeds of W3 were collected on December 8, 2018, from the fence line of a sorghum field in Croppa Creek, New South Wales (NSW), Australia (S27°36.980; E152°12.486). W8 seeds were collected on November 12, 2018, from a wheat field in Wagga Wagga, NSW (S34°53.092; E147°04.555). Seeds of S3 were collected on March 25, 2019, from the fence line of a cotton field in Griffith, NSW (S34°30.019; E146°08.611). Seeds of S6 were collected on March 26, 2019, from a cotton field in Griffith, NSW (S34°43.417; E146°03.409). Plants were still green, suggesting that these plants were germinated in the last summer season. In Australia, winter, spring, summer, and autumn seasons are from June to August, September to November, December to February, and March to May, respectively. Seeds were stored at room temperature (25 ± 2 C) until used for experiments.

General Protocol

Pot trials were conducted during the spring and summer seasons of 2021 to 2022 at the weed science research facility of the University

of Queensland, Gatton, Australia. All trials were conducted two times. Plastic pots (14 cm diameter) filled with potting mix (Centenary Landscape, Queensland, Australia) were used in this study. Seeds (10 to 12 per pot) were planted at 0.5 cm depth, and seedlings were thinned immediately after emergence to keep five plants per pot. Pots were kept on benches under natural temperature and light conditions. The average minimum and maximum temperatures during the experiment were 17.7 C and 30.1 C, respectively. Irrigation was applied two times every day using an automated sprinkler system, but no fertilizer was added to the pots.

Herbicides were applied at the 4- to 5-leaf stage of rigid ryegrass. All herbicide treatments were applied in a spray cabinet with a track sprayer fitted with TeeJet® XR110015 flat-fan nozzles (TeeJet® Technologies, Wheaton, IL, USA). The sprayer delivered a spray volume of 108 L ha⁻¹. After treatment, pots were returned to the outdoor area. At 28 d after herbicide application, plant survival and biomass data were recorded. Plants were considered alive if they had at least 1 new leaf. Surviving plants from each pot were harvested together from the base and dried in an oven at 70 C for 72 h. Biomass of dry plants was weighed.

Experiment 1: Performance of POST Herbicides

Using the procedures described earlier, five plants per pot of four rigid ryegrass accessions were established in pots and grown outdoors to the 4- to 5-leaf stage. The performance of 11 commonly used POST herbicides was evaluated on rigid ryegrass accessions (Table 1). There were also unsprayed control treatments for each accession. The selected herbicides are commonly used in Australian farming systems to control rigid ryegrass. Each treatment was replicated four times.

Experiment 2: Dose-Response Experiment

Experiment 1 identified differential responses of summer-emerging and winter-emerging accessions of rigid ryegrass to glufosinate, glyphosate, haloxypop, and pinoxaden. To further examine these differences, these accessions were screened in dose-response pot studies using the procedures described earlier. As winter-emerging accessions (W3 and W8) were effectively controlled at the field rates of glufosinate, glyphosate, haloxypop, and pinoxaden in Experiment 1, these accessions were screened

Table 2. Herbicide doses used for four accessions of rigid ryegrass in Experiment 2.^a

Herbicide	Accessions	Doses	
		g ai/ae ha ⁻¹	
Glufosinate	S, S6	0, 187.5, 375, 750 (1X), 1500, 3000, 6000	
	W3, W8	0, 46.9, 93.8, 187.5, 375, 750 (1X), 1500	
Glyphosate	S3, S6	0, 113.5, 227, 454 (1X), 908, 1816, 3632	
	W3, W8	0, 28.4, 56.8, 113.5, 227, 454 (1X), 908	
Haloxypop	S3, S6	0, 13, 26, 52 (1X), 104, 208, 416	
	W3, W8	0, 3.3, 6.5, 13, 26, 52 (1X), 104	
*Imazamox (33 g L ⁻¹) + imazapyr (15 g L ⁻¹)	S3, S6, W3, W8	0, 9, 18, 36 (1X), 72, 144, 288	
Iodosulfuron	S3, S6, W3, W8	0, 2.5, 5, 10 (1X), 20, 40, 80	
Mesosulfuron	S3, S6, W3, W8	0, 0.8, 1.5, 3 (1X), 6, 12, 24	
Pinoxaden	S3, S6	0, 7.5, 15, 30 (1X), 60, 120, 240	
	W3, W8	0, 1.9, 3.8, 7.5, 15, 30 (1X), 60	

^aRecommended doses are shown with 1X. An asterisk indicates a commercial mixture.

Table 3. Effect of herbicide treatments on survival and biomass of four accessions of rigid ryegrass when sprayed during summer months (Experiment 1).^{a,b}

Herbicide	Survival				Biomass			
	S3	S6	W3	W8	S3	S6	W3	W8
	%				g pot ⁻¹			
Control	100	100	100	100	2.21	3.35	2.11	1.81
Butroxydim	0	0	0	0	0	0	0	0
Clethodim	0	0	0	0	0	0	0	0
Glufosinate	75	70	20	15	0.47	0.45	0.03	0.02
Glyphosate	85	85	0	5	2.20	2.80	0	0.03
Haloxypop	50	55	0	10	1.17	1.15	0	0.81
*Imazamox (33 g L ⁻¹) + imazapyr (15 g L ⁻¹)	0	10	5	5	0	0.02	0.01	0.16
Iodosulfuron	25	75	40	20	0.23	1.72	1.04	0.51
Mesosulfuron	5	10	0	5	0.11	0.13	0	0.27
Pinoxaden	30	45	0	0	0.70	2.79	0	0
Paraquat	0	0	0	0	0	0	0	0
*Paraquat (125 g L ⁻¹) + amitrole (250 g L ⁻¹)	0	0	0	0	0	0	0	0
LSD _{0.05}	19	16	12	16	0.53	0.98	0.34	0.60

^aAn asterisk indicates a commercial mixture.

^bAbbreviation: LSD_{0.05}, least significant difference at the 5% level of significance.

at the seven rates 0X, 0.0625X, 0.125X, 0.25X, 0.5X, 1X, and 2X (Table 2). Summer-emerging accessions (S3 and S6) were not controlled at the field rates of glufosinate, glyphosate, haloxypop, and pinoxaden in Experiment 1; these accessions were screened at the seven rates 0X, 0.25X, 0.5X, 1X, 2X, 4X, and 8X (Table 2). Summer- and winter-emerging accessions responded similarly to imazamox (33 g L⁻¹) + imazapyr (15 g L⁻¹) (a commercial mixture), iodosulfuron, and mesosulfuron in Experiment 1. Therefore rates were kept the same (0X, 0.25X, 0.5X, 1X, 2X, 4X, and 8X) for these three herbicides. Herbicides that provided complete control of the four accessions of rigid ryegrass in Experiment 1 were not included in the dose-response experiment.

Statistical Analyses

Experiments were conducted in a randomized complete block design with four replications in Experiment 1 and three replications in Experiment 2. A pot with five plants was considered an experimental unit (one pot representing a replication). All experiments were repeated within a week after the completion of the first run. Data were subjected to analysis of variance (ANOVA) (Genstat 2021) to test for treatment by experimental run interaction. There was no interaction, and therefore data were pooled over

the two experimental runs. In Experiment 1, each accession was analyzed separately, and means of survival and biomass were separated at $P \leq 0.05$ using Fisher's protected least significant difference test. In Experiment 2, seedling survival and biomass (percent of nontreated control) data were regressed against herbicide dose using a three-parameter logistic model using SigmaPlot 14.5 (Systat Software Inc., Palo Alto, CA, USA):

$$Y = a / [1 + (x/H_{50})^b]$$

In this model, Y is the survival or biomass data at herbicide dose x , a is the maximum seedling survival or biomass, H_{50} is the herbicide dose (g ae or ai ha⁻¹) required for a 50% reduction in seedling survival (LD₅₀) or biomass (GR₅₀), and b is the slope.

Results and Discussion

Experiment 1: Performance of POST Herbicides

Summer-emerging accessions (S3 and S6) of rigid ryegrass were more poorly controlled by glufosinate, glyphosate, haloxypop, and pinoxaden herbicide treatments than were winter-emerging accessions (W3 and W8) (Table 3). Glyphosate-, haloxypop-, and pinoxaden-resistant rigid ryegrass accessions have been

Table 4. Estimated herbicide doses required for a 50% reduction in accession survival (LD_{50}) and plant biomass production (GR_{50}) of rigid ryegrass accessions (Experiment 2).^a

Herbicide	Accession	LD_{50} —g ai/ae ha ⁻¹ —	GR_{50} —g ai/ae ha ⁻¹ —
Glufosinate	S3	1,053 (63)	682 (77)
	S6	1,133 (136)	1,119 (116)
	W3	497 (218)	322 (82)
	W8	399 (109)	266 (56)
Glyphosate	S3	894 (35)	260 (74)
	S6	567 (122)	1,207 (688)
	W3	83 (15)	95 (5)
	W8	132 (13)	95 (22)
Haloxyfop	S3	7.4 (2.9)	40.8 (9.0)
	S6	13.3 (4.3)	138.6 (30.2)
	W3	1.3 (0.9)	5.2 (1.0)
	W8	2.4 (0.1)	3.1 (0.1)
*Imazamox (33 g L ⁻¹) + imazapyr (15 g L ⁻¹)	S3	11.2 (0.7)	13.4 (1.4)
	S6	18.5 (3.3)	12.4 (1.0)
	W3	14.1 (0.6)	8.4 (0.2)
	W8	8.9 (0.5)	10.0 (1.4)
Iodosulfuron	S3	5.9 (1.8)	6.7 (0.7)
	S6	20.2 (3.7)	22.9 (5.7)
	W3	14.8 (3.6)	10.1 (1.4)
	W8	8.3 (1.6)	7.0 (1.9)
Mesosulfuron	S3	0.6 (0.1)	0.4 (0.1)
	S6	0.7 (0.1)	0.9 (0.2)
	W3	0.7 (0.1)	0.7 (0.1)
	W8	0.6 (0.1)	0.6 (0.4)
Pinoxaden	S3	11.8 (1.1)	22.0 (2.6)
	S6	12.9 (5.4)	54.7 (30.7)
	W3	1.9 (0.1)	2.5 (0.1)
	W8	1.7 (0.3)	2.0 (0.2)

^aValues in parentheses are standard errors. An asterisk indicates a commercial mixture.

reported in Australia (Boutsalis et al. 2012; Heap 2022; Owen and Powles 2010), but glufosinate-resistant rigid ryegrass has not previously been reported in Australia. The first case of glufosinate-resistant rigid ryegrass was reported in Greece in 2018 (Travlos et al. 2018). Although 70% to 75% of seedlings of S3 and S6 accessions survived the field rate of glufosinate (750 g ai ha⁻¹), the growth of these seedlings was severely affected, as indicated by the 79% to 87% reductions in biomass compared to the nontreated control plants (Table 3). In the nontreated control treatment, the biomass of S3, S6, W3, and W8 accessions was 2.2, 3.4, 2.1, and 1.8 g pot⁻¹, respectively.

The summer- and winter-emerging rigid ryegrass accessions were susceptible to widely used ACCase-inhibiting herbicides with apparent low-level resistance to ALS-inhibiting herbicides. Butroxydim, clethodim, paraquat, and paraquat + amitrole provided complete control of all four accessions of rigid ryegrass (Table 3). The susceptibility of all four accessions to butroxydim and clethodim is unusual given the high frequencies of resistance to these herbicides occurring in rigid ryegrass accessions in Australian cropping systems (Boutsalis et al. 2012; Broster et al. 2013; Owen et al. 2014; Saini et al. 2015). There was low-level survival ($\leq 10\%$) in three of the rigid ryegrass accessions following imazamox + imazapyr and mesosulfuron treatments (Table 3). Although there was survival, the plants were severely affected by these herbicide treatments, as indicated by the biomass reductions of at least 80%. The iodosulfuron treatment resulted in 20% to 75% survival of rigid ryegrass accessions, with the growth of these seedlings reduced by 50% to 90% (Table 3).




Figure 1. Effect of glufosinate dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

Experiment 2: Dose-Response Experiment

Glufosinate

The dose-response assay revealed that a glufosinate rate of 500 and 400 g ai ha⁻¹ resulted in 50% mortality (LD_{50}) in W3 and W8 accessions, respectively, whereas the same mortality was achieved with 1,050 and 1,130 g ai ha⁻¹ in S3 and S6, respectively (Table 4; Figure 1). The GR_{50} values for the winter accessions (W3 and W8) were 270 to 320 g ai ha⁻¹, whereas these values for the summer-emerging accessions were 680 to 1,120 g ai ha⁻¹.

Glufosinate is a nonselective, POST herbicide that is registered for fallow and preseedling weed control in Australian grain production systems. Although glufosinate-resistant rigid ryegrass has not been reported in Australia, there are reports that rigid ryegrass is not well controlled by glufosinate (Kumaratilake et al. 2002). Environmental factors, such as temperature and relative humidity, can greatly affect the efficacy of glufosinate (Kumaratilake and Preston 2005; Ramsey et al. 2002). However, these parameters were not studied in the current study. In Australia, glufosinate usage is expected to increase to manage glyphosate-resistant weed species (Chauhan et al. 2021). Summer-emerging accessions were collected from the cotton-growing fields and fence lines, and the poor control of these accessions with glufosinate suggests that the use of XtendFlex™ (Bayer CropScience, Hawthorn East, Victoria, Australia) cotton (resistant to glyphosate, glufosinate, and dicamba) in the future (Werth et al. 2021) may not provide effective control of these accessions.




Figure 2. Effect of glyphosate dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

Glyphosate

The dose–response trial confirmed poor control of S3 and S6 accessions of rigid ryegrass by glyphosate. The LD₅₀ value for S3 and S6 accessions were 890 and 570 g ai·ha⁻¹, respectively, whereas those values for W3 and W8 accessions were 80 and 130 g ai·ha⁻¹, respectively (Table 4; Figure 2). The GR₅₀ values for S3 and S6 were 260 and 1,210 g ai·ha⁻¹, respectively—greater than for the winter accessions (95 g ai·ha⁻¹).

Studies have reported glyphosate resistance in several weed species, including rigid ryegrass, in Australia and other parts of the world (Desai et al. 2020; Heap 2022; Owen and Powles 2010). The first case of glyphosate-resistant rigid ryegrass was reported more than 20 years ago in Australia (Powles et al. 1998; Pratley et al. 1999), and since then, several accessions of glyphosate-resistant rigid ryegrass have been reported in several countries (Fernandez-Moreno et al. 2017). Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot] and perennial ryegrass (*Lolium perenne* L.), closely related species of rigid ryegrass, have also evolved resistance to glyphosate (Singh et al. 2020; Yannicari et al. 2016).

Glyphosate is relied on for broad-spectrum preplanting and fallow weed control in conservation agriculture systems as well as for selective weed control in glyphosate-resistant crops grown in Australia (Duke and Powles 2008). Therefore the increasing frequency of glyphosate-resistant rigid ryegrass accessions is of concern to Australian growers. These summer-emerging accessions were collected from glyphosate-resistant cotton fields, suggesting that the use of this herbicide in cotton increases the selection pressure for resistance evolution. With the expected




Figure 3. Effect of haloxyfop dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

increasing problem of summer-emerging rigid ryegrass, it will be very important to monitor the evolution of glyphosate resistance in this weed species as it occurs in summer fallows and glyphosate-resistant cotton crops.

Haloxyfop

Although 30% of the seedlings of S6 survived the 8X rate of haloxyfop, the LD₅₀ value was only 13 g ai·ha⁻¹ (Table 4; Figure 3). The GR₅₀ value for this accession was 140 g ai·ha⁻¹. The GR₅₀ value for S3 (another summer-emerging accession) was 40 g ai·ha⁻¹, whereas the GR₅₀ values for the winter-emerging accessions were 3 to 5 g ai·ha⁻¹.

Haloxyfop is used to control grass weeds in fallow and broadleaf crops (Chauhan et al. 2021). Although there are several cases of winter-emerging rigid ryegrass accessions resistant to haloxyfop, this is the first study showing poor control of summer-emerging accessions. The summer-emerging accessions were collected from cotton fields and edges, and growers use haloxyfop to control grass weeds in cotton (from the crop's second leaf to before the onset of flowering). Therefore a possible reason for the poor control could be the selection pressure for this herbicide, as herbicides from this mode of action group are prone to the risk of developing herbicide resistance. It is possible that these accessions evolved resistance to ACCase-inhibiting herbicides prior to evolving to be summer-emerging biotypes. Resistance to some ACCase-inhibiting herbicides can occur because of a decreased rate of herbicide activation, an increased rate of herbicide detoxification, sequestration of the herbicide, or overproduction of ACCase activity (Saini




Figure 4. Effect of the commercial mixture of imazamox (33 g L^{-1}) + imazapyr (15 g L^{-1}) dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate (36 g ai ha^{-1}) of the commercial mixture; $24.75 \text{ g ai ha}^{-1}$ of imazamox + $11.25 \text{ g ai ha}^{-1}$ of imazapyr of the herbicide.

et al. 2015). In such situations, growers need to use other effective ACCase-inhibiting herbicides (e.g., clethodim, butoxydim) in rotation or mixture.

Imazamox (33 g L⁻¹) + imazapyr (15 g L⁻¹)

The response of the summer- and winter-emerging accessions of rigid ryegrass was similar to the application of imazamox + imazapyr (Table 4; Figure 4). The LD_{50} values for the commercial mixture ranged from 9 (imazamox at 6.2 g ai ha^{-1} and imazapyr at 2.8 g ai ha^{-1}) to $18.5 \text{ g ai ha}^{-1}$ (imazamox at $12.7 \text{ g ai ha}^{-1}$ and imazapyr at 5.8 g ai ha^{-1}), whereas the GR_{50} values for different accessions were 8.4 (imazamox at 5.8 g ai ha^{-1} and imazapyr at 2.6 g ai ha^{-1}) to $13.4 \text{ g ai ha}^{-1}$ (imazamox at 9.2 g ai ha^{-1} and imazapyr at 4.2 g ai ha^{-1}).

Imazamox + imazapyr, a commercial mixture of ALS-inhibiting imidazolinone herbicides, is used for the control of certain annual grass and broadleaf weeds in Clearfield® (BASF Australia, Southbank, Victoria, Australia) crops and igrowth® (Pacific Seeds, Toowoomba, Queensland, Australia) herbicide-resistant grain sorghum. Although the LD_{50} and GR_{50} values for different accessions of rigid ryegrass were well below half the field rate of this herbicide, up to 6% of seedlings survived the 4X rate of this herbicide. These results suggest monitoring the escaped seedlings and the evolution of resistance to this herbicide mixture as the area under Clearfield® and igrowth® herbicide-resistant sorghum increases. After applications of these imidazolinone herbicides,




Figure 5. Effect of iodosulfuron dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

there is a requirement for significant intervals for planting crops, such as chickpea (*Cicer arietinum* L.; 10 mo) and cotton (22 mo).

Iodosulfuron

Accessions responded differently to different doses of iodosulfuron. S6 and W3 accessions had greater LD_{50} and GR_{50} values than the field rate (10 g ai ha^{-1}) of iodosulfuron. The LD_{50} value for S6 and W3 accessions were 20 and 15 g ai ha^{-1} , respectively, whereas these values for S3 and W8 were 6 and 8 g ai ha^{-1} , respectively (Table 4; Figure 5). Similarly, the GR_{50} values for S6 and W3 were 23 and 10 g ai ha^{-1} , respectively, whereas the GR_{50} value for S3 and W8 was 7 g ai ha^{-1} .

Iodosulfuron, an ALS-inhibiting herbicide, is recommended for the control of rigid ryegrass in wheat. In Australia, only one case of iodosulfuron-resistant rigid ryegrass has been reported, from South Australia (Heap 2022). The current study found two accessions (S6 and W3) resistant to iodosulfuron. Recently, iodosulfuron-resistant rigid ryegrass was also reported in Spain (Torra et al. 2021). The study reported the evolution of multiple and cross-resistance to ALS- and ACCase-inhibiting herbicides by mechanisms consistent with the presence of both target-site and non-target-site resistance mechanisms.

Mesosulfuron

All four accessions were found to be highly susceptible to the field rate of mesosulfuron. The LD_{50} values for different accessions ranged from 0.6 to 0.7 g ai ha^{-1} , whereas the GR_{50} values ranged from 0.4 to 0.9 g ai ha^{-1} (Table 4; Figure 6).




Figure 6. Effect of mesosulfuron dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

Mesosulfuron, an ALS-inhibiting herbicide, is used to control rigid ryegrass in wheat. Mesosulfuron-resistant rigid ryegrass has been reported in France, Israel, and Spain (Heap 2022; Torra et al. 2021). In the United States, a dose-response assay revealed 37-fold resistance in Italian ryegrass to mesosulfuron compared with a susceptible standard (Singh et al. 2020). The widespread use of sulfonylurea herbicides in Australian crop production systems may result in the evolution of mesosulfuron-resistant rigid ryegrass; therefore the use of this herbicide needs to be managed using a diversified weed management system. After an application of mesosulfuron, there is a need to wait for 9 to 12 mo if growers want to plant barley, chickpea, or cotton.

Pinoxaden

Summer- and winter-emerging accessions of rigid ryegrass responded differently to different rates of pinoxaden. The LD₅₀ values for summer- and winter-emerging accessions were 12 to 13 and 2 g ai ha⁻¹, respectively (Table 4; Figure 7). The GR₅₀ values were much greater than the LD₅₀ values. The GR₅₀ values for S3 and S6 accessions were 22 and 55 g ai ha⁻¹, respectively, whereas the GR₅₀ values for W3 and W8 accessions were 3 and 2 g ai ha⁻¹, respectively.

Pinoxaden, an ACCase-inhibiting herbicide, is used to control rigid ryegrass and *Avena* species in wheat and barley crops in Australia. Although pinoxaden was first registered in Australia in 2006, 30% to 40% of the fields surveyed in 2003 and 2005 had pinoxaden-resistant rigid ryegrass accessions (Boutsalis et al. 2012). These observations suggest cross-resistance in rigid




Figure 7. Effect of pinoxaden dose on survival (A) and biomass (percent of nontreated control) (B) of four accessions (S3, S6, W3, and W8) of rigid ryegrass. Plants were sprayed at the 4- to 5-leaf stage of each accession. The vertical dashed line represents the 1X rate of the herbicide.

ryegrass to herbicides that were not previously used. As the summer-emerging accessions of rigid ryegrass were collected from cotton fields and edges, it is possible that these accessions never received an application of pinoxaden but were exposed to haloxypfop, another ACCase-inhibiting herbicide. Italian ryegrass has also been found to have a 23-fold resistance to pinoxaden in the Texas Blacklands of the United States (Singh et al. 2020).

The two winter-emerging accessions were better controlled by POST herbicides compared to the two summer-emerging accessions. The summer-emerging accessions could be resistant to the herbicides; however, this cannot be concluded in this study, as there was no susceptible control of summer-emerging rigid ryegrass. Future studies should include herbicide-susceptible accessions of summer-emerging rigid ryegrass. This study was conducted outdoors during the spring and summer seasons. Although the weather conditions were optimum for summer-emerging accessions, they may not have been representative for winter-emerging accessions. Therefore future research should include only summer-emerging accessions in the spring and summer months.

This study revealed poor control of summer-emerging accessions (S3 and S6) of rigid ryegrass by multiple herbicides (glyphosate, glufosinate, haloxypfop, and pinoxaden). The S6 accession had the highest GR₅₀ value for these herbicides, and it was also poorly controlled by iodosulfuron. Both summer-emerging accessions were collected from cotton fields and edges, suggesting that glyphosate-resistant cotton or the future availability of glyphosate- and glufosinate-resistant cotton in Australia (e.g., XtendFlex™) may not provide effective control of these summer-emerging

accessions. Haloxypop is commonly used to control dominant summer annual grass weed species, such as feather fingergrass (*Chloris virgata* Sw.) and jungle rice [*Echinochloa colona* (L.) Link], in cotton. While currently there are no reported cases of haloxypop-resistant feather fingergrass and jungle rice in Australia, the results presented here suggest that summer-emerging rigid ryegrass accessions with evolved resistance may become more problematic in these production systems. At present, summer-emerging accessions of rigid ryegrass are mainly present along crop edges or fence lines (Thompson and Chauhan 2022); therefore preventive measures, such as field hygiene, clean water channels, or clean field boundaries, could prove very effective in delaying the spread of summer-emerging rigid ryegrass.

In Australia, annual sowthistle (*Sonchus oleraceus* L.) is another example that was once considered to be only a winter weed but is now present in winter as well as summer cropping systems (Manalil et al. 2017). As rigid ryegrass is an obligate outcrossing species with high genetic diversity (Charmet et al. 1996), stopping the spread of multiple-herbicide-resistant summer-emerging rigid ryegrass is critical. The occurrence of rigid ryegrass as a summer-emerging weed in summer cropping systems is problematic; however, the evolution of resistance in these accessions to herbicides frequently relied on for summer annual weed control is a significant threat to summer cropping systems. Further research should investigate the management of herbicide resistance in summer-emerging annual ryegrass accessions.

Acknowledgments. The authors thank the Grains Research and Development Corporation (GRDC) for investing in this research. The authors declare no conflicts of interest.

References

- Beckie HJ, Jasieniuk M (2021) *Lolium rigidum* and *Lolium multiflorum*. Pages 261–283 in Chauhan BS, ed. *Biology and Management of Problematic Crop Weed Species*. New York, NY: Academic Press
- Boutsalis P, Gill GS, Preston C (2012) Incidence of herbicide resistance in rigid ryegrass (*Lolium rigidum*) across southeastern Australia. *Weed Technol* 26:391–398
- Broster JC, Koetz EA, Wu H (2013) Herbicide resistance levels in annual ryegrass (*Lolium rigidum* Gaud.) and wild oat (*Avena* spp.) in southwestern New South Wales. *Plant Protect Q* 28:126–132
- Charmet G, Balfourier F, Chatard V (1996) Taxonomic relationships and interspecific hybridization in the genus *Lolium* (grasses). *Gen Res Crop Evol* 43:319–327
- Chauhan BS, Congreve M, Mahajan G (2021) Management options for large plants of glyphosate-resistant feather fingergrass (*Chloris virgata*) in Australian fallow conditions. *PLoS ONE* 16:e0261788
- Chauhan BS, Gill G, Preston C (2007) Effect of seeding systems and dinitroaniline herbicides on emergence and control of rigid ryegrass (*Lolium rigidum*) in wheat. *Weed Technol* 21:53–58
- Desai HS, Thompson M, Chauhan BS (2020) Target-site resistance to glyphosate in *Chloris virgata* biotypes and alternative herbicide options for its control. *Agronomy* 10:1266
- Duke SO, Powles SB (2008) Glyphosate: a once-in-a-century herbicide. *Pest Manag Sci* 64:319–325
- Fernandez-Moreno PT, Rojano-Delgado AM, Menendez J, De Prado R (2017) First case of multiple resistance to glyphosate and PPO-inhibiting herbicides in rigid ryegrass (*Lolium rigidum*) in Spain. *Weed Sci* 65:690–698
- Genstat (2021) Genstat for Windows. 21st ed. Hemel Hempstead, UK: VSN International.
- Heap I (2022) International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org/>. Accessed: March 30, 2022
- Kumaratilake AR, Lorraine-Colwill DF, Preston C (2002) A comparative study of glufosinate efficacy in rigid ryegrass (*Lolium rigidum*) and sterile oat (*Avena sterilis*). *Weed Sci* 50:560–566
- Kumaratilake AR, Preston C (2005) Low temperature reduces glufosinate activity and translocation in wild radish (*Raphanus raphanistrum*). *Weed Sci* 53:10–16
- Lemerle D, Tang H, Murray G, Morris S, Tang H (1996) Survey of weeds and diseases in cereal crops in the southern wheat belt of New South Wales. *Aust J Exp Agric* 36:545–554
- Lewellyn RS, Ronning D, Ouzman J, Walker S, Mayfield A, Clarke M (2016) Impact of Weeds on Australian Grain Production: The Cost of Weeds to Australian Grain Growers and the Adoption of Weed Management and Tillage Practices. Canberra, Australia: CSIRO. 112 p
- Manalil S, Werth J, Jackson R, Chauhan BS, Preston C (2017) An assessment of weed flora 14 years after the introduction of glyphosate-tolerant cotton in Australia. *Crop Past Sci* 68:773–780
- Owen MJ, Martinez NJ, Powles SB (2014) Multiple herbicide-resistant *Lolium rigidum* (annual ryegrass) now dominates across the Western Australian grain belt. *Weed Res* 54:314–324
- Owen MJ, Powles SB (2010) Glyphosate-resistant rigid ryegrass (*Lolium rigidum*) populations in the Western Australian grain belt. *Weed Technol* 24:44–49
- Powles SB, Lorraine-Colwill DF, Dellow J, Preston C (1998) Evolved resistance to glyphosate in rigid ryegrass (*Lolium rigidum*) in Australia. *Weed Sci* 46:604–607
- Pratley J, Urwin N, Stanton R, Baines P, Broster J, Cullis K, Schafer D, Bohn J, Krueger R (1999) Resistance to glyphosate in *Lolium rigidum*. I. Bioevaluation. *Weed Sci* 47:405–411
- Ramsey RJL, Stephenson GR, Hall JC (2002) Effect of relative humidity on the uptake, translocation, and efficacy of glufosinate ammonium in wild oat (*Avena fatua*). *Pestic Biochem Physiol* 73:1–8
- Saini RK, Malone J, Preston C, Gill G (2015) Target enzyme-based resistance to clethodim in *Lolium rigidum* populations in Australia. *Weed Sci* 63: 946–953
- Singh V, Maity A, Abugho S, Swart J, Drake D, Bagavathiannan M (2020) Multiple herbicide-resistant *Lolium* spp. is prevalent in wheat production in Texas Blacklands. *Weed Technol* 34:652–660
- Thompson M, Chauhan BS (2022) Changing seasonality of *Lolium rigidum* (annual ryegrass) in southeastern Australia. *Front Agron* 4
- Thompson M, Mahajan G, Chauhan BS (2021) Seed germination ecology of southeastern Australian rigid ryegrass (*Lolium rigidum*) populations. *Weed Sci* 69:454–460
- Torra J, Montull JM, Taberner A, Onkokesung N, Boonham N, Edwards R (2021) Target-site and non-target-site resistance mechanisms confer multiple and cross-resistance to ALS and ACCase inhibiting herbicides in *Lolium rigidum* from Spain. *Front Plant Sci* 12. <https://doi.org/10.3389/fpls.2021.625138>
- Travlos IS, Cheimona N, De Prado R, Jhala AJ, Chachalis D, Tani E (2018) First case of glufosinate-resistant rigid ryegrass (*Lolium rigidum* Gaud.) in Greece. *Agronomy* 8:35
- Werth J, Thornby D, Keenan M, Hereward J, Chauhan BS (2021) Effectiveness of glufosinate, dicamba, and clethodim on glyphosate-resistant and -susceptible populations of five key weeds in Australian cotton systems. *Weed Technol* 35:967–973
- Yannicari M, Vila-Aiub M, Istilart C, Acciari H, Castro AM (2016) Glyphosate resistance in perennial ryegrass (*Lolium perenne* L.) is associated with a fitness penalty. *Weed Sci* 64:71–79