

## Ca-BEARING RECTORITE FROM TOOHO MINE, JAPAN

**Key Words**—Allevardite, Beidellite, Interstratification, Margarite, Rectorite, X-ray powder diffraction.

Rectorite (alleverdite) was recognized by Caillère *et al.* (1950) and Brindley (1956) as a mica-type mineral with sheets of water molecules separating the 2:1 mica-like layers. Brown and Weir (1963) showed that rectorite and alleverdite are the same mineral that consists of pairs of mica-like (high charge, non-swelling) and montmorillonite-like (low charge, swelling) layers. Kodama (1966) concluded that rectorite consists of a regular alternation of paragonite-like layers and expandable layers having beidellitic and montmorillonitic compositions. Weaver and Pollard (1975) reported that the mica-like layers generally contain interlayer Na (paragonite), but in some samples K is the predominant cation.

A Ca-bearing rectorite is reported here, which appears to be an interstratified Ca-paragonite/beidellite, from the Tooho mine, Aichi Prefecture, Japan.

## MATERIAL AND METHODS

The specimen originated from the Tooho mine ("resekii" deposit), Aichi Prefecture, Japan. This deposit was formed by hydrothermal alteration of andesitic country rock. The dominant clay minerals in this deposit are pyrophyllite, interstratified mica/smectite, and mica, which are concentrated in central, intermediate, and marginal zones, respectively. The rectorite material occurs in a part of the zone of interstratified mica/smectite that has been altered by later hydrothermal activity to a brittle clayey material containing small veinlets of dickite, fluorite, and Li-bearing tosudite (Nishiyama *et al.*, 1975).

The <2- $\mu\text{m}$  fraction of the specimen was used for this study. The impurities in the specimen are 7% quartz, 5% tosudite, 3% topaz, 2% pyrophyllite, 1% dickite, and ~0.2% fluorite. Quantitative estimations of the impurities except fluorite were carried out by X-ray powder diffraction intensity measurement using the method of known additions (Brindley, 1961). The amount of fluorite was estimated from the fluorine analysis of the specimen using a specific ion electrode (Ingram, 1970), after correcting for the fluorine in topaz.

## RESULTS AND DISCUSSIONS

*X-ray powder diffraction*

X-ray powder diffraction (XPD) patterns of an oriented specimen on a glass slide show a regular series of basal reflections (Table 1). The 001 spacing of the mineral is 24.83 Å at 58% relative humidity (RH), and the sequence of the basal reflections (standard deviation,  $\sigma = 0.18$ ) was as good as those of other rectorites (Table 2); therefore this mineral can be described as a 1:1 regular interstratification resembling rectorite.


After the mineral was treated with  $\text{Mg}^{2+}$  and ethylene glycol (EG), better integral series of the basal reflections having d-spacings of 24.40 and 26.50 Å, respectively, were observed. This means that each expandable layer in the mineral behaved homogeneously. These interlayer characteristics resemble those of rectorite rather than the Funyu material described by Shimoda *et al.* (1974) in which the integral series became less integral after such treatments (Table 2). After KCl treatment,

the basal spacing was reduced to 22.23 Å with less regularity ( $\sigma = 0.28$ ) (compare the data for rectorite, Kodama, 1966). With EG treatment of the  $\text{K}^+$ -saturated specimen, two phases were observed with d-spacings of 26.7 (about 60%) and 23.8 Å (about 40%), which correspond respectively to two and one layers of ethylene glycol in the interlayer position.

Table 3 shows the observed and calculated F values of the XRD basal reflections of the specimen before and after various treatments. Relatively good agreement was obtained. One-dimension Fourier syntheses calculated from the observed F values show the levels of the component layers of the mineral. The levels were measured from the midpoint of the octahedral sheet width on the electron density curves (Nishiyama and Oinuma, 1978) (Table 4). The d-spacing of the mica layer was 9.64 Å for both the untreated and EG-treated specimens and 9.70 Å for the specimen heated at 500°C for 1 hr. These d-spacings are similar to those of rectorite, 9.52, 9.50, and 9.62 Å for the untreated, EG-treated, and 560°C-treated alleverdites (rectorite), respectively (Brindley, 1956), rather than to the mica layer of the Funyu sample whose d-spacings were 9.94 and 9.99 Å for untreated and EG-treated materials, respectively.


*Chemical composition and structural properties*

The chemical composition of the Tooho mineral, corrected for mineral impurities and estimated on an ignited weight basis, is listed in Table 5. From these data the following structural formula can be calculated:




Cations exchanged by 1 N  $\text{NH}_4^+$  solution amounted to 0.01%  $\text{K}_2\text{O}$  and 0.83%  $\text{CaO}$  on an ignited weight basis ( $\text{Ca}_{0.16}\text{K}_{0.002}$  on the basis of one  $\text{O}_{20}(\text{OH})_4$  structural unit).

Assuming that all interlayer cations in the expandable layer of the mineral are exchangeable and that the cation ratio in the octahedral sheets of the expandable and non-expandable layers remains constant, the structural formula of the mica component of the mineral can be written as



and that of the expandable component as



These structural formulae show that the mica layer belongs to the margarite (Ca-mica) and the paragonite (Na-mica) series rather than muscovite (K-mica). This conclusion is not contradicted by the XRD data; the thickness of the mica layer and  $b_0$  of the mineral (9.64 and 8.916 Å) match more closely those of paragonite (9.66 and 8.91 Å) or margarite (9.71 and 8.92 Å) than those of muscovite (10.0 and 9.04 Å).

The structural formula of the mineral shows that the expandable component is an Al-rich smectite group. The layer charge of the expandable layer was calculated to be  $-0.33$  per  $\text{O}_{10}(\text{OH})_2$ , and is predominantly due to tetrahedral substitution. Therefore, the term beidellitic layer appears to be appropriate for the expandable component in this mineral. Schultz (1969) reported that after saturation with  $\text{K}^+$  and heating at 300°C for 30 min, dioctahedral smectites with a total net

Table 1. X-ray powder diffraction data for Ca-rectorite before and after various treatments.<sup>1</sup>

00ℓ	Untreated			Ethylene glycol			KCl			KCl + EG			500°C × 1 hr		
	d (Å)	d × ℓ (Å)	I <sup>2</sup>	d (Å)	d × ℓ (Å)	I <sup>2</sup>	d (Å)	d × ℓ (Å)	I <sup>2</sup>	d (Å)	d × ℓ (Å)	I <sup>2</sup>	d (Å)	d × ℓ (Å)	I <sup>2</sup>
1	24.54	25.54	19,957	26.77	26.77	23,248	22.08	22.08	1879	26.29	26.29	3561	20.08	22.08	1272
2	12.27	24.54	6328	13.29	26.58	3546	10.91	21.83	963	13.19	26.38	467	9.88	19.76	1252
3	—	—	—	8.84	26.52	468	—	—	—	?	—	—	6.558	19.67	114
4	6.281	24.12	44	6.63	26.52	250	5.504	22.01	41	6.707	26.83	44	4.861	19.44	664
5	4.996	24.98	989	5.292	26.46	430	4.483	22.41	30	5.356	26.62	37	3.866	19.33	52
6	—	—	—	4.428	26.56	173	—	—	—	4.745	23.73	53	—	—	—
7	3.547	24.82	124	3.785	26.49	66	3.186	22.30	276	—	—	—	3.209	19.25	1950
8	3.099	24.79	1440	3.308	26.46	1030	—	—	—	3.858	27.00	24	—	—	—
9	2.722	24.49	11	2.945	26.50	298	2.535	22.81	—	3.351	23.46	362 (Q)	—	—	—
10	2.488	24.88	13	2.648	26.48	45	—	—	8	3.351	26.80	60	—	—	—
11	2.264	24.90	22	—	—	—	2.006	22.06	28	2.978	26.80	—	1.925	19.25	194
12	2.066	24.79	83	2.206	26.47	13	1.860	22.32	10	—	—	—	—	—	—
13	1.917	24.92	190	2.035	26.45	117	—	—	—	1.975	23.70	14	1.602	19.22	22
14	1.787	25.01	5	1.891	26.47	136	—	—	—	—	—	—	—	—	—
15	1.665	24.97	3	1.764	26.46	7	—	—	—	—	—	—	—	—	—
16	1.544	24.70	22	1.661	26.57	4	—	—	—	—	—	—	—	—	—
17	—	—	—	1.557	26.46	19	—	—	—	1.923	23.93	24	—	1.376	19.26
18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
19	—	—	—	1.392	26.44	2	—	—	—	—	—	—	—	—	—
20	—	—	—	1.322	26.44	18	—	—	—	—	—	—	—	—	—
21	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
22	—	—	—	1.200	26.40	11	—	—	—	—	—	—	—	—	—

<sup>1</sup> Relative humidity 49–58%.

<sup>2</sup> Peak height.




Figure 1. Electron micrograph of the Tooho Ca-rectorite.

layer charge of less than 0.85/unit cell ( $0.425/O_{10}(OH)_2$ ) reexpanded to 17 Å; those with net layer charge of more than 0.85/unit cell did not. After  $K^+$ -saturation and EG treatment, the mineral in this study showed two peaks at 26.7 and 23.8 Å, meaning that the beidellitic layer consists of two distinct interlayers which are characterized by less and more layer charges than  $0.43/O_{10}(OH)_2$ . Simultaneously, the basal reflections became less rational after such treatments, and some complicated interstratified structures may have due to the heterogeneous distribution of layer charges in the beidellitic layers.

The predominant interlayer cation of the beidellitic layer was  $Ca^{2+}$ , a fact verified by 001 spacing of the mineral of 24.83 Å at 58% RH and 23.5 Å at 35% RH, similar to the 24.5 Å spacing of Ca-saturated rectorite at 50% RH (Kodama, 1966). Furthermore, Matsuda and Nagasawa (1977) showed that the Ca-saturated expandable layer of a regularly interstratified mica/smectite changes from two interlayers of water molecules to one with decreasing RH from 30% to 10%.

Table 3. Observed and calculated structure factors of the (00 $l$ ) reflections of the Ca-rectorite before and after various treatments.

00 $l$	Untreated		500°C × 1 hr		Ethylene glycol	
	F obs	F cal	F obs	F cal	F obs	F cal
1	66.8	85.6	14.9	13.7	68.1	89.4
2	74.4	-72.0	29.9	-27.8	54.9	-57.2
3	—	10.9	13.7	13.3	29.9	-24.0
4	12.3	1.3	45.0	-31.3	29.2	22.4
5	73.8	-43.6	16.0	12.5	48.2	-28.7
6	—	10.5	119.3	-95.4	36.8	-26.3
7	37.4	-33.4	—	12.7	26.8	31.3
8	147.6	-107.1	—	-2.1	122.8	-103.9
9	14.8	17.2	—	9.8	75.0	-67.6
10	18.0	13.1	67.4	93.1	32.6	22.9
11	26.0	-31.9	—	8.6	—	21.1
12	56.0	60.5	28.2	29.1	21.7	-26.7
13	92.8	83.2	—	7.2	71.4	50.7
14	15.6	27.2	30.0	-48.1	84.2	94.7
15	13.1	3.7	—	—	21.2	32.2
16	41.2	49.2	—	—	15.9	-11.8
17	—	—	—	—	40.4	32.4
18	—	—	—	—	—	8.0
19	—	—	—	—	14.7	-30.8
20	—	—	—	—	47.1	-35.8
21	—	—	—	—	—	7.5
22	—	—	—	—	41.5	71.8

#### Infrared and differential thermal analysis

The OH-stretching absorption band of the Tooho specimen was observed at 3643  $cm^{-1}$  which is the same value as other rectorites. The absorption bands between 825 and 690  $cm^{-1}$  of the Tooho specimen, however, are distinct from rectorites. The absorption bands of quartz at 800, 782, and 695  $cm^{-1}$  and of other mineral impurities obscured the absorption bands of the Ca-rectorite in these regions, but it is likely that the rectorite bands at 808 and 730  $cm^{-1}$  are ambiguous or are shifted

Table 2. Averaged basal spacings (Å) of the Ca-rectorite and other related minerals.

	Untreated	MgCl <sub>2</sub>	KCl	500°C	EG	MgCl <sub>2</sub> + EG	KCl + EG		
A	$\bar{x}$	24.83	24.40	22.23	19.50	26.50	26.81	26.71	23.80
	$\sigma$	0.18	0.12	0.28	0.28	0.07	0.14	0.24	—
	n	13	10	8	8	19	13	8	4
B	$\bar{x}$	24.63			19.24 <sup>1</sup>	26.48			
	$\sigma$	0.12			0.13	0.03			
	n	20			13	16			
C	$\bar{x}$	23.87	24.61	22.3	19.3 <sup>2</sup>	26.72 <sup>3</sup>			
	( $\sigma$ )	0.17	0.03	1.r. <sup>4</sup>	1.r. <sup>4</sup>				
	n								
D	$\bar{x}$	24.86	24.92 <sup>5</sup>	22.34	19.5 <sup>5</sup>	26.74	26.60 <sup>5</sup>	26.86	
	$\sigma$	0.15		0.25		0.27		0.55	
	n	11		9		13		8	

A = Ca-bearing rectorite; B = allevardite (rectorite) (Brindley, 1956); C = rectorite (Kodama, 1966); D = Funyu sample (Shimoda *et al.*, 1974);  $\sigma = [\sum(x_i - \bar{x})^2/n]^{1/2}$ ; n = numbers of measured basal reflections.

<sup>1</sup> 550°C.

<sup>2</sup> 600°C.

<sup>3</sup> Shimoda *et al.* (1974).

<sup>4</sup> Less rational series of diffuse reflections.

<sup>5</sup> Matsuda and Nagasawa (1977).

Table 4. Lattice spacings of the d(001) and component layers of the Ca-rectorite and related minerals.

Treatment	Ca-rectorite			Allevardite (rectorite) <sup>1</sup>			Funiyu sample		
	d(001) (Å)	Component layer		d(001) (Å)	Component layer		d(001) (Å)	Component layer	
		Mica	Exp. <sup>2</sup> (Å)		Mica	Exp. <sup>2</sup> (Å)		Mica	Exp. <sup>2</sup> (Å)
Untreated	24.83	9.64	15.19	24.62	9.52	15.10	24.86	9.94	14.92
EG	26.50	9.64	16.86	26.48	9.50	16.98	26.74	9.99	16.75
500°C	19.50	9.70	9.80	19.24 <sup>3</sup>	9.62	9.62			

<sup>1</sup> Brindley (1956).<sup>2</sup> Expandable layer.<sup>3</sup> 560°C.

to lower frequencies in this material. The band at 825 cm<sup>-1</sup>, which is a shoulder of the 808 cm<sup>-1</sup> band in rectorites, is distinct in this mineral.

Endothermic reactions were noted by differential thermal analysis for the Tooho specimen at 105°, 178°, and 557°C due to dehydration. Smectites having monovalent interlayer cations (Na<sup>+</sup>, K<sup>+</sup>) show one endothermic peak at low temperature, whereas, those having divalent interlayer cations (Ca<sup>2+</sup>, Mg<sup>2+</sup>, and/or Li<sup>+</sup>) show two endothermic peaks (Hendricks *et al.*, 1940). Therefore the two peaks at 105° and 178°C of the Tooho specimen are probably due to the dehydration of the smectite (beidellite) layer wherein Ca<sup>2+</sup> is the interlayer cation.

#### Electron microscopy

A transmission electron micrograph (Figure 1) shows a very thin flaky habit of the mineral. Folded ribbons which are characteristic of allevardite and also of rectorite (Brown and Weir, 1963) are rare in this sample.

#### CONCLUSIONS

The mineral from the Tooho "roseki" deposit, Aichi Prefecture, Japan is a regular 1:1 interstratification of mica and expandable layers. The mica layer belongs to a margarite-paragonite series, whereas the expandable layer is similar to bei-

dellite with Ca<sup>2+</sup> exchange cations. Because the name rectorite has generally been given to regular 1:1 interstratifications of dioctahedral paragonite and smectite, it is convenient to describe the Tooho mineral as a Ca-bearing rectorite. This Ca-bearing rectorite has not previously been reported in natural deposits; however, Eberl (1978) synthesized hydrothermally a Ca-rectorite from a Wyoming bentonite at 400°C.

#### ACKNOWLEDGMENTS

The first author thanks Mrs. A. Kamijo who presented a JEM-T7S electron microscope to our Toyo University. We are indebted to Mr. T. Hatta, University of Tsukuba, for the determination of fluorine.

Natural Science Laboratory  
Toho University  
Hakusan 5-28-20, Bunkyo-ku  
Tokyo 112, Japan

TSUTOMU NISHIYAMA

Institute of Geoscience  
University of Tsukuba  
Sakuramura, Niihari-gun  
Ibaraki Prefecture, Japan

SUSUMU SHIMODA

#### REFERENCES

Table 5. Chemical compositions of the Tooho specimen the Ca-rectorite and related minerals.

	1 (%)	2 <sup>1</sup> (%)	3 <sup>1</sup> (%)	4 <sup>1</sup> (%)	5 <sup>1</sup> (%)
SiO <sub>2</sub>	47.46	51.98	54.09	54.11	52.22
TiO <sub>2</sub>			0.36	0.01	0.16
Al <sub>2</sub> O <sub>3</sub>	31.52	39.35	38.48	40.38	38.81
Fe <sub>2</sub> O <sub>3</sub>	1.08	1.44	0.78	0.15	0.86
FeO			0.66		
MgO	0.19	0.25	0.42	0.78	
CaO	2.44	3.46	1.19	0.52	2.49
Li <sub>2</sub> O	0.07	0.03			
Na <sub>2</sub> O	1.68	2.43	2.69	3.87	2.42
K <sub>2</sub> O	0.76	1.08	1.31	0.29	3.02
H <sub>2</sub> O+	7.11				
H <sub>2</sub> O-	7.20				
F	0.70				
Total	100.21	100.02	99.98	100.23	99.98

1 = Tooho specimen containing impurities. 2 = Ca-bearing rectorite corrected for other mineral impurities. 3 = Allevardite (rectorite) (Hénin *et al.*, 1954 and Brindley, 1956). 4 = Rectorite (Kodama, 1966). 5 = Funiyu sample (Shimoda *et al.*, 1974).

<sup>1</sup> On an ignited weight basis.

- Brindley, G. W. (1956) Allevardite, a swelling double-layer mica mineral: *Amer. Mineral.* **41**, 91–103.
- Brindley, G. W. (1961) Quantitative analysis of clay mixtures: in *X-ray Identification and Crystal Structures of Clay Minerals*, 2nd ed., G. Brown, ed., Mineral. Society, London, 489–516.
- Brown, G. and Weir, A. H. (1963) The identity of rectorite and allevardite: in *Proc. Int. Clay Conf., Stockholm, 1963*, vol. 1, I. Th. Rosenquist and P. Graff-Petersen, eds., Pergamon Press, Oxford, 27–35.
- Caillère, S., Mathieu-Scicaud, A., and Hénin, S. (1950) Nouvel essai d'identification du minéral de La Table près Allevard, l'allevardite: *Bull. Soc. Fr. Mineral. Cristallogr.* **73**, 193–202.
- Eberl, D. D. (1978) The reaction of montmorillonite to mixed-layer clay: the effect of interlayer alkali and alkaline earth cations: *Geochim. Cosmochim. Acta* **42**, 1–7.
- Hendricks, S. B., Nelson, R. A., and Alexander, L. T. (1940) Hydration mechanism of the clay mineral montmorillonite, saturated with various cations: *J. Amer. Chem. Soc.* **62**, 1457–1464.
- Hénin, S., Esquevin, J., and Caillère, S. (1954) Sur la fibrosité de certain minéraux de nature montmorillonitique: *Bull. Soc. Fr. Mineral. Cristallogr.* **77**, 491–499.
- Ingram, B. L. (1970) Determination of fluorine in silicate rocks without separation of aluminum using a specific ion electrode: *Anal. Chem.* **42**, 1825–1827.
- Kodama, H. (1966) The nature of the component layers of rectorite: *Amer. Mineral.* **51**, 1035–1055.
- Matsuda, T. and Nagasawa, K. (1977) Comparisons of char-

- acteristics of expansible components between regularly interstratified mica-smectite and chlorite-smectite: *J. Mineral. Soc. Japan* 13 (spec. issue), 111–118 (in Japanese).
- Nishiyama, T., Shimoda, S., Shimosaka, K., and Kanaoka, S. (1975) Lithium-bearing tosudite: *Clays & Clay Minerals* 23, 337–342.
- Nishiyama, T. and Oiuma, K. (1978) One-dimensional Fourier synthesis of clay minerals. Part 1. Dioctahedral mica, montmorillonite and 1:1 regular interstratified mica-montmorillonite: *J. Toyo Univ. Gen. Ed. (Nat. Sci.)* No. 21, 1–38.
- Shimoda, S., Kohyama, N., and Ishikawa, Y. (1974) Some problems on the origin of interstratified mica clay minerals: *J. Mineral. Soc. Japan* 11 (spec. issue), 169–179 (in Japanese).
- Schultz, L. G. (1969) Lithium and potassium absorption, dehydroxylation temperature, and structural water content of aluminous smectites: *Clays & Clay Minerals* 17, 115–149.
- Weaver, C. E. and Pollard, L. D. (1975) *The Chemistry of Clay Minerals*: Elsevier, Amsterdam, 107–109.

(Received 16 June 1980; accepted 9 December 1980)

## ANNOUNCEMENT

### THE CLAY MINERALS SOCIETY

18th Annual Meeting — 30th Annual Clay Minerals Conference

University of Illinois — Urbana, Illinois

October 4 to 8, 1981

- SCIENTIFIC PROGRAM:** Symposium on the Transformation of Smectite to Illite  
General Sessions  
Poster Sessions
- FIELD TRIP:** Sedimentary Structures in Shale (including a stop at the Fithian cyclothem locality) and Soil Formation on Glacial Till (October 8)
- COSPONSORS:** Department of Geology and Illinois Geological Survey
- FURTHER INFORMATION:** Dr. John Hower  
Conference Chairman  
Department of Geology  
University of Illinois  
Urbana, Illinois 61801
- Dr. D. D. Eberl  
Program Chairman  
Department of Geology  
University of Illinois  
Urbana, Illinois 61801