

ON FACTORIALS EXPRESSIBLE AS SUMS OF AT MOST THREE FIBONACCI NUMBERS

FLORIAN LUCA¹ AND SAMIR SIKSEK²

¹*Instituto de Matemáticas, Universidad Nacional Autónoma de México,
CP 58089, Morelia, Michoacán, México* (fluca@matmor.unam.mx)

²*Mathematics Institute, University of Warwick,
Coventry CV4 7AL, UK* (s.siksek@warwick.ac.uk)

(Received 3 September 2008)

Abstract In this paper, we determine all the factorials that are a sum of at most three Fibonacci numbers.

Keywords: diophantine equation; Fibonacci numbers; factorials

2010 *Mathematics subject classification:* Primary 11D85

1. Introduction

Diophantine equations involving factorials have a long history. For example, in 1876 Brocard [6] asked for the integral solutions of $n! + 1 = x^2$; this was asked again (apparently independently) by Ramanujan [17] in 1913. The Brocard–Ramanujan equation is still an unsolved problem today; see D25 in Guy’s book [13]. Other Diophantine equations involving factorials have proved more tractable. For example, Erdős and Obláth [11] showed that the equation $x^p + y^p = n!$ has no solutions with x, y coprime and $p > 2$. Many have considered equations of the form $P(x) = n!$, where P is a polynomial; the best results so far appear to be those of Berend and Harmse [1], who show that there are only finitely many solutions if P has an irreducible factor of relatively large degree.

Diophantine equations involving Fibonacci numbers have been no less popular, as documented in [13, D25] and in the historical sections of [8] and [7]. Moreover, there have been several papers attacking Diophantine equations that involve both factorials and Fibonacci numbers. For example, in [12] it is shown that if k is fixed, then there are only finitely many positive integers n such that

$$F_n = m_1! + m_2! + \cdots + m_k!$$

holds for some positive integers m_1, \dots, m_k , and all solutions of the above equation with $k \leq 2$ have been determined. It is conjectured in [12] that if $m_1 < m_2 < \cdots < m_k$ holds in the above equation, then k itself must be bounded. Some results on this problem

can be found in [3]. In [16], it was shown that the largest solution of the Diophantine equation

$$F_{n_1} F_{n_2} \cdots F_{n_k} = m_1! m_2! \cdots m_\ell!$$

with $1 \leq n_1 < \cdots < n_k$ and $2 \leq m_1 \leq \cdots \leq m_\ell$ is $F_1 F_2 F_3 F_4 F_5 F_6 F_8 F_{10} F_{12} = 11!$.

In this paper we prove the following result.

Theorem 1.1. *Let $(F_m)_{m \geq 0}$ be the Fibonacci sequence given by $F_0 = 0$, $F_1 = 1$ and $F_{m+2} = F_{m+1} + F_m$ for all $m \geq 0$. The only factorials expressible as the sum of at most three Fibonacci numbers are*

$$\begin{aligned} 0! = 1! = F_1 = F_2, \quad 2! = F_1 + F_1 = F_2 + F_1 = F_2 + F_2 = F_3, \\ 3! = F_4 + F_4 = F_5 + F_1 = F_5 + F_2 = F_3 + F_3 + F_3 = F_4 + F_3 + F_1 = F_4 + F_3 + F_2, \\ 4! = F_8 + F_4 = F_6 + F_6 + F_6 = F_7 + F_6 + F_4 = F_8 + F_3 + F_1 = F_8 + F_3 + F_2, \\ 6! = F_{15} + F_{10} + F_{10} = F_{15} + F_{11} + F_8. \end{aligned}$$

It is not hard to show that every positive integer N has a representation, called the *Zeckendorf decomposition*, of the form $N = F_{n_1} + F_{n_2} + \cdots + F_{n_s}$, where $n_i - n_{i+1} \geq 2$, and that, up to identifying F_1 with F_2 , this representation is unique. Our problem is therefore related to the Zeckendorf decomposition of factorials. Denote by $Z(N)$ the number s of Fibonacci numbers appearing in the Zeckendorf decomposition of N .

Conjecture. $Z(n!)$ tends to infinity with n .

We are unable to prove our conjecture, but our Theorem 1.1 determines all positive integers n such that $Z(n!) \leq 3$.

It is appropriate to point out some analogous results to our conjecture that appear in [15]. Let $b \geq 2$ be a positive integer. For a positive integer N let $s_b(N)$ be the sum of the base b digits of n . In [15], it is shown that the inequality $s_b(n!) \gg \log n$ holds for all positive integers n , where the implied constant depends on b . Thus, the complexity of representing $n!$ in base b grows as n tends to infinity. The method of proof is elementary and it is based on the observation that $n!$ is a multiple of $b^m - 1$ for all $m = 1, 2, \dots, \lfloor \log n / \log b \rfloor$.

Particular Diophantine equations of the form

$$a_1^{x_1} + \cdots + a_k^{x_k} = n!,$$

where a_1, \dots, a_k are given positive integers and x_1, \dots, x_k, n are non-negative integer unknowns, have been studied in [10]. For example, all the solutions of the Diophantine equation

$$2^{x_1} + 3^{x_2} + 5^{x_3} + 7^{x_4} + 11^{x_5} = n! \tag{1.1}$$

have $n \leq 6$.

For the purpose of the present paper, as $F_0 = 0$, it suffices to determine all solutions to the following Diophantine equation:

$$F_x + F_y + F_z = n!, \quad x, y, z \geq 0, \quad n \geq 1. \tag{1.2}$$

Before doing this, we explain very briefly why our method for solving this equation is far more complicated than the method for solving (1.1) in [10]. To solve (1.1), all we have to do is find a positive integer M such that the congruence

$$2^{x_1} + 3^{x_2} + 5^{x_3} + 7^{x_4} + 11^{x_5} \equiv 0 \pmod{M}$$

has no solutions. Once this is done, we know that, for any solution to (1.1), $M \nmid n!$, giving a bound on n . This elementary idea cannot be used for (1.2); for example, $F_0 + F_{-2} + F_1 = 0$, and so the congruence $F_x + F_y + F_z \equiv 0 \pmod{M}$ has solutions for all M .

Our strategy for (1.2) is as follows. Let $(L_m)_{m \geq 0}$ be the Lucas sequence defined by $L_0 = 2$, $L_1 = 1$ and $L_{m+2} = L_{m+1} + L_m$ for all $m \geq 0$. Let $m \geq 6$ be an even integer such that $L_{m/2} \leq n$. We compute the first few terms of an expansion of F_x as an ‘ $F_{m/2}$ -adic’ power series, in a way that is very similar to Strassman’s Theorem (see [9, pp. 59–73]), except that we do not require $F_{m/2}$ to be prime. From this, we deduce congruence conditions modulo m and modulo $F_{m/2}$ on the unknowns x, y, z in (1.2); the idea here is reminiscent of Skolem’s method (see [9, pp. 228–231] and [4, pp. 290–300]). We use the Chebyshev θ -function to combine the information obtained from all even $m \geq 6$ with $L_{m/2} \leq n$. For n very large, this shows that x, y, z are too large compared with n for Equation (1.2) to hold, and so gives a bound on n . Our initial bound obtained in this way is $n \leq L_{501} < 5.045 \times 10^{104}$. An iterative argument, using the same information derived from the ‘Strassman’ expansion, is applied 50 times to reduce the bound to $n \leq L_{37} = 54\,018\,521$. The proof is completed using a sieving argument.

2. Inequalities

In this section, we gather some inequalities that will be useful later.

Lemma 2.1. *For all integers $n \geq 2$,*

$$\log(n!) < (n + \frac{1}{2}) \log n - n + 1 \leq n \log n. \tag{2.1}$$

Proof. By Stirling’s formula,

$$n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\lambda_n},$$

where

$$\frac{1}{12n + 1} < \lambda_n < \frac{1}{12n}.$$

Hence,

$$\log(n!) \leq (n + \frac{1}{2}) \log n - n + \frac{1}{2} \log(2\pi) + \lambda_n.$$

But $\frac{1}{2} \log(2\pi) + \lambda_n \leq \frac{1}{2} \log(2\pi) + \frac{1}{24} < 1$, leading to the first inequality in the statement of the lemma. To obtain the second, we need to show that

$$n \geq \frac{1}{2} \log n + 1$$

holds for all $n \geq 2$. This is in fact true for $n = 2$, and so is true for all real $n \geq 2$ since

$$\frac{d}{dt} \left(t - \frac{\log t}{2} - 1 \right) = 1 - \frac{1}{2t} > 0 \quad \text{for all } t \geq 2.$$

□

We write $\alpha = \frac{1}{2}(1 + \sqrt{5})$ and $\beta = \frac{1}{2}(1 - \sqrt{5})$ for the two roots of the characteristic equation $\lambda^2 - \lambda - 1 = 0$ of the Fibonacci sequence. It is well known that the *Binet formula*

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}$$

holds for all $n \geq 0$. We will find it convenient to extend the Fibonacci sequence to negative subscripts either using Binet's formula directly or by defining $F_{n-2} = F_n - F_{n-1}$ for all $n = 1, 0, -1, \dots$. The Binet formula for the Lucas numbers is

$$L_n = \alpha^n + \beta^n \quad \text{for all } n \geq 0.$$

As with the Fibonacci numbers, we will sometimes make use of negative subscripted Lucas numbers. It is easy to see that if $n \geq 0$, then $F_{-n} = (-1)^{n-1}F_n$ and $L_{-n} = (-1)^n L_n$.

Here are a few inequalities involving the Fibonacci and Lucas numbers.

Lemma 2.2. *Any solution to the Diophantine equation (1.2) with $n \geq 3$ satisfies*

$$\max(x, y, z) \leq \frac{n \log n}{\log \alpha} - 1.$$

Proof. From $F_x \leq n!$ and the Binet formula, we obtain that $\alpha^x \leq \sqrt{5}n! + 1$. Hence,

$$\begin{aligned} x \log \alpha &\leq \log(n!) + \log(\sqrt{5}) + \log(1 + 1/(\sqrt{5}n!)) \\ &\leq (n + \frac{1}{2}) \log n - n + 1 + \log(\sqrt{5}) + \log(1 + 1/(6\sqrt{5})). \end{aligned}$$

The lemma follows from the inequality

$$n \geq \frac{1}{2} \log n + 1 + \log \alpha + \log(\sqrt{5}) + \log(1 + 1/(6\sqrt{5})),$$

which is easily established for $n \geq 3$ by modifying the argument at the end of the proof of Lemma 2.1. □

Lemma 2.3. *Let $y \geq 1$. If $x \geq 2.079 \log y + 2.441$, then $F_x \geq y$. Moreover, if $y \geq 200$ and $x \leq 2.076 \log y$, then $L_x \leq y$.*

Proof. For the first part, note that $2.079 \log \alpha = 1.0004\dots$ to four decimal places. Thus,

$$\begin{aligned} x \log \alpha &\geq \log y + 2.441 \log \alpha \\ &\geq \log \sqrt{5}y + \log(1 + 1/\sqrt{5}) \\ &\geq \log \sqrt{5}y + \log(1 + 1/(\sqrt{5}y)). \end{aligned}$$

Hence, $\alpha^x \geq \sqrt{5}y + 1$, giving $F_x \geq y$.

For the second part, observe that $2.076 \log \alpha = 0.998996 \dots$ to six decimal places. So, $x \log \alpha \leq 0.999 \log y$, and therefore

$$L_x \leq \alpha^x + 1 \leq y^{0.999} + 1 \leq y \left(\frac{1}{200^{0.001}} + \frac{1}{200} \right) < y.$$

□

Lemma 2.4. For $x \geq 0$, we have $\log(1 + x) \leq x$. For $0 \leq x \leq \frac{1}{2}$, we have $\log(1 - x) \geq -2x$.

2.1. The Chebyshev function

We shall need some estimates involving the Chebyshev function

$$\theta(x) = \sum_{p \leq x} \log p,$$

where the sum is taken over all primes less than or equal to x . It is well known that $\theta(x)/x \rightarrow 1$ as $x \rightarrow \infty$. Here, we need lower estimates for this ratio for small values of x .

Proposition 2.5. For all real $x \geq 1$, $\theta(x) \leq 1.001102x$. Moreover,

if $10 \leq x \leq 20$,	then $\theta(x)/x \geq 0.4861$,
if $20 \leq x \leq 30$,	then $\theta(x)/x \geq 0.6628$,
if $30 \leq x \leq 40$,	then $\theta(x)/x \geq 0.7033$,
if $40 \leq x \leq 50$,	then $\theta(x)/x \geq 0.7228$,
if $50 \leq x \leq 500$,	then $\theta(x)/x \geq 0.7615$,
if $500 \leq x \leq 1000$,	then $\theta(x)/x \geq 0.9194$,
if $x \geq 1000$,	then $\theta(x)/x \geq 0.9456$.

Proof. Theorem 6 of [18] gives $\theta(x) < 1.001102x$ if $0 < x$, and $\theta(x) \geq 0.998684x$ if $x \geq 1319007$. To obtain the lower bounds claimed by the proposition, we used a simple MAGMA [5] script to determine the infima of $\theta(x)/x$ in the finite ranges above as well as in the range $1000 \leq x \leq 2 \times 10^6$. Note that over the interval $[p, p']$, where p, p' are primes, the infimum of $\theta(x)/x$ is $\theta(p)/p'$. □

3. Elementary lemmas

We shall also need the following elementary properties of the Fibonacci and Lucas numbers. Properties (3.1)–(3.3) are well known (see, for example, [14]) and can be proved immediately using the Binet formulae for the Fibonacci and Lucas numbers. For integers n ,

$$F_{2n} = F_n L_n, \quad L_{2n} = 5F_n^2 + 2(-1)^n, \quad L_n^2 = 5F_n^2 + 4(-1)^n. \quad (3.1)$$

For all pairs of integers m and n ,

$$2F_{m+n} = F_m L_n + F_n L_m, \quad 2L_{m+n} = L_m L_n + 5F_m F_n. \quad (3.2)$$

If m and n have the same parity, then

$$F_m + F_n = \begin{cases} F_{(m+n)/2}L_{(m-n)/2} & \text{if } m \equiv n \pmod{4}, \\ F_{(m-n)/2}L_{(m+n)/2} & \text{if } m \not\equiv n \pmod{4}. \end{cases} \tag{3.3}$$

Lemma 3.1. *Let m be a non-zero integer. Then*

$$F_{\lambda m}^2 \equiv (-1)^{(\lambda+1)m} \lambda^2 F_m^2 \pmod{F_m^4}.$$

Moreover, if m is even, then

$$F_{\lambda m} \equiv (-1)^{m(\lambda+1)/2} \lambda F_m \pmod{F_{m/2}^3}.$$

Proof. Define

$$H_\lambda = \left(\frac{F_{\lambda m}}{F_m} \right)^2 = \frac{\alpha^{2\lambda m} + \beta^{2\lambda m} - 2(-1)^{\lambda m}}{(\alpha^m - \beta^m)^2}.$$

This is a ternary recurrence sequence with characteristic polynomial

$$(X - \alpha^{2m})(X - \beta^{2m})(X - (-1)^m) = (X^2 - L_{2m}X + 1)(X - (-1)^m).$$

However, from (3.1) we have $L_{2m} \equiv 2(-1)^m \pmod{F_m^2}$. Hence,

$$H_{\lambda+3} \equiv 3(-1)^m H_{\lambda+2} - 3H_{\lambda+2} + (-1)^m H_\lambda \pmod{F_m^2}.$$

Moreover, $H_0 = 0$, $H_1 = 1$ and $H_2 = L_m^2 \equiv 4(-1)^m \pmod{F_m^2}$, again by (3.1). An easy induction shows that

$$H_\lambda \equiv (-1)^{(\lambda+1)m} \lambda^2 \pmod{F_m^2},$$

and multiplying by F_m^2 completes the proof of the first part of the lemma.

The proof of the second part is similar, but easier, using the binary recurrence sequence of general term $G_\lambda = F_{\lambda m}/F_m$. □

Lemma 3.2. *Let m be a non-zero even integer. Then*

$$F_{x_0+2\lambda m} \equiv F_{x_0} \pmod{F_m}, \tag{3.4}$$

and

$$2F_{x_0+\lambda m} \equiv 2(-1)^{m\lambda/2} F_{x_0} + 2(-1)^{m(\lambda+1)/2} F_m L_{x_0} \lambda + 5F_m^2 F_{x_0} \lambda^2 \pmod{F_{m/2}^3}. \tag{3.5}$$

Proof. Using (3.2), we see that

$$\begin{aligned} 2F_{x_0+2\lambda m} &= F_{x_0} L_{2\lambda m} + L_{x_0} F_{2\lambda m} \\ &= F_{x_0} (2 + 5F_{\lambda m}^2) + L_{x_0} F_{2\lambda m}. \end{aligned}$$

However, F_m divides $F_{\lambda m}$ and $F_{2\lambda m}$. If $2 \nmid F_m$, then (3.4) follows. Suppose now that $2 \mid F_m$. Then $2 \mid L_m$, and we note that $2F_m$ divides both $F_{\lambda m}^2$ and $L_m F_m = F_{2m} \mid F_{2\lambda m}$. Hence, $2F_{x_0+2\lambda m} \equiv 2F_{x_0} \pmod{2F_m}$. This completes the proof of (3.4).

We now drop the requirement that 2 divides F_m and we move on to prove (3.5). To this aim, we combine (3.1) and (3.2) with Lemma 3.1 to obtain

$$\begin{aligned} 2F_{x_0+\lambda m} &= F_{x_0}L_{\lambda m} + L_{x_0}F_{\lambda m} \\ &= F_{x_0}(2(-1)^{m/2} + 5F_{\lambda m}^2) + L_{x_0}F_{\lambda m} \\ &\equiv F_{x_0}(2(-1)^{m/2} + 5F_m^2\lambda^2) + (-1)^{m(\lambda+1)/2}F_mL_{x_0}\lambda \pmod{F_{m/2}^3}, \end{aligned}$$

which gives (3.5). □

4. Some congruences

The following two results are useful in applying the ‘Strassman procedure’ alluded to in § 1.

Lemma 4.1. *Let $m \geq 4$ be even and let $-m < x_0, y_0 \leq m$ be integers such that x_0 is odd, y_0 is even and $F_{x_0} + F_{y_0} \equiv 0 \pmod{F_m}$. Then $(x_0, y_0) \in \{(\pm(m-1), m-2), (\pm 1, 2)\}$.*

Proof. Since $x_0, y_0 \in (-m, m]$ and $m \geq 4$, it follows that if either F_{x_0} or F_{y_0} is negative, then it is less than F_m in absolute value. Thus, $F_{x_0} + F_{y_0} \in \{-F_m, 0, F_m, 2F_m\}$.

The case $F_{x_0} + F_{y_0} = 2F_m$ is impossible since $F_{x_0} \leq F_m, F_{y_0} \leq F_m$ and at least one of these two inequalities is strict because x_0 is not equal to y_0 .

Assume that $F_{x_0} + F_{y_0} = F_m$. Since both x_0 and y_0 are at most m , it follows that $F_{x_0} \geq 0$ and $F_{y_0} \geq 0$. Furthermore, both these inequalities are in fact strict since equality is achieved only in the case $y_0 = 0$, leading to $F_{x_0} = F_m$, so $x_0 = m$, which in turn is impossible because x_0 must be odd. Thus, both F_{x_0} and F_{y_0} are positive and less than F_m . If $\max\{F_{x_0}, F_{y_0}\} \leq F_{m-2}$, then

$$F_{m-3} + 2F_{m-2} = F_m = F_{x_0} + F_{y_0} < 2F_{m-2},$$

leading to $F_{m-3} < 0$, which is impossible. So one of F_{x_0} or F_{y_0} equals F_{m-1} , and therefore the other one is $F_m - F_{m-1} = F_{m-2}$. By parity arguments, we get that $F_{y_0} = F_{m-2}$ and $F_{x_0} = F_{m-1}$; so $(x_0, y_0) = (\pm(m-1), m-2)$.

Assume that $F_{x_0} + F_{y_0} = 0$. Then $F_{x_0} = -F_{y_0} = F_{-y_0}$, since y_0 is even. Since x_0 is odd, F_{x_0} is positive, so $y_0 < 0$. If $y_0 = -2$, we then get $F_{x_0} = F_2 = 1$, leading to $x_0 = \pm 1$. This gives us the possibilities $(x_0, y_0) = (\pm 1, -2)$. Finally, if $y_0 \leq -4$, then $x_0 = \pm y_0$, which is false since x_0 must be odd.

Assume that $F_{x_0} + F_{y_0} = -F_m$. Then $F_m \geq |F_{y_0}| = |-F_m - F_{x_0}| = F_m + F_{x_0} > F_m$, again because $F_{x_0} > 0$, but this last inequality is false. This completes the proof of this lemma. □

Lemma 4.2. *Let $m \geq 6$ be even and let $-m < x_0, y_0, z_0 \leq m$ be integers satisfying $F_{x_0} + F_{y_0} + F_{z_0} \equiv 0 \pmod{F_m}$. Then*

- (i) either all three x_0, y_0, z_0 are even and their sum is a multiple of m , or
- (ii) up to symmetries, (x_0, y_0, z_0) equals $(a, \pm(a+1), -a-2)$ for some even integer a , or
- (iii) up to symmetries, (x_0, y_0, z_0) equals one of $(\pm 3, -2, -2), (\pm 1, \pm 3, -4), (m, \pm 1, -2), (m, \pm(m-1), m-2), (\pm(m-1), \pm(m-3), m-4), (m-2, m-2, \pm(m-3))$ or $(0, \pm(m-1), m-2)$.

Proof. Assume that one of $x_0, y_0, z_0 \in \{0, m\}$. Say $x_0 \in \{0, m\}$. Then $F_{y_0} + F_{z_0} \equiv 0 \pmod{F_m}$. If y_0 and z_0 have distinct parities, then, by Lemma 4.1, we get that, up to symmetries,

$$(x_0, y_0, z_0) \in \{(m, \pm(m-1), m-2), (m, \pm 1, -2)\}.$$

Assume now that y_0 and z_0 have the same parities. If $y_0 \in \{0, m\}$, then $F_m \mid F_{z_0}$, so that $z_0 \in \{0, m\}$ and $x_0 + y_0 + z_0 \equiv 0 \pmod{m}$. Assume now that neither of y_0, z_0 is in $\{0, m\}$. If they are even, then $|F_{y_0}| \leq F_{m-2}$ and $|F_{z_0}| \leq F_{m-2}$, therefore $|F_{y_0} + F_{z_0}| \leq 2F_{m-2} < F_m$, so $F_{y_0} + F_{z_0} = 0$, leading to $F_{y_0} = -F_{z_0} = F_{-z_0}$. Since both y_0 and z_0 are even, we get that $y_0 = -z_0$, so indeed all three numbers x_0, y_0, z_0 are even and $x_0 + y_0 + z_0 \equiv 0 \pmod{m}$. If y_0 and z_0 are odd, then we may replace y_0 and z_0 by their absolute values and note that F_{y_0} and F_{z_0} are both positive. Assume that $y_0 \geq z_0$. If $y_0 \leq m-3$, then $F_{y_0} + F_{z_0} \leq 2F_{m-3} < F_m$, which is impossible. If $y_0 = m-1$, then $F_{z_0} = F_m - F_{m-1} = F_{m-2}$, leading to $z_0 = m-2$ (because $m-2 \geq 4$), which is false since z_0 is odd.

From now on, we assume that $x_0, y_0, z_0 \in (-m, m)$ and that none of them is zero. Then $|F_{x_0} + F_{y_0} + F_{z_0}| \leq 3F_{m-1} < 2F_m$, so that $F_{x_0} + F_{y_0} + F_{z_0} \in \{0, \pm F_m\}$. Assume that $F_{x_0} + F_{y_0} + F_{z_0} = 0$. Since none of these numbers is zero, it follows that at least one of them is negative. Say z_0 is such that F_{z_0} is negative and has the largest absolute value (among the negative numbers from the set $\{F_{x_0}, F_{y_0}, F_{z_0}\}$). Then z_0 is even and negative and $F_{x_0} + F_{y_0} = F_{-z_0}$. Assume first that F_{x_0} and F_{y_0} are positive. Then $|y_0| < |z_0|$. If $|y_0| = |z_0| - 1 = -z_0 - 1$, we then get that $F_{x_0} = F_{-z_0-2}$. If $|z_0| \geq 6$, then $x_0 = -z_0 - 2$. Putting $x_0 = a$, we get that $x_0 = a, y_0 = \pm(a+1), z_0 = -a-2$. Thus, we obtain the possibility

$$(x_0, y_0, z_0) = (a, \pm(a+1), -a-2), \quad \text{for some even } a.$$

If $z_0 = -4$, then $|y_0| = 3$ and $F_{x_0} = F_2$, and therefore $x_0 \in \{\pm 1, 2\}$. The case $x_0 = 2$ is part of the previous parametric family with $a = 2$, while, for $x_0 = \pm 1$, we obtain the possibilities

$$(x_0, y_0, z_0) = (\pm 1, \pm 3, -4).$$

Continue to assume that $F_{x_0} + F_{y_0} = F_{-z_0}$ but that one of F_{x_0} and F_{y_0} is negative. Say $F_{y_0} < 0$. Then y_0 is even and negative. Thus, $F_{x_0} = F_{|y_0|} + F_{|z_0|}$. If $|z_0| = 2$, then $|y_0| = 2$ and we get that $F_{x_0} = 2$, so $x_0 = \pm 3$. Thus, we obtain the possibility

$$(x_0, y_0, z_0) = (\pm 3, -2, 2).$$

If $|z_0| \geq 4$, then either $F_{|y_0|} + F_{|z_0|} = 2F_{|z_0|} \in (F_{|z_0|+1}, F_{|z_0|+2})$, so this is not a Fibonacci number, or $|y_0| \leq |z_0| - 2$, therefore $F_{|z_0|} < F_{|z_0|} + F_{|y_0|} \leq F_{|z_0|} + F_{|z_0|-2} < F_{|z_0|+1}$, so $F_{|z_0|} + F_{|y_0|}$ is not a Fibonacci number either.

Now assume that $F_{x_0} + F_{y_0} + F_{z_0} = F_m$. If $\max\{F_{x_0}, F_{y_0}, F_{z_0}\} \leq F_{m-3}$, we then get that $F_{x_0} + F_{y_0} + F_{z_0} \leq 3F_{m-3} < F_m$, which is impossible. So, let us assume that $F_{x_0} \in \{F_{m-1}, F_{m-2}\}$ and that $F_{x_0} \geq F_{y_0} \geq F_{z_0}$. If $F_{x_0} = F_{m-1}$, then $F_{y_0} + F_{z_0} = F_{m-2}$. If $F_{y_0} = F_{m-1}$ also, then $F_{z_0} = F_{m-2} - F_{m-1} = -F_{m-3}$, which is impossible since $m - 3$ is odd. Clearly, $F_{y_0} \neq F_{m-2}$, since the contrary leads to $F_{z_0} = 0$. If $F_{y_0} = F_{m-3}$, then $F_{z_0} = F_{m-2} - F_{m-3} = F_{m-4}$. Thus, we have obtained the possibilities

$$(x_0, y_0, z_0) = (\pm(m - 1), \pm(m - 3), m - 4).$$

If $F_{y_0} \leq F_{m-4}$, then $F_{z_0} = F_{m-2} - F_{y_0} \geq F_{m-3} > F_{m-4} \geq F_{y_0}$, which is impossible. This takes care of the case when $F_{x_0} = F_{m-1}$. Assume now that $F_{x_0} = F_{m-2}$. Then $F_{y_0} + F_{z_0} = F_{m-1}$. If $F_{y_0} \leq F_{m-3}$, then $F_{y_0} + F_{z_0} \leq 2F_{m-3} < F_{m-1}$, which is a contradiction. Thus, $F_{y_0} = F_{m-2}$, giving $F_{z_0} = F_{m-3}$. Hence,

$$(x_0, y_0, z_0) = (m - 2, m - 2, \pm(m - 3)).$$

Now assume that $F_{x_0} + F_{y_0} + F_{z_0} = -F_m = F_{-m}$. If at most two of the Fibonacci numbers involved are negative, then they are in absolute value less than or equal to F_{m-2} (because their indices are even). Thus, $F_m = |F_{x_0} + F_{y_0} + F_{z_0}| < 2F_{m-2}$, which is false. Consequently, all three Fibonacci numbers are negative, so all their indices are negative and even. Changing (x_0, y_0, z_0) to $(-x_0, -y_0, -z_0) = (|x_0|, |y_0|, |z_0|)$, we get a solution to

$$F_{|x_0|} + F_{|y_0|} + F_{|z_0|} = F_m.$$

If at most two of the Fibonacci numbers involved are less than or equal to F_{m-2} , then

$$F_m = F_{|x_0|} + F_{|y_0|} + F_{|z_0|} \leq 2F_{m-2} + F_{m-4} < F_m,$$

which is impossible, while if all three of them are F_{m-2} , we then get $3F_{m-2} = F_m$, which is also impossible for $m \geq 6$ since the left-hand side is in fact larger than the right-hand side.

This completes the proof of Lemma 4.2. □

5. Skolem’s method

In this section we show—using an argument similar to Skolem’s method—that if x, y, z, n is a solution of the Diophantine equation (1.2), then certain linear forms in x, y, z are multiples of m or $F_{m/2}$ for all even integers $m \geq 6$ such that F_m is not too large with respect to n . Throughout this section we study the equation

$$F_x + F_y + F_z = n!$$

in non-negative integers x, y, z with $n \geq 7$. From now on we make the following convention. If precisely two of the unknowns x, y, z are even, then we shall suppose that these are x and z . If exactly one of them is even, we shall suppose that it is x .

Lemma 5.1. *Let x, y, z, n be as above. Let $m \geq 6$ be an even integer such that both F_m and $2F_{m/2}^2$ divide $n!$ (a sufficient condition for both of these to divide $n!$ is $L_{m/2} \leq n$). Then*

- (a) *not all of x, y, z are odd,*
- (b) *if x, y, z are all even, m divides $x + y + z$,*
- (c) *if x is even and y, z are odd, m divides $x + 4$,*
- (d) *if x, z are both even and y is odd, either m divides $x + z + 2$ or $F_{m/2}$ divides $3x \pm 4y + 3z$; moreover, this latter expression is non-zero.*

Proof. Let $m \geq 6$ be an even integer. First, we prove the observation that $L_{m/2} \leq n$ implies that both F_m and $2F_{m/2}^2$ divide $n!$. Thus, suppose that $L_{m/2} \leq n$. Then $F_{m/2} < L_{m/2}$ and $F_m = F_{m/2}L_{m/2}$. Hence, F_m divides $n!$. Clearly, $2F_{m/2}^2$ divides $n!$ for $m = 6$. Suppose $m \geq 8$, so $F_{m/2} < \frac{1}{2}L_{m/2}$. Thus, $2F_{m/2}^2$ divides $n!$. This proves the observation.

From now on we drop the condition that $L_{m/2} \leq n$ but assume that both F_m and $2F_{m/2}^2$ divide $n!$. Write

$$x = x_0 + 2\lambda m, \quad y = y_0 + 2\mu m, \quad z = z_0 + 2\epsilon m,$$

where $-m < x_0, y_0, z_0 \leq m$. By (3.4), $F_{x_0} + F_{y_0} + F_{z_0} \equiv 0 \pmod{F_m}$. Lemma 4.2 gives a number of possibilities for x_0, y_0, z_0 . Clearly, x, y, z have the same parities, respectively, as x_0, y_0, z_0 . By examining the possibilities in Lemma 4.2, we see that x, y, z are not all odd, and that if x, y, z are all even, then m divides $x + y + z$.

Suppose that two of x, y, z are odd and one is even. By our convention above, x must be even. Then, from Lemma 4.2, we see that (x, y, z) is congruent modulo m to one of

$$(-4, \pm 1, \pm 3), (-4, \pm 3, \pm 1),$$

showing that m divides $x + 4$.

There now only remains the case in which precisely two of x, y, z are even and one is odd. By our convention, x, z are even and y is odd. From Lemma 4.2, we see that (x, y, z) is congruent modulo $2m$ to one of

$$\begin{aligned} &(a, \pm(a+1), -a-2), (-a-2, \pm(a+1), a), (m, \pm(m-1), m-2), \\ &(m-2, \pm(m-1), m), (0, \pm(m-1), m-2), (m-2, \pm(m-1), 0), \\ &(m, \pm 1, -2), (-2, \pm 1, m), \end{aligned}$$

or to one of $(-2, \pm 3, -2), (m-2, \pm(m-3), m-2)$. In all but the last two cases, m divides $x + z + 2$.

It remains to consider the case where (x, y, z) is congruent modulo $2m$ to one of $(-2, \pm 3, -2), (m-2, \pm(m-3), m-2)$. Note here that $x \equiv z \pmod{4}$ and that $(x, y, z) \equiv (-2, \pm 3, -2) \pmod{m}$. We now write

$$x = -2 + \lambda_1 m, \quad y = \pm 3 + \mu_1 m, \quad z = -2 + \epsilon_1 m.$$

Observe that λ_1, μ_1 and ϵ_1 have the same parity. Moreover, $F_{-2} + F_{\pm 3} + F_{-2} = 0$. It follows from (3.5) that

$$2n! \equiv \pm 2F_m(L_{-2}\lambda_1 + L_{\pm 3}\mu_1 + L_{-2}\epsilon_1) + 5F_m^2(F_{-2}\lambda_1^2 + F_{\pm 3}\mu_1^2 + F_{-2}\epsilon_1^2) \pmod{F_{m/2}^3}.$$

Here, our observation that λ_1, μ_1 and ϵ_1 have the same parity is crucial. We now consider two subcases. The first is $3 \nmid m$. This means that $F_{m/2}, L_{m/2}$ are odd and coprime. Recall that $2F_{m/2}^2$ divides $n!$. From $F_m = F_{m/2}L_{m/2}$ and the coprimality of $F_{m/2}$ and $L_{m/2}$, we obtain that $F_{m/2}$ divides $L_{-2}\lambda_1 + L_{\pm 3}\mu_1 + L_{-2}\epsilon_1 = 3\lambda_1 \pm 4\mu_1 + 3\epsilon_1$. Thus, $3x \pm 4y + 3z = m(3\lambda_1 \pm 4\mu_1 + 3\epsilon_1)$ is divisible by $F_{m/2}$. The second subcase is $3 \mid m$. Hence, $F_{m/2}$ and $L_{m/2}$ are both even, and their greatest common divisor is 2. We now obtain that $\frac{1}{2}F_{m/2}$ divides $3\lambda_1 \pm 4\mu_1 + 3\epsilon_1$. But m is even. Thus, $3x \pm 4y + 3z$ is divisible by $F_{m/2}$ in this case as well.

All that it remains to prove is that the expression $3x \pm 4y + 3z$ does not vanish. This is clearly true for $3x + 4y + 3z$. Suppose that $3x - 4y + 3z = 0$. Recall our observation above that $x \equiv z \pmod{4}$. Then $y = \frac{3}{4}(x + z)$, and using (3.3) we obtain

$$n! = F_x + F_z + F_{3(x+z)/4} = F_{(x+z)/2}L_{(x-z)/2} + F_{3(x+z)/4}.$$

The right-hand side is divisible by $F_{(x+z)/4}$, and so this divides $n!$. If $\frac{1}{4}(x + z) \leq 12$, then we can list all the solutions. Hence, suppose that $\frac{1}{4}(x + z) > 12$. By the Primitive Divisor Theorem [2], $F_{(x+z)/4}$ has some prime divisor p such that $p \equiv \pm 1 \pmod{\frac{1}{4}(x + z)}$. But $p \mid n!$, which gives that $p \leq n$. Thus,

$$\frac{1}{4}(x + z) \leq n + 1,$$

and so $x \leq 4n + 4, y \leq 3n + 3$ and $z \leq 4n + 4$. However, $F_{\max\{x,y,z\}} \geq \frac{1}{3}n!$, giving a contradiction for $n \geq 7$. □

6. Bounds on n when x is even and y, z have the same parity

6.1. Case I: x, y, z are all even

Let us suppose that $n \geq 200$. In this case, we know from Lemma 5.1 that all even $m \geq 6$ with $L_{m/2} \leq n$ satisfy $m \mid (x + y + z)$.

Let p run through the integers

$$3 \leq p \leq 2.076 \log n.$$

By Lemma 2.3, $L_p \leq n$, and so $2p$ divides $x + y + z$. Thus, by Lemma 2.2,

$$\theta(2.076 \log n) \leq \log(x + y + z) \leq \log\left(\frac{3n \log n}{\log \alpha}\right).$$

The first bound that we prove for n is $n \leq L_{31}$. Suppose that $n \geq L_{31} + 1$. Then $2.076 \log n > 30$, and so, by Proposition 2.5,

$$\theta(2.076 \log n) \geq 0.7033 \times 2.076 \log n > 1.46 \log n.$$

Hence,

$$1.46 \log n < \log \left(\frac{3n \log n}{\log \alpha} \right) = \log n + \log \log n + \log(3/\log \alpha).$$

This is impossible for $n \geq L_{31} + 1$. Thus, $n \leq L_{31}$.

Hence,

$$x + y + z \leq \frac{3n \log n}{\log \alpha} \leq 279\,962\,456.$$

Suppose that $n \geq L_{19}$. Then $x + y + z$ is divisible by $2p$ for all integers $3 \leq p \leq 19$. However,

$$\text{lcm}(6, 8, \dots, 38) = 465\,585\,120 > 279\,962\,456 \geq x + y + z,$$

giving a contradiction. Thus, $n \leq L_{19}$. Repeating the argument shows that $n \leq L_{13}$ and finally that $n \leq L_{11} = 199$. This contradicts our initial assumption that $n \geq 200$, and so $n \leq 199$.

6.2. Case II: x is even and y and z are odd

In this case we know from Lemma 5.1 that all even $m \geq 6$ with $L_{m/2} \leq n$ satisfy $m \mid (x + 4)$. A similar argument to the one above now shows that $n \leq 199$.

7. Bound for n when x, z are even and y is odd

7.1. An initial bound

Suppose that $n \geq 200$. Let $0 < \gamma < 1$ be a real number to be chosen later. Let p be a prime satisfying

$$2.079\gamma \log n + 2.441 \leq p \leq 2.076 \log n.$$

By Lemma 2.3, we have that $L_p \leq n$ and $F_p \geq n^\gamma$. We know, by Lemma 5.1 applied to $m = 2p$, that either $2p$ divides $x + z + 2$ or F_p divides one of the (non-zero) expressions $3x \pm 4y + 3z$.

From Lemma 2.2,

$$x + z + 2 \leq \frac{2n \log n}{\log \alpha}, \quad |3x \pm 4y + 3z| \leq \frac{10n \log n}{\log \alpha}.$$

Suppose that k is a positive integer satisfying

$$n^{\gamma(k+1)} > \frac{10n \log n}{\log \alpha}. \quad (7.1)$$

Then at most k of the numbers F_p for the primes p in the given range divide $3x + 4y + 3z$, and at most another k of these divide $3x - 4y + 3z$. Note that here we are making use of the fact that the F_p are coprime as p runs through the primes; this is a consequence of the well-known property $\gcd(F_u, F_v) = F_{\gcd(u,v)}$ for all integers u and v .

It follows that for all but at most $2k$ primes p in the range above, $2p$ divides $x + z + 2$. Hence,

$$\begin{aligned} &\theta(2.076 \log n) - \theta(2.079\gamma \log n + 2.441) - 2k \log(2.076 \log n) \\ &\leq \log(x + z + 2) \leq \log\left(\frac{2n \log n}{\log \alpha}\right) \leq \log n + \log \log n + 1.425. \end{aligned} \quad (7.2)$$

Now suppose that k and γ are fixed and that n is very large. Recall that for large x , $\theta(x) = x + o(x)$ as $x \rightarrow \infty$. Thus, the above inequalities give

$$(1.076 - 2.079\gamma) \log n \leq o(\log n) \quad \text{as } n \rightarrow \infty,$$

showing that n must in fact be bounded provided that γ is small enough. We use this idea to obtain an explicit bound for n .

We first show that $n \leq L_{501}$. So, suppose that $n \geq L_{501} + 1$. We let $k = 2$ and $\gamma = 0.35$. It is easy to show that (7.1) holds. Moreover,

$$2.076 \log n \geq 500,$$

and so, by Proposition 2.5, we have

$$\theta(2.076 \log n) \geq 0.9194 \times 2.076 \log n > 1.908 \log n$$

and

$$\theta(2.079\gamma \log n + 2.441) \leq 1.001102(2.079\gamma \log n + 2.441) < 0.729 \log n + 2.444.$$

Equation (7.2) gives

$$0.188 \log n \leq 5 \log \log n + 6.791.$$

This is impossible for $n \geq L_{501} + 1$. Hence, $n \leq L_{501} < 5.045 \times 10^{104}$.

7.2. A recursive procedure for reducing the bound

We now give an iterative argument that will be used repeatedly to reduce the above bound. Our argument is reminiscent of that given at the end of §6 but is substantially more complicated. Write

$$\mathcal{E} = x + z + 2, \quad \mathcal{F} = |(3x + 4y + 3z)(3x - 4y + 3z)|.$$

For a positive integer $b \geq 2$ we put

$$C_{\mathcal{E},b} = \frac{2L_b \log L_b}{\log \alpha}, \quad C_{\mathcal{F},b} = \frac{60L_b^2 (\log L_b)^2}{(\log \alpha)^2}.$$

Lemma 7.1. *If $n \leq L_b$, then $\mathcal{E} \leq C_{\mathcal{E},b}$ and $\mathcal{F} \leq C_{\mathcal{F},b}$.*

Proof. This follows from Lemma 2.2. □

For positive integers u and a with $2 \leq u \leq a$ define

$$H_a(u) = \text{lcm}\{F_v : 3 \leq v \leq a, u \mid v\}.$$

For $2 \leq u_1 < u_2 < \dots < u_n \leq a$ define

$$H_a(u_1, \dots, u_n) = \text{lcm}(H_a(u_1), \dots, H_a(u_n)).$$

Lemma 7.2. *Suppose that $n \geq L_a$ and that $2 \leq u \leq a$. If $u \nmid \mathcal{E}$, then $H_a(u) \mid \mathcal{F}$. If $2 \leq u_1 < u_2 < \dots < u_n \leq a$ and all $u_i \nmid \mathcal{E}$, then $H_a(u_1, \dots, u_n)$ divides \mathcal{F} .*

Proof. Let v be an integer satisfying $3 \leq v \leq a$ and $u \mid v$. Write $m = 2v$. Then m is an even integer satisfying $m \geq 6$ and $L_{m/2} \leq L_a \leq n$. By Lemma 5.1, either m divides \mathcal{E} or $F_v = F_{m/2}$ divides \mathcal{F} . But $u \mid v \mid m$ and $u \nmid \mathcal{E}$. Thus, $m \nmid \mathcal{E}$, and so F_v divides \mathcal{F} as required. \square

Now let $2 \leq a \leq b$. Put

$$\mathcal{P}_a = \{u : 2 \leq u \leq a, u \text{ is a prime power}\}.$$

We define a sequence of subsets of the powerset of \mathcal{P}_a as

$$S_0(a, b) = \{\emptyset\}, \quad S_1(a, b) = \{\{u\} : 2 \leq u \leq a, H_{u,a} \leq C_{\mathcal{F},b}\},$$

and, for $k \geq 1$, we define $S_{k+1}(a, b)$ to be the set of $(k+1)$ -tuples $\{u_1, \dots, u_{k+1}\}$ satisfying

- (i) $2 \leq u_1 < u_2 < \dots < u_{k+1} \leq a$,
- (ii) $\{u_1, \dots, u_k\} \in S_k(a, b)$,
- (iii) $\{u_{k+1}\} \in S_1(a, b)$,
- (iv) $H_a(u_1, \dots, u_{k+1}) \leq C_{\mathcal{F},b}$.

We put

$$S(a, b) = \bigcup_{k \geq 0} S_k(a, b).$$

Lemma 7.3. *Let $2 \leq a \leq b$ and suppose that $L_a \leq n \leq L_b$. Let $V = \{u \in \mathcal{P}_a : u \mid \mathcal{E}\}$. Then $\mathcal{P}_a \setminus V \in S(a, b)$.*

Proof. Write $\mathcal{P}_a \setminus V = \{u_1, \dots, u_j\}$, where $u_1 < \dots < u_j$. No u_i divides \mathcal{E} , and so, by Lemma 7.2, $H_a(u_1, \dots, u_j)$ divides \mathcal{F} . By Lemma 7.1, we have $H_a(u_1, \dots, u_j) \leq C_{\mathcal{F},b}$. Clearly, for each $k \leq j-1$, $H_a(u_1, \dots, u_{k-1}) \leq C_{\mathcal{F},b}$ and $H_a(u_k) \leq C_{\mathcal{F},b}$. This shows inductively that $\{u_1, \dots, u_k\} \in S_k(a, b)$ for $k = 1, \dots, j$. Thus, $\mathcal{P}_a \setminus V \in S_j(a, b) \subseteq S(a, b)$. \square

Lemma 7.4. *Let $2 \leq a \leq b$. Suppose that for each $U \in S(a, b)$ we have*

$$\text{lcm}(\mathcal{P}_a \setminus U) > C_{\mathcal{E},b}.$$

Then there is no solution to the Diophantine equation (1.2) with x, z even, y odd and $L_a \leq n \leq L_b$.

Proof. Suppose that $L_a \leq n \leq L_b$. Let $V = \{u \in \mathcal{P}_a : u \mid \mathcal{E}\}$ and let $U = \mathcal{P}_a \setminus V$. By Lemma 7.3, $U \in S(a, b)$. Moreover,

$$\text{lcm}(\mathcal{P}_a \setminus U) = \text{lcm}(V) \mid \mathcal{E}.$$

However, by Lemma 7.1, $\mathcal{E} \leq C_{\mathcal{E},b}$. This gives a contradiction. □

We have shown previously that $n \leq L_{501}$. We shall apply Lemma 7.4 to repeatedly reduce this bound on n . First we let $a = 490$ and $b = 501$. We used a simple MAGMA script to compute \mathcal{P}_a and $S_k(a, b)$. We found that \mathcal{P}_a has 112 elements, $S_1(a, b)$ has 84 elements, $S_2(a, b)$ has 2565 elements, $S_3(a, b)$ has 8609 elements, $S_4(a, b)$ has 16 elements and $S_k(a, b) = \emptyset$ for $k \geq 5$. Altogether, $S(a, b)$ has 11 275 elements. We check the criterion of Lemma 7.4 and find that it holds for all $U \in S(a, b)$. Thus, there are no solutions to (1.2) with x, z even, y odd, and $L_{490} \leq n \leq L_{501}$. This shows that $n \leq L_{490}$. Repeating the above argument another 50 times shows that $n \leq L_{37} = 54\,018\,521$.

8. The final sieve

We know from the previous three sections that all solutions of the Diophantine equation (1.2) satisfy $n \leq 54\,018\,521$. In this section, we shall determine all solutions to (1.2) with $n \leq 6 \times 10^7$ and thus complete the proof of Theorem 1.1.

Lemma 8.1. *In (1.2), suppose that $x \geq y \geq z$. Then*

$$\frac{\log n!}{\log \alpha} + C_1 \leq x \leq \frac{\log n!}{\log \alpha} + C_2, \tag{8.1}$$

where

$$C_1 = \frac{\log(\frac{1}{3}\sqrt{5}) - 1/\sqrt{5}}{\log \alpha}, \quad C_2 = \frac{\log(\sqrt{5}) + 1/(6\sqrt{5})}{\log \alpha}.$$

Proof. The lemma is easily checked for $n \leq 2$, so suppose $n \geq 3$. Clearly,

$$\frac{1}{3}n! \leq F_x \leq n!,$$

and so

$$\frac{1}{3}\sqrt{5}n! - 1 \leq \alpha^x \leq \sqrt{5}n! + 1.$$

Taking logarithms, we find that

$$x \log \alpha \leq \log n! + \log \sqrt{5} + \log(1 + 1/(n!\sqrt{5})) \leq \log n! + \log \sqrt{5} + 1/(6\sqrt{5})$$

by using Lemma 2.4 and the fact that $n \geq 3$. Moreover,

$$x \log \alpha \geq \log n! + \log(\frac{1}{3}\sqrt{5}) + \log(1 - 3/(n!\sqrt{5})) \geq \log n! + \log(\frac{1}{3}\sqrt{5}) - 1/\sqrt{5},$$

again using Lemma 2.4 and $n \geq 3$. This completes the proof. □

For the purpose of searching for all solutions to (1.2), we may without loss of generality suppose that $x \geq y \geq z$. The last lemma above gives, for each n , an interval containing at most three integers x .

Now let

$$l_1 = F_{43} = 433\,494\,437, \quad l_2 = F_{47} = 2\,971\,215\,073.$$

Both these Fibonacci numbers are prime and they have been chosen because the period of the Fibonacci sequence modulo each l_i is particularly small; the periods are 172 and 188, respectively. Let

$$T_1 = \{F_u + F_v \pmod{l_1} : 0 \leq u \leq 171\}$$

and

$$T_2 = \{F_u + F_v \pmod{l_2} : 0 \leq u \leq 187\}.$$

Using MAGMA, we find that T_1 has 2821 elements and T_2 has 3453 elements. Thus,

$$\#T_1/l_1 \approx 6.5 \times 10^{-6}, \quad \#T_2/l_2 \approx 1.16 \times 10^{-6}.$$

Now our MAGMA program for determining the solutions of (1.2) with $n \leq 6 \times 10^7$ is as follows. For each n we need to compute three quantities. The first is $\log(n!)/\log(\alpha)$, the second is $n! \pmod{l_1}$ and the third is $n! \pmod{l_2}$. Knowing these for $n = k - 1$ quickly gives these for $n = k$. For each n , we determine the integers x in the interval (8.1). For each x , we compute $F_x \pmod{l_1}$ and $F_x \pmod{l_2}$. If $n! - F_x$ modulo l_1 does not belong to T_1 , or $n! - F_x$ modulo l_2 does not belong to T_2 , then we know that there is no solution to Equation (1.2) with the given values of n and x . Computing $F_x \pmod{l_i}$ can be done in $O(\log x) = O(\log n)$ steps as it involves only computing α^x modulo l_i , and so it is very fast. Our script took less than six hours to run on a dual core 3.00 GHz Opteron and produced only the following pairs of values of (x, n) for which $n! - F_x$ belongs to T_i modulo l_i ($i = 1, 2$):

$$(0, 1), (1, 1), (0, 2), (1, 2), (2, 2), (3, 2), (3, 3), (4, 3), (5, 3), \\ (6, 4), (7, 4), (8, 4), (9, 5), (11, 5), (14, 6), (15, 6).$$

From this, we easily recover our list of solutions in Theorem 1.1.

Note that the probability of a random integer belonging modulo l_1 to T_1 and modulo l_2 to T_2 is less than 10^{-11} . Since the possibilities for (x, n) are most $3 \times 6 \times 10^7 < 2 \times 10^8$, it is not at all surprising that our sieve found only pairs of (x, n) for which there are solutions (n, x, y, z) to (1.2).

Acknowledgements. F.L. is supported by Grant CONACyT 46755. S.S. is supported by a grant from the UK Engineering and Physical Sciences Research Council and by a Marie Curie International Reintegration Grant (MIRG-CT-2006-044530).

References

1. D. BEREND AND J. E. HARMSE, On polynomial-factorial Diophantine equations, *Trans. Am. Math. Soc.* **358**(4) (2006), 1741–1779.
2. YU. BILU, G. HANROT AND P. M. VOUTIER, Existence of primitive divisors of Lucas and Lehmer numbers, *J. Reine Angew. Math.* **539** (2001), 75–122.
3. M. BOLLMAN, Sums of consecutive factorials in the Fibonacci sequence, *Congr. Numer.* **194** (2009), 77–83.
4. Z. I. BOREVICH AND I. R. SHAFAREVICH, *Number theory* (Academic Press, 1966).
5. W. BOSMA, J. CANNON AND C. PLAYOUST, The MAGMA algebra system I: the user language, *J. Symb. Computat.* **24** (1997), 235–265 (see also www.maths.usyd.edu.au)
6. H. BROCARD, Question 166, *Nouv. Corresp. Math.* **2** (1876), 287.
7. Y. BUGEAUD, F. LUCAS, M. MIGNOTTE AND S. SIKSEK, Perfect powers from products of terms in Lucas sequences, *J. Reine Angew. Math.* **611** (2007), 109–129.
8. Y. BUGEAUD, M. MIGNOTTE AND S. SIKSEK, Classical and modular approaches to exponential Diophantine equations, I, Fibonacci and Lucas perfect powers, *Annals Math.* **163** (2006), 969–1018.
9. J. W. S. CASSELS, *Local fields*, London Mathematical Society Student Texts, Volume 3 (Cambridge University Press, 1986).
10. M. CIPU, F. LUCA AND M. MIGNOTTE, Solutions of the Diophantine equation $au^x + bv^y + cw^z = n!$, *Annals Sci. Math. Québec* **31** (2007), 127–137.
11. P. ERDŐS AND R. OBLÁTH, Über diophantische Gleichungen der Form $n! = x^p \pm y^p$ und $n! \pm m! = x^p$, *Acta Sci. Math. (Szeged)* **8** (1937), 241–255.
12. G. GROSSMAN AND F. LUCA, Sums of factorials in binary recurrence sequences, *J. Number Theory* **93**(2) (2002), 87–107.
13. R. K. GUY, *Unsolved problems in number theory*, 3rd edn (Springer, 2004).
14. R. KNOTT, Fibonacci numbers and the golden section, www.mcs.surrey.ac.uk/Personal/R.Knott/Fibonacci/.
15. F. LUCA, The number of nonzero digits of $n!$, *Can. Math. Bull.* **45**(1) (2002), 115–118.
16. F. LUCA AND P. STĂNICĂ, $F_1 F_2 F_3 F_4 F_5 F_6 F_8 F_{10} F_{12} = 11!$, *Portugaliae Math.* **63** (2006), 251–260.
17. S. RAMANUJAN, Question 469, *J. Indian Math. Soc.* **5** (1913), 59.
18. J. B. ROSSER AND L. SCHOENFELD, Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$, *Math. Comp.* **29** (1975), 243–269.