

A SUFFICIENT CONDITION FOR AN EXTREME COVERING OF n -SPACE BY SPHERES

T. J. DICKSON

(Received 29 August 1966)

1

The problem of finding the most economical coverings of n -dimensional Euclidean space by equal spheres whose centres form a lattice, which is equivalent to a problem concerning the inhomogeneous minima of positive definite quadratic forms, has been discussed recently by Barnes and Dickson [1]. The reader is referred to [1] for a complete background on the problem. Terms and notations used will be as in that paper.

Among other results, Barnes and Dickson proved that if f is an extreme form in the interior of a Voronoï cone Δ then (i) every extreme form in Δ is a multiple of f and (ii) f and Δ have the same group of automorphisms. It is the main object of this paper to extend these results to the following necessary and sufficient condition for extremeness.

THEOREM. *If \mathcal{F} is the set of all forms in a Voronoï cone Δ which are invariant under all automorphisms of Δ , and f is a form in the interior of a Voronoï cone Δ then f is extreme if and only if $f \in \mathcal{F}$ and*

$$(1.1) \quad \mu(f) = \min_{g \in \mathcal{F}} \mu(g).$$

§§ 2 and 3 are concerned with proving this result and § 4 with applying it to the known Voronoï cones.

In § 5 the case where the only form satisfying (1.1) lies on the boundary of the cone is discussed.

2

The following two lemmas are necessary for the proof of the above theorem.

LEMMA 1. *If f is an interior form, ϕ any quadratic form, then the inequality*

$$\mu(f + \varepsilon\phi) \leq \mu(f)$$

holds for all sufficiently small ε (of either sign) if and only if ϕ is a multiple of f .

This result appeared in [1] but was submerged in the proof of Theorem 2 of that paper. It was not stated as a separate result at that stage.

This lemma means that in the interior of a Voronoï cone the function $\mu(f)$ cannot have a local maximum on a line i.e. the maximum of $\mu(f)$ on a line segment must occur at one or both of the end points.

LEMMA 2. *If k forms f_0, f_1, \dots, f_{k-1} are in the interior of the same Voronoï cone and are not all multiples of the same form, then*

$$(2.1) \quad \mu(f_0 + f_1 + \dots + f_{k-1}) < \max_i \mu(f_i).$$

PROOF. The lemma will be proved by induction on k .

(i) $k = 2$.

Consider the forms

$$f_t(x) = (1-t)f_0(x) + tf_1(x)$$

for $0 \leq t \leq 1$. If the maximum value of $\mu(f_t)$ is attained at a point, f_{t_1} say, for which $0 < t_1 < 1$, then

$$\mu(f_{t_1 + \varepsilon(f_0 - f_1)}) \leq \mu(f_{t_1})$$

for all sufficiently small ε of either sign.

This is impossible, by Lemma 1, as f_0 is not a multiple of f_1 and so $\mu(f_t) < \max_{i=0,1} \mu(f_i)$ for all $0 < t < 1$.

Taking $t = \frac{1}{2}$ and remembering that $\mu(\lambda f) = \mu(f)(\lambda > 0)$, we have

$$(2.2) \quad \mu(f_0 + f_1) < \max_{i=0,1} \mu(f_i)$$

(ii) Assume (2.1) true for $k \leq K$ and consider case $k = K + 1$.

Without loss of generality we can assume that f_0, f_1, \dots, f_{K-1} are not all multiples of the same form and f_K is not a multiple of $f_0 + f_1 + \dots + f_{K-1}$.

By assumption

$$(2.3) \quad \mu(f_0 + f_1 + \dots + f_{K-1}) < \max_{i=0, \dots, K-1} \mu(f_i)$$

and, from (2.2),

$$(2.4) \quad \mu(f_0 + \dots + f_K) < \max\{\mu(f_K), \mu(f_0 + \dots + f_{K-1})\}.$$

(2.3) and (2.4) together give (2.1) and so (2.1) holds for all k .

3. Proof of the Theorem

(i) Let f be an extreme interior form of a Voronoï cone Δ .

From the results of Barnes and Dickson [1], $f \in \mathcal{F}$ and obviously

$$(3.1) \quad \mu(f) \leq \mu(g)$$

for all $g \in \mathcal{F}$ sufficiently close to f .

Let f_0 be a form in \mathcal{F} such that $\mu(f_0) = \min_{g \in \mathcal{F}} \mu(g)$ and consider line segment of forms

$$f_t(x) = tf(x) + (1-t)f_0(x)$$

for $0 \leq t \leq 1$.

By (3.1) and the definition of f_0 , the maximum value of $\mu(f_t)$ is attained at a point f_{t_1} for which $0 < t_1 < 1$ and so

$$\mu(f_{t_1} + \varepsilon(f_0 - f)) \leq \mu(f_{t_1})$$

for all sufficiently small ε of either sign. By lemma 1, $f_0 - f$ is a multiple of f_{t_1} i.e. f_0 is a multiple of f .

Thus $\mu(f) = \min_{g \in \mathcal{F}} \mu(g)$.

(ii) Let f be an interior form of Δ such that $f \in \mathcal{F}$ and

$$(3.2) \quad \mu(f) = \min_{g \in \mathcal{F}} \mu(g).$$

Let there be another form f_0 in Δ for which $\mu(f_0) < \mu(f)$.

Obviously $f_0 \notin \mathcal{F}$.

Consider the forms obtained by applying *all* automorphisms of Δ to f_0 and let these forms be $f_i (i = 0, \dots, k-1)$.

As $f_0 \notin \mathcal{F}$, not all the f_i are multiples of the same form.

Now $\mu(f_i) = \mu(f_0)$ for $(i = 0, \dots, k-1)$, so, by Lemma 2

$$\mu(f_0 + f_1 + \dots + f_{k-1}) < \mu(f_0) < \mu(f).$$

But $f_0 + f_1 + \dots + f_{k-1} \in \mathcal{F}$ and we have a contradiction to (3.2).

Thus $\mu(g) \geq \mu(f)$ for all g in Δ and hence f is extreme.

The proof is now complete.

In (i) of the above proof, if f_0 had been defined as any form for which

$$\mu(f_0) = \min_{g \in \Delta} \mu(g)$$

then the same argument would give that f_0 is a multiple of f . Thus we have:

COROLLARY. *If f is an extreme interior form of Δ then*

$$\mu(g) > \mu(f)$$

for all g in Δ not a multiple of f .

4. Applications

Dickson extracted all the extreme quaternary forms [2] by finding the set \mathcal{F} for each cone and minimizing $\mu(f)$ over \mathcal{F} in each case, i.e. finding the forms satisfying (1.1). However, as this condition had not been shown to be sufficient for extremeness, another result of [1] had to be used

to prove that the forms were in fact extreme and the uniqueness of the extreme form in any Voronoï cone used to conclude that they were the only extreme forms. With the above theorem, however, we see that we can conclude immediately that the respective forms are extreme. A similar comment applies to the treatment of the principal domain in [1] by Barnes and Dickson.

The following discussion of the known Voronoï cones shows how easily the extreme forms can be extracted using the above theorem.

(i) *The principal domain*

For every $n \geq 2$, the principal domain Δ is the set of forms expressible in the form

$$\phi(x) = \sum_{\substack{i,j=0 \\ i < j}}^n \rho_{ij}(x_i - x_j)^2, \quad x_0 = 0, \quad \rho_{ij} \geq 0 \quad (0 \leq i < j \leq n).$$

The group $G(\Delta)$ of automorphisms of Δ has order $2(n+1)!$ and is transitive on the edge forms $(x_i - x_j)^2$ ($0 \leq i < j \leq n$) of Δ . $G(\Delta)$ is in fact generated by (i) all permutations of x_1, x_2, \dots, x_n ; (ii) $x_i \rightarrow -x_i$ ($1 \leq i \leq n$); (iii) $x_1 \rightarrow x_1, x_i \rightarrow x_1 - x_i$ ($2 \leq i \leq n$).

So the set \mathcal{F} as defined in the above theorem contains only multiples of

$$f(x) = \sum_{\substack{i,j=0 \\ i < j}}^n (x_i - x_j)^2 = n \sum_{i=1}^n x_i^2 - 2 \sum_{1 \leq i < j \leq n} x_i x_j.$$

As this is an interior form then by the above theorem it is extreme and there are no other extreme forms (including boundary forms) in Δ .

(ii) *The cone Δ'' of quaternary forms*

Δ'' is the set of forms expressible in the form

$$\begin{aligned} \phi(x) = & \rho_1 x_1^2 + \rho_2 x_2^2 + \rho_3 x_3^2 + \rho_4 x_4^2 + \rho_{12} \omega + \rho_{13} (x_1 - x_3)^2 \\ & + \rho_{14} (x_1 - x_4)^2 + \rho_{23} (x_2 - x_3)^2 + \rho_{24} (x_2 - x_4)^2 + \rho_{34} (x_1 + x_2 - x_3 - x_4)^2 \end{aligned}$$

where $\rho_i \geq 0, \rho_{ij} \geq 0$ for all i, j and

$$\omega = 2x_1^2 + 2x_2^2 + 2x_3^2 + 2x_4^2 + 2x_1 x_2 - 2x_1 x_3 - 2x_1 x_4 - 2x_2 x_3 - 2x_2 x_4.$$

The group $G(\Delta'')$ of automorphisms of Δ'' contains the transformations (i) $x_1 \leftrightarrow x_2$ (ii) $x_3 \leftrightarrow x_4$ (iii) $x_1 \rightarrow x_1 - x_3, x_2 \rightarrow x_4, x_3 \rightarrow x_1, x_4 \rightarrow x_4 - x_2$. The family of forms of Δ'' (neglecting boundary forms) which are invariant under these transformations are those expressible in the form

$$\begin{aligned} f(x) = & x_1^2 + x_2^2 + x_3^2 + x_4^2 + (x_1 - x_3)^2 + (x_1 - x_4)^2 + (x_2 - x_3)^2 + (x_2 - x_4)^2 \\ & + (x_1 + x_2 - x_3 - x_4)^2 + \alpha \omega \quad (\alpha > 0). \end{aligned}$$

As ω is not equivalent to the other edge forms of Δ'' , then this is the set \mathcal{F} (neglecting boundary forms).

By using Voronoi's table of vertices for the cone Δ'' [3p. 173], as in [2], we obtain that

$$\mu(f) = \frac{2(2-3\alpha+3\alpha^2-\alpha^3)}{(1+\alpha)^{\frac{3}{2}}(2-\alpha)^{\frac{1}{2}}}$$

for all f in \mathcal{F} .

Solving $\partial\mu(f)/\partial\alpha = 0$, we obtain $\alpha = \frac{1}{2}(5-\sqrt{13})$.

As this value of α gives an interior form of Δ'' then $f(\mathbf{x})$ is extreme when $\alpha = \frac{1}{2}(5-\sqrt{13})$ and this is the only extreme form in Δ'' (neglecting multiples).

(iii) *The cone Δ' .*

In a similar manner the form $f_3(\mathbf{x})$ of [2] can be found to be the extreme form in the cone Δ' of quaternary forms. Consider, however, the general cone Δ' [3, p. 159].

For all $n \geq 4$, the cone Δ' is the set of forms expressible in the form

$$\phi(\mathbf{x}) = \sum_{i=1}^n \rho_i x_i^2 + \sum_{\substack{i < j \\ ij \neq 12}} \rho_{ij} (x_i - x_j)^2 + \rho_{12} \omega,$$

where $\rho_i \geq 0$, $\rho_{ij} \geq 0$ for all i, j and

$$\omega = (n-2)(x_1^2 + x_2^2) + 2 \sum_{i=3}^n x_i^2 + 2x_1x_2 - 2 \sum_{i=3}^n (x_1 + x_2)x_i.$$

It can be shown rather laboriously, making use once again of the above theorem, that an interior form in this cone is extreme if and only if it is a multiple of

$$f(\mathbf{x}) = (n-1-\alpha)(x_1^2 + x_2^2) + (2 + (n-2)\beta) \sum_{i=3}^n x_i^2 + 2\alpha x_1x_2 - 2 \sum_{i=3}^n (x_1 + x_2)x_i - 2\beta \sum_{i < j} x_i x_j,$$

where α, β satisfy

$$(4.1) \quad \begin{cases} 0 < \alpha < 1, \\ \beta > 0, \\ n-2-2\alpha-\alpha\beta > 0, \end{cases}$$

$$(4.2) \quad \begin{aligned} 6(\beta+2)(1-\alpha)\{(n^2-n-1) + (1-2n)\alpha\} \\ = n(n-1)(n-2)\beta^2 + 6n(n-2)\beta + 12(n-2), \end{aligned}$$

$$(4.3) \quad \begin{aligned} 3(1-\alpha)^2\{(n-1)\beta + 2(2n-1)\} \\ = n(n-1)^2\beta^2 - n(n-1)(n-6)\beta - 6n^2 + 24n - 12. \end{aligned}$$

5. Boundary forms

All the extreme forms found so far are interior forms, so there has been no results yet which disprove the conjecture made previously that all extreme forms are interior forms.

For each Voronoï cone, the set \mathcal{F} is non-empty and closed (the form $\sum_i \phi_i(\mathbf{x})$ where $\phi_i(\mathbf{x})$ ($i = 1, \dots, N$) are all the edge forms of the cone, must belong to \mathcal{F}). Thus there will be at least one form f_0 in \mathcal{F} for which

$$(5.1) \quad \mu(f_0) = \min_{g \in \mathcal{F}} \mu(g).$$

We already know that if there is an *interior* form f_0 in \mathcal{F} satisfying (5.1) then (neglecting multiples of f_0),

- (i) f_0 is the only form in \mathcal{F} satisfying (5.1)
- (ii) f_0 is extreme and
- (iii) f_0 is the only extreme form in Δ , boundary forms included.

If, however, there is no interior form f_0 satisfying (5.1) i.e. the minimum of $\mu(f)$ on \mathcal{F} is attained on the boundary then this implies that there are no extreme interior forms, but, as yet, it implies nothing about the existence or non-existence of extreme boundary forms.

Now, it can be shown that Lemma 1 can be extended to include the case where f is a boundary form and $f + \varepsilon\varphi$ is also a boundary form for all sufficiently small ε . Using this the general theory can also be extended to include boundary forms. It is found that when there is a boundary form f_0 in \mathcal{F} satisfying (5.1) then once again this is unique and also is still the only *possible* extreme form. However, in this case it is not necessarily extreme and will only be extreme if it satisfies (5.1) for all the cones in which it lies. We can summarize all cases as follows:

For any cone Δ , there will be a unique (up to multiplication by a constant) form $f_0 \in \mathcal{F}$ satisfying (5.1) which is the only possible extreme form in Δ . f_0 is extreme if and only if it is the unique form found in the same manner for all cones containing f_0 .

The case where f_0 occurs on the boundary is definitely possible. The author has examined the general cone Δ' for higher values of n and found that there are no solutions of (4.2) and (4.3) which satisfy the conditions (4.1) for $n \geq 14$. This means that there are no interior extreme forms in Δ' for these values of n . The author has also found that for these values of n the form f_0 as defined above lies on that boundary of Δ' which is common with the principal domain. As the principal domain does not contain any boundary extreme forms then we know that f_0 is not extreme and hence Δ' does *not* contain an extreme form, boundary or interior, for $n \geq 14$.

Whether it is possible for f_0 to occur on the boundary of a cone and satisfy the above condition for extremeness, is still a matter for conjecture.

References

- [1] Barnes, E. S. and Dickson, T. J.: 'Extreme coverings of n -space by spheres', *J. Aust. Math. Soc.* 7 (1967), 115—127.
- [2] Dickson T. J.: 'The extreme coverings of 4-space by spheres', *J. Aust. Math. Soc.* 7 (1967), 490—496.
- [3] Voronoi, G.: 'Recherches sur les paralleloedres primitifs (Part 2) *J. reine angew. Math.* 136 (1909) 67—181.

The University of Western Australia