

Synthetic Stellar libraries and SSP simulations in the Gaia Era

Antonella Vallenari¹, Rosanna Sordo¹, Rosaria Tantalo², France Allard³, Ronny Blomme⁴, Jean-Claud Bouret⁵, Ines Brott⁶, Yves Fremat⁴, Christophe Martayan⁴, Yassine Damerdji⁷, Bengt Edvardsson⁸, Eric Josselin⁹, Bertrand Plez⁹, Oleg Kochukhov¹⁰, Mary Kontizas¹¹, Andreas Schweitzer¹², Jean Zorec⁴, Paraskevi Tsalmantza¹³, Ulisse Munari¹ & Tenay Saguner^{1,2}

¹INAF - Osservatorio Astronomico di Padova ²Astronomy Department, Padova University

³CRAL-ENS Lyon (France) ⁴Royal Observatory, Brussels (Belgium) ⁵CNRS Marseille (France) ⁶Sterrenkundig Instituut Utrecht (The Netherlands) ⁷Univ. De Liege (Belgium) ⁸Mid Sweden University Sundsvall (Sweden) ⁹GRAAL Montpellier (France) ¹⁰Uppsala University (Sweden) ¹¹Athens University (Greece) ¹²Hamburger Sternwarte, Hamburg, (Germany) ¹³MPI für Astronomie, Heidelberg, (Germany)

Abstract. The Gaia mission will obtain accurate positions, parallaxes and proper motions for 10^9 object all over the sky. In addition, it will collect low resolution spectroscopy in the optical range for $\sim 10^9$ objects, stars, galaxies, and QSOs. Parameters of those objects are expected to be part of the final Catalog. Complete and up-to-date libraries of synthetic stellar spectra are needed to train the algorithms to classify this huge amount of data. Here we focus on the use of the synthetic libraries of spectra calculated by the Gaia community to derive grids of Single Stellar Populations as building blocks of population synthesis models.

Keywords. Gaia; stars: synthetic spectra; population synthesis

1. Gaia Stellar and Single Stellar Population libraries

Large and up-to-date libraries of spectra are of fundamental importance for the Gaia mission (to be launched early 2012). They are used as templates to train supervised algorithms for an automatic classification of the Gaia spectra (Bailer-Jones *et al.* 2008). State-of-the-art synthetic stellar libraries are calculated by the Gaia community with an homogeneous coverage of parameter space at 0.1 nm sampling in the optical range (300-1100 nm) for stars spanning the most different types, from M to O, from A-peculiar to Emission lines to White Dwarfs (Bouret *et al.* 2008, Shulyak *et al.* 2004, Gustafsson *et al.* 2008, Alvarez *et al.* 1998, Brott *et al.* 2005, Allard *et al.* 2000, Martayan *et al.* 2008, Hindson *et al.* 2008). More detail can be found in Sordo *et al.* (2009). In addition, we make use of the Munari *et al.* (2005) synthetic stellar library, which is based on Kurucz's codes and allows to investigate the effects of $[\alpha/\text{Fe}]$ variations up to relatively high T_{eff} . This library (in its solar-scaled version) was extended in the UV, by the UVBlue library (Rodríguez-Merino *et al.* 2005) which uses the Kurucz's model atmospheres and covers the 85-470 nm spectral range. Here we focus on the use of these spectra to built synthetic libraries of single stellar populations (SSPs) to be used in the population synthesis study of star clusters and galaxies (we quote among others Vallenari *et al.* 2008, Tsalmantza *et al.* 2007, 2009). SSPs are derived as described in Tantalo (2005). We calculate three sets at changing parameters: a) *Optical-UV solar-scaled SSPs*: using Padova isochrones by Bertelli *et al.* (1994) together with the Munari *et al.* (2005) library extended by UVBlue library. The SSPs have $[\alpha/\text{Fe}] = +0.0$; b) *Optical solar-scaled SSPs*: the Padova isochrones

by Bertelli *et al.* (1994) are coupled with the Gaia libraries. The SSPs have $[\alpha/\text{Fe}] = +0.0$; c) Optical α -enhanced SSPs: Teramo isochrones by Pietrinferni *et al.* (2006) are coupled with the Munari *et al.* (2005) library. The SSPs have $[\alpha/\text{Fe}] = +0.4$. Extensive tests on the results (Lick indexes calculation, comparison with spectra of observed clusters) will be presented in Vallenari *et al.* (2009) (in preparation). The comparison of Lick indexes calculated from our SSP library (α -enhanced version) with the observational data on Galactic and M31 clusters from Trager *et al.* (1998) shows a good agreement (see Fig. 1).




Figure 1. Lick indexes $\langle \text{Fe} \rangle$ vs $[\text{Mg}/\text{Fe}]$ calculated using Munari *et al.* (2005) library, BASTI tracks, $[\text{Fe}/\text{H}] = -2.0$ (red), -1.5 (blue), -0.5 (green), $[\alpha/\text{Fe}] = +0.4$ are compared with Trager *et al.* (1998) data on globular and M31 clusters (open squares)

References

- Alvarez, R. & Plez, B. 1998, *AJ* 330, 1109
 Allard, F., Hauschildt, P. H., & Schweitzer, A. 2000, *ApJ*, 539, 366
 Bailer-Jones, C. A. L., Smith, K. W., Tiede, C., Sordo, R., & Vallenari, A. 2008, *MNRAS*, 391, 1838
 Bertelli, G., Bressan, A., Chiosi, C., Fagotto, F., & Nasi, E. 1994, *A&AS*, 106, 275
 Bouret, J.-C. *et al.* 2008, in preparation
 Brott, I. & Hauschildt, P. H. 2005, *ESA SP-576*, 565
 Castanheira, B. G. *et al.* 2006, *A&A*, 450, 331
 Gustafsson, B., Edvardsson, B., Eriksson, K. *et al.* 2008, *A&A*, 486, 951
 Hindson, L., Napiwotzki, R., Heber, U., & Lemke, M. 2008, *ASP-CS* 392, 163
 Martayan, C. *et al.* 2008, *SF2A-2008*, 499
 Munari, U., Sordo, R., Castelli, F., & Zwitter, T. 2005, *A&A* 442, 1127
 Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2006, *ApJ*, 642, 797
 Rodríguez-Merino, L. H., Chavez, M., Bertone, E., & Buzzoni, A. 2005, *ApJ*, 626, 411
 Shulyak *et al.* 2004, *A&A*, 428, 993
 Sordo, R. *et al.* 2009, *MemSAI*, 80, 103
 Tantalo, R. 2005, The Initial Mass Function 50 Years Later, 327, 235
 Trager, S. C., Worthey, G., Faber, S. M., Burstein, D., & Gonzalez, J. J. 1998 *ApJS*, 116, 1
 Tsalmantza, P. *et al.* 2007, *A&A*, 470, 761
 Tsalmantza, P. *et al.* 2009, arXiv:0907.1671
 Vallenari, A. & Sordo, R. 2008, arXiv:0812.0293