

## THE HECKE ALGEBRA ON THE COHOMOLOGY OF $\Gamma_0(p_0)$

XIANGDONG WANG

### § 1. Introduction

Let  $p_0$  be a prime,  $p_0 > 3$  and  $\Gamma_0(p_0)$ ,  $\Gamma_1(p_0)$ , as usual, the congruence subgroups of  $\Gamma = PSL_2(\mathbb{Z})$ .

$$\Gamma_0(p_0) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma \mid c \equiv 0 \pmod{p_0} \right\},$$

$$\Gamma_1(p_0) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(p_0) \mid d \equiv 1 \pmod{p_0} \right\}.$$

Denote

$$\mathcal{A} = \left\{ r = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, \gcd(a, b, c, d) = 1, \det(r) \not\equiv 0 \pmod{p_0} \right\},$$

$$\mathcal{A}_0 = \left\{ r = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{A} \mid c \equiv 0 \pmod{p_0} \right\},$$

$$\mathcal{A}_1 = \left\{ r = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{A}_0 \mid d \equiv 1 \pmod{p_0} \right\}$$

with  $\mathcal{A}_1 \subset \mathcal{A}_0 \subset \mathcal{A}$  and  $\mathcal{A}_0/\mathcal{A}_1 \cong (\mathbb{Z}/p_0)^*$ . Let  $R = \mathbb{Z}[\frac{1}{6}]$ . We consider the following  $R$ -module  $M_n = \{\sum_{v=0}^n a_v x^v y^{n-v} \mid a_v \in R\}$ . The semigroup  $\mathcal{A}$  acts on  $M_n$  via

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} x^v y^{n-v} = (ax + cy)^v (bx + dy)^{n-v}.$$

Let  $\eta: \Gamma_0(p_0)/\Gamma_1(p_0) \cong (\mathbb{Z}/p_0)^* \rightarrow R^*$  be the Legendre-symbol. We extend  $\eta$  to  $\mathcal{A}_0$  such that  $\eta$  acts trivially on  $\mathcal{A}_1$ , i.e.  $\eta$  is a character from  $\mathcal{A}_0/\mathcal{A}_1$  to  $R^*$ . Denote by  $R_\eta$  the  $R$ -module of rank 1 with a  $\mathcal{A}_0$ -operation given by  $s_0 \cdot 1 = \eta(s_0) \cdot 1$ ,  $\forall s_0 \in \mathcal{A}_0$ . Set  $M_{n,\eta} = M_n \otimes R_\eta$ . This is then a  $R[\mathcal{A}_0]$ -module. The goal of the present paper is to investigate the Hecke algebra on the cohomology group  $H^*(\Gamma_0(p_0), M_{n,\eta})$ . Let  $S_k(\Gamma_0(p_0), \eta)$ , as usual, be the

---

Received February 26, 1990.

cuspidal forms with the weight  $k$ . Then the Eichler-Shimura theorem says that the following sequence

$$0 \rightarrow S_{n+2}(\Gamma_0(p_0), \eta) \oplus \overline{S_{n+2}(\Gamma_0(p_0), \eta)} \rightarrow H^1(\Gamma_0(p_0), M_{n,\eta} \otimes \mathbb{C}) \xrightarrow{r^*} \bigoplus_s H^1(\Gamma_0(p_0)_s, M_{n,\eta} \otimes \mathbb{C}) \rightarrow 0$$

is exact, where  $s$  runs over cusps of  $\Gamma_0(p_0)$  and  $\Gamma_0(p_0)_s := \{r \in \Gamma_0(p_0) \mid r \cdot s = s\} = \langle T_s \rangle$  is an infinite cyclic group. It is well known that  $\Gamma_0(p_0)$  has two cusps  $0, \infty$ . The dimension of

$$H^1(\Gamma_0(p_0)_s, M_{n,\eta} \otimes \mathbb{C}) \cong M_{n,\eta} / (1 - T_s)M_{n,\eta}$$

is 1, which follows in particular that

$$\dim(H^1(\Gamma_0(p_0), M_{n,\eta} \otimes \mathbb{C})) = 2 \dim(S_{n+2}(\Gamma_0(p_0), \eta)) + 2$$

(cf. [Hab] p. 284). By the above identification, we see that the study of the Hecke algebra on the cuspidal forms is equivalent to that on the cohomology  $H^1(\Gamma_0(p_0), M_{n,\eta})$ , see Chap. 1 in [Hab] for more details and backgrounds. Applying the Shapiro lemma to the cohomology group of  $\Gamma_0(p_0)$  we get in Section 5 a basis for the cohomology  $H^1(\Gamma_0(p_0), M_{n,\eta})$ . Using this basis we obtain an algorithm that can be used to compute the Hecke operator  $T_l$  on the cohomology  $H^1(\Gamma_0(p_0), M_{n,\eta})$ . Finally the characteristic polynomials of  $T_2, T_3, T_5$  and  $T_7$  are given in Table 1 for small  $p_0$  and  $n$ .

### § 2. The Shapiro-Lemma

In order to determine the cohomology of  $\Gamma_0(p_0)$ , we first recall the Shapiro-Lemma. Denote by  $W_{n,\eta}$  the induced module of  $M_{n,\eta}$  on  $\Gamma$ :

$$W_{n,\eta} = \text{Ind}_{\Gamma_0(p_0)}^{\Gamma} M_{n,\eta} = \{f: \Gamma \rightarrow M_{n,\eta} \mid f(r_0 r) = r_0 \cdot f(r), \forall r_0 \in \Gamma_0(p_0)\}$$

The operation of  $\Gamma$  on  $W_{n,\eta}$  is defined by  $(a \cdot f)(r) := f(ra)$ ,  $a, r \in \Gamma$ . We extend now this operation to an operation of  $\mathcal{A}$  on  $W_{n,\eta}$ . For  $a \in \mathcal{A}$ ,  $r \in \Gamma$ , there exist always  $a' \in \mathcal{A}_0$ ,  $r' \in \Gamma$ , such that  $ra = a'r'$ . We define  $(a \cdot f)(r) := a' \cdot f(r')$ . It is obvious that this definition coincides with the above definition if  $a \in \Gamma$ . Now on the cohomology groups

$$H^1(\Gamma_0(p_0), M_{n,\eta}) \quad \text{and} \quad H^1(\Gamma, W_{n,\eta})$$

we can define the Hecke algebra (cf. [Hab] Chap. 1). By the Shapiro-Lemma (cf. [Bro] or [AS] § 1) there is a canonical isomorphism between

$$H^1(\Gamma_0(p_0), M_{n,\eta}) \cong H^1(\Gamma, W_{n,\eta})$$

as modules under the Hecke algebra.

**§ 3. The dimension of the cohomology  $H^1(\Gamma, W_{n,\gamma})$**

To get started, we consider the  $\Gamma$ -module  $W_{n,\gamma}$ . Let

$$a_i = \begin{pmatrix} 0 & -1 \\ 1 & i \end{pmatrix}, \quad i = 0, 1, \dots, p_0 - 1, \quad a_{p_0} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$\{a_i\}$  is then a set of representatives of  $\Gamma$  with respect to  $\Gamma_0(p_0)$ :

$$\Gamma = \bigcup_{i=0}^{p_0} \Gamma_0(p_0)a_i.$$

An element  $f \in W_{n,\gamma}$  is uniquely determined by the values  $f(a_0), f(a_1), \dots, f(a_{p_0})$  by using the condition  $f(r_0r) = r_0f(r)$ . The dimension of  $W_{n,\gamma}$  over  $R$  is  $(p_0 + 1) \cdot \dim(M_{n,\gamma}) = (p_0 + 1)(n + 1)$ . In other words,  $W_{n,\gamma}$  is generated by the elements  $(w_0, w_1, \dots, w_{p_0})$  with  $w_i \in M_{n,\gamma}$ .

Now we consider the cohomology  $H^1(\Gamma, W_{n,\gamma})$ . The structure of cohomology  $H^1(\Gamma, W_{n,\gamma})$  is well known (cf. [Wan] § 1):

$$H^1(\Gamma, W_{n,\gamma}) \cong W_{n,\gamma} / (W_{n,\gamma}^S + W_{n,\gamma}^Q)$$

where  $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ,  $Q = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$  and  $W_{n,\gamma}^r := \{w \in W_{n,\gamma} \mid r.w = w\}$  for  $r \in \Gamma$ .

We begin with the description of  $W_{n,\gamma}^S$ . It is easy to show that

$$\begin{cases} a_0S = a_{p_0} \\ a_iS = S_i a_j, \quad i \cdot j \equiv -1 \pmod{p_0}, \quad S_i = \begin{pmatrix} -j & -1 \\ 1 + ij & i \end{pmatrix} \in \Gamma_0(p_0) \\ a_{p_0}S = a_0 \end{cases}$$

and by the definition we obtain

$$\begin{cases} (S.f)(a_0) = f(a_{p_0}) \\ (S.f)(a_i) = S_i.f(a_j), \quad i = 1, \dots, p_0 - 1 \\ (S.f)(a_{p_0}) = f(a_0). \end{cases}$$

Therefore,  $W_{n,\gamma}^S$  has the expression:

$$\begin{aligned} W_{n,\gamma}^S &= \{f \in W_{n,\gamma} \mid f(a_0) = f(a_{p_0}), f(a_i) = S_i.f(a_j)\} \\ &= \{(w_0, \dots, w_{p_0}) \in M_{n,\gamma} \times \dots \times M_{n,\gamma} \mid w_0 = w_{p_0}, w_i = S_i.w_j\} \end{aligned}$$

Let  $T = SQ = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . One shows immediately that

$$\begin{cases} a_i T = a_{i+1}, & i = 0, 1, \dots, p_0 - 2 \\ a_{p_0-1} T = \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix} a_0 \\ a_{p_0} T = T a_{p_0} \end{cases}$$

and

$$\begin{cases} a_0 Q = T a_{p_0} \\ a_1 Q = T^{-1} a_0 \\ a_i Q = S_i a_{j+1}, & i = 2, 3, \dots, p_0 - 1 \\ a_{p_0} Q = a_1 \end{cases}$$

from which it follows

$$W_{n,\gamma}^T = \left\{ (w_0, \dots, w_{p_0}) \in M_{n,\gamma} \times \dots \times M_{n,\gamma} \mid w_0 = \dots = w_{p_0-1} \right. \\ \left. = \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix} w_0, w_{p_0} = T w_{p_0} \right\}$$

$$W_{n,\gamma}^Q = \{ (w_0, \dots, w_{p_0}) \in M_{n,\gamma} \times \dots \times M_{n,\gamma} \mid T w_{p_0} = w_0, w_1 = w_{p_0}, S_i w_{j+1} = w_i \}.$$

For the purpose of determining the dimension of  $H^1(\Gamma, W_{n,\gamma})$  we show now

3.1 LEMMA.  $W_{n,\gamma}^S \cap W_{n,\gamma}^Q = \{0\}$ .

*Proof.* Let  $f = (w_0, \dots, w_{p_0}) \in W_{n,\gamma}^S \cap W_{n,\gamma}^Q$ . It implies that  $f \in W_{n,\gamma}^T$  i.e.,

$$w_0 = w_1 = \dots = w_{p_0-1} = \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix} w_0, \text{ and } w_{p_0} \in M_{n,\gamma}^T.$$

Hence it follows that  $w_{p_0} = ax^n, w_0 = by^n$  for some  $a, b$ . For  $f \in W_{n,\gamma}^S$  we have  $w_0 = w_{p_0}$ , i.e.  $ax^n = by^n$ , which implies that  $a = b = 0$ .  $\square$

Therefore, the dimension of the cohomology  $W_{n,\gamma}$  is

$$\dim(H^1(\Gamma, W_{n,\gamma})) = \dim(W_{n,\gamma}) - \dim(W_{n,\gamma}^S) - \dim(W_{n,\gamma}^Q).$$

Now we compute the dimensions of  $W_{n,\gamma}^S$  and  $W_{n,\gamma}^Q$ .

Let  $\nu_2, \nu_3$  the number of  $\Gamma_0(p_0)$ -inequivalent elliptic points of the order 2, 3 respectively.

$$\nu_2 = 0 \text{ or } 2 \equiv p_0 + 1 \pmod{4}, \quad \nu_3 = 0 \text{ or } 2 \equiv p_0 + 1 \pmod{3}$$

It is obvious that

$$\nu_2 = 2 \Leftrightarrow p_0 \equiv 1 \pmod{4} \Leftrightarrow \eta(-1) = 1 \Leftrightarrow \text{there is a } i_0 \text{ with } i_0^2 \equiv -1 \pmod{p_0}.$$

In that case one has  $\eta(i_0) = i_0^{(p_0-1)/2} = (-1)^{(p_0-1)/4}$  and  $a_{i_0} S = S_{i_0} a_{i_0}$ . Furthermore it is easy to show that

$$\begin{aligned} \nu_3 = 2 &\Leftrightarrow p \equiv 1 \pmod 3 \Leftrightarrow 6 \mid p_0 - 1 \Leftrightarrow \text{there is a } i_0 \text{ of order 6 in } (\mathbb{Z}/p_0)^* \\ &\Leftrightarrow i_0^3 \equiv -1 \pmod{p_0} \Leftrightarrow i_0(i_0 - 1) \equiv -1 \pmod{p_0}. \end{aligned}$$

It follows that  $a_{i_0}Q = S_{i_0}a_{i_0}$ . Since  $(i_0 - 1)^2 \equiv -i_0$  one has  $\eta(i_0) = \eta(-1)\eta(i_0 - 1)^2 = \eta(-1) = (-1)^{(p_0-1)/2}$ .

3.2 LEMMA.

$$\dim(W_{n,\eta}^S) = 2\left[\frac{p_0 + 1}{4}\right](n + 1) + 2d_s$$

$$\dim(W_{n,\eta}^Q) = \left[\frac{p_0 + 1}{3}\right](n + 1) + 2d_q$$

where

$$d_s = \begin{cases} 0 & p_0 \equiv 3 \pmod 4 \\ 2\left[\frac{n}{4}\right] + 1 & p_0 \equiv 1 \pmod 8, \\ 2\left[\frac{n + 2}{4}\right] & p_0 \equiv 5 \pmod 8 \end{cases}, \quad d_q = \begin{cases} 0 & p_0 \equiv 2 \pmod 3 \\ 2\left[\frac{n}{6}\right] + 1 & p_0 \equiv 1 \pmod{12}. \\ 2\left[\frac{n + 3}{6}\right] & p_0 \equiv 7 \pmod{12} \end{cases}$$

In particular,

$$\begin{aligned} \dim(H^1(\Gamma, W_{n,\eta})) &= \left(p_0 + 1 - 2\left[\frac{p_0 + 1}{4}\right] - \left[\frac{p_0 + 1}{3}\right]\right)(n + 1) \\ &\quad - 2d_s - 2d_q \end{aligned}$$

$$\begin{aligned} \dim(S_{n+2}(\Gamma_0(p_0), \eta)) &= \frac{1}{2}\left(p_0 + 1 - 2\left[\frac{p_0 + 1}{4}\right] - \left[\frac{p_0 + 1}{3}\right]\right)(n + 1) \\ &\quad - d_s - d_q - 1. \end{aligned}$$

*Proof.* For  $f = (w_0, \dots, w_{p_0}) \in W_{n,\eta}^S$  we have  $w_i = S_i w_j$  and  $S_j = S_i^{-1}$ . If  $j \neq i$ , then  $w_j$  is uniquely determined by  $w_i$ . The number of such pair  $(i, j)$  is  $2[(p_0 + 1)/4]$ . If  $j = i$ , that means  $p_0 \equiv 1 \pmod 4$ , one has  $w \in \text{Ker}(1 - S_i)$ . We calculate the dimension of  $\text{Ker}(1 - S_i)$ . Let  $m \otimes 1 \in M_{n,\eta}$ , then  $S_i(m \otimes 1) = \eta(i)(S_i m \otimes 1)$ . For  $S_i = \begin{pmatrix} -i & -1 \\ 1 + i^2 & i \end{pmatrix}$  there is a regular matrix  $P$  with  $S_i = PSP^{-1}$ . It follows that

$$d_s = \dim(M_{n,\eta}^{S_i}) = \dim(\text{Ker}(1 - S_i)) = \dim(\text{Ker}(1 - \eta(i)S)).$$

For  $p_0 \equiv 1 \pmod 8$  one has  $\eta(i) = (-1)^{(p_0-1)/4} = 1$ . The dimension of  $\text{Ker}(1 - S)$  can be easily determined,  $\dim \text{Ker}(1 - S) = 2[n/4] + 1$ . Since there are two  $i$  with  $i^2 \equiv -1$ , dimension of  $W_{n,\eta}^S$  has the expression:

$$\dim(W_{n,\eta}^S) = 2\left[\frac{p_0 + 1}{4}\right](n + 1) + 4\left[\frac{n}{4}\right] + 2$$

The other cases can be proved in the same manner. □

**§ 4. The dimension of  $H^i(\Gamma, W_{n,\eta})_{\pm}$**

Let  $\Gamma_{\infty} = \langle T \rangle$  be the stabilizer of the cusp  $\infty$  in  $\Gamma$ . We have an exact sequence:

$$0 \rightarrow H^0(\Gamma_{\infty}, W_{n,\eta}) \rightarrow H_c^1(\Gamma, W_{n,\eta}) \rightarrow H^1(\Gamma, W_{n,\eta}) \rightarrow H^1(\Gamma_{\infty}, W_{n,\eta}) \rightarrow \dots$$

where  $H_c^i(\cdot, \cdot)$  is the cohomology with the compact support, referring to [Hab] Chap. 1 for details and backgrounds. It has been shown in [Wan] § 1 that the cohomology

$$H^1(\Gamma_{\infty}, W_{n,\eta}) \cong W_{n,\eta}/(1 - T)W_{n,\eta}.$$

4.1 LEMMA.

$$H^1(\Gamma_{\infty}, W_{n,\eta} \otimes \mathbb{Q}) \cong \mathbb{Q}\phi_0 + \mathbb{Q}\phi_{\infty}$$

where  $\phi_0(T) = (x^n, 0, \dots, 0) \in W_{n,\eta}$ ,  $\phi_{\infty}(T) = (0, \dots, 0, y^n) \in W_{n,\eta}$ .

*Proof.* For each  $w = (w_0, \dots, w_{p_0}) \in W_{n,\eta}$ , we consider the equation

$$(*) \quad w = a(x^n, 0, \dots, 0) + b(0, \dots, 0, y^n) + (T - 1)v$$

with  $v = (v_0, \dots, v_{p_0}) \in W_{n,\eta}$ , which means:

$$\begin{aligned} w_0 &= ax^n + v_1 - v_0 \\ w_i &= v_{i+1} - v_i, \quad 0 < i < p_0 - 1 \\ w_{p_0-1} &= \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix} v_0 - v_{p_0-1} \\ w_{p_0} &= by^n + (T - 1)v_{p_0}, \end{aligned}$$

it follows that

$$\left(1 - \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix}\right)v_0 = ax^n - \sum_{j=0}^{p_0-1} w_j.$$

We take  $a$  as the coefficient of  $x^n$  in  $\sum_{j=0}^{p_0-1} w_j$  and  $b$  as the coefficient of  $y^n$  in  $w_{p_0}$ . The equations

$$\begin{aligned} \left(1 - \begin{pmatrix} 1 & 0 \\ -p_0 & 1 \end{pmatrix}\right)v_0 &= c_0y^n + c_1xy^{n-1} + \dots + c_{n-1}x^{n-1}y \\ (1 - T)v_{p_0} &= d_1xy^{n-1} + d_2x^2y^{n-2} + \dots + d_nx^n \end{aligned}$$

are always solvable in  $M_{n,\eta} \otimes \mathbb{Q}$  for any  $c_0, \dots, c_{n-1}, d_1, \dots, d_n \in \mathbb{Q}$ . Therefore the equation (\*) is solvable in  $M_{n,\eta} \otimes \mathbb{Q}$ .  $\square$

Let  $\varepsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ . We define for a cocycle  $\phi \in Z^1(\Gamma, W_{n,\eta})$

$$(\varepsilon\phi)(r) := \varepsilon\phi(\varepsilon^{-1}r\varepsilon) \quad \forall r \in \Gamma.$$

It induces an automorphism of the order 2 on the cohomologies (cf. [Wan] §1). Hence we obtain two exact sequences:

$$0 \rightarrow H^0(\Gamma_\infty, W_{n,\eta})_+ \rightarrow H^1_c(\Gamma, W_{n,\eta})_+ \rightarrow H^1(\Gamma, W_{n,\eta})_+ \xrightarrow{r^*} H^1(\Gamma_\infty, W_{n,\eta})_+ \rightarrow \dots$$

$$0 \rightarrow H^0(\Gamma_\infty, W_{n,\eta})_- \rightarrow H^1_c(\Gamma, W_{n,\eta})_- \rightarrow H^1(\Gamma, W_{n,\eta})_- \xrightarrow{r^*} H^1(\Gamma_\infty, W_{n,\eta})_- \rightarrow \dots$$

where  $H^1(\Gamma, W_{n,\eta})_\pm := \{\phi \in H^1(\Gamma, W_{n,\eta}) \mid \varepsilon.\phi = \pm\phi\}$ . Since

$$\begin{cases} a_0\varepsilon = \varepsilon a_0 \\ a_i\varepsilon = E.a_{p_0-i}, \quad i = 1, 2, \dots, p_0 - 1 \\ a_{p_0}\varepsilon = \varepsilon a_{p_0} \end{cases}$$

where  $E := \begin{pmatrix} -1 & 0 \\ p_0 & 1 \end{pmatrix}$ . The operation of  $\varepsilon$  on  $W_{n,\eta}$  is

$$\begin{cases} (\varepsilon u)(a_0) = \varepsilon.u(a_0) \\ (\varepsilon u)(a_i) = E.u(a_{p_0-i}) \\ (\varepsilon u)(a_{p_0}) = \varepsilon.u(a_{p_0}). \end{cases}$$

In particular, it follows that

$$\varepsilon.\phi_\infty(T) = \phi_\infty(T), \quad \varepsilon.\phi_0(T) = (-1)^n\phi_0(T).$$

4.2 LEMMA.

- a.  $\phi_\infty \in H^1(\Gamma_\infty, W_{n,\eta})_-$
- b.  $\phi_0 \in H^1(\Gamma_\infty, W_{n,\eta})_-$  for  $n$  even;  $\phi_0 \in H^1(\Gamma, W_{n,\eta})_+$  for  $n$  odd.

*Proof.*

$$\begin{aligned} (\varepsilon\phi_\infty)(T) &= \varepsilon.\phi_\infty(\varepsilon^{-1}T\varepsilon) = \varepsilon.\phi_\infty(T^{-1}) = -\varepsilon T^{-1}\phi_\infty(T) = -T\varepsilon.\phi_\infty(T) \\ &= -\varepsilon.\phi_\infty(T) + (1 - T)\varepsilon.\phi_\infty(T) \sim -\varepsilon.\phi_\infty(T) = -\phi_\infty(T). \end{aligned}$$

It means that  $\varepsilon.\phi_\infty = -\phi_\infty$ . (b) can be proved in the same way.  $\square$

By applying the Eichler-Shimura isomorphism, together with the observation above, we obtain

4.3 COROLLARY.

a. For  $n$  even we have

$$\dim(H^1(\Gamma, W_{n,\eta})_-) = \frac{1}{2} \dim(H^1(\Gamma, W_{n,\eta})) + 1$$

$$\dim(H^1(\Gamma, W_{n,\eta})_+) = \frac{1}{2} \dim(H^1(\Gamma, W_{n,\eta})) - 1.$$

b. For  $n$  odd we have

$$\dim(H^1(\Gamma, W_{n,\eta})_-) = \frac{1}{2} \dim(H^1(\Gamma, W_{n,\eta}))$$

$$\dim(H^1(\Gamma, W_{n,\eta})_+) = \frac{1}{2} \dim(H^1(\Gamma, W_{n,\eta})).$$

§ 5. The basis of  $H^1(\Gamma, W_{n,\eta})$

It is well known that

$$H^1(\Gamma, W_{n,\eta}) \cong W_{n,\eta}/(W_{n,\eta}^S + W_{n,\eta}^Q).$$

Our goal in this section is to choose a subset  $V$  of  $W_{n,\eta}$  such that  $W_{n,\eta} = W_{n,\eta}^S \oplus W_{n,\eta}^Q \oplus V$ . Since the group  $\Gamma$  is generated by  $S, Q$  with the relations  $S^2 = 1, Q^3 = 1$  (cf. [Ser]), the cohomology

$$H^1(\Gamma, W_{n,\eta}) = \frac{\{(\phi(S), \phi(Q)) \mid \phi(S) \in (1 - S)W_{n,\eta}, \phi(Q) \in (1 - Q)W_{n,\eta}\}}{\{((1 - S)u, (1 - Q)u) \mid u \in W_{n,\eta}\}}$$

$$\cong \frac{\{\phi(Q) \mid \phi(S) = 0, \phi(Q) \in (1 - Q)W_{n,\eta}\}}{\{(1 - Q)u \mid u \in W_{n,\eta}^S\}}$$

$$\cong \{(1 - Q)v \mid v \in V\},$$

i.e, every class  $\phi \in H^1(\Gamma, W_{n,\eta})$  has the form

$$\begin{cases} \phi(S) = 0 \\ \phi(Q) = (1 - Q)u, \quad u \in V. \end{cases}$$

Defining by  $\alpha_i$  (resp.  $\beta_i$ ) the permutation of  $\{0, 1, \dots, p_0\}$  induced by the operation of  $S$  (resp.  $Q$ ) on  $\{a_0, a_1, \dots, a_{p_0}\}$ . We have (cf. § 3)

$$\alpha_i \cdot i \equiv -1 \pmod{p_0}, \quad 0 < i < p_0$$

$$\beta_i = \alpha_i + 1, \quad 1 < i < p_0$$

5.1 DEFINITION. For  $i, j, k \in \{1, 2, \dots, p_0 - 1\}$

a. A pair  $(i, j)$  is called a  $\alpha$ -pair if  $j = \alpha_i, i = \alpha_j$ , or equivalently,  $i \cdot j \equiv -1 \pmod{p_0}$ ;



b. A triple  $(i, j, k)$  is called a  $\beta$ -triple if  $j = \beta_i, k = \beta_j, i = \beta_k$ , or equivalently,  $i \cdot j \cdot k \equiv -1 \pmod{p_0}$ ;

c. Let  $B$  be a subset of  $\{1, 2, \dots, p_0 - 1\}$ . We denote by  $\langle B \rangle$  the subset of  $\{1, 2, \dots, p_0 - 1\}$  determined by the following conditions:

- i.  $B \subset \langle B \rangle$ ;
  - ii. if  $(i, j)$  is an  $\alpha$ -pair and  $j \in \langle B \rangle$  then  $i \in \langle B \rangle$ ;
  - iii. if  $(i, j, k)$  is a  $\beta$ -triple and  $j, k \in \langle B \rangle$  then  $i \in \langle B \rangle$ ;
- d. A subset  $B$  of  $\{1, 2, \dots, p_0 - 1\}$  is called a basis set if it satisfies:
- i.  $\langle B \rangle = \{1, 2, \dots, p_0 - 1\}$ ;
  - ii.  $\forall i \in B, \langle B \setminus \{i\} \rangle \neq \{1, 2, \dots, p_0 - 1\}$ .

It follows immediately from the definition that the number of the  $\alpha$ -pair is  $2[(p_0 + 1)/4] - 1$  and the number of the  $\beta$ -triple is  $[(p_0 + 1)/3] - 1$ . Therefore the number of the elements in  $B$  is

$$\begin{aligned} \#B &= (p_0 - 1) - \left(2\left[\frac{p_0 + 1}{4}\right] - 1\right) - \left(\left[\frac{p_0 + 1}{3}\right] - 1\right) \\ &= p_0 + 1 - 2\left[\frac{p_0 + 1}{4}\right] - \left[\frac{p_0 + 1}{3}\right]. \end{aligned}$$


We define inductively two series of subsets of  $\{1, 2, \dots, p_0 - 1\}$ .

$$\begin{aligned} L_1 &= \{1\} \\ M_r &= \{\alpha_i \mid i \in L_r\} \setminus L_r, \quad r > 0 \\ L_{r+1} &= \{j = \beta_i, \beta_j \mid i \in M_r\} \setminus M_r \end{aligned}$$

5.2 EXAMPLE.  $p_0 = 13$ . In that case  $\nu_2 = 2, \nu_3 = 2$ . The permutations of  $\{a_0, a_1, \dots, a_{p_0}\}$  induced by the operation of  $S$  and  $Q$  are:

$i$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
$S$	13	12	6	4	3	5	2	11	8	10	9	7	1	0
$Q$	13	0	7	5	4	6	3	12	9	11	10	8	2	1

The sets  $L_r$  and  $M_r$  can be described by the diagram:




5.3 LEMMA.

- a.  $\{1, 2, \dots, p_0 - 1\} = \bigcup_{r=1}^N (L_r \cup M_r)$  for some  $N < p_0$ ;
- b. For each  $i \in L_r$ , there exists a  $j \in L_r$  with  $i \cdot j \equiv 1 \pmod{p_0}$ ;
- c. For each  $i \in M_r$ , there exists a  $j \in M_r$  with  $i \cdot j \equiv 1 \pmod{p_0}$ .

*Proof.* a. Assume that  $a$  is the smallest element in  $\{1, 2, \dots, p_0 - 1\}$  with the property  $a \notin \bigcup_{r=1}^\infty (L_r \cup M_r)$ . Let  $a = \beta_b$  for some  $b \in \{1, 2, \dots, p_0 - 1\}$ . Then  $a = \beta_b = \alpha_b + 1$  and  $\alpha_b < a$ . By the assumption it implies  $\alpha_b \in \bigcup_{r=1}^\infty (L_r \cup M_r)$ , which follow that  $b \in \bigcup_{r=1}^\infty (L_r \cup M_r)$  and  $a \in \bigcup_{r=1}^\infty (L_r \cup M_r)$  by the definition of  $\langle B \rangle$ . It contradicts the assumption.

b. We prove the assertion by the induction. The assertion for  $r = 1$  is obvious. Let  $a$  be an element in  $L_{r+1}$ , then there is an element  $c \in M_r$  such that  $a = \beta_c$  or  $c = \beta_a$ . We treat only the case  $a = \beta_c$ . Let  $b = \beta_a \in L_{r+1}$  and  $d = \alpha_c \in L_r$ . By the induction assumption there is a  $e \in L_r$  with  $d \cdot e \equiv 1 \pmod{p_0}$ . Let  $f = \alpha_e$ , we see immediately that  $f \cdot c \equiv 1 \pmod{p_0}$ . Let  $g = \beta_f$ ,  $h = \beta_g \in L_{r+1}$ , we look at the following diagram:



and assert that  $a \cdot h \equiv 1 \pmod{p_0}$ . Indeed,

$$\begin{aligned}
 a &= \beta_c = \alpha_c + 1 = d + 1 \equiv (d + 1) \cdot (-ef) \equiv (e + 1) \cdot (-f) \\
 &\equiv 1 - f = 1 - \beta_h = -\alpha_h
 \end{aligned}$$

i.e.,  $a \cdot h \equiv -\alpha_h \cdot h \equiv 1$ .

- c. It follows immediately from (b).

5.4 LEMMA. *There is a basis set  $B$  with the property: if  $a \in B$  then  $p_0 - a \in B$ .*


The proof of the lemma presents in fact an algorithm to compute the basis set  $B$ .

*Proof.* First note that  $\langle L_r \rangle \subset \langle M_r \rangle \subset \langle L_{r+1} \rangle$ .

*Case 1:* If  $a, \alpha_a \in L_r$  and  $a \notin \langle B \rangle$ , there is an elements  $b \in L_r$  with  $ab \equiv 1$ , which yields  $\alpha_a \cdot \alpha_b \equiv 1$ . Since  $(a + \alpha_b)b = ab + \alpha_b \cdot b \equiv 1 +$

$= 0$ , one has  $a + \alpha_b = p_0$  and  $\{a, b, \alpha_a, \alpha_b\} \subset \langle \{a, \alpha_b\} \rangle$ . Hence we add  $a, \alpha_b$  to  $B$ .

*Case 2.*  $(a, b, c)$  is a  $\beta$ -triple,  $a, b \in M_r, c \in L_{r+1}$  and  $a, b \notin B$ . For  $a, b \in M_r$  there are  $d, e \in M_r$  with  $ad \equiv 1, be \equiv 1$ . We consider the following diagram:




one verifies trivially that  $a + d_1 = p_0$  and

$$\{a, b, c, d, e, f, a_1, b_1, d_1, e_1\} \subset \langle \{a, d_1, c, f\} \rangle.$$

Therefore we add  $a, d_1$  to  $B$ .

*Case 3:*  $(a, b, c)$  is a  $\beta$ -triple,  $a, b, c \in M_r$  and  $a, b, c \notin \langle B \rangle$ . There are  $d, e, f \in M_r$  with  $ad \equiv 1, be \equiv 1, cf \equiv 1$ . We consider the following diagram:



It is obvious, that  $a + d_1 = p_0, b + e_1 = p_0$  and

$$\{a, b, c, d, e, f, a_1, b_1, c_1, d_1, e_1, f_1\} \subset \langle \{a, b, d_1, e_1\} \rangle.$$

We add thus  $a, b, d_1, e_1$  to  $B$ .

In such a way we obtain a basis set  $B$ . □

In the example 5.2 we can take the basis set  $B = \{5, 8, 4, 9\}$ .

We study now the cohomology  $H^1(\Gamma, W_{n,\eta}) = W_{n,\eta}/(W_{n,\eta}^S + W_{n,\eta}^Q)$ . Let  $B$  be a basis set. Then each element  $(0, w_1, \dots, w_{p_0-1}, 0) \in W_{n,\eta}$  is congruent mod  $W_{n,\eta}^S + W_{n,\eta}^Q$  to an element  $g = (v_0, \dots, v_{p_0}) \in W_{n,\eta}$  with  $v_i = 0$  for  $i \in B$ .

If  $\nu_2 = 2$ , there is a  $i_0 \in B$  such that  $i_0^2 \equiv -1$ . If  $w_{i_0} \in \text{Ker}(1 - S_{i_0}) = M_{n,\eta}^{S_{i_0}}$ , then  $(0, \dots, 0, w_{i_0}, 0, \dots, 0) \in W_{n,\eta}^S$ . Therefore

$$\begin{aligned} \{(0, \dots, w_{i_0}, \dots, 0) \mid w_{i_0} \in M_{n,\eta}\} / (W_{n,\eta}^S + W_{n,\eta}^Q) \\ \cong \{(0, \dots, v_{i_0}, \dots, 0) \mid v_{i_0} \in M_{n,\eta} / M_{n,\eta}^{S_{i_0}}\}. \end{aligned}$$

Similarly, if  $\nu_3 = 2$  and  $i_0 \in B$ ,  $i_0^3 \equiv -1$ , then

$$\begin{aligned} \{(0, \dots, w_{i_0}, \dots, 0) \mid w_{i_0} \in M_{n,\eta}\} / (W_{n,\eta}^S + W_{n,\eta}^Q) \\ \cong \{(0, \dots, v_{i_0}, \dots, 0) \mid v_{i_0} \in M_{n,\eta} / M_{n,\eta}^{S_{i_0}}\}. \end{aligned}$$

Now we consider the index 0,  $p_0$ . Since

$$(0, \dots, 0, w_{p_0}) = (-w_{p_0}, 0, \dots, 0) \text{ mod } W_{n,\eta}^S + W_{n,\eta}^Q,$$

we need only to consider only the index 0. Let

$$(w_0, 0, \dots, 0) = \underbrace{(a, 0, \dots, 0, a)}_{\in W_{n,\eta}^S} + \underbrace{(Tb, b, 0, \dots, 0, b)}_{\in W_{n,\eta}^Q} + (0, c, 0, \dots, 0)$$

for some  $a, b, c$ , then  $b = -a, c = a, (1 - T)a = w_0$ . The equation  $(1 - T)a = w_0$  can be solved only for  $w_0 = c_1xy^{n-1} + c_2x^2y^{n-2} + \dots + c_nx^n$ . Therefore the element  $(y^n, 0, \dots, 0)$  is linear independent to

$$\begin{aligned} \{(0, \dots, 0, v_i, 0, \dots, 0) \mid i \in B, v_i \in M_{n,\eta} \text{ or } M_{n,\eta} / M_{n,\eta}^{S_{i_0}} \\ \text{if } i^2 \equiv -1 \text{ or } i^3 \equiv -1\} \text{ mod } W_{n,\eta}^S + W_{n,\eta}^Q. \end{aligned}$$

On the other hand,

$$(0, x^n, 0, \dots, 0) = (-x^n, 0, \dots, 0, -x^n) + (Tx^n, x^n, 0, \dots, 0, x^n) \in W_{n,\eta}^S + W_{n,\eta}^Q$$

and  $(0, x^n, 0, \dots, 0)$  can be represented by the elements of

$$\{(0, \dots, 0, v_i, 0, \dots, 0) \mid i \in B, v_i \in M_{n,\eta} \text{ or } M_{n,\eta} / M_{n,\eta}^{S_{i_0}} \text{ if } i^2 \equiv -1 \text{ or } i^3 \equiv -1\},$$

which implies that the elements of

$$\{(0, \dots, 0, v_i, 0, \dots, 0) \mid i \in B, v_i \in M_{n,\eta} \text{ or } M_{n,\eta} / M_{n,\eta}^{S_{i_0}} \text{ if } i^2 \equiv -1 \text{ or } i^3 \equiv -1\}$$

are linear dependent mod  $W_{n,\eta}^S + W_{n,\eta}^Q$ . A basis of  $H^1(\Gamma, W_{n,\eta})$  is then  $(y^n, 0, \dots, 0)$  and

$$\{(0, \dots, 0, v_i, 0, \dots, 0) \mid i \in B, v_i \in M_{n,\eta} \text{ or } M_{n,\eta}/M_{n,\eta}^{S_i} \\ \text{if } i^2 \equiv -1 \text{ or } i^3 \equiv -1 \pmod{\sim},$$

where the relation  $\sim$  is given by the equation

$$(0, x^n, 0, \dots, 0) \equiv 0 \pmod{W_{n,\eta}^S + W_{n,\eta}^Q}$$

**§ 6. The basis of  $H^1(\Gamma, W_{n,\eta})_{\pm}$**

We shall first deal with the operation of  $\varepsilon$  on  $H^1(\Gamma, W_{n,\eta})$ . From the definition in § 4 we have for a class  $\phi \in H^1(\Gamma, W_{n,\eta})$ ,  $\phi(S) = 0$ ,  $\phi(Q) = (1 - Q)u$ ,

$$\begin{aligned} (\varepsilon\phi)(S) &= \varepsilon.\phi(\varepsilon S\varepsilon) = \varepsilon.\phi(S^{-1}) = 0 \\ (\varepsilon\phi)(Q) &= \varepsilon.\phi(\varepsilon Q\varepsilon) = \varepsilon.\phi(SQ^{-1}S) = -\varepsilon SQ^{-1}\phi(Q) = -\varepsilon SQ^{-1}(1 - Q)u \\ &= (1 - Q)\varepsilon Su = (1 - Q)S\varepsilon u. \end{aligned}$$

If  $\phi \in H^1(\Gamma, W_{n,\eta})_-$ , i.e.  $\varepsilon\phi + \phi = 0$ , it follows that  $S\varepsilon u + u \in W_{n,\eta}^S + W_{n,\eta}^Q$ . Since  $S\varepsilon u + u = (S + 1)\varepsilon u + u - \varepsilon u$  and  $(S + 1)\varepsilon u \in W_{n,\eta}^S$ , we obtain

$$\phi \in H^1(\Gamma, W_{n,\eta})_- \iff u - \varepsilon u \in W_{n,\eta}^S + W_{n,\eta}^Q.$$

Similarly,

$$\phi \in H^1(\Gamma, W_{n,\eta})_+ \iff u + \varepsilon u \in W_{n,\eta}^S + W_{n,\eta}^Q.$$

In order to determine a basis of  $H^1(\Gamma, W_{n,\eta})_-$  we consider the vector space

$$U := \{u = (u_0, \dots, u_{p_0}) \in W_{n,\eta} \mid u - \varepsilon.u = 0\}.$$

$U$  has a basis consisting of the elements  $(u_0, \dots, u_{p_0})$  which satisfy one of the following conditions (cf. § 4):

1.  $\begin{cases} u_0 = x^j y^{n-j}, & j \text{ even} \\ u_i = 0, & i > 0 \end{cases}$
2.  $\begin{cases} u_{p_0} = x^j y^{n-j}, & j \text{ even} \\ u_i = 0, & i < p_0 \end{cases}$
3.  $\begin{cases} u_i = x^j y^{n-j} \\ u_{p_0-i} = E.u_i \\ u_k = 0, & k \neq i, p_0 - i. \end{cases}$

In particular, the classes  $\phi \in H^1(\Gamma, W_{n,\eta})$ ,  $\phi(S) = 0$ ,  $\phi(Q) = (1 - Q)u$  are classes in  $H^1(\Gamma, W_{n,\eta})_-$  for  $n$  even, where  $u = (u_0, \dots, u_{p_0}) \in W_{n,\eta}$  with

$$1. \begin{cases} u_0 = y^n \\ u_j = 0, \quad j > 0 \end{cases}$$

or

$$2. \begin{cases} u_i \in W_{n,\eta} \text{ or } M_{n,\eta}/M_{n,\eta}^{S_i}, \quad i \in B, i < p_0/2 \\ u_{p_0-i} = E.u_i \\ u_j = 0, \quad j \neq i, p_0 - i. \end{cases}$$

The number of the above classes is

$$1 + \frac{1}{2} *B \dim (M_{n,\eta}) - d_s - d_Q = \dim (H^1(\Gamma, W_{n,\eta})_-).$$

By using the fact that the basis set  $B$  consists of the pair  $(i_1, i_2)$  with  $i_1 + i_2 = p_0$  we find that the above classes generate the cohomology  $H^1(\Gamma, W_{n,\eta})_-$ . Therefore this set of classes is a basis of  $H^1(\Gamma, W_{n,\eta})_-$  for  $n$  even.

Similarly, we choose a basis of  $H^1(\Gamma, W_{n,\eta})_+$  for  $n$  odd:  $\begin{cases} \phi(S) = 0 \\ \phi(Q) = (1 - Q)u \end{cases}$  with

$$\begin{cases} u_i \in M_{n,\eta} \text{ or } M_{n,\eta}/M_{n,\eta}^{S_i}, \quad i \in B, i < p_0/2 \\ u_{p_0-i} = -E.u_i \\ u_j = 0, \quad j \neq i, p_0 - i. \end{cases}$$

6.1. *Remark.* In general it is very difficult to determine the basis of  $H^1(\Gamma, W_{n,\eta})_+$  for  $n$  even, because the dimension of  $H^1(\Gamma, W_{n,\eta})_+$  is  $\frac{1}{2} \dim (H^1(\Gamma, W_{n,\eta})) - 1$ , and the dimension of the vector space generated by the set

$$\begin{cases} u_i \in M_{n,\eta} \text{ or } M_{n,\eta}/M_{n,\eta}^{S_i}, \quad i \in B, i < p_0/2 \\ u_{p_0-i} = E.u_i \end{cases}$$

is  $\frac{1}{2} \dim (H^1(\Gamma, W_{n,\eta}))$ . It implies that there is a relation between the above elements. The case  $H^1(\Gamma, W_{n,\eta})_-$  for  $n$  odd is similar.

We are now interested in the boundary map  $r^*$  on the basis.

6.2 LEMMA. For a class  $\phi \in H^1(\Gamma, W_{n,\eta})$  with  $\phi(S) = 0, \phi(Q) = (1 - Q)u,$

- a. if  $u = (y^n, 0, \dots, 0)$  then  $r^*\phi = \phi_\infty;$
- b. if  $u = (0, \dots, 0, u_i, 0, \dots, 0), 0 < i < p_0$  then  $r^*\phi = a\phi_0$  for some  $a.$

*Proof.* a.

$$\begin{aligned} (r^*\phi)(T) &= \phi(T) = S\phi(Q) = S(1 - Q)u = (S - T)u \\ &= (S - 1)u + (1 - T)u \sim (S - 1)u = (-y^n, 0, \dots, 0, y^n). \end{aligned}$$

The solution of the equation (\*) in § 4.1 is  $a = 0, b = 1$ , i.e.,  $r^*\phi = \phi_\infty$ .

b.  $r^*\phi(T) \sim (S - 1)u = (0, \dots, -u_i, 0, \dots, S_i^{-1}u_i, 0, \dots, 0)$ . It is obvious that  $b = 0$  (cf. the proof of § 4.1). Hence  $r^*\phi = a\phi_0$  for some  $a$ .  $\square$

**§ 7. The Hecke operator  $T_l$  on  $H^1(\Gamma, W_{n,\gamma})$**

To get started, we recall the definition of the Hecke operator  $T_l$  on  $H^1(\Gamma, W_{n,\gamma})$ , where  $l$  is a prime,  $l \neq p_0$ . Let

$$b_i = \begin{pmatrix} 1 & i \\ 0 & l \end{pmatrix}, \quad i = 0, 1, \dots, l - 1 \quad \text{and} \quad b_l = \begin{pmatrix} l & 0 \\ 0 & 1 \end{pmatrix},$$

they are a complete set of representatives of  $\Gamma \begin{pmatrix} l & 0 \\ 0 & 1 \end{pmatrix} \Gamma$  with respect to  $\Gamma$ :

$$\Gamma \begin{pmatrix} l & 0 \\ 0 & 1 \end{pmatrix} \Gamma = \bigcup_{i=0}^l \Gamma b_i$$

For each  $r \in \Gamma$  there is a  $s_i \in \Gamma$  such that  $b_i r = s_i b_j$  for some  $j$ . Define for a cocycle  $f \in Z^1(\Gamma, W_{n,\gamma})$

$$(T_l f)(r) := \sum_{i=0}^l b'_i f(s_i)$$

where  $b'_i := \det(b_i) b_i^{-1}$ .

All this is discussed in more detail in [AS] § 1 or [Wan] § 1.2.

7.1. EXAMPLE.  $l = 2, p_0 = 5, n = 4$

For  $l = 2$  the representatives are

$$b_0 = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \quad b_1 = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}, \quad b_2 = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}.$$

A simple calculation shows that

$$\begin{cases} b_0 S = S b_2 \\ b_1 S = S Q^{-1} S Q S b_1 \\ b_2 S = S b_0 \end{cases} \quad \begin{cases} b_0 T = b_1 \\ b_1 T = T b_0 \\ b_2 T = T^2 b_2 \end{cases} \quad \begin{cases} b_0 Q = Q S Q b_2 \\ b_1 Q = S Q^{-1} S Q^{-1} b_0 \\ b_2 Q = S b_1 \end{cases}$$

By the definition we get for a class  $\phi \in H^1(\Gamma, W_{n,\gamma})$

$$(T_2 \phi)(S) = b'_0 \phi(S) + b'_1 \phi(S Q^{-1} S Q S) + b'_2 \phi(S) = (S - 1) b'_1 S Q^{-1} \phi(Q)$$

$$(T_2 \phi)(Q) = b'_0 \phi(Q S Q) + b'_1 \phi(S Q^{-1} S Q^{-1}) + b'_2 \phi(S) = (1 - Q)(b'_0 + b'_0 S Q) \phi(Q).$$

Hence the cocycle  $T_2 \phi$  is cohomology to

$$T_2\phi \sim \begin{cases} (T_2\phi)(S) = 0 \\ (T_2\phi)(Q) = (1 - Q)(b'_0 + b'_0SQ + b'_1SQ^{-1})\phi(Q). \end{cases}$$

It is easy to see that

$$\begin{aligned} (1 - Q)(b'_0 + b'_0QS + b'_1SQ^{-1}) &= (1 - Q)(b'_0 + (Q + Q^2)b'_2Q^{-1}) \\ &= (1 - Q)(b'_0 - b'_2Q^{-1}), \end{aligned}$$

we obtain then

$$(T_2\phi)(Q) = (1 - Q)(b'_0 - b'_2Q^{-1})\phi(Q).$$

For  $p_0 = 5$  we choose a basis set  $B = \{2, 3\}$ . The basis of  $H^1(\Gamma, W_{n,\eta})_-$  is then  $(y^n, 0, 0, 0, 0, 0)$  and  $(0, 0, w_2, Ew_2, 0, 0)$   $w_2 \in M_{n,\eta}/M_{n,\eta}^{S_2}$ . For  $n = 5$  the numerical computation shows that  $M_{n,\eta}/M_{n,\eta}^{S_2} = Rv_1 + Rv_2 + Rv_3$  with

$$\begin{aligned} v_1 &= x^4 - 8x^3y + 24x^2y^2 - 32xy^3 + 16y^4 \\ v_2 &= x^3y - 6x^2y^2 + 12xy^3 - 8y^4 \\ v_3 &= x^2y^2 - 4xy^3 + 4y^4. \end{aligned}$$

Let  $v_0 = y^4$ , then the basis of  $H^1(\Gamma, W_{n,\eta})_-$  is  $\phi_i, i = 0, 1, 2, 3$  with  $\phi_i(S) = 0, \phi_i(Q) = (1 - Q)v_i$ . The operation of  $T_2$  is

$$T_2(v_0, v_1, v_2, v_3) = (v_0, v_1, v_2, v_3) = \begin{pmatrix} -31 & 0 & 0 & 0 \\ * & 31 & 0 & 0 \\ * & 0 & -10 & 18 \\ * & 0 & -8 & 10 \end{pmatrix}.$$

The characteristic polynomial of  $T_2$  on  $H^1(\Gamma, W_{n,\eta})_-$  is

$$\chi_2(x) = (x + 31)(x - 31)(x^2 + 44).$$

The factors  $(x + 31)$  and  $(x - 31)$  come from the operation of  $T_2$  on the boundary cohomology  $H^1(\Gamma_\infty, W_{n,\eta} \otimes \mathbb{Q}) \cong \mathbb{Q}\phi_0 + \mathbb{Q}\phi_\infty$ . More precise,

$$T_2\phi_\infty = -31\phi_\infty, \quad T_2\phi_0 = 31\phi_0.$$

Therefore the characteristic polynomial of  $T_2$  on  $S_6(\Gamma_0(p_0), \eta)$  is  $x^2 + 44$ . The numerical computations of  $T_2, T_3, T_5$  and  $T_7$  for small  $p_0$  and  $n$  are given in the table 1.

**7.2 Remark.** The space  $S_{n+2}(\Gamma_0(p_0), \eta)$  carries the Petersson product, a non-degenerate Hermitian product on  $S_{n+2}(\Gamma_0(p_0), \eta)$ . If  ${}^t$  denotes ‘‘transpose’’ with respect to this product, then  $T_i^t = \eta(l)T_i$ . Let now  $\lambda$  be an eigenvalue of  $T_i$ , we have then  $\bar{\lambda} = \eta(l)\lambda$  (cf. [Rib] § 1). Therefore,



if  $\eta(l) = -1$ , then  $\lambda = ia$  with  $a \in \mathbb{R}$ . If  $\eta(l) = 1$ ,  $\lambda \in \mathbb{R}$ .

(1)  $p_0 \equiv 1 \pmod{4}$ . In that case the dimension of  $S_{n+2}(\Gamma_0(p_0), \eta)$  is even.

i.  $\eta(l) = -1$ . The characteristic polynomial of  $T_l$  is

$$\begin{aligned} \chi_l(x) &= (x - ia_1)(x + ia_1)(x - ia_2)(x + ia_2) \cdots (x - ia_r)(x + ia_r) \\ &= (x^2 + a_1^2)(x^2 + a_2^2) \cdots (x^2 + a_r^2) \\ &= x^{2r} + b_1x^{2r-2} + \cdots + b_r \end{aligned}$$

with  $b_1, \dots, b_r \geq 0$ .

ii.  $\eta(l) = 1$ . The characteristic polynomial of  $T_l$  is

$$\chi_l(x) = g(x)^2$$

for some polynomial  $g(x)$ . The roots of  $g(x)$  are all real.

(2)  $p_0 \equiv 3 \pmod{4}$ . In that case the dimension of  $S_{n+2}(\Gamma_0(p_0), \eta)$  is odd.

i.  $\eta(l) = -1$ . There are zero eigenvalues. The characteristic polynomial is

$$\chi_l(x) = x^h(x^{2s} + b_1x^{2s-2} + \cdots + b_s)$$

where  $h$  is the class number of the field  $\mathbb{Q}(\sqrt{-p_0})$ .

ii.  $\eta(l) = +1$ . The characteristic polynomial is

$$\chi_l(x) = g(x)^2 \cdot f(x)$$

where  $f(x)$  is a polynomial generated by the Theta series and  $\deg(f(x)) = h$  (cf. [Shi]).

The results in the table 1 confirm the remark above.

ACKNOWLEDGEMENTS. The author is grateful for the support received from the DFG during the preparation of this paper.

Table 1

The characteristic polynomials of the Hecke operators  $T_2, T_3, T_5$ , and  $T_7$  on the cusp forms  $S_k(\Gamma_0(p_0), \eta)$ , where  $\eta$  is the Legendre symbol.

---


$$P_0=5, K=N+2: \quad \text{ETA}(2, P_0)=-1, \text{ETA}(3, P_0)=-1, \text{ETA}(7, P_0)=-1$$


---

N=4

$$T_2:=X^2+44$$

$$T_3:=X^2+396$$

$$T_7:=X^2+3564$$

N=6

$$T_2:=X^2+116$$

$$T3 := X^2 + 1044$$

$$T7 := X^2 + 176436$$

$$N = 8$$

$$T2 := X^4 + 1708 * X^2 + 1216$$

$$T3 := X^4 + 33552 * X^2 + 45529776$$

$$T7 := X^4 + 104167728 * X^2 + 2144749073480496$$

$$N = 10$$

$$T2 := X^4 + 4132 * X^2 + 2496256$$

$$T3 := X^4 + 341568 * X^2 + 18385718256$$

$$T7 := X^4 + 4904976672 * X^2 + 2087691277621558896$$

$$N = 12$$

$$T2 := X^6 + 41052 * X^4 + 440779968 * X^2 + 617678127104$$

$$T3 := X^6 + 8329788 * X^4 + 17708569483248 * X^2 + 1517182687182390336$$

$$T7 := X^6 + 213997084092 * X^4 + 10526623838205776341488 * X^2 \\ + 46528027403146207719038230676544$$

$$N = 14$$

$$T2 := X^8 + 117588 * X^6 + 2455515648 * X^4 + 4160982695936$$

$$T3 := X^8 + 48755052 * X^6 + 160831293357168 * X^4 + 79914543281387267904$$

$$T7 := X^8 + 8435989101708 * X^6 + 21799671533824901950559088 * X^4 \\ + 17560391031732483266163471186728360256$$

$$N = 16$$

$$T2 := X^8 + 813836 * X^6 + 197805587136 * X^4 + 15212877148553216 * X^2 \\ + 338022604671796903936$$

$$T3 := X^8 + 634018824 * X^6 + 123866741829162816 * X^4$$

$$+ 8052359188906852344353664 * X^2 + 62556794360183564540341578775296$$

$$T7 := X^8 + 1358809234759656 * X^6 + 571583885437582806176526269376 * X^4$$

$$+ 71743645253248677409589367384237677906875776 * X^2$$

$$+ 1225649387103886247126536790871068024121055114558759170816$$

$$N = 18$$

$$T2 := X^8 + 2907524 * X^6 + 2568216374016 * X^4 + 678867689422782464 * X^2 \\ + 8301849147532531204096$$

$$T3 := X^8 + 4476366576 * X^6 + 6998614044948851616 * X^4$$

$$+ 4394102925151257527276257536 * X^2$$

$$+ 859178610673769519506507390330864896$$

$$T7 := X^8 + 51160209747400944 * X^6 + 649955449858844816462059274614176 * X^4$$

$$+ 1364277688497122259242343356898905016125205537024 * X^2$$

$$+ 669240116784884405332807029722912360484202369263457687836124416$$

$N = 20$

$$\begin{aligned} T2: &= X^{10} + 15122620 * X^8 + 74461069946560 * X^6 + 143355636201404579840 * X^4 \\ &\quad + 92050796042892961151713280 * X^2 + 14584363461253989437721829965824 \\ T3: &= X^{10} + 75700218780 * X^8 + 1690290073124929870560 * X^6 \\ &\quad + 10589033423492535098094901061760 * X^4 \\ &\quad + 10613905392864453389568881849143800910080 * X^2 \\ &\quad + 2839805815981800681177617222924350898397211646976 \\ T7: &= X^{10} + 2623942726584980220 * X^8 \\ &\quad + 2393834166138243432310096381198875360 * X^6 \\ &\quad + 897532555190091115792311471245276662831526251090803840 * X^4 \\ &\quad + 12130074515248198130994387822394541291034854198094242424652328 \\ &\quad \quad 8877748480 * X^2 \\ &\quad + 46127042768695709673188041565524085282943142185101532687761730 \\ &\quad \quad 59937738127467213792820224 \end{aligned}$$

$N = 22$

$$\begin{aligned} T2: &= X^{10} + 62579380 * X^8 + 1269587477762560 * X^6 + 9620767823712245596160 * X^4 \\ &\quad + 19648398991934117012339425280 * X^2 \\ &\quad + 3574276364739503586982992256434176 \\ T3: &= X^{10} + 565341209820 * X^8 + 118033406092349714504160 * X^6 \\ &\quad + 10931210697192722327499640220787840 * X^4 \\ &\quad + 406914738264133623534685754882233338060775680 * X^2 \\ &\quad + 2922982673270172565978306559380807929420812626129050624 \\ T7: &= X^{10} + 111196555384787994780 * X^8 \\ &\quad + 3435262712787547437076787484432246075360 * X^6 \\ &\quad + 36412038333453087389178656736733867773560385038722769196160 * X^4 \\ &\quad + 85586684250837585052810715708244193791504968303062585347065682 \\ &\quad \quad 3315813705484 * X^2 \\ &\quad + 34879917834200075143347515459724686022028645166264732887044408 \\ &\quad \quad 1898307411589887020773579776 \end{aligned}$$

$P0 = 7, K = N + 2: \quad \eta(2, P0) = 1, \eta(3, P0) = -1, \eta(5, P0) = -1$

$N = 1$

$$\begin{aligned} T2: &= X + 3 \\ T3: &= X \\ T5: &= X \end{aligned}$$

$N = 3$

$$T2: = X - 1$$

$$T3:=X$$

$$T5:=X$$

$$N=5$$

$$T2:=(X+8)^2*(X-9)$$

$$T3:=X*(X^2+2040)$$

$$T5:=X*(X^2+2040)$$

$$N=7$$

$$T2:=(X^2-16*X-120)^2*(X+31)$$

$$T3:=X*(X^4+17184*X^2+40430880)$$

$$T5:=X*(X^4+1809120*X^2+736852788000)$$

$$N=9$$

$$T2:=(X^2+24*X-592)^2*(X-57)$$

$$T3:=X*(X^4+132480*X^2+4381776000)$$

$$T5:=X*(X^4+11349120*X^2+25531635024000)$$

$$N=11$$

$$T2:=(X^3-10216*X+172800)^2*(X+47)$$

$$T3:=X*(X^6+2434704*X^4+1858882957920*X^2+429665499302054400)$$

$$T5:=X*(X^6+1290415440*X^4+544550093091324000*X^2+75252114900743951016000000)$$

$$N=13$$

$$T2:=(X^4-88*X^3-49600*X^2+3161344*X+199833600)^2*(X+87)$$

$$T3:=X*(X^8+32897856*X^6+307339393288320*X^4+678298556041314969600*X^2+3197232909629570972160000)$$

$$T5:=X*(X^8+30327873600*X^6+303490459358455478400*X^4+1203282796541403639170914560000*X^2+1643994907570049884150368794126400000000)$$

$$N=15$$

$$T2:=(X^4+272*X^3-98776*X^2-15713792*X+773514240)^2*(X-449)$$

$$T3:=X*(X^8+193153824*X^6+13542540815792160*X^4+407914538508420139929600*X^2+4459777119693624095941077504000)$$

$$T5:=X*(X^8+579368436960*X^6+81730362131262670356000*X^4+2948421249394654853254317120000000*X^2+1938958613271787722241837348802664000000000)$$

$$N=17$$

$$T2:=(X^5-456*X^4-716336*X^3+195823104*X^2+124785737728*X-13438656184320)^2*(X+999)$$

$$T3:=X*(X^{10}+2541979176*X^8+1981194676580514240*X^6$$

$$\begin{aligned}
& + 470805560399816932850265600 * X^4 \\
& + 33914967955417991795516068276224000 * X^2 \\
& + 463598587189134022224773827601838489600000) \\
T5 := & X * (X^{10} + 26205473373480 * X^8 + 229513487290145010811339200 * X^6 \\
& + 771146143064265537788863097464793280000 * X^4 \\
& + 727302726371763893278482096922796468827756800000000 * X^2 \\
& + 258396135501073534080222733876861109352500261616000000000000)
\end{aligned}$$

---


$$P0 = 11, K = N + 2: \quad \text{ETA}(2, P0) = -1, \text{ETA}(3, P0) = 1, \text{ETA}(5, P0) = 1, \\
\text{ETA}(7, P0) = -1$$


---

N = 1

$$\begin{aligned}
T2 & := X \\
T3 & := X + 5 \\
T5 & := X + 1 \\
T7 & := X
\end{aligned}$$

N = 3

$$\begin{aligned}
T2 & := X * (X^2 + 30) \\
T3 & := (X + 3)^2 * (X - 7) \\
T5 & := (X - 31)^2 * (X + 49) \\
T7 & := X * (X^2 + 3000)
\end{aligned}$$

N = 5

$$\begin{aligned}
T2 & := X * (X^4 + 270 * X^2 + 16680) \\
T3 & := (X^2 - 12 * X - 1509)^2 * (X - 10) \\
T5 & := (X + 65)^4 * (X - 74) \\
T7 & := X * (X^4 + 393000 * X^2 + 38537472000)
\end{aligned}$$

N = 7

$$\begin{aligned}
T2 & := X * (X^6 + 1374 * X^4 + 436560 * X^2 + 40320000) \\
T3 & := (X^3 + 18 * X^2 - 6285 * X - 201150)^2 * (X + 113) \\
T5 & := (X^3 + 224 * X^2 - 525475 * X + 31988350)^2 * (X - 1151) \\
T7 & := X * (X^6 + 22327704 * X^4 + 102738589578240 * X^2 + 134544048242688000000)
\end{aligned}$$

N = 9

$$\begin{aligned}
T2 & := X * (X^8 + 6030 * X^6 + 11712120 * X^4 + 7669330560 * X^2 + 564269690880) \\
T3 & := (X^4 + 201 * X^3 - 98919 * X^2 - 1150929 * X + 1149750126)^2 * (X - 475) \\
T5 & := (X^4 - 1215 * X^3 - 21311915 * X^2 - 2265218325 * X + 17429871112150)^2 \\
& \quad * (X + 3001) \\
T7 & := X * (X^8 + 767889840 * X^6 + 102582267767649600 * X^4 \\
& \quad + 1566249894398109763584000 * X^2 + 6330325858079634845966794752000)
\end{aligned}$$

$N=11$

$$\begin{aligned} T2: &= X*(X^{10} + 30654*X^8 + 318945120*X^6 + 1305642637440*X^4 \\ &\quad + 2049564619929600*X^2 + 957721368231936000) \\ T3: &= (X^5 - 1218*X^4 - 775914*X^3 + 838214892*X^2 + 189020241225*X \\ &\quad + 120422340866250)^2*(X + 1358) \\ T5: &= (X^5 - 13246*X^4 - 413004050*X^3 + 7878939523400*X^2 \\ &\quad - 32298230888024375*X + 12308222362848968750)^2*(X + 25774) \\ T7: &= X*(X^{10} + 72369291504*X^8 + 1579588871009845139520*X^6 \\ &\quad + 12964051646785030759215833088000*X^4 \\ &\quad + 37709819138673185762480264655566929920000*X^2 \\ &\quad + 23187441850664232142842389272548747887247360000000) \end{aligned}$$

$$P0=13, K=N+2: \quad \text{ETA}(2, P0)=-1, \text{ETA}(3, P0)=1, \text{ETA}(5, P0)=-1, \\ \text{ETA}(7, P0)=-1$$

$N=2$

$$\begin{aligned} T2: &= X^2 + 9 \\ T3: &= (X + 1)^2 \\ T5: &= X^2 + 81 \\ T7: &= X^2 + 225 \end{aligned}$$

$N=4$

$$\begin{aligned} T2: &= X^6 + 161*X^4 + 5856*X^2 + 18864 \\ T3: &= (X^3 - 8*X^2 - 549*X + 4068)^2 \\ T5: &= X^6 + 8018*X^4 + 13754433*X^2 + 2485690416 \\ T7: &= X^6 + 82950*X^4 + 1662348177*X^2 + 423560602764 \end{aligned}$$

$N=6$

$$\begin{aligned} T2: &= X^6 + 449*X^4 + 37224*X^2 + 205776 \\ T3: &= (X^3 + 28*X^2 - 2601*X - 71748)^2 \\ T5: &= X^6 + 243506*X^4 + 1206410625*X^2 + 93756690000 \\ T7: &= X^6 + 847206*X^4 + 231424342425*X^2 + 20471634652072500 \end{aligned}$$

$N=8$

$$\begin{aligned} T2: &= X^{10} + 3841*X^8 + 5134480*X^6 + 2823572208*X^4 + 614223235584*X^2 \\ &\quad + 43308450164736 \\ T3: &= (X^5 + X^4 - 66033*X^3 + 1260423*X^2 + 530326440*X + 14266185264)^2 \\ T5: &= X^{10} + 14820283*X^8 + 74785768290163*X^6 + 146559998245698565881*X^4 \\ &\quad + 87330504466586448091944000*X^2 + 12065478109519129517166006240000 \\ T7: &= X^{10} + 252125259*X^8 + 23724928789729587*X^6 \\ &\quad + 1025407325324195954977977*X^4 \end{aligned}$$

$$+ 19661129805887483504404526084736 * X^2$$

$$+ 121307703706137674344780717867862132400$$

N=10

$$T2 := X^{12} + 18433 * X^{10} + 121088056 * X^8 + 340607607312 * X^6 + 380893885719552 * X^4$$

$$+ 134825856231997440 * X^2 + 1497425476589715456$$

$$T3 := (X^6 + 244 * X^5 - 665334 * X^4 - 129598956 * X^3 + 109163403621 * X^2$$

$$+ 14522233287672 * X - 255121008509808)^2$$

$$T5 := X^{12} + 289917556 * X^{10} + 32326953002900950 * X^8$$

$$+ 1726712418063587931532500 * X^6$$

$$+ 44108094881553049831926298640625 * X^4$$

$$+ 430033290962195234920750132329450000000 * X^2$$

$$+ 10886105645673774994569770130605197500000000$$

$$T7 := X^{12} + 13650769356 * X^{10} + 64465836700280921262 * X^8$$

$$+ 139418894150631875357617076028 * X^6$$

$$+ 143785268511480525150168789070017931401 * X^4$$

$$+ 63753954827004609548776322006655133952858100000 * X^2$$

$$+ 9054507376401194828902343707676292621596570213493750000$$

$$P0=17, K=N+2: \quad \text{ETA}(2, P0)=1, \quad \text{ETA}(3, P0)=-1, \quad \text{ETA}(5, P0)=-1,$$

$$\text{ETA}(7, P0)=-1$$

N=2

$$T2 := (X^2 + X - 8)^2$$

$$T3 := X^4 + 74 * X^2 + 1072$$

$$T5 := X^4 + 480 * X^2 + 38592$$

$$T7 := X^4 + 530 * X^2 + 68608$$

N=4

$$T2 := (X^3 + X^2 - 68 * X - 36)^2$$

$$T3 := X^6 + 668 * X^4 + 145216 * X^2 + 10185984$$

$$T5 := X^6 + 9488 * X^4 + 8442048 * X^2 + 40743936$$

$$T7 := X^6 + 71708 * X^4 + 104887424 * X^2 + 2346850713600$$

N=6

$$T2 := (X^5 + 9 * X^4 - 452 * X^3 - 2988 * X^2 + 27904 * X + 83616)^2$$

$$T3 := X^{10} + 16832 * X^8 + 93191572 * X^6 + 192821327856 * X^4 + 116860780245888 * X^2$$

$$+ 9421474370420736$$

$$T5 := X^{10} + 351440 * X^8 + 44989957632 * X^6 + 2580556932172800 * X^4$$

$$+ 65876023734658560000 * X^2 + 602974359706927104000000$$

$$T7 := X^{10} + 4233136 * X^8 + 5824132863636 * X^6 + 2871845375371443376 * X^4$$

$$+ 497996015831560956471424 * X^2 + 14856566017369895192889851904$$

$N = 8$

$$T2 := (X^6 - 15 * X^5 - 1892 * X^4 + 20460 * X^3 + 770176 * X^2 - 3195840 * X - 6636441)^2$$

$$T3 := X^{12} + 122690 * X^{10} + 5157152560 * X^8 + 87983684680032 * X^6 \\ + 612743619071665152 * X^4 + 1335826553351738886144 * X^2 \\ + 203949399568932198678528$$

$$T5 := X^{12} + 13939648 * X^{10} + 67854209805568 * X^8 + 136905662805154384896 * X^6 \\ + 103030638845234136672153600 * X^4 \\ + 23873047875692895959460126720000 * X^2 \\ + 213202160733331266611086098432000000$$

$$T7 := X^{12} + 181444282 * X^{10} + 8551923317087424 * X^8 \\ 145015964651608425915232 * X^6 + 922072716536803810905054408448 * X^4 \\ + 2318685324256549381944604148484046848 * X^2 \\ + 1967676585788509591285949532270066715852800$$

---


$$P0 = 19, K = N + 2: \quad \text{ETA}(2, P0) = -1, \text{ETA}(3, P0) = -1, \text{ETA}(5, P0) = 1, \\ \text{ETA}(7, P0) = 1$$


---

$N = 1$

$$T2 := X * (X^2 + 13)$$

$$T3 := X * (X^2 + 13)$$

$$T5 := (X - 4)^2 * (X + 9)$$

$$T7 := (X + 5)^3$$

$N = 3$

$$T2 := X * (X^4 + 35 * X^2 + 142)$$

$$T3 := X * (X^4 + 301 * X^2 + 5112)$$

$$T5 := (X^2 + 21 * X + 92)^2 * (X - 31)$$

$$T7 := (X^2 - 68 * X + 499)^2 * (X + 73)$$

$N = 5$

$$T2 := X * (X^8 + 483 * X^6 + 75582 * X^4 + 4242376 * X^2 + 71047680)$$

$$T3 := X * (X^8 + 3442 * X^6 + 4292649 * X^4 + 2281096296 * X^2 + 432254085120)$$

$$T5 := (X^4 - 54 * X^3 - 49415 * X^2 + 3367200 * X + 292006000)^2 * (X + 54)$$

$$T7 := (X^4 + 70 * X^3 - 157380 * X^2 - 29481334 * X - 1276939885)^2 * (X - 610)$$

$N = 7$

$$T2 := X * (X^{12} + 2323 * X^{10} + 2010462 * X^8 + 803113072 * X^6 + 150633270400 * X^4 \\ + 12173735396352 * X^2 + 333034797957120)$$

$$T3 := X * (X^{12} + 59719 * X^{10} + 1354569075 * X^8 + 14270784462117 * X^6 \\ + 66670855305320376 * X^4 + 99071703704871505152 * X^2)$$



$$+ 33664128506976532561920)$$

$$T5:=(X^6-4*X^5-1446203*X^4+95652050*X^3+409166434600*X^2-102103842940000*X+6563900254320000)^2*(X+289)$$

$$T7:=(X^6-1843*X^5-25578196*X^4+37453164210*X^3+157007096825425*X^2+139069305605381375*X-21933742012221418750)^2*(X-527)$$

$$P0=23, K=N+2: \quad \text{ETA}(2, P0)=1, \quad \text{ETA}(3, P0)=1, \quad \text{ETA}(5, P0)=-1, \\ \text{ETA}(7, P0)=-1$$

$$N=1$$

$$T2:=X^3-12*X+7$$

$$T3:=X^3-27*X+38$$

$$T5:=X^3$$

$$T7:=X^3$$

$$N=3$$

$$T2:=(X^2+4*X-2)^2*(X^3-48*X+79)$$

$$T3:=(X^2+6*X-45)^2*(X^3-243*X+14)$$

$$T5:=X^3*(X^4+2556*X^2+1270188)$$

$$T7:=X^3*(X^4+11988*X+31754700)$$

$$N=5$$

$$T2:=(X^4-4*X^3-162*X^2+920*X+832)^2*(X+7)*(X^2-7*X-143)$$

$$T3:=(X^4-15*X^3-957*X^2+13293*X^2-12870)^2*(X+38) \\ *(X^2-38*X-743)$$

$$T5:=X^3*(X^8+95100*X^6+3184494300*X^4+44006549508000*X^2 \\ +214214641502400000)$$

$$T7:=X^3*(X^8+660492*X^6+152231816700*X^4+13982809796769600*X^2 \\ +400834240321008384000)$$

$$N=7$$

$$T2:=(X^6+4*X^5-850*X^4-3248*X^3+147872*X^2+268672*X-5317760)^2 \\ *(X^3-768*X+1951)$$

$$T3:=(X^6+36*X^5-20508*X^4-1030644*X^3+86837139*X^2+5371429140*X \\ +55514443500)^2*(X^3-19683*X+1062686)$$

$$T5:=X^3*(X^{12}+3434556*X^{10}+4503520431468*X^8+2817283398730424640*X^6 \\ +847955819735403719760000*X^4+103782437973914306469472512000*X^2 \\ +2208782254549937077079536204800000)$$

$$T7:=X^3*(X^{12}+40494132*X^{10}+605958970060332*X^8 \\ +4096821152215401422400*X^6+12087187496206708701149510400*X^4 \\ +11540311691303117336118557810688000*X^2)$$

$$+ 252607388911566511898618438444236800000)$$

---


$$P_0 = 29, K = N + 2: \quad \text{ETA}(2, P_0) = -1, \text{ETA}(3, P_0) = -1, \text{ETA}(5, P_0) = 1, \\ \text{ETA}(7, P_0) = 1$$


---

$$N = 2$$

$$T_2 := X^6 + 38X^4 + 301X^2 + 560$$

$$T_3 := X^6 + 61X^4 + 791X^2 + 875$$

$$T_5 := (X^3 - 11X^2 - 133X + 1071)^2$$

$$T_7 := (X^3 + 14X^2 - 108X - 1192)^2$$

$$N = 4$$

$$T_2 := X^{12} + 278X^8 + 28285 + 1260472X^6 + 22944832X^4 + 140087936X^2 + 966400$$

$$T_3 := X^{12} + 2245X^{10} + 1884878X^8 + 715200530X^6 + 112977325989X^4 \\ + 4281127461369X^2 + 46577165867100$$

$$T_5 := (X^6 - 23X^5 - 12280X^4 + 235866X^3 + 33953337X^2 - 384523443X \\ - 2627317458)^2$$

$$T_7 := (X^6 - 10X^5 - 76080X^4 + 1925088X^3 + 1377655664X^2 - 73626194400X \\ - 519034134784)^2$$

$$N = 6$$

$$T_2 := X^{16} + 1382X^{14} + 744077X^{12} + 200869632X^{10} + 28931822432X^8 \\ + 2155663113216X^6 + 71710495842560X^4 + 663330761523200X^2 \\ + 590388176896000$$

$$T_3 := X^{16} + 22051X^{14} + 187767701X^{12} + 793510274339X^{10} \\ + 1809033803032599X^8 + 2281021494195869649X^6 \\ + 1527214705246483000335X^4 + 458931418705423915202025X^2 \\ + 30465487147014831010162500$$

$$T_5 := (X^8 + 99X^7 - 276993X^6 - 31299849X^5 + 17584369885X^4 \\ + 1686634037625X^2 - 196943514064875X^2 - 14966521618921875X \\ - 200278684287731250)^2$$

$$T_7 := (X^8 - 330X^7 - 3716228X^6 + 233875960X^5 + 3438911219312X^4 \\ + 1091616310004000X^3 - 293827217111058624X^2 \\ - 89873092347162858880X + 8119186526578407384064)^2$$

---


$$P_0 = 31, K = N + 2: \quad \text{ETA}(2, P_0) = 1, \text{ETA}(3, P_0) = -1, \text{ETA}(5, P_0) = 1, \\ \text{ETA}(7, P_0) = 1$$


---

$$N = 1$$

$$T_2 := (X + 1)^2(X^3 - 12X + 15)$$

$$T_3 := X^3(X^2 + 26)$$

$$T5:=(X-2)^2*(X^3-75*X+246)$$

$$T7:=(X-8)^2*(X^3-147X+*430)$$

N=3

$$T2:=(X^3+X^2-30*X+6)^2*(X^3-48*X-97)$$

$$T3:=X^3*(X^6+398*X^4+49236*X^2+1934136)$$

$$T5:=(X^3+4*X^2-291*X+1014)^2*(X^3-1875*X-29266)$$

$$T7:=(X^3+66*X^2-1005*X-31688)*(X^3-7203*X^2+50398)$$

N=5

$$T2:=(X^6+X^5-222*X^4-370*X^3+9416*X^2+13440*X-90624)^2*(X+15) \\ *(X^2-15*X+33)$$

$$T3:=X^3*(X^{12}+7208*X^{10}+19859688*X^8+26566749360*X^6+17884354852944*X^4 \\ +5570285336959680*X^2+590986232936064000)$$

$$T5:=(X^6+73*X^5-51615*X^4-3624325*X^3+522398750*X^2+25671172500*X \\ -103336)^2*(X^2-246*X+13641)*(X+246)$$

$$T7:=(X^6-3*X^5-207897*X^4-2308819*X^3+13269144858*X^2+215614693848*X \\ -247)^2*(X^2-430*X-168047)*(X+430)$$

P0=37, K=N+2:  $\text{ETA}(2, P0)=-1$ ,  $\text{ETA}(3, P0)=1$ ,  $\text{ETA}(5, P0)=-1$ ,  
 $\text{ETA}(7, P0)=1$

N=2

$$T2:=X^8+50*X^6+709*X^4+3000*X^2+1764$$

$$T3:=(X^4+3*X^3-50*X^2-57*X-427)^2$$

$$T5:=X^8+431*X^6+29521*X^4+588072*X^2+2039184$$

$$T7:=(X^4-2*X^3-587*X^2+2460*X+53892)^2$$

N=4

$$T2:=X^{16}+390*X^{14}+60701*X^{12}+4799932*X^{10}+203487156*X^8+4519465040*X^6 \\ +48993644736*X^4+211923220224*X^2+178006118400$$

$$T3:=(X^8+9*X^7-1280*X^6-11016*X^5+422488*X^4+2751084*X^3-25673805*X^2 \\ -30714957*X+141986196)^2$$

$$T5:=X^{16}+31026*X^{14}+373650779*X^{12}+2220056867434*X^{10} \\ +6834316986168825*X^8+10475224449621004436*X^6 \\ +6885539411711705092656*X^4+1110302609356408225416384*X^2 \\ +19726944242324026399110144$$

$$T7:=(X^8-95*X^7-54561*X^6+2410919*X^5+907038560*X^4+534484632*X^3 \\ -349499585616*X^2-2731120576272*X+3409346511153792)^2$$

$$P_0=41, K=N+2: \quad \text{ETA}(2, P_0)=1, \text{ETA}(3, P_0)=-1, \text{ETA}(5, P_0)=1, \\ \text{ETA}(7, P_0)=-1$$

$$N=2$$

$$T_2:=(X^5+3X^4-25X^2-51X^2+104X+32)^2$$

$$T_3:=X^{10}+180X^8+10910X^6+276172X^4+2531856X^2+524672$$

$$T_5:=(X^5+2X^4-282X^3-1400X^2+9016X+43904)^2$$

$$T_7:=X^{10}+1912X^8+1274822X^6+344662636X^4+30875879696X^2 \\ +87767656832$$

$$P_0=43, K=N+2: \quad \text{ETA}(2, P_0)=-1, \text{ETA}(3, P_0)=-1, \text{ETA}(5, P_0)=-1, \\ \text{ETA}(7, P_0)=-1$$

$$N=1$$

$$T_2:=X*(X^6+20X^4+121X^2+214)$$

$$T_3:=X*(X^6+45X^4+431X^2+214)$$

$$T_5:=X*(X^6+117X^4+3863X^2+25894)$$

$$T_7:=X*(X^6+150X^4+4896X^2+3424)$$

$$P_0=47, K=N+2: \quad \text{ETA}(2, P_0)=1, \text{ETA}(3, P_0)=1, \text{ETA}(5, P_0)=-1, \\ \text{ETA}(7, P_0)=1$$

$$N=1$$

$$T_2:=(X+1)^2*(X^5-20X^3+80X-17)$$

$$T_3:=(X+2)^2*(X^5-45X^3+405X-298)$$

$$T_5:=X^5*(X^2+78)$$

$$T_7:=(X+4)^2*(X^5-245X^3+12005X-31922)$$

$$N=3$$

$$T_2:=(X^5+X^4-40X^3+12X^2+300X-316)^2*(X^5-80X^3+1280X+1759)$$

$$T_3:=(X^5-4X^4-207X^3+576X^2+7803X+9558)^2*(X^5-405X^3+32805X \\ +29294)$$

$$T_5:=X^5(X^{10}+5490X^8+10917588X^6+9407020248X^4+3230761626000X^2 \\ +270690407718048)$$

$$T_7:=(X^5-14X^4-2905X^3-45230X^2-141377X+94796)^2*(X^5-12005X^3 \\ +28824005X-45406386)$$

$$P_0=53, K=N+2: \quad \text{ETA}(2, P_0)=-1, \text{ETA}(3, P_0)=1, \text{ETA}(5, P_0)=-1, \\ \text{ETA}(7, P_0)=1$$

$$N=2$$

$$T_2:=X^{12}+80X^{10}+2356X^8+30996X^6+176575X^4+393232X^2+285376$$

$$T3 := X^{12} + 215 * X^{10} + 16178 * X^8 + 505118 * X^6 + 5738621 * X^4 + 15503831 * X^2 + 673036$$

$$T5 := X^{12} + 789 * X^{10} + 196604 * X^8 + 18690640 * X^6 + 682399088 * X^4 + 6573121072 * X^2 + 2960741056$$

$$T7 := (X^6 - 12 * X^5 - 1052 * X^4 + 10868 * X^3 + 215348 * X^2 - 624840 * X - 9386656)^2$$

## REFERENCE

- [AS] A. Ash and G. Stevens, Cohomology of arithmetic groups and congruences between systems of Hecke eigenvalues, *J. reine angew. Math.*, **356** (1986), 192–220.
- [Bro] K. Brown, Cohomology of groups, GTM 87, Springer Verlag (1982).
- [Hab] K. Haberland, Perioden von Modulformen einer Variabler und Gruppenkohomologie I, II, III, *Math. Nachr.*, **112** (1983), 245–315.
- [Rib] K. A. Ribet, Galois representations attached to eigenform with nebentypus, in: *Lecture Notes in Math.*, **601**, Springer Verlag.
- [Ser] J.-P. Serre, A course in arithmetic, GTM 7, Springer Verlag (1973).
- [Shi] G. Shimura, On elliptic curves with complex multiplication as factor of the Jacobians of modular function fields, *Nagoya Math. J.*, **43** (1971), 199–208.
- [Wan] X.-D. Wang, Die Eisensteinklasse in  $H^1(SL_2(\mathbb{Z}), M_n(\mathbb{Z}))$  und die Arithmetik spezieller Werte von L-Funktionen, *Bonner Math. Schriften*, **202**, Bonn (1989).

*Mathematisches Institut  
der Universität Bonn  
Berlingstr. 1  
D-5300 Bonn  
Germany*