

Conclusions: HBO treatment to prevent the development persistent neuropsychologic disturbances, and to decrease the incidence of delayed neuropsychological sequelae. The use of HBO posing detoxication, antihypoxic and metabolic effects in the complex treatment of drug - addiction to improve the outcomes. Thus, HBO may be used as a method of a metabolic brain protect.

P03.436

CLINICAL SYMPTOMS AMONG FEMALE AND MALE DRUG-NAÏVE SCHIZOPHRENIC PATIENTS

B.J. Havaki-Kontaxaki, V.P. Kontaxakis*, S.S. Stamouli, C.T. Kollias, G.N. Christodoulou. *Department of Psychiatry, University of Athens, Greece*

Background: Gender has been recognized to be an important factor in the expression of schizophrenic illness. The aim of this study is to identify differences between female and male drug-naïve first-episode schizophrenics regarding their clinical symptomatology and psychosocial functioning.

Methods: Twenty-five drug-naïve first-episode schizophrenic patients (14 women, 11 men) consecutively admitted at Eginition Hospital, Athens, were included in the study. Their mean age was 27.8 (± 6.8) years. Patients were assessed on admission (during the first week) using the Positive and Negative Syndrome scale (PANSS, Kay 1991) and the Global Assessment of Functioning Scale (GAF, APA 1987). There were no statistically significant differences between female and male schizophrenic patients regarding age (27.8 vs 27.7), education (years of schooling, 12.2 vs 12.0), employment status (unemployment, 78% vs 55%), way of admission (voluntarily, 93% vs 91%).

Results: Female and male schizophrenics did not differ in terms of the total PANSS score (113.7 vs 117.8), the PANSS-positive subscale score (27.3 vs 28.9), the PANSS-negative subscale score (27.8 vs 29.4), the PANSS-general psychopathology subscale score (58.6 vs 61.3) and the GAF score (32.2 vs 29.6). However, women more often presented with hostility (79% vs 18%, $p < 0.01$) and depression (86% vs 45%, $p < 0.05$) while men with grandiosity (21% vs 63%, $p < 0.05$).

P03.437

ARE REGIONAL SUICIDE RATES MIRRORED IN SUBSEQUENT NATIONAL POLITICAL CHANGES?

M. Voracek*, A.K. Formann, G. Sonneck. *University of Vienna, Austria*

Recently, political changes in Austria have received worldwide attention. We subjected the voting pattern in the preceding general elections in Austria (autumn 1999) to a suicide-epidemiological analysis. Previous research studied short-term effects of political changes on suicide occurrence. Regional suicide rates were found to decrease during U.S. Presidential campaigns and elections, and regional parasuicide and suicide rates were found to decrease after U.K. general elections. Reversing the question we investigated, whether regional suicide rates in Austria were related to the subsequent national political changes. For the 121 political districts in Austria, weighted by their size (measured by persons entitled to vote), we regressed 7 predictor variables (percentage of gains/losses of the 5 important political parties in Austria in the 1999 general election, as compared to the 1995 general election, '99-to-'95-changes of voters' turnout, and proportion of invalid voting) on average standardized suicide rates (SMR) for the preceding period (1988-94). Stepwise weighted multiple regression analysis lead to

a significant model, including 3 of the predictor variables. The '95-to-'99-changes for two oppositional parties and the changes in the proportion of invalid voting together accounted for no less than one quarter of the variability in preceding district-level suicide rates. Generally, pronounced caution should be exercised in interpreting this pattern as representing cause-effect relationships. Thus, additional analyses aiming to rule out possible confounding variables in the statistical model are presented. Nevertheless, this result gives some preliminary evidence that variations in preceding regional suicide incidence (which at first may seem of little direct concern to voting behavior) actually may be mirrored in subsequent changes in voting behavior (proportion of invalid voting, gains/losses of oppositional parties). This finding can be discussed and understood in terms of the classic Durkheimian concept of societal anomie.

P03.438

AN ACID TEST FOR NOSTRADAMUS: TOTAL ECLIPSE OF THE SUN 1999 AND SUICIDE INCIDENCE

M. Voracek*, G. Sonneck. *University of Vienna, Austria*

Apart from the millennium, the total eclipse of the sun on Wednesday, August 11th, 1999, was definitely the single most important mass-media event in Middle Europe that year. In Austria, extensive and in-depth mass-media coverage, announcing and hailing the coming event, started months before, leading to a collective experience of almost spiritual dimensions in the population. Approx. 40% of the Austrian working population were on leave that day, and approx. 750,000 of Austria's population (8 million inhabitants) traveled into the narrow strip (approx. 110 km wide) of eclipse totality, causing severe traffic jam and public transport disruptions. In advance of the event, several concerns of possible individual and collective psychiatric affections as well were published (Gratton & Line, 1999). However, any data concerning an eventual short-term impact of total eclipses of the sun on national suicide incidence are lacking. Given the circumstances of the event as described above, we took the opportunity to address this question. Austrian individual records suicide data for 1999 will be available from Statistics Austria in May 2000. The results of a comparison of suicide incidence at and around the day of the total eclipse with a preceding control period (suicide incidence on and around August 11th in the years 1984-98) will be presented. We hypothesize that a "Nostradamian" view (apocalyptic/bad omen) of the event under scrutiny will not withstand this acid test: in a line with broad empirical evidence upon the timing of suicide (cf. Voracek & Sonneck, 1999) we expect a significant decrease in Austrian suicide incidence before and at August 11th 1999. Furthermore, that decrease should have been more pronounced for regions within the narrow strip of total visibility of the eclipse.

P03.439

SUICIDAL BEHAVIOR IN PRISON INMATES: PROSPECTIVE AUSTRIAN NATIONAL DATA 1996-99

M. Voracek*, O. Bernecker, G. Sonneck. *University of Vienna, Austria*

It is widely known that prison inmates represent a special, high-risk subpopulation for suicidal behavior and suicidal acts. Typically, investigations of prison suicide are based on retrospective research designs (cumulative case records). For apparent reasons, reliability, validity, and generalizability of suicide-epidemiological findings from such retrospective research designs are clearly limited. In Austria, within an ongoing surveillance system and prospective research design which started in 1996, extensive data of all suicidal