

Weak Explicit Matching for Level Zero Discrete Series of Unit Groups of p -Adic Simple Algebras

Allan J. Silberger and Ernst-Wilhelm Zink

Abstract. Let F be a p -adic local field and let A_i^\times be the unit group of a central simple F -algebra A_i of reduced degree $n > 1$ ($i = 1, 2$). Let $\mathcal{R}^2(A_i^\times)$ denote the set of irreducible discrete series representations of A_i^\times . The “Abstract Matching Theorem” asserts the existence of a bijection, the “Jacquet-Langlands” map, $\mathcal{JL}_{A_2, A_1} : \mathcal{R}^2(A_1^\times) \rightarrow \mathcal{R}^2(A_2^\times)$ which, up to known sign, preserves character values for regular elliptic elements. This paper addresses the question of explicitly describing the map \mathcal{JL} , but only for “level zero” representations. We prove that the restriction $\mathcal{JL}_{A_2, A_1} : \mathcal{R}_0^2(A_1^\times) \rightarrow \mathcal{R}_0^2(A_2^\times)$ is a bijection of level zero discrete series (Proposition 3.2) and we give a parameterization of the set of unramified twist classes of level zero discrete series which does not depend upon the algebra A_i and is invariant under \mathcal{JL}_{A_2, A_1} (Theorem 4.1).

0 Introduction

This paper gives the Jacquet-Langlands correspondence (\mathcal{JL}) explicitly for level zero discrete series up to unramified twist (see Theorem 4.1). In particular, we prove that \mathcal{JL} restricts to a bijective correspondence of level zero discrete series (see Proposition 3.2). The subtler question of giving \mathcal{JL} precisely on each inertial class is left to our next paper, which depends upon this one. In this paper we use the Abstract Matching Theorem (AMT) (see Section 0.3), so our results are of the nature, “AMT implies that \mathcal{JL} gives this correspondence”.

0.1 Some Structure and Notation

This section fixes some notation and terminology. Our notation will be consistent with that of [GSZ]. Equation numbers are always in parentheses.

Let F be a p -adic local field and $n > 1$ an integer. For $d \geq 1$ let $D := D_d$ denote a central F -division algebra of index d and let $A := M_m(D)$ a central simple F -algebra of reduced degree $n := dm$.

Let $o := o_F$ denote the ring of integers of F , ϖ_F denotes a prime element of o , and $\mathfrak{p}_F := \varpi_F o$ the maximal ideal of o . Let $k := k_F$ denote the residual field of F with $q := |k|$ its order.

Let $O := O_D$ denote the ring of integers of D , $\varpi := \varpi_D$ a prime element of O such that $\varpi^d = \varpi_F$, and let $\mathfrak{p} := \mathfrak{p}_D = \varpi O$ be the prime ideal of O . We fix a maximal unramified field extension $F_d \subset D$ which is normalized by ϖ . The residual field

Received by the editors February 6, 2002.
AMS subject classification: 22E50, 11R39.
©Canadian Mathematical Society 2003.

$k_D := O/\mathfrak{p}$ is of order q^d and may be identified with k_d , the residual field of F_d . More generally, for $\ell \geq 1$ we write F_ℓ for an unramified extension of F of degree ℓ , o_ℓ for its ring of integers, and k_ℓ for a finite field extension of k of degree ℓ .

We write $X(k_\ell^\times)$ for the group of multiplicative characters of k_ℓ and $X_t(F_\ell^\times)$ for the group of tame multiplicative characters of F_ℓ . Thus $\chi \in X_t(F_\ell^\times)$ has the *reduction* $\bar{\chi} \in X(k_\ell^\times)$ (see (2.3)).

For any ordered partition s_1, \dots, s_r of m we call the block diagonal subgroup $M = \text{GL}_{s_1} \times \dots \times \text{GL}_{s_r}$ a *standard Levi subgroup* of GL_m and we call the parabolic subgroup $P = M \ltimes U$ which contains the upper triangular subgroup of GL_m a *standard parabolic subgroup* of GL_m . When necessary we write $G(R)$ to denote the group of R points of the algebraic group with respect to the ring R . Usually we abuse notation to identify algebraic subgroups of GL_m with their D -points.

For a hereditary order $\mathfrak{A} \subset A = M_m(D)$ we write $\mathfrak{P} := \mathfrak{P}_{\mathfrak{A}}$ for its Jacobson radical. Fix the maximal hereditary order $\mathfrak{A}_1 = M_m(O) \subset A$ and write $\mathfrak{P}_1 := \mathfrak{P}_{\mathfrak{A}_1} = M_m(\mathfrak{p})$. We call \mathfrak{A} *standard* if $\mathfrak{A} \subset \mathfrak{A}_1$ (hence $\mathfrak{P}_1 \subset \mathfrak{P}_{\mathfrak{A}}$) and if $(\mathfrak{A}/\mathfrak{P}_1)^\times$ is a standard parabolic subgroup of $\text{GL}_m(k_D)$. Most of the time we consider standard principal orders \mathfrak{A}_r , with Jacobson radical \mathfrak{P}_r , which are determined by the period $r \mid m$ and which are such that $(\mathfrak{A}_r/\mathfrak{P}_r)^\times$ is the standard Levi subgroup $[\text{GL}_s(k_D)]^r$ of $\text{GL}_m(k_D)$.

0.2 The Category of Level Zero Representations

Consider the Bernstein spectrum $\Omega(A^\times)$, the set of A^\times conjugacy classes $[M, \pi]$ of cuspidal pairs in which M is a standard Levi subgroup of A^\times and π is an irreducible supercuspidal representation of M . Each irreducible smooth representation Π of A^\times has a well defined supercuspidal support $\mathcal{CS}(\Pi) \in \Omega(A^\times)$ and thus we have a surjective finite to one map

$$\mathcal{CS}: \text{Irr}(A^\times) \longrightarrow \Omega(A^\times).$$

The partition of $\Omega(A^\times)$ into connected components Ω pulls back to a partition $\text{Irr}(A^\times) = \bigsqcup_{\Omega} \text{Irr}(\Omega)$, where $\text{Irr}(\Omega) := \{\Pi : \mathcal{CS}(\Pi) \in \Omega\}$. This partition gives rise to the Bernstein decomposition

$$(0.1) \quad \mathcal{M}(A^\times) = \prod_{\Omega} \mathcal{M}(\Omega)$$

of the category of smooth representations of A^\times into subcategories $\mathcal{M}(\Omega)$. The set of objects of $\mathcal{M}(\Omega)$ consists of those representations which have all irreducible subquotients in $\text{Irr}(\Omega)$.

Let $\mathcal{C} := \mathcal{C}(D)$ be a set of representatives for the set of unramified twist classes of irreducible unitary supercuspidal representations of $\text{GL}_s(D)$ for all $s \geq 1$, and let $\text{Div}^+(\mathcal{C})$ be the set of effective divisors on \mathcal{C} . For every $\mathcal{D} = \sum_{\pi \in \mathcal{C}} m_\pi \pi \in \text{Div}^+(\mathcal{C})$ we consider the cuspidal pair

$$\left(M_{\mathcal{D}} = \prod_{\pi \in \mathcal{C}} \text{GL}_{d(\pi)}(D)^{\times m_\pi}, \pi_{\mathcal{D}} = \bigotimes_{\pi \in \mathcal{C}} \pi^{\otimes m_\pi} \right)$$

and the connected component $\Omega(\mathcal{D})$ generated by the A^\times orbit of $(M_{\mathcal{D}}, \pi_{\mathcal{D}})$. The mapping

$$\mathcal{D} = \sum m_\pi \pi \longmapsto \Omega_{\mathcal{D}} = \text{the connected component of } [M_{\mathcal{D}}, \pi_{\mathcal{D}}] \in \Omega(A^\times)$$

defines a one-one correspondence between the set of divisors of degree $d(\mathcal{D}) = \sum m_\pi d(\pi) = m$ and the set of connected components of $\Omega(A^\times)$.

Let $\mathfrak{A}_1, \mathfrak{B}_1$ as in Section 0.1. Then the unit representation $1_{1+\mathfrak{B}_1}$ is a type for A^\times and the subcategory $\mathcal{M}_0(A^\times)$ of level zero representations of A^\times has as its set of objects the set of representations (Π, V) which are generated by the set of $(1 + \mathfrak{B}_1)$ -fixed vectors in V . It follows either from [BD, Corollaire 3.9] or, more explicitly, from [GSZ, 5.5] that $\mathcal{M}_0(A^\times)$ is closed under forming subquotients. Moreover,

$$(0.2) \quad \mathcal{M}_0(A^\times) = \bigoplus_{\Omega \text{ level zero}} \mathcal{M}(\Omega)$$

is a finite partial sum of (0.1). (For a general result see [BK2, beginning of Section 4, especially Theorem 4.3].) In the decomposition (0.2) the level zero components may be further represented as $\Omega = \Omega_{\mathcal{D}}$ for divisors $\mathcal{D} = \sum m_\pi \pi$ such that all π from the support have $(1 + \mathfrak{B}_1)$ -fixed vectors [GSZ, 5.5].

More precisely, consider the residual field k_D and the set $\mathcal{C}(k_D)$ of irreducible cuspidal representations of $GL_s(k_D)$ for all $s \geq 1$. With respect to the action of $\text{Gal}(k_D|k)$ on matrix elements in $GL_s(k_D)$, hence on the set of cuspidal representations, let $\tilde{\mathcal{C}}(k_D)$ denote the set of $\text{Gal}(k_D|k)$ orbits in $\mathcal{C}(k_D)$. Using [GSZ, (24) and 5.5(ii)], we have the bijection

$$(0.3) \quad \tilde{\mathcal{C}}(k_D) \longleftrightarrow \mathcal{C}_0 \quad [\sigma] \longmapsto \pi_{[\sigma]}$$

onto the subset of level zero unramified twist class representatives $\mathcal{C}_0 \subset \mathcal{C}$. The bijection is given by choosing one representative $\pi_{[\sigma]}$ from the uniquely determined unramified twist class of irreducible representations of $GL_s(D)$ which contain the representation σ of $GL_s(k_D)$. From (0.3) we obtain a bijection of divisors $\Delta \mapsto \mathcal{D}$, $\text{Div}^+(\tilde{\mathcal{C}}(k_D)) \xrightarrow{\sim} \text{Div}^+(\mathcal{C}_0)$ (see [GSZ, 5.3]). We write $\Omega_{\mathcal{D}} = \Omega_\Delta$ if \mathcal{D} corresponds to Δ and note that (0.2) becomes $\mathcal{M}_0(A^\times) = \bigoplus_{\Delta} \mathcal{M}(\Omega_\Delta)$, where Δ runs over the finite set of degree m divisors on $\tilde{\mathcal{C}}(k_D)$.

0.3 The Abstract Matching Theorem (AMT)

The Abstract Matching Theorem (AMT) of Deligne/Kazhdan/Vigneras [DKV], Rogawski [Ro], and Badulescu [Ba] asserts that for any two central simple F -algebras $A_1 \cong M_{m_1}(D_{d_1}), A_2 \cong M_{m_2}(D_{d_2})$ of reduced degree $n = d_1 m_1 = d_2 m_2$ there exists a bijective mapping, the ‘‘Jacquet-Langlands correspondence’’,

$$(0.4) \quad \mathcal{J}\mathcal{L}_{A_2, A_1} : \mathcal{R}^2(A_1^\times) \longrightarrow \mathcal{R}^2(A_2^\times) \quad \Pi^{A_2} := \mathcal{J}\mathcal{L}_{A_2, A_1}(\Pi^{A_1}),$$

where $\mathcal{R}^2(A_i^\times)$ denotes the set of discrete series representations of the unit group $A_i^\times = GL_{m_i}(D_{d_i})$ ($i = 1, 2$), such that the characters $\Theta_{\Pi^{A_i}}$ satisfy

$$(0.5) \quad (-1)^{m_1-1} \Theta_{\Pi^{A_1}}(x) = (-1)^{m_2-1} \Theta_{\Pi^{A_2}}(x)$$

for all regular elliptic x . (We assume an identification of the regular elliptic conjugacy classes among all the unit groups of reduced degree n simple F -algebras.)

0.4 The Weak Explicit Matching Theorem

Let $\mathcal{R}_0^2(A^\times) \subset \mathcal{R}^2(A^\times)$ denote the set of level zero discrete series representations and let Ω_Δ denote a level zero Bernstein component. In Proposition 2.8 we prove:

- (i) $\text{Irr}(\Omega_\Delta) \cap \mathcal{R}_0^2(A^\times) \neq \emptyset \iff \Delta = r[\sigma] \in \text{Div}^+(\tilde{\mathcal{C}}(k_D))$ is a simple degree m divisor, *i.e.*, $rd([\sigma]) = m$.
- (ii) For Δ a simple divisor as in (i) let $\mathcal{S}_\Delta^A := \text{Irr}(\Omega_\Delta) \cap \mathcal{R}_0^2(A^\times)$. Then \mathcal{S}_Δ^A is comprised of precisely one unramified twist class of discrete series representations.

We say that two representations $\Pi, \Pi' \in \mathcal{R}^2(A^\times)$ are *inertially equivalent* and write $\Pi \sim \Pi'$ if there exists an unramified character μ of A^\times such that $\Pi = \Pi' \otimes \mu$.

To prove (ii) as well as a certain multiplicity one statement for discrete series representations we reduce to the unramified split case where these facts are well known. For this we use our description of the Hecke algebras of level zero types [GSZ] and establish a reformulation of [BK1, (7.7.5)] for our context. The Appendix is devoted to these reformulations of Bushnell/Kutzko’s work.

Now let A_1, A_2 be as in Section 0.3; in particular, assume that $d_1 m_1 = d_2 m_2 = n$. To give a natural identification of the simple degree m_1 divisors on $\tilde{\mathcal{C}}(k_{d_1})$ with the simple degree m_2 divisors on $\tilde{\mathcal{C}}(k_{d_2})$ we use Proposition 1.1, which constructs out of Green’s parameterization of the cuspidal representations of the general linear group over a finite field a natural bijection

$$(0.6) \quad \langle \phi \rangle \setminus X(k_n^\times) \leftrightarrow \text{the set of simple degree } m \text{ divisors on } \tilde{\mathcal{C}}(k_d)[\bar{\chi}] \mapsto \Delta([\bar{\chi}])$$

for any factorization $n = dm$. The left side consists of the set of $\text{Gal}(k_n|k)$ orbits of characters of the multiplicative group k_n^\times . Using (0.6) we may set $\mathcal{S}_\chi^A := \mathcal{S}_{\Delta([\bar{\chi}]})^A$ and this is meaningful for any algebra $A = M_m(D_d)$ of reduced degree n .

Our main result, the “Weak Explicit Matching Theorem”, see Theorem 4.1, asserts that the Jacquet-Langlands correspondence (0.4) restricts to a bijection

$$\mathcal{JL}_{A_2, A_1} : \mathcal{R}_0^2(A_1^\times) \longrightarrow \mathcal{R}_0^2(A_2^\times)$$

of level zero discrete series sets such that $\mathcal{S}_\chi^{A_1}$ maps to $\mathcal{S}_\chi^{A_2}$ for all $[\bar{\chi}] \in \langle \phi \rangle \setminus X(k_n^\times)$. We actually prove a slightly finer result: Since AMT implies that \mathcal{JL}_{A_2, A_1} preserves central characters, we have partitions of $\mathcal{S}_\chi^{A_1}, \mathcal{S}_\chi^{A_2}$ into finite subsets $\mathcal{S}_\chi^{A_1}, \mathcal{S}_\chi^{A_2}$, respectively, of cardinality $e := n/f, f := |[\bar{\chi}]|$, where $\chi \in X_i(F_n^\times)$ has the reduction $\bar{\chi}$ and $\chi_F := \chi|_{F^\times}$ is the common central character of all the representations in $\mathcal{S}_\chi^{A_1} \cup \mathcal{S}_\chi^{A_2}$. We prove that $\mathcal{JL}(\mathcal{S}_\chi^{A_1}) = \mathcal{S}_\chi^{A_2}$.

Let us conclude this section by explaining why Theorem 4.1 is called “explicit”. Let $[\bar{\chi}] \in \langle \phi \rangle \setminus X(k_n^\times)$ correspond to the simple degree m divisor $r[\sigma]$ on $\tilde{\mathcal{C}}(k_d)$ (σ constructed via the correspondence (0.6); *cf.* Proposition 1.1) and let $A = M_m(D_d)$. Then we give two characterizations of the representations $\Pi \in \mathcal{S}_\chi^A$, which distinguish this class from all other discrete series representations of A^\times , each characterization being sufficient to determine the inertial class of $\Pi \in \mathcal{R}_0^2(A^\times)$:

- (a) Let \mathfrak{A}_r be the standard principal order of period r in A ; thus $\mathfrak{A}_r^\times / (1 + \mathfrak{P}_r) \cong [\mathrm{GL}_s(k_d)]^r$ ($m = rs$). Then $\Pi \in \mathcal{S}_{\bar{\chi}}^A$ if and only if Π contains the lift of $\sigma^{\otimes r}$ to an irreducible representation of \mathfrak{A}_r^\times .
- (b) Let M be the standard block diagonal Levi subgroup $[\mathrm{GL}_s(D)]^r$ of A^\times . Then $\Pi \in \mathcal{S}_{\bar{\chi}}^A$ if and only if the supercuspidal support of Π is the conjugacy class of an unramified twist of the representation $\pi_{[\sigma]}^{\otimes r}$ of M . By [GSZ, 5.1], the supercuspidal representation $\pi_{[\sigma]}$ of $\mathrm{GL}_s(D)$ is “explicitly” given by induction from a compact mod center subgroup of A^\times .

Finally, Theorem 4.1 is “weak” in the sense that it is explicit only up to the inertial ambiguities which remain to be resolved.

1 Parameters for the Set of Level Zero Simple Components of $\Omega(A^\times)$

In Proposition 1.1 we prove (0.6), which gives a natural parameterization for the set of Bernstein components corresponding to level zero simple divisors. Only such components can have level zero discrete series subquotients associated to them.

Proposition 1.1 *Let $k_n|k$ be a degree n extension of finite fields with generating automorphism ϕ and let $n := dm$. Then Green’s parameterization of the cuspidal representations of finite general linear groups in terms of orbits of regular characters [Gr] induces a natural bijection*


$$(1.1) \quad \mathcal{G}_m: \langle \phi \rangle \backslash X(k_n^\times) \xrightarrow{\sim} \bigsqcup_{s|m} \mathrm{GL}_s(k_d)_{\mathrm{cusp}}^\wedge / \langle \phi \rangle$$

between the set of $\langle \phi \rangle$ -orbits in the group of characters $X(k_n^\times)$ and the set of orbits of cuspidal representations of the groups $\mathrm{GL}_s(k_d)$ for all $s|m$. If $[\bar{\chi}]$ is a character orbit of length f and $f \mid n$, then the corresponding Galois orbit of cuspidal representations $[\sigma]$ consists of representations of $\mathrm{GL}_{f'}(k_d)$ ($f' := f/(d, f)$) and the length of $[\sigma]$ is (d, f) . In particular, $||[\bar{\chi}]|| = d(\sigma)||[\sigma]|| = f'(d, f)$, $d(\sigma) = f'$ being the degree of the divisor σ .

Remarks

- (i) For $m = 1$ and $d = n$ the correspondence \mathcal{G}_1 is the identity. For $m = n$ and $d = 1$, since there is no Galois action on the right side of (1.1), we see that \mathcal{G}_n gives the usual Green’s parameterization of the cuspidal representations of $\mathrm{GL}_s(k)$ for $s \mid n$.
- (ii) On the right in (1.1) it would be enough to let ϕ be a generator of the Galois group $\mathrm{Gal}(k_d|k)$.

Proof Given a $\langle \phi \rangle$ -orbit $[\bar{\chi}] \subset X(k_n^\times)$ of length f we write $\text{lcm}(d, f) = df'$, and we use the following diagram of finite fields:



Since $f \mid df'$ and $df' \mid n$, for any $\bar{\chi} \in X(k_n^\times)$ such that $[\bar{\chi}] = f$, we have a unique k_d -regular character $\bar{\chi}_{df'} \in X(k_{df'}^\times)$ such that $\bar{\chi}_{df'} \circ N_{k_n|k_{df'}} = \bar{\chi}$. Therefore, by [Gr], we also have the unique cuspidal representation $\sigma = \sigma(\bar{\chi}_{df'})$ of $\text{GL}_{f'}(k_d)$ which has the character values $\theta_\sigma(x) = (-1)^{f'-1} \sum_{\bar{\eta} \in \text{Gal}(k_{df'}|k_d) \cdot \bar{\chi}_{df'}} \bar{\eta}(x)$ for k_d -regular $x \in k_{df'}^\times$. Observing that the bijection of Green $\langle \phi^d \rangle \setminus X(k_{df'}^\times)_{k_d\text{-reg}} \xrightarrow{\sim} \text{GL}_{f'}(k_d)_{\text{cusp}}^\wedge$, which maps orbits of k_d -regular characters in $X(k_{df'}^\times)$ to cuspidal representations of $\text{GL}_{f'}(k_d)$, is compatible with the action of $\langle \phi \rangle$ on both sides, we see that the $\langle \phi \rangle$ -orbit $[\sigma(\bar{\chi}_{df'})]$ has length (d, f) . Conversely, an orbit $[\sigma] \in \text{GL}_s(k_d)_{\text{cusp}}^\wedge / \langle \phi \rangle$ determines an orbit of k_d -regular characters $[\bar{\psi}_\sigma] \in \langle \phi \rangle \setminus X(k_{df'}^\times)$ and a unique orbit $[\bar{\chi}_\sigma] \subset X(k_n^\times)$ such that $\bar{\chi}_\sigma = \bar{\psi}_\sigma \circ N_{k_n|k_{df'}}$. These orbits are of the same length; specifically, we have $|[\bar{\psi}_\sigma]| = |[\bar{\chi}_\sigma]| = s|[\sigma]|$. ■

For $\bar{\chi} \in X(k_n^\times)$ we write $e := n/f$ and $e' := (e, m)$. From the equation $dm = ef$ it follows that $m = e'f'$. Via \mathcal{G}_m the character orbit $[\bar{\chi}]$ determines the simple divisor $e'[\sigma]$, which has the degree m .

Now consider the central simple algebra $A = M_m(D)$, $D := D_d$ a central division algebra of index d over F . From [GSZ, (24)] (see (0.3)) we know how to associate to $[\sigma]$ a unitary supercuspidal representation $\pi_{[\sigma]} \in \mathcal{C}(D)$ such that $d(\pi_{[\sigma]}) = d([\sigma])$. Thus $[\bar{\chi}]$ determines both a degree m divisor $\mathcal{D}_m([\bar{\chi}]) = e'\pi_{[\sigma]} \in \text{Div}(\mathcal{C})$ and a simple level zero component $\Omega_{[\bar{\chi}]} := \Omega_{\mathcal{D}_m([\bar{\chi}])} \subset \Omega(A^\times)$.

Proposition 1.2 *The mapping $\langle \phi \rangle \setminus X(k_n^\times) \ni [\bar{\chi}] \mapsto \Omega_{[\bar{\chi}]} = \Omega_{\mathcal{D}_m([\bar{\chi}])}$ gives a bijection from $\langle \phi \rangle \setminus X(k_n^\times)$ to the set of simple level zero connected components of $\Omega(A^\times)$. If $[\bar{\chi}]$ corresponds to the degree m divisor $e'[\sigma]$, then $(\mathfrak{A}_{e'}^\times, \sigma^{\otimes e'})$ is a type for the component $\Omega_{[\bar{\chi}]}$. If $\Pi \in \text{Irr}(\Omega_{[\bar{\chi}]})$ is an irreducible representation with cuspidal support in $\Omega_{[\bar{\chi}]}$, then its central character ω_Π is a tame character of F^\times with $\bar{\omega}_\Pi = \bar{\chi}|_{k^\times} \in X(k^\times)$.*

Proof The proof follows from [GSZ, 5.4 and 5.5]. In particular, the central character ω_Π is tame because Π is a level zero representation and the reduction $\bar{\omega}_\Pi$ is constructed by restricting to the units of F^\times and factoring mod the principal units. It is obviously determined by the type. From Proposition 1.1 we obtain $\bar{\omega}_\Pi = (\bar{\chi}_{df'})^{e'}|_{k^\times} = \bar{\chi}|_{k^\times}$. ■

2 A Parameterization of the Set of Inertial Classes $\mathcal{R}_0^2(A^\times)/\sim$

We shall continue to use a “bar” over a Greek letter to denote a multiplicative character of a finite field; e.g. if χ is a tame multiplicative character of a p -adic field, then $\bar{\chi}$ usually will denote its “reduction”, i.e., the reduction of its restriction to the group of units.

Proposition 2.1 *Let $\Omega = \Omega(\mathcal{D})$ be a level zero Bernstein component and assume that \mathcal{D} is not a simple divisor. Then $\mathcal{R}^2(A^\times) \cap \text{Irr}(\Omega) = \emptyset$.*

Proof Let $\Omega \subset \Omega(A^\times)$ be a level zero connected component. Assume that $\Omega = \Omega(\mathcal{D})$, where the degree m divisor \mathcal{D} is not simple. By [GSZ, 5.3 and 5.4], the types $(\mathfrak{A}^\times, \tau)$ associated to Ω all have the same divisor $\Delta(\tau) = \sum_{[\sigma]} r_{[\sigma]}(\tau)[\sigma]$ with $r_{[\sigma]} > 0$ for at least two distinct orbits $[\sigma]$. Then we consider the proper Levi subgroup $\tilde{M} = \prod_{[\sigma]} \text{GL}_{r_{[\sigma]}(\tau) \cdot d(\sigma)}(D) \subset A^\times$. Using the Hecke algebra isomorphism [GSZ, 1.9] we obtain from [BK2, (8.4)] that the parabolic induction from \tilde{M} to G is an equivalence between the categories $\mathcal{M}_{\tilde{M}}(\Omega)$ and $\mathcal{M}_G(\Omega)$, where $\Omega = \Omega(\mathcal{D})$ is interpreted as a connected component of the Bernstein spectrum $\Omega(\tilde{M})$ in the obvious way. This implies that $\Pi \in \text{Irr}(\Omega)$ is always parabolically induced from \tilde{M} and therefore Π cannot be a discrete series. Indeed either by AMT (or, to give a local argument, by Harish-Chandra’s Selberg principle) the character of a discrete series representation cannot completely vanish on the regular elliptic set, whereas the character of a parabolically induced representation does vanish for regular elliptic elements. ■

Corollary 2.2 *If $\mathcal{R}_0^2(A^\times) \cap \text{Irr}(\Omega) \neq \emptyset$, then $\Omega = \Omega_{[\bar{\chi}]}$ for some Galois orbit $[\bar{\chi}] \in X(k_n^\times)$.*

Proof From Proposition 2.1 we know that, if $\mathcal{R}_0^2(A^\times) \cap \Omega \neq \emptyset$, then $\Omega = \Omega(\mathcal{D})$ for some simple divisor \mathcal{D} . From Proposition 1.2 we have a parameterization of the set of all simple level zero components such that each is of the form $\Omega_{[\bar{\chi}]}$ for some $\bar{\chi} \in X(k_n^\times)$. ■

Definition 2.3 We write $\mathcal{S}_{\bar{\chi}}^A \subset \text{Irr}(\Omega_{[\bar{\chi}]})$ for the set of irreducible discrete series representations which have supercuspidal support in $\Omega_{[\bar{\chi}]}$.

In the following we are working toward Proposition 2.8, toward proving that in the level zero case $\mathcal{S}_{\bar{\chi}}^A = \mathcal{R}_0^2(A^\times) \cap \text{Irr}(\Omega_{[\bar{\chi}]})$ is not empty and consists of a single unramified twist class. As already mentioned in Section 0.4 our proof of these assertions makes use of level zero types and their Hecke algebras; it does not depend upon AMT. The notation is as explained in the paragraph prior to Proposition 1.2.

Proposition 2.4 *Assume that $\mathcal{D}_m([\bar{\chi}]) = e' \pi_{[\sigma]}$, i.e., $\Omega_{[\bar{\chi}]} \subset \Omega(A^\times)$ is the connected component which contains the cuspidal pair $(M, \pi_{[\sigma]}^{\otimes e'})$ with respect to the Levi subgroup $M = [\text{GL}_{f'}(D)]^{\times e'} \subset A^\times$. For any $\Pi \in \mathcal{S}_{\bar{\chi}}^A$ the normalized Jacquet functor $r_{M, A^\times}(\Pi)$ is irreducible; specifically, $r_{M, A^\times}(\Pi) = \pi_{[\sigma]}^{\otimes e'} \cdot \psi$ for some unramified character ψ of M which is not unitary.*

Proof The type $(\mathfrak{A}_{e'}^\times, \tau)$ for $\Omega_{[\bar{\chi}]}$ is a cover of $(M \cap \mathfrak{A}_{e'}^\times, \sigma^{\otimes e'})$, which is a type for the supercuspidal component $\Omega_{[\bar{\chi}]}(M) \subset \Omega(M)$ (see [GSZ, 5.3 and its proof]). Therefore, by [BK2, (7.9)(ii)], we have a natural isomorphism of complex vector spaces

$$(2.1) \quad \text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \Pi) \cong \text{Hom}_{\mathfrak{A}_{e'}^\times \cap M}(\tau, r_{M, A^\times}(\Pi)).$$

Let ω denote the central character of Π . Then we also have the natural isomorphism

$$(2.2) \quad \text{Hom}_{\mathfrak{A}_{e'}^\times \langle \varpi_F \rangle}(\tau_\omega, \Pi) \cong \text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \Pi).$$

Since Π is irreducible, $\text{Hom}_{\mathfrak{A}_{e'}^\times \langle \varpi_F \rangle}(\tau_\omega, \Pi)$ is a finite-dimensional simple module over the Hecke algebra $\mathcal{H}(A^\times, \mathfrak{A}_{e'}^\times \langle \varpi_F \rangle, \tau_\omega)$. Since our type has the form $\tau = \sigma^{\otimes e'}$ the considerations of the appendix can be applied (see the text before (a.2)). In particular we may apply Proposition A9 with $(K, \rho) = (\mathfrak{A}_{e'}^\times \langle \varpi_F \rangle, \tau_\omega)$ because Π is a discrete series representation. This implies that the modules in (2.2) are complex vector spaces of dimension one.

Considering now the right side of (2.1), we see that $r_{M, A^\times}(\Pi)$ contains the type $(M \cap \mathfrak{A}_{e'}^\times, \tau)$ with multiplicity one and therefore $r_{M, A^\times}(\Pi)$ contains an irreducible component $\pi_{[\sigma]}^{\otimes e'} \cdot \psi$ and indeed only one. The character ψ cannot be unitary, since if it were unitary, then Π would be a subrepresentation of a unitarily induced representation and therefore not a discrete series representation. Moreover, any other component of $r_{M, A^\times}(\Pi)$ has to be supercuspidal and conjugate to $\pi_{[\sigma]}^{\otimes e'} \cdot \psi$ under the normalizer of M , i.e., it has to be of the form $\pi_{[\sigma]}^{\otimes e'} \cdot \psi'$ (see e.g. [BR, III.2 Theorem 18]). Any such component contains $\tau|_{M \cap \mathfrak{A}_{e'}^\times}$. By [BK2, (7.9)] this is impossible since τ occurs as a component of $\Pi|_{\mathfrak{A}_{e'}^\times}$ only once. Thus $r_{M, A^\times}(\Pi) = \pi_{[\sigma]}^{\otimes e'} \cdot \psi$ and is irreducible. ■

Remark 2.5 Since $r_{M, A^\times}(\Pi)$ is irreducible, it belongs to the supercuspidal support of Π . Therefore, we have $r_{M, A^\times}(\Pi) = (\pi_{[\sigma]} \tilde{\eta}')^{\otimes e'} \psi_1$, where ψ_1 is a positive, real-valued character and $\tilde{\eta}'$ is a unitary unramified character.

Let $(\mathfrak{A}_{e'}^\times, \tau)$ be the type such that $\tau = \sigma^{\otimes e'}$ and set $\tau^{\nu_1, \dots, \nu_{e'}} := \sigma^{\phi^{\nu_1}} \otimes \dots \otimes \sigma^{\phi^{\nu_{e'}}}$. The set of tuples $(\nu_1, \dots, \nu_{e'}) \in (\mathbb{Z}/(d, f)\mathbb{Z})^{e'}$ parameterizes the $(d, f)^{e'}$ conjugates of τ .

Corollary 2.6 Let $(\Pi, V) \in S_{\bar{\chi}}^A$ and let $V^{1+\mathfrak{F}_{e'}}$ denote the subspace of V consisting of all $(1 + \mathfrak{F}_{e'})$ -fixed vectors, where $\mathfrak{F}_{e'}$ denotes the Jacobson radical of $\mathfrak{A}_{e'}$. As a representation of $\mathfrak{A}_{e'}^\times$ the space $V^{1+\mathfrak{F}_{e'}}$ decomposes as $\bigoplus \tau^{\nu_1, \dots, \nu_{e'}}$, where, for $0 \leq \nu_1, \dots, \nu_{e'} < (d, f)$, each representation occurs with multiplicity one.

Proof From [GSZ, 5.5(i)] we know that each of the above types occurs and with the same multiplicity. As we have seen in the proof of Proposition 2.4, τ occurs simply in $\Pi|_{\mathfrak{A}_{e'}^\times}$. Therefore, the same is true for each of the types $\tau^{\nu_1, \dots, \nu_{e'}}$. ■

We consider the unramified extension $F_n|F$ of degree n and we write $X_t(F_n^\times)$ for the set of tamely ramified characters of F_n^\times . Then we have a reduction map

$$(2.3) \quad X_t(F_n^\times) \ni \chi \longmapsto \bar{\chi} \in X(k_n^\times),$$

where $\bar{\chi}$ denotes $\chi|_{o_n^\times}$ regarded as a character of k_n^\times . Here we identify k_n with the residual field of F_n and we use the assumption that χ is a tame character. Under these assumptions $\bar{\chi}$ is well defined; we call $\bar{\chi}$ the reduction of χ . Noting that F_n has a prime element in F , we see that (2.3) is compatible with the Galois action of $\text{Gal}(F_n|F) \cong \text{Gal}(k_n|k) = \langle \phi \rangle$, the lengths of orbits being preserved. Moreover, the character $\chi \in X_t(F_n^\times)$ is uniquely determined by its reduction $\bar{\chi} \in X(k_n^\times)$ and its restriction $\chi_F \in X_t(F^\times)$. For $\Pi \in \mathcal{S}_\chi^A$ it follows from Definition 2.3 and from the last statement of Proposition 1.2 that the central character $\omega_\Pi \in X_t(F^\times)$ has the reduction $\bar{\omega}_\Pi = \bar{\chi}|_{k^\times}$. From this we obtain a partition $\mathcal{S}_\chi^A = \bigcup_\chi \mathcal{S}_\chi^A$, where $\chi \in X_t(F_n^\times)$ runs over all tame characters which have the prescribed reduction $\bar{\chi}$:

Definition 2.7 For $\chi \in X_t(F_n^\times)$ set $\mathcal{S}_\chi^A := \{\Pi \in \mathcal{S}_\chi^A : \omega_\Pi = \chi_F\}$.

Like $\mathcal{S}_{\bar{\chi}}^A$, cf. Definition 2.3, the definition of \mathcal{S}_χ^A depends only upon the $\langle \phi \rangle$ -orbit of χ .

Proposition 2.8 If $\chi \in X_t(F_n^\times)$ generates a $\langle \phi \rangle$ -orbit $[\chi]$ of length f , then \mathcal{S}_χ^A is a set of order $e = n/f$. In particular, the group of unramified characters of order dividing n acts transitively on \mathcal{S}_χ^A . The set \mathcal{S}_χ^A is a single unramified twist class.

Proof We note that \mathcal{S}_χ^A consists of all $\Pi \in \mathcal{S}_\chi^A$ which have a fixed central character. By Proposition 1.2 the representations contained in \mathcal{S}_χ^A admit the type $(\mathfrak{A}_{e'}^\times, \tau)$, where $\tau = \sigma^{\otimes e'}$. Therefore, we may apply Corollary A10 with $r = e'$ and $l = (d, f)$, since Proposition 1.1 implies that $(d, f) = |[\sigma]|$. Since $dr/l = de'/(d, f) = e$, it follows that \mathcal{S}_χ^A consists of e representations. If $\eta \in X(F^\times)$ and $\bar{\eta} = \eta \circ \text{Nrd}_{A|F}$, then we know that $\omega_{\Pi \otimes \bar{\eta}} = \omega_\Pi \otimes \eta^n$, which implies that the group of unramified characters η of order dividing n acts on each set \mathcal{S}_χ^A . Therefore, it suffices to show that, for $\Pi \in \mathcal{S}_\chi^A$, we have $\Pi \otimes \bar{\chi}_F = \Pi$ if and only if $\chi_F^f = 1$. If this is so, then, since $n/f = e$, it will follow that all e inequivalent discrete series representations with the same central character lie in a single unramified twist class. Since the group of unramified characters of A^\times transitively permutes the set of all \mathcal{S}_χ^A for which the reduction of χ is $\bar{\chi}$, it will follow that, if \mathcal{S}_χ^A is only one unramified twist class for each χ , then the same is true for $\mathcal{S}_{\bar{\chi}}^A$.

Let us proceed with the proof that for an unramified character $\eta \in X(F^\times)$ we have $\Pi \otimes \bar{\eta} = \Pi$ if and only if $\eta^f = 1$. From Remark 2.5 and the comments above Proposition 1.2, which explain how to construct a degree m simple divisor from a cuspidal divisor $e'[\sigma]$, where $e'd(\sigma) = e'f' = m$, it follows that the supercuspidal support of $\Pi \in \mathcal{S}_\chi^A$ has the form $\pi_{[\sigma]}^{\otimes e'} \cdot \psi$, where ψ is some non-unitary unramified character of $M = (\text{GL}_{f'}(D))^{\times e'}$. By Jacquet's subrepresentation theorem $\Pi \subset \text{Ind}_P^{A^\times} (\pi_{[\sigma]}^{\otimes e'} \cdot \psi)$, where $P = M \rtimes U_P$ is a parabolic subgroup of A^\times and the induction is normalized.

Thus, $\Pi \cdot \tilde{\eta}$ is contained in $\text{Ind}_P^{A^\times} (\pi_{[\sigma]}^{\otimes e'} \cdot \psi \cdot \tilde{\eta}|_M)$. From the irreducibility of the Jacquet module we see that $\Pi \otimes \tilde{\eta} = \Pi$ implies that

$$(2.4) \quad \pi_{[\sigma]}^{\otimes e'} = \pi_{[\sigma]}^{\otimes e'} \tilde{\eta}|_M.$$

Furthermore $\text{Nrd}_{A|F}|_M = (\text{Nrd}_{f'})^{\otimes e'}$, where $\text{Nrd}_{f'}$ denotes the reduced norm mapping $M_{f'}(D)|_F$. Therefore from (2.4) we obtain $\pi_{[\sigma]} = \pi_{[\sigma]} \cdot (\eta \circ \text{Nrd}_{f'})$ for the supercuspidal representation $\pi_{[\sigma]}$ of $\text{GL}_{f'}(D)$. With this justification we reduce to the supercuspidal case $\Pi = \pi_{[\sigma]}$, $f' = m$, and $e' = 1$. In this case from [GSZ, 5.1] we know that $\Pi = \text{cInd}(\tilde{\sigma})$, where $\tilde{\sigma}$ is an extension of σ to $\langle \varpi^l \rangle \mathfrak{A}_1^\times$. Here and in [GSZ] $l = |[\sigma]|$ (see Section A.1). Moreover we have seen (*loc. cit.*) that $\tilde{\sigma} \mapsto \text{cInd}(\tilde{\sigma})$ is a bijection between the possible extensions of σ and the unramified twists of $\pi_{[\sigma]} \in \mathcal{C}_0$ (see (0.3)). Therefore $\tilde{\eta} \otimes \text{cInd}(\tilde{\sigma}) = \text{cInd}(\tilde{\eta} \otimes \tilde{\sigma})$ also equals $\text{cInd}(\tilde{\sigma})$ if and only if $\tilde{\eta} \otimes \tilde{\sigma} = \tilde{\sigma}$, so we must consider representations of $\langle \varpi^l \rangle \times \mathfrak{A}_1^\times$. Note that $l = |[\sigma]| = (d, f)$, since σ corresponds to $[\bar{\chi}]$ of length f (see Proposition 1.1). The reduced norm $\text{Nrd}_{A|F}$ induces an injection $\langle \varpi^l \rangle \mathfrak{A}_1^\times / \mathfrak{A}_1^\times \hookrightarrow F^\times / \mathfrak{o}_F^\times$ with image of index $ml = f'(d, f) = f$. Therefore $\tilde{\eta} \otimes \tilde{\sigma} = \tilde{\sigma}$, if and only if $\eta^f = 1$. ■

Writing $\mathcal{R}_0^2(A^\times) / \sim$ for the set of inertial equivalence classes of level zero discrete series representations, we now have the bijection

$$(2.5) \quad \langle \phi \rangle \setminus X(k_n^\times) \ni \bar{\chi} \longmapsto \mathcal{S}_\chi^A \in \mathcal{R}_0^2(A^\times) / \sim.$$

In concluding this section, we lift (2.5) to a bijection

$$\text{Gal}(F_n|F) \setminus X_t(F_n^\times) \ni [\chi] \longmapsto \mathcal{S}_\chi^A \in \mathcal{R}_0^2(A^\times) / \approx,$$

where we set $\Pi_1 \approx \Pi_2$ if $\Pi_1 = \Pi_2 \otimes \eta$ for some unramified character η of A^\times which is trivial on the center of A^\times . To formulate our result, which summarizes what has been proved to this point, we also extend $(\mathfrak{A}_{e'}^\times, \tau)$ to $(\mathfrak{A}_{e'}^\times F^\times, \tau_{\chi_F})$, τ_{χ_F} being the extension of $\tau = \sigma^{\otimes e'}$ to $\mathfrak{A}_{e'}^\times \langle \varpi_F \rangle = \mathfrak{A}_{e'}^\times F^\times$ such that $\chi_F 1_\tau$ is the restriction of τ_{χ_F} to F^\times .

Theorem 2.9 *Let $\chi \in X_t(F_n^\times)$ be a unitary character generating the $\text{Gal}(F_n|F)$ -orbit $[\chi]$ of length f and set $\chi_F := \chi|_{F^\times}$.*

- (i) *There are precisely $e = n/f$ discrete series representations Π_1, \dots, Π_e of A^\times with central character $\omega_{\Pi_i} = \chi_F$ such that $\Pi_i \in \mathcal{S}_\chi^A$, i.e., such that $\Pi_i|_{\mathfrak{A}_{e'}^\times F^\times} \supset \tau_{\chi_F}$. This set of representations comprises an equivalence class $\mathcal{S}_\chi^A \in \mathcal{R}_0^2(A^\times) / \approx$. The finite set \mathcal{S}_χ^A consists of the set of all elements in the inertial class \mathcal{S}_χ^A with the central character χ_F .*
- (ii) *The representations $\Pi_i \in \mathcal{S}_\chi^A$ have supercuspidal support in $\Omega_{[\bar{\chi}]}$. In particular, these representations are supercuspidal if and only if $e' = 1$.*
- (iii) *The map $\chi \mapsto \mathcal{S}_\chi^A$ induces a bijection $\text{Gal}(k_n|k) \setminus X_t(F_n^\times) \leftrightarrow \mathcal{R}_0^2(A^\times) / \approx$ which lifts the bijection (2.5).*

Remark 2.10 In the case $A = D_n$ the injection

$$(2.6) \quad F_n^\times / (1 + \mathfrak{p}_{F_n}) \xrightarrow{\sim} O_{D_n}^\times F^\times / (1 + \mathfrak{p}_{D_n}) \hookrightarrow D_n^\times / (1 + \mathfrak{p}_{D_n})$$

gives $\tau_\chi = \chi \in X_t(F_n^\times)$ as a character of $O_{D_n}^\times F^\times / (1 + \mathfrak{p}_{D_n})$. Thus we may regard $\mathbb{S}_\chi^{D_n}$ as the set of irreducible components of the representation $\text{Ind}_{O_{D_n}^\times F^\times}^{D_n^\times}(\chi)$.

3 The Characters $\mathcal{R}_0^2(A^\times)$ at Very Regular Elements

Let $(\Pi, V) \in \mathcal{R}_0^2(A^\times)$ and assume that the supercuspidal support of Π lies in $\Omega_{[\bar{\chi}]}$. Let $(\mathfrak{A}_{e'}^\times, \tau)$, where $\tau = \sigma^{\otimes e'}$, be the corresponding type. In Corollary 2.6 we considered the $\mathfrak{A}_{e'}^\times$ representation on $V^{1+\mathfrak{P}_{e'}}$. Now we want to study the \mathfrak{A}_1^\times representation on $V^{1+\mathfrak{P}_1}$. As always $\mathcal{M}(G)$ denotes the category of (smooth) representations of a group G .

Lemma 3.1

- (i) *The Jacquet restriction mapping $r_{\mathfrak{A}_1^\times, \mathfrak{A}_{e'}^\times} : \mathcal{M}(\mathfrak{A}_1^\times) \rightarrow \mathcal{M}(\mathfrak{A}_{e'}^\times)$ induces a bijection between the set of irreducible constituents of $V^{1+\mathfrak{P}_1}$ and the set of $\mathfrak{S}_{e'}$ -orbits of irreducible constituents of $V^{1+\mathfrak{P}_{e'}}$. In particular, $V^{1+\mathfrak{P}_1}$ decomposes simply.*
- (ii) *If an $\mathfrak{S}_{e'}$ -orbit in $V^{1+\mathfrak{P}_{e'}}$ has length greater than one, then the corresponding constituent of $V^{1+\mathfrak{P}_1}$ is induced from a proper parabolic subgroup of \mathfrak{A}_1^\times and its character vanishes on the regular elliptic set of \mathfrak{A}_1^\times .*
- (iii) *If an orbit has length equal one, then the corresponding constituent of $V^{1+\mathfrak{P}_1}$ is a generalized Steinberg representation with cuspidal support $(\sigma^{\phi'})^{\otimes e'}$.*

Proof (i) Since $(\mathfrak{A}_{e'}^\times, \tau)$ is a type in (Π, V) and any cuspidal level zero type of (Π, V) corresponds, up to conjugacy, to a representation of $\mathfrak{A}_{e'}^\times$, the mapping $r := r_{\mathfrak{A}_1^\times, \mathfrak{A}_{e'}^\times}$ does not annihilate any irreducible \mathfrak{A}_1^\times -submodule of $V^{1+\mathfrak{P}_1}$. Obviously, we may regard $V^{1+\mathfrak{P}_{e'}} = r(V^{1+\mathfrak{P}_1})$. Therefore, since, by Corollary 2.6, the $\mathfrak{A}_{e'}^\times$ -module $V^{1+\mathfrak{P}_{e'}}$ decomposes simply, it follows that the \mathfrak{A}_1^\times -module $V^{1+\mathfrak{P}_1}$ decomposes simply too. Let π be an irreducible \mathfrak{A}_1^\times representation occurring in $V^{1+\mathfrak{P}_1}$. Every constituent of $r(\pi)$ occurs with multiplicity one and if π' is any other constituent of $V^{1+\mathfrak{P}_1}$, then $r(\pi)$ and $r(\pi')$ are disjoint. By Frobenius reciprocity a constituent τ' of $V^{1+\mathfrak{P}_{e'}}$ occurs in $r(\pi)$ if and only if $\pi \subset \text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau'$. For $i = 1, 2$ let $\tau_i = \sigma_1^i \otimes \dots \otimes \sigma_{e'}^i$ be a constituent of $V^{1+\mathfrak{P}_{e'}}$. Let $T_i := \{\sigma_1^i, \dots, \sigma_{e'}^i\}$ be the corresponding multiset, $i = 1, 2$. If $T_1 \neq T_2$ (i.e., the two representations τ_1 and τ_2 are not just rearrangements of the same tensor factors), then, as is well known, it follows from the Bruhat decomposition of the group \mathfrak{A}_1^\times , the two induced representations $\text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_i$ are disjoint. Therefore the set of constituents of $r(\pi)$ lies in a single $\mathfrak{S}_{e'}$ -orbit, i.e., π has associated to it by r a single multiset T . On the other hand, assume that $T_1 = T_2$ and assume that $\sigma_k^1 = \sigma_k^2$ for all $k \neq j, j+1$ and that $\sigma_j^i = \sigma_{j+1}^{i'}$ for $i \neq i'$ and $1 \leq i, i' \leq 2$. Let $\mathfrak{U}^j \subset M_{e'}(M_{f'}(O_D))$ be the standard hereditary order which differs from $\mathfrak{A}_{e'}$ only in that the block $j+1, j$ is allowed to consist of arbitrary matrices in $M_{f'}(O_D)$. Then $\mu := \text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_1 =$

$\text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_2$ is irreducible and, by transitivity of induction, $\text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_1 = \text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_2$; if $\pi \subset \text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \tau_i$, then $\mu \subset r_{\mathfrak{A}_1^\times, \mathfrak{A}_{e'}^\times}(\pi)$. Since $r_{\mathfrak{A}_1^\times, \mathfrak{A}_{e'}^\times}(\mu) = \tau_1 + \tau_2$, it follows that $r(\pi)$ contains τ_1 if and only if it contains τ_2 . Since the set of transpositions of the form $j \leftrightarrow j + 1$ ($1 \leq j < e'$) generate $\mathfrak{S}_{e'}$, it follows that the set of constituents $r(\pi)$ consists of one full $\mathfrak{S}_{e'}$ -orbit of constituents of $V^{1+\mathfrak{P}_{e'}}$.

(ii) First, if a representation of $\mathfrak{A}_1^\times = \text{GL}_m(k_d)$ is induced from a proper parabolic subgroup of $\text{GL}_m(k_d)$, then certainly its character vanishes on the regular elliptic set of $\text{GL}_m(k_d)$. To prove all of (ii) it is enough to check that the intertwining number of the induced representation $\text{Ind}_{\mathfrak{A}_{e'}^\times}^{\mathfrak{A}_1^\times} \sigma_1^{\otimes r_1} \otimes \dots \otimes \sigma_\ell^{\otimes r_\ell}$, where $\sigma_1, \dots, \sigma_\ell$ are inequivalent cuspidal representations of $\text{GL}_{f_j}(k_d)$ and $r_1 + \dots + r_\ell = e'$, is equal to the product $\prod_{j=1}^\ell |\mathfrak{S}_{r_j}|$. We omit this argument (see e.g. [DM, p. 58, 6.5]), which implies that the correct intertwining number is already obtained after inducing to $\prod_{j=1}^\ell \text{GL}_{r_j, f_j}(k_d)$; by transitivity of induction each component of the representation of $\prod_{j=1}^\ell \text{GL}_{r_j, f_j}(k_d)$ induces an irreducible representation of $\text{GL}_m(k_d)$, since further induction does not produce more intertwining.

(iii) If the irreducible constituent τ of $V^{1+\mathfrak{P}_{e'}}$ is fixed by $\mathfrak{S}_{e'}$, then from Corollary 2.6 and from (i) we conclude that $\tau = (\sigma^{\phi^\nu})^{\otimes e'}$ and $\tau = r(\pi)$ for some constituent π of $V^{1+\mathfrak{P}_1}$. Let ω be the central character of $(\Pi, V) \in \mathcal{R}_0^2(A^\times)$. Then obviously $\text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \pi) \subseteq \text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \Pi) = \text{Hom}_{\mathfrak{A}_{e'}^\times \langle \varpi_F \rangle}(\tau_\omega, \Pi)$. Again τ has the special form on which the considerations of the appendix apply (see the text before (a.2)). Therefore we may argue as below (2.2) that the right side of this inclusion has complex dimension one. This, together with the fact that the inclusion is non-zero, implies that the inclusion is an isomorphism of one-dimensional complex vector spaces. Now let $\mathcal{H} = \mathcal{H}(A^\times, \mathfrak{A}_{e'}^\times \langle \varpi_F \rangle, \tau_\omega)$ ($\omega = \omega_\Pi$), \mathcal{H}^2 the completion of \mathcal{H} (see (a.11) and subsequent remarks), $\mathcal{H}_0 = \mathcal{H}(\mathfrak{A}_1^\times \langle \varpi_F \rangle, \mathfrak{A}_{e'}^\times \langle \varpi_F \rangle, \tau_\omega)$, and $\bar{\mathcal{H}}_0$ the reduction of \mathcal{H}_0 modulo $(1 + \mathfrak{P}_1) \langle \varpi_F \rangle$. Since Π is discrete series, we see from Proposition A7 that the \mathcal{H}^{opp} -module $\text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \Pi)$ can be embedded into \mathcal{H}^2 . The module structure on $H = \text{Hom}_{\mathfrak{A}_{e'}^\times}(\tau, \Pi)$ carries over to a convolution action on a subspace of \mathcal{H}^2 . In the unramified split case this embedding sends H to the space $\mathbb{C} \cdot e_{\text{St}}$, where e_{St} denotes the \mathcal{J} bi-invariant matrix coefficient of a Steinberg representation. Via our isometric isomorphisms we bring this function back to a function, also to be denoted e_{St} , for the general case. For $\varphi \in \mathcal{H}$ we have $\varphi * e_{\text{St}} = \lambda(\varphi)e_{\text{St}}$, where $\lambda: \mathcal{H} \rightarrow \mathbb{C}$ is the character of \mathcal{H} associated to our one-dimensional module. By restriction we obtain a character λ_0 of \mathcal{H}_0 . Since convolution by $\varphi \in \mathcal{H}_0$ commutes with restriction to $\mathfrak{A}_1^\times \langle \varpi_F \rangle$, this character can also be represented by $\varphi * e_{\text{St},0} = \lambda_0(\varphi)e_{\text{St},0}$, where $e_{\text{St},0}$ denotes the restriction of e_{St} to $\mathfrak{A}_1^\times \langle \varpi_F \rangle$. Now we can pass to the reduction $\bar{\mathcal{H}}_0$ and observe that $\bar{\varphi} * e_{\text{st}} = \bar{\lambda}_0(\bar{\varphi})e_{\text{st}}$, where e_{st} denotes the reduction of $e_{\text{St},0}$. Since e_{st} is the Steinberg idempotent in $\bar{\mathcal{H}}_0$, we see from [SZ, 4.2] that the character $\bar{\lambda}_0$ which we obtained by restriction and reduction corresponds to a generalized Steinberg representation of the finite field group. This implies that the constituent π of $V^{1+\mathfrak{P}_1}$ is the generalized Steinberg representation of \mathfrak{A}_1^\times with cuspidal support $\tau = (\sigma^{\phi^\nu})^{\otimes e'}$. ■

Proposition 3.2 *The map $\mathcal{J}_{\mathcal{L}_{A,D_n}}: \mathcal{R}_0^2(D_n^\times) \rightarrow \mathcal{R}_0^2(A^\times)$ is a bijection.*

Proof based on AMT For the proof we want to compare the characters $\Pi^{D_n} \in S_\chi^{D_n}$ and $\Pi^A \in S_\chi^A$ at very regular elements.

First we recall that very regular elements $x \in A^\times$ are defined by the following properties:

- (i) $F[x]|F$ is an unramified extension of degree n in A ,
- (ii) x is a unit of $F[x]$, and
- (iii) the reduction \bar{x} of x modulo the prime ideal of $F[x]$ generates the residue class field extension $k_n|k$.

Let $x \in A^\times$ be very regular; we may assume that $x \in \mathfrak{A}_1^\times \subset A^\times$. Let $\varphi_{F,x} \in o[T]$ be the minimal polynomial of x over F and let $\bar{\varphi}_{F,x} \in k[T]$ be the reduction modulo \mathfrak{p}_F . Obviously for $\bar{x} = x \pmod{\mathfrak{P}_1}$ we have $\bar{\varphi}_{F,x}(\bar{x}) = \overline{\varphi_{F,x}(x)} = 0 \in \mathfrak{A}_1/\mathfrak{P}_1$. Therefore $\bar{\varphi}_{F,x'} = \bar{\varphi}_{F,x}$ for all $x' \in x(1 + \mathfrak{P}_1) = x + \mathfrak{P}_1$, i.e., the whole coset consists of very regular elements.

Lemma 3.3 *Let $x, x' \in \mathfrak{A}_1^\times$ be very regular. Then $\bar{\varphi}_{F,x'} = \bar{\varphi}_{F,x}$ if and only if $yx'y^{-1} \in x(1 + \mathfrak{P}_1)$ for some $y \in \langle \varpi_D \rangle \rtimes \mathfrak{A}_1^\times$.*

Proof It suffices to prove “only if”. By Skolem/Noether, $\bar{\varphi}_{F,x'} = \bar{\varphi}_{F,x}$ if and only if $\bar{x}, \bar{x}' \in (\mathfrak{A}_1/\mathfrak{P}_1)^\times = \text{GL}_m(k_D)$ are conjugate up to an action of the Galois group $\text{Gal}(k_D|k)$. Since conjugation by a prime element $\varpi_D \in D$ which normalizes the maximal unramified field extension $F_d \subset D$ generates $\text{Gal}(k_d|k)$, there exists $y \in \langle \varpi_D \rangle \mathfrak{A}_1^\times$ such that $\bar{y}\bar{x}'\bar{y}^{-1} = \bar{x}$. But this pulls back to $yx'y^{-1} \in x(1 + \mathfrak{P}_1)$. ■

Let $\Pi^A = \mathcal{J}_{\mathcal{L}_{A,D_n}}(\Pi^{D_n})$, where Π^{D_n} is a level zero representation. Recall that the character Θ_Π of an admissible representation Π of a p -adic reductive group G is defined as a linear functional on the space of complex-valued, compactly supported, locally constant test functions defined on G . By Harish-Chandra’s result [HC, Section 4, Cor.], Θ_Π is represented by a locally constant function on the regular set of G .

For the following we use AMT:

Lemma 3.4 *Let $x \in \mathfrak{A}_1^\times$ be very regular. The character Θ_{Π^A} is represented on the coset $x(1 + \mathfrak{P}_1)$ by a constant function.*

Proof Since the character $\Theta_{\Pi^{D_n}}$ is a level zero character, its value at any very regular element x depends only on the reduction \bar{x} and only on the reduction of the minimal polynomial $\bar{\varphi}_{F,x}$. By AMT the same is true for the values of Θ_{Π^A} at very regular $x \in \mathfrak{A}_1^\times$. From Lemma 3.3 it therefore follows that Θ_{Π^A} is constant on cosets $x(1 + \mathfrak{P}_1)$ of very regular $x \in \mathfrak{A}_1^\times$. ■

Lemma 3.5 *Let $x \in \mathfrak{A}_1^\times$ be very regular. The character Θ_{Π^A} is represented on the coset $x(1 + \mathfrak{P}_1)$ by the finite sum $\Sigma := \sum_\theta \theta$, where θ is summed over the irreducible level zero characters of \mathfrak{A}_1^\times which occur in $\Pi^A|_{\mathfrak{A}_1^\times}$ and each is counted once. Moreover, the two constant functions $\Theta_{\Pi^A}(u)$ and $\Sigma(u)$ on $x(1 + \mathfrak{P}_1)$ are identically equal.*

Proof Since Σ is a sum of level zero characters of \mathfrak{A}_1^\times , it is constant on $x(1 + \mathfrak{P}_1)$. Let ρ denote the characteristic function of the coset $x(1 + \mathfrak{P}_1)$ and let “vol” denote the volume of this open compact set. Then $\Sigma(x) = \Sigma(\rho) / \text{vol}$. On the other hand from Harish Chandra’s result (*loc. cit.*) combined with Lemma 3.4 we see that Θ_{Π^A} has on $x(1 + \mathfrak{P}_1)$ the constant value $\Theta_{\Pi^A}(x) = \Theta_{\Pi^A}(\rho) / \text{vol}$. But $\Theta_{\Pi^A}(\rho) = \text{tr}(\Pi^A(\rho))$ is the trace of the finite rank operator $\Pi^A(\rho)$. By Schur orthogonality on the compact group \mathfrak{A}_1^\times only level zero components of $\Pi^A|_{\mathfrak{A}_1^\times}$ contribute to that trace. The multiplicity one assertion in Lemma 3.1 (i) implies that $\Theta_{\Pi^A}(\rho) = \Sigma(\rho)$. ■

Let us complete the proof of Proposition 3.2. From Remark 2.10 it is easy to see that the character of $\Pi^{D_n} \in S_{\chi}^{D_n}$ at a very regular argument $x \in \mathfrak{A}_1^\times$ is $\Theta_{\Pi^{D_n}}(x) = \sum_{\bar{\eta} \in [\bar{\chi}]} \bar{\eta}(\bar{x})$. This is the division algebra case of (3.4) below. According to [SZ, 1.1 (ii)], no character sum of the form $\sum_{\bar{\eta} \in [\bar{\chi}]} \bar{\eta}(\bar{x})$ on k_n^\times is identically zero on the set of regular elements. Therefore, there exists a very regular element $x \in D_n^\times$ such that $\Theta_{\Pi^{D_n}}(x) \neq 0$. By AMT $\Theta_{\Pi^A}(x) = \Theta_{\mathcal{J}\mathcal{L}(\Pi^{D_n})}(x) \neq 0$ (we identify regular elliptic conjugacy classes of D_n^\times and A^\times with the same minimal polynomials). Again by Schur orthogonality we may conclude that $\Pi^A|_{\mathfrak{A}_1^\times}$ contains level zero components, so Π^A is a level zero representation. Therefore $\mathcal{J}\mathcal{L}_{A,D_n}$ gives an injection of $\mathcal{R}_0(D_n^\times)$ into $\mathcal{R}_0^2(A^\times)$. By AMT this injection preserves the central character, *i.e.*, $\mathcal{J}\mathcal{L}_{A,D_n}$ takes $\mathcal{R}_0(D_n^\times, \omega)$ into $\mathcal{R}_0^2(A^\times, \omega)$, where ω is a fixed central character, *i.e.*, a fixed character of F^\times . On the other hand, for all A we know that $\mathcal{R}_0^2(A^\times, \omega) = \bigcup_{\chi \in X_r(F_n^\times)} \mathcal{S}_\chi^A$, where $\chi \in X_r(F_n^\times)$ runs over all characters such that $\chi_F = \omega$. From Theorem 2.9 (i) we see that $\mathcal{R}_0^2(D_n^\times, \omega)$ and $\mathcal{R}_0^2(A^\times, \omega)$ are finite sets of the same cardinality, so $\mathcal{J}\mathcal{L}_{A,D_n}$ is surjective too. ■

Proposition 3.6 For $\Pi \in \mathcal{S}_\chi^A$ and $x \in A^\times$ very regular

$$(3.1) \quad \Theta_{\Pi^A}(x) = (-1)^{m-1} \sum_{\bar{\eta} \in [\bar{\chi}]} \bar{\eta}(\bar{x}).$$

Proof Let $\Pi^A \in \mathcal{S}_\chi^A$ and assume that $x \in A^\times$ is very regular. Without loss of generality we may assume that $x \in \mathfrak{A}_1^\times$. Then we know from Lemma 3.3 combined with AMT that Θ_{Π^A} is constant on $x(1 + \mathfrak{P}_1)$. By Lemma 3.4, $\Theta_\Pi = \Theta_{V^{1+\mathfrak{P}_1}}$ at very regular elements. We pass to \mathfrak{A}_1^\times . Using Lemma 3.1 (ii), we see that it is enough to compute the character at very regular elements by summing the characters of the generalized Steinberg components occurring in $V^{1+\mathfrak{P}_1}$. Thus, by Lemma 3.1 (iii), we have a single Galois orbit of GS representations of $\mathfrak{A}_1^\times = \text{GL}_m(k_d)$ corresponding to the $\mathfrak{S}_{e'}$ -orbits of length one in $V^{1+\mathfrak{P}_{e'}}$, *i.e.*, corresponding to the set of representations $\tau^i = \sigma^{\phi^i \otimes e'}$, $0 \leq i < (d, f)$. From [SZ, 6.1] we see that the character θ_i of the GS representation of $\text{GL}_m(k_d)$ which has cuspidal support $\tau^i = \sigma^{\phi^i \otimes e'}$ is $\theta_i(\bar{x}) = (-1)^{m-1} \sum \bar{\eta}(\bar{x})$ for $\bar{x} \in k_n$ regular with respect to the degree m extension $k_n|k_d$, where $\bar{\eta}$ runs over a $\text{Gal}(k_n|k_d)$ -orbit of characters of k_n^\times corresponding to σ^{ϕ^i} of $\text{GL}_{f'}(k_d)$. (In terms of (1.1) this refers to the case $k = k_d$.) Therefore we obtain

$$\Theta_{\Pi^A}(x) = (-1)^{m-1} \sum_{\bar{\chi}' \in \text{Gal}(k_d|k) \cdot \bar{\chi}} \sum_{\bar{\eta} \in \text{Gal}(k_n|k_d) \cdot \bar{\chi}'} \bar{\eta}(\bar{x}) = (-1)^{m-1} \sum_{\bar{\eta} \in \text{Gal}(k_n|k) \cdot \bar{\chi}} \bar{\eta}(\bar{x}). \quad \blacksquare$$

4 Weak Explicit Matching Theorem

Now we prove our main result.

Weak Explicit Matching Theorem 4.1 *Let $\chi \in X_t(F_n^\times)$, let D_n be a central division algebra of index n over F , and let A be a central simple F -algebra of reduced degree n . Then $\mathcal{J}\mathcal{L}_{A,D_n}(\mathcal{S}_\chi^{D_n}) = \mathcal{S}_\chi^A$ (cf. Definition 2.7), i.e., the Jacquet-Langlands map $\mathcal{J}\mathcal{L}_{A,D_n}$ defines a bijection of $\mathcal{S}_\chi^{D_n}$ to \mathcal{S}_χ^A and of $\mathcal{S}_{\bar{\chi}}^{D_n}$ to $\mathcal{S}_{\bar{\chi}}^A$.*

Proof Let $\Pi^A \in \mathcal{S}_\chi^A$ and set $\Pi^{D_n} := \mathcal{J}\mathcal{L}_{D_n,A}(\Pi^A)$. The idea of the proof is to compare the characters Θ_{Π^A} and $\Theta_{\Pi^{D_n}}$ on the corresponding sets of very regular elements x . From (3.4) and (0.5) we obtain

$$(4.1) \quad \Theta_{\Pi^{D_n}}(x) = \sum_{\bar{\eta} \in [\bar{\chi}]} \bar{\eta}(x)$$

for x very regular. On the other hand, Proposition 3.2 implies that Π^{D_n} is a level zero representation of D_n^\times , and this implies that $\Pi^{D_n} \in \mathcal{S}_{\chi'}^{D_n}$ for some $\chi' \in X_t(F_n^\times)$. Applying (3.4) to $\Pi^{D_n} \in \mathcal{S}_{\chi'}^{D_n}$ we obtain

$$(4.2) \quad \Theta_{\Pi^{D_n}}(x) = \sum_{\bar{\eta}' \in [\bar{\chi}']} \bar{\eta}'(x),$$

again for all very regular elements x . Thus the character sums (4.1) and (4.2) are equal for very regular arguments. By [SZ, 1.1 (i)] this implies that $[\bar{\chi}] = [\bar{\chi}']$. Moreover, $\chi_F = \chi'_F$ because Π^A and $\Pi^{D_n} = \mathcal{J}\mathcal{L}_{D_n,A}(\Pi^A)$ have the same central character (AMT). Therefore, since $\chi \in X_t(F_n^\times)$ is determined by its restriction $\chi|_{F^\times}$ and its reduction $\bar{\chi}$, $[\chi] = [\chi'] \in \langle \phi \rangle \backslash X_t(F_n^\times)$. Thus $\mathcal{J}\mathcal{L}_{D_n,A}$ maps \mathcal{S}_χ^A into $\mathcal{S}_{\bar{\chi}}^{D_n}$. Since these finite sets contain the same number of elements, by Proposition 2.8, and since, by AMT, $\mathcal{J}\mathcal{L}_{D_n,A}$ is injective, we conclude that $\mathcal{J}\mathcal{L}_{D_n,A}(\mathcal{S}_\chi^A) = \mathcal{S}_{\bar{\chi}}^{D_n}$. ■

Remark 4.2 Since the inertial class of discrete series representations $\mathcal{S}_{\bar{\chi}}^A$ corresponds to a single Bernstein component $\Omega_{[\bar{\chi}]}$ and since associated supercuspidal supports $(M, \pi_{[\sigma]}^{\otimes e'} \cdot \psi)$ also correspond to the same orbit $[\bar{\chi}]$, Theorem 4.1 implies that, up to unramified twist, the Jacquet-Langlands map commutes with parabolic induction.

A Appendix: Mod Center Hecke Algebras and $\mathcal{R}_0^2(A^\times)$

In order to justify assertions needed for the proofs of Propositions 2.4 and 2.8 we reformulate some parts of [BK1] and extend some of the results of [GSZ]. We review some of the connections between finite-dimensional mod center Hecke algebra modules and discrete series representations.

A.1 The Setting

We keep the notation and re-introduce some of the context of [GSZ]. From [GSZ, Part 4] we have the Hecke algebra $\mathcal{H}(r, z)$ of type \tilde{A}_{r-1} (cf. [BK1, (5.4)]). Thus we have

$\tilde{W} = W \rtimes \langle h \rangle$, an extension of the Coxeter group (W, S) , where S is a set of cardinality r and \tilde{W} is an extension of W by the infinite cyclic group $\langle h \rangle$. The group $\langle h \rangle$, acting on the normal subgroup W by conjugation, stabilizes $S \subset W$ and cyclically permutes the elements of S : $hs_i h^{-1} = s_{i-1}$ ($r - 1 \geq i > 0$) and $hs_0 h^{-1} = s_{r-1}$. The group \tilde{W} is equipped with a length function l which extends the length function on W and satisfies $l(h^i) = 0$ for all $i \in \mathbb{Z}$.

Fact A.1 For any $z \in \mathbb{C}^\times$ the vector space $\mathbb{C}[\tilde{W}]$ has one and only one \mathbb{C} -algebra structure $\mathcal{H}(\tilde{W}, z)$ such that:

- (i) $[s]^2 = (z - 1)[s] + z[1]$ for $s \in S$ and
- (ii) $[w_1] \cdot [w_2] = [w_1 w_2]$ whenever $l(w_1 w_2) = l(w_1) + l(w_2)$ ($w_1, w_2 \in \tilde{W}$).

Remarks (i) From (ii) we have $[h] \cdot [w] = [hw]$ and $[w] \cdot [h] = [wh]$ for all $w \in \tilde{W}$. The algebra

$$\mathcal{H}(\tilde{W}, z) = \mathcal{H}(W, z) \otimes_{\mathbb{C}} \mathbb{C}[\langle [h] \rangle]$$

is a twisted tensor product of the algebra for the Coxeter group W with the group ring $\mathbb{C}[\langle [h] \rangle]$:

$$(a.1) \quad ([w_1] \otimes [h^i])([w_2] \otimes [h^j]) = [w_1][h^i w_2 h^{-i}] \otimes [h^{i+j}]$$

for all $w_1, w_2 \in W$ and $i, j \in \mathbb{Z}$.

(ii) The center of $\mathcal{H}(\tilde{W}, z)$ is the subalgebra $\mathcal{Z} := \mathbb{C}[\langle [h^r] \rangle]$.

Now consider $A = M_m(D)$ and the distinguished pair (F_d, ϖ) , where $F_d|F$ is a maximal unramified extension of F in D and ϖ denotes a prime element of D which normalizes F_d ; thus $\varpi^d = \varpi_F$, a prime element of F . The group \tilde{W}_A denotes the extended relative affine Weyl group of A^\times ; concretely, \tilde{W}_A consists of all monomial matrices with entries which are either zero or powers of ϖ .

Let τ be the irreducible unitary representation of the principal order unit group \mathfrak{A}_r^\times which is lifted from the cuspidal representation $\sigma^{\otimes r}$ of $\overline{\mathfrak{A}}_r^\times = \text{GL}_s(k_D)^{\times r}$ ($rs = m$, $k_D = k_d$). Let l be the cardinality of the Galois orbit $\text{Gal}(k_d|k) \cdot \sigma$. Let \tilde{W} denote the set of monomial $m \times m$ matrices of the form $w = T \otimes I_s$, where T is an $r \times r$ monomial matrix with non-zero entries which are powers of ϖ^l and I_s denotes the $s \times s$ identity matrix. Let W be the subgroup of \tilde{W} consisting of all elements with reduced norm ± 1 and let

$$(a.2) \quad h := \begin{bmatrix} & I_{(r-1)s} \\ \varpi^l I_s & \end{bmatrix}.$$

Then the support of the Hecke algebra $\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ of operator-valued τ -spherical functions consists of the union of the set of double cosets $\mathfrak{A}_r^\times w \mathfrak{A}_r^\times$ over $w \in \tilde{W}$, where $\tilde{W} = W \rtimes \langle h \rangle \subset \tilde{W}_A$ (see [GSZ, 2.1]).

Slightly restating [GSZ, (4.2)], we have:

Fact A.2 Fix any non-trivial function $\mathcal{F} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ which has support on $\mathfrak{A}_r^\times h \mathfrak{A}_r^\times$. Then there is a uniquely determined isomorphism of algebras

$$\Phi_{\mathcal{F}}: \mathcal{H}(\tilde{W}, q^{ds}) \xrightarrow{\sim} \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$$

such that:

- (i) For all $w \in \tilde{W}$ the element $[w] \in \mathcal{H}(\tilde{W}, q^{ds})$ is mapped to a function φ_w which has support on $\mathfrak{A}_r^\times w \mathfrak{A}_r^\times$.
- (ii) $\varphi_h = \Phi_{\mathcal{F}}([h]) = \mathcal{F}$.

A.2 Algebra Involutions and Isomorphisms of Hilbert Spaces

Let \mathcal{W} be a representation space for the irreducible unitary representation τ and write $\langle \cdot, \cdot \rangle_{\mathcal{W}}$ for the inner product on \mathcal{W} . For $F \in \text{End}_{\mathbb{C}}(\mathcal{W})$ let F^* denote the adjoint of F with respect to $\langle \cdot, \cdot \rangle_{\mathcal{W}}$ and define the involution \star on $\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ by setting

$$(a.3) \quad \varphi^\star(g) := \varphi(g^{-1})^\star$$

for all $g \in A^\times, \varphi \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$.

Clearly, the involution \star defined by (a.3) sends a function with support on $\mathfrak{A}_r^\times w \mathfrak{A}_r^\times$ ($w \in \tilde{W}$) to a function with support on $\mathfrak{A}_r^\times w^{-1} \mathfrak{A}_r^\times$. Moreover, $(\lambda\varphi)^\star = \bar{\lambda}\varphi^\star$ for $\varphi \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau), \lambda \in \mathbb{C}$, and $(\varphi_1 \star \varphi_2)^\star = \varphi_2^\star \star \varphi_1^\star$ for all $\varphi_1, \varphi_2 \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$.

For the algebra $\mathcal{H}(\tilde{W}, z)$ we assume that $0 < z \in \mathbb{R}$ and define the involution \star such that $(\lambda[w])^\star := \bar{\lambda}[w^{-1}]$ ($\lambda \in \mathbb{C}, w \in \tilde{W}$) and such that $(x + y)^\star = x^\star + y^\star$ ($x, y \in \mathcal{H}(\tilde{W}, z)$). Again we have an anti-commutative conjugate linear involution.

Next we define inner products on the algebras $\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ and $\mathcal{H}(\tilde{W}, z)$:
 On $\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ we set

$$\begin{aligned} h(\varphi_1, \varphi_2) &:= (\dim \tau)^{-1} \int_{A^\times} \text{tr}_{\mathcal{W}}(\varphi_1(g) \cdot \varphi_2(g)^\star) dg \\ &= (\dim \tau)^{-1} \text{tr}_{\mathcal{W}}(\varphi_1 \star \varphi_2^\star)(1_{A^\times}). \end{aligned}$$

On $\mathcal{H}(\tilde{W}, z)$ ($0 < z \in \mathbb{R}$) we define the inner product $\langle \cdot, \cdot \rangle$ by assuming that the set \tilde{W} is an orthogonal basis for $\mathcal{H}(\tilde{W}, z)$ and setting $\langle [w], [w] \rangle = z^{l(w)}$. In particular, this implies that $\langle [h^i], [h^i] \rangle = 1$ for any $i \in \mathbb{Z}$.

Let $\mathcal{F} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ be a function supported on $\mathfrak{A}_r^\times h \mathfrak{A}_r^\times$. We know that $\mathcal{F} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)^\times$. By Fact A.2, there is a unique isomorphism

$$\Phi_{\mathcal{F}}: \mathcal{H}(\tilde{W}, q^{ds}) \xrightarrow{\sim} \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$$

such that $\Phi_{\mathcal{F}}([h]) = \mathcal{F}$. Bushnell/Kutzko [BK1, (5.6.16)] call $\Phi_{\mathcal{F}}$ a *unitary* isomorphism if $\mathcal{F}^\star = \mathcal{F}^{-1}$.

On pp. 193–195 of [BK1], beginning at (5.6.17), Bushnell/Kutzko prove:

Proposition A.3 Assume that $\Phi_{\mathcal{F}}$ is a unitary homomorphism. Then:

- (i) $\Phi_{\mathcal{F}}$ is a \star -homomorphism, i.e., $\Phi_{\mathcal{F}}(x^{\star}) = \Phi_{\mathcal{F}}(x)^{\star}$ for all $x \in \mathcal{H}(\tilde{W}, q^{ds})$.
- (ii) If the Haar measure on A^{\times} is normalized such that $\int_{\mathfrak{A}_r^{\times}} dx = 1$, then $\Phi_{\mathcal{F}}$ is an isometry of pre-hilbert spaces, i.e., $h(\Phi_{\mathcal{F}}(x), \Phi_{\mathcal{F}}(y)) = \langle x, y \rangle$ for all $x, y \in \mathcal{H}(\tilde{W}, q^{ds})$.

Proof (i) Since $\Phi_{\mathcal{F}}$ is linear and both the involutions \star are conjugate linear, it follows that, in order to show that $\Phi_{\mathcal{F}}$ is a \star -homomorphism, it is enough to show that $\Phi_{\mathcal{F}}([w]^{\star}) = \Phi_{\mathcal{F}}([w])^{\star}$ for all $w \in \tilde{W}$. Indeed, since $([w_1] \cdot [w_2])^{\star} = [w_2]^{\star} \cdot [w_1]^{\star}$ for all $w_1, w_2 \in \tilde{W}$ and $(f_1 \star f_2)^{\star} = f_2^{\star} \star f_1^{\star}$ for all $f_1, f_2 \in \mathcal{H}(A^{\times}, \mathfrak{A}_r, \tau)$, it suffices to show that $\Phi_{\mathcal{F}}([s]^{\star}) = \Phi_{\mathcal{F}}([s])^{\star}$ for all $s \in S$ and that $\Phi_{\mathcal{F}}([h]^{\star}) = \Phi_{\mathcal{F}}([h])^{\star}$. Since $[s]^{\star} = [s]$, we must show that $\Phi_{\mathcal{F}}([s])^{\star} = \Phi_{\mathcal{F}}([s])$. Let $f := f_s$ denote the obviously unique function in $\mathcal{H}(A^{\times}, \mathfrak{A}_r^{\times}, \tau)$ with support $\mathfrak{A}_r^{\times} s \mathfrak{A}_r^{\times}$ such that $f \star f = q^{ds} 1_{\mathcal{H}} + (q^{ds} - 1)f$. Since $\Phi_{\mathcal{F}}([s])$ has support in $\mathfrak{A}_r^{\times} s \mathfrak{A}_r^{\times}$ and $[s]^2 = q^{ds}[1] + (q^{ds} - 1)[s]$, it follows that $\Phi_{\mathcal{F}}([s]) = f$. Thus we must show that $f^{\star} = f$. On the one hand, we know that $\text{tr}(f \star f)(1) = q^{ds} \dim(\tau) > 0$. Since $s = s^{-1}$, f^{\star} also has support $\mathfrak{A}_r^{\times} s \mathfrak{A}_r^{\times}$. Thus, $f^{\star} = cf$ with $c \in \mathbb{C}^{\times}$; since $f^{\star\star} = f$ and \star is conjugate linear, we have $|c| = 1$. Since $f(x)f(x)^{\star}$ is a positive operator on \mathcal{W} for every $x \in \mathfrak{A}_r^{\times} s \mathfrak{A}_r^{\times}$, we see that

$$\text{tr}(f \star f^{\star})(1) = \text{tr} \left(\int_{A^{\times}} f(x)f(x)^{\star} dx \right) > 0.$$

Therefore, $c = 1$ and $f = f^{\star}$. Finally, if $\Phi_{\mathcal{F}}([h]) = \mathcal{F}$, where $\mathcal{F}^{\star} = \mathcal{F}^{-1}$, then $\Phi_{\mathcal{F}}([h])^{\star} = \Phi_{\mathcal{F}}([h^{-1}])$, since $[h] \star [h^{-1}] = 1$ and $\Phi_{\mathcal{F}}$ is an isomorphism. Since $[h]^{\star} = [h^{-1}]$, we have shown that $\Phi_{\mathcal{F}}([h])^{\star} = \Phi_{\mathcal{F}}([h]^{\star})$, which completes the proof of (i).

(ii) We begin with the following:

Lemma A.4 For all $w_1, w_2 \in \tilde{W}$

$$(a.4) \quad \langle [w_1], [w_2] \rangle = \langle [w_1] \cdot [w_2]^{\star}, 1 \rangle.$$

Proof First we observe that any $w \in \tilde{W}$ can be written in the form $w' h^i = s_{i_1} \cdots s_{i_{l(w)}} h^i$, where $i \in \mathbb{Z}$ is unique and $w' = s_{i_1} \cdots s_{i_{l(w)}}$ is a reduced expression for $w' \in W$. If $w_1 = w'_1 h^i$ and $w_2 = w'_2 h^{i'}$ with $i \neq i'$, then both sides of (a.4) are zero, since $[w]^{\star} = [w^{-1}]$ and we may apply (a.1). Thus we need check only the case $i = i'$ and it is clear that without loss of generality we may assume $i = 0$, i.e., $w_1, w_2 \in W$. Now we argue by induction on $l(w_2)$, assuming that $l(w_1)$ is arbitrary. The case $l(w_2) = 0$, i.e., $w_2 = 1$ is trivial, so we assume $l(w_2) > 0$ and that (a.4) is true for all w_1 and w'_2 with $l(w'_2) < l(w_2)$. Let $w_2 = w'_2 s$ with $s \in S$ and $l(w'_2) < l(w_2)$. By associativity in $\mathcal{H}(r, z)$ we have

$$\langle [w_1] \cdot [w_2]^{\star}, 1 \rangle = \langle [w_1] \cdot [w'_2 s]^{\star}, 1 \rangle = \langle [w_1] \cdot [s w'_2{}^{-1}], 1 \rangle = \langle ([w_1] \cdot [s]) \cdot [w'_2{}^{-1}], 1 \rangle.$$

There are two cases to consider: Assume $l(w_1 s) > l(w_1)$. Then $[w_1] \cdot [s] = [w_1 s]$. In this case, $\langle [w_1], [w_2] \rangle = 0$, since $l(w_2 s) = l(w'_2) < l(w_2)$ implies that $w_1 \neq w_2$. Moreover, $\langle [w_1 s] \cdot [w'_2]^{\star}, 1 \rangle = \langle [w_1 s], [w'_2] \rangle$, since $l(w'_2) < l(w_2)$. Again it is clear

that $\langle [w_1s], [w'_2] \rangle = 0$, since $w_1s \neq w'_2$, i.e., $l(w_1s \cdot s) = l(w_1) < l(w_1s)$ and $l(w'_2 \cdot s) = l(w_2) > l(w'_2)$. Thus (a.4) holds in the case that $l(w_1s) > l(w_1)$. Now assume that $w_1s = w'_1$ and that $l(w'_1) < l(w_1)$. In this case,

$$\begin{aligned} \langle ([w_1] \cdot [s]) \cdot [w'_2{}^{-1}], 1 \rangle &= \langle [w'_1](z[1] + (z - 1)[s]) \cdot [w'_2{}^{-1}], 1 \rangle \\ &= z\langle [w'_1] \cdot [w'_2{}^{-1}], 1 \rangle + (z - 1)\langle [w'_1] \cdot [s] \cdot [w'_2{}^{-1}], 1 \rangle, \end{aligned}$$

using the distributivity and linearity of the multiplication in $\mathcal{H}(r, z)$. Now we have $\langle [w'_1] \cdot [s] \cdot [w'_2{}^{-1}], 1 \rangle = \langle [w_1] \cdot [w'_2]{}^*, 1 \rangle = \langle [w_1], [w'_2] \rangle = 0$, since $w_1 \neq w'_2$, and we may use induction ($l(w'_2) < l(w_2)$). Also by induction we have

$$z\langle [w'_1] \cdot [w'_2{}^{-1}], 1 \rangle = z\langle w'_1, w'_2 \rangle = z \cdot z^{l(w'_2)} \delta_{w'_1, w'_2} = z^{l(w_2)} \delta_{w_1, w_2}. \quad \blacksquare$$

Let us complete the proof of Proposition A.3 (ii): It is enough to prove the case $x = [w_1]$ and $y = [w_2]$ with $w_1, w_2 \in \tilde{W}$ and to observe that then (ii) follows via linearity. Writing $\varphi_w := \Phi_{\mathcal{F}}([w])$, we have

$$(\dim \tau)h(\varphi_w, \varphi_w) = \text{tr}_{\mathcal{W}}((\varphi_w * \varphi_w^*)(1)) = \text{tr}_{\mathcal{W}}(\Phi_{\mathcal{F}}([w] \cdot [w]^*)(1)).$$

From (a.4) and the orthogonality of the basis \tilde{W} of $\mathcal{H}(\tilde{W}, q^{ds})$ we obtain

$$[w] \cdot [w]^* = \langle [w], [w] \rangle 1 + \sum_{w \neq 1} c_w [w].$$

Thus $\Phi_{\mathcal{F}}([w] \cdot [w]^*)(1) = \langle [w], [w] \rangle \Phi_{\mathcal{F}}(1)(1) = \langle [w], [w] \rangle \mu(\mathfrak{A}_r^\times)^{-1} \tau(1)$. This gives

$$h(\Phi_{\mathcal{F}}([w]), \Phi_{\mathcal{F}}([w])) = \langle [w], [w] \rangle \mu(\mathfrak{A}_r^\times)^{-1},$$

which proves (ii), assuming that the measure $\mu(\mathfrak{A}_r^\times) = 1$. \blacksquare

We must still prove that \mathcal{F} may be chosen such that $\Phi_{\mathcal{F}}$ is unitary:

Lemma A.5 *There exists a function $\mathcal{F} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ with support $\mathfrak{A}_r^\times h \mathfrak{A}_r^\times$ such that $\mathcal{F}^* = \mathcal{F}^{-1}$.*

Proof [GSZ, 1.1] implies that the function \mathcal{F} has the value $\mathcal{F}(h) = J$, where $J \in \text{Hom}_{\mathfrak{A}_r^\times \cap h^{-1} \mathfrak{A}_r^\times h}(\tau, \tau^h)$ is an intertwining operator. In [GSZ, 4.3] we saw that if $\tilde{\mathcal{F}} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ has support $\mathfrak{A}_r^\times h^{-1} \mathfrak{A}_r^\times$ and if $\tilde{\mathcal{F}}(h^{-1}) = J^{-1}$, then $\mathcal{F} * \tilde{\mathcal{F}} = \lambda 1_{\mathcal{H}}$, where $\lambda = \mu(\mathfrak{A}_r^\times) \mu(\mathfrak{A}_r^\times h \mathfrak{A}_r^\times)$. Thus if we replace J by $J_1 = \lambda^{-1/2} J$ as the value of $\mathcal{F}(h)$ and J^{-1} by $J_2 = \lambda^{-1/2} J^{-1}$ as the value of $\tilde{\mathcal{F}}(h^{-1})$, then $\mathcal{F}^{-1} = \tilde{\mathcal{F}}$. Since $\mathcal{F}^*(h^{-1}) = \mathcal{F}(h)^* = J_1^*$, it suffices to show that $J \in \text{End}_{\mathbb{C}}(\mathcal{W})$ may be chosen to be a unitary operator, since in this case $J_1^* = \lambda^{-1/2} J^* = \lambda^{-1/2} J^{-1} = J_2$, i.e., in this case, $\mathcal{F}^* = \tilde{\mathcal{F}} = \mathcal{F}^{-1}$.

Let us show that J can be chosen to be a unitary operator in $\text{End}_{\mathbb{C}}(\mathcal{W})$. By the definition of h (see (a.2)) we know that $h^{dr/l} = \varpi_F 1_{A^\times}$, a central element of \tilde{W} (l is the length of the Galois orbit $\text{Gal}(k_d|k) \cdot \sigma$). The representation τ of $M(k_D)$ extends as a unitary representation to the group $M(k_D) \rtimes C$, where $C = \langle h \rangle \text{ mod } h^{dr/l}$, since C is a finite cyclic group and $h^{dr/l}$ is central. Clearly, the generator h of C must under this extension map to an element of $\text{Hom}_{\mathfrak{A}_r^\times \cap h^{-1} \mathfrak{A}_r^\times h}(\tau, \tau^h) = \text{Hom}_{M(k_D)}(\tau, \tau^h)$. Thus J can be chosen to be unitary. \blacksquare

A.3 Discrete Series and Hecke Algebras with a Central Character

To reduce the description of discrete series in the general case to the unramified split case we reformulate some parts of [BK1, (6.1)].

Let ω be a character of $\langle \varpi_F \rangle$ and let τ_ω denote the extension of τ by ω to $\mathfrak{A}_r^\times \langle \varpi_F \rangle = \mathfrak{A}_r^\times F^\times$. We consider the projection maps:

$$(a.5) \quad \mathcal{H}(\tilde{W}, q^{ds}) \longrightarrow \bar{\mathcal{H}}(\tilde{W}, q^{ds}) := \mathcal{H}(\tilde{W}, q^{ds}) / ([\varpi_F] - [1])$$

and

$$(a.6) \quad \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau) \longrightarrow \mathcal{H}(A^\times, \mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega), \quad f \longmapsto \bar{f} := Pf,$$

where $Pf(x) := \sum_{i \in \mathbb{Z}} \omega(\varpi_F^{-i}) f(\varpi_F^i x)$. It is easy to see that the projection operator P is a homomorphism of algebras, that the quotient algebra $\mathcal{H}(A^\times, \mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega)$ has the structure of convolution algebra with respect to the product

$$(\bar{f}_1 * \bar{f}_2)(x) = \int_{A^\times / \langle \varpi_F \rangle} \bar{f}_1(y) \bar{f}_2(y^{-1}x) dy.$$

The unit element is:

$$\bar{e}(x) = \begin{cases} \mu(\mathfrak{A}_r^\times)^{-1} \tau_\omega(x) & \text{for } x \in \mathfrak{A}_r^\times \langle \varpi_F \rangle \\ 0 & \text{otherwise.} \end{cases}$$

We fix the normalization $\mu(\mathfrak{A}_r^\times) = 1$.

Lemma A.6 *Let $J \in \text{End}_{\mathbb{C}}(\mathcal{W}_\tau)$ be constructed as in the second half of the proof of Lemma A.5 and choose $\mathcal{F} \in \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ with support $\mathfrak{A}_r^\times h \mathfrak{A}_r^\times$ such that*

$$\mathcal{F}(h) := \frac{1}{\sqrt{\mu}} J\zeta,$$

where $\mu = \mu(\mathfrak{A}_r^\times h \mathfrak{A}_r^\times)$ and ζ is a root of unity chosen such that $\zeta^{dr/l} = \omega(\varpi_F)$. Then the isomorphism $\Phi_{\mathcal{F}}$ of Fact A.2 induces an isomorphism

$$\Phi_{\bar{\mathcal{F}}}: \bar{\mathcal{H}}(\tilde{W}, q^{ds}) \longrightarrow \mathcal{H}(A^\times, \mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega),$$

which implies the commutative diagram

$$\begin{array}{ccc} \mathcal{H}(\tilde{W}, q^{ds}) & \xrightarrow{\Phi_{\mathcal{F}}} & \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau) \\ (a.5) \downarrow & & \downarrow (a.6) \\ \bar{\mathcal{H}}(\tilde{W}, q^{ds}) & \xrightarrow{\Phi_{\bar{\mathcal{F}}}} & \mathcal{H}(A^\times, \mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega) \end{array}$$

in which the horizontal arrows are the indicated isomorphisms and the numbers on the vertical arrows are the projection homomorphisms (a.5) and (a.6) defined previously.

Proof Since $h^r = \varpi^l 1_{A^\times}$, an operator which normalizes \mathfrak{A}_r^\times , we have

$$\mu(\mathfrak{A}_r^\times \cdot h^r \cdot \mathfrak{A}_r^\times) = \mu(h^r \mathfrak{A}_r^\times) = 1.$$

Therefore, since l is the length of the Galois orbit of σ , $\mathcal{F}^r(h^r) = J^r \zeta^r$ intertwines the representation τ of \mathfrak{A}_r^\times and, in particular, since by hypothesis $\zeta^{dr/l} = \omega(\varpi_F)$,

$$\mathcal{F}^{dr/l}(\varpi_F 1_{A^\times}) = \mathcal{F}^{dr/l}(h^{dr/l}) = J^{dr/l} \zeta^{dr/l} = \omega(\varpi_F) 1_\tau = \bar{e}(\varpi_F 1_{A^\times}).$$

Thus $\bar{\mathcal{F}}^{dr/l} = \bar{e}$. We leave it to the reader to verify that the isomorphism of the affine Weyl group algebra $\Phi_{\mathcal{F}}|_{\mathcal{H}(W, q^{ds})}$ passes to the quotient and induces an isomorphism $\Phi_{\bar{\mathcal{F}}}|_{\bar{\mathcal{H}}(W, q^{ds})}$ such that the resulting subdiagram is commutative. In fact the vertical arrows for the affine Weyl group algebras are isomorphisms as well as the horizontal arrows.

We must show that the vertical algebra homomorphisms from the isomorphic cyclic group algebras $\mathbb{C}[\langle [h] \rangle]$ and $\mathbb{C}[\langle \mathcal{F} \rangle]$ involve the same polynomial relations, *i.e.*, that

$$\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau) / (\mathcal{F}^{dr/l} - e) \cong \mathcal{H}(A^\times, \mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega).$$

Since $[h] \mapsto \Phi_{\mathcal{F}}([h]) = \mathcal{F}$ generates a homomorphism of cyclic groups, it suffices to show that no smaller power \mathcal{F}^ℓ of \mathcal{F} maps to a scalar multiple of \bar{e} . For this we note that $\bar{\mathcal{F}}^\ell$ a root of the identity, *i.e.*, a root of $\bar{\mathcal{F}}^{dr/l} = \bar{e}$, implies that $\ell \mid \frac{dr}{l}$. We therefore assume that $\ell \mid \frac{dr}{l}$. If $r \nmid \ell$, then the support of \mathcal{F}^ℓ is a double coset $\mathfrak{A}_r^\times h^\ell \mathfrak{A}_r^\times$ such that h^ℓ does not normalize \mathfrak{A}_r^\times ; in this case, h^ℓ cannot centralize the representation τ , so \mathcal{F}^ℓ cannot be a multiple of the identity element \bar{e} . Assume that $r \mid \ell$. In this case the support of the function \mathcal{F}^ℓ is the double coset $\mathfrak{A}_r^\times h^\ell$ and $h^\ell = \varpi^{\ell/r} 1_{A^\times}$; since $r \mid \ell$, h^ℓ normalizes \mathfrak{A}_r^\times and the double coset supported at h^ℓ is the same as a one-sided coset. We see also that $J^\ell \zeta^\ell = \mathcal{F}^\ell(h^\ell) = \mathcal{F}(\varpi^{\ell/r} \cdot 1_{A^\times})$. To complete the proof of Lemma A.6 it is sufficient to show that the operator J , which intertwines the representation τ , cannot centralize τ . In this case, \mathcal{F}^ℓ cannot be a scalar multiple of \bar{e} .

Let us show that J^ℓ has to be conjugate to a non-trivial permutation matrix, which implies that there exists $x \in \mathfrak{A}_r^\times$ such that $J^\ell \tau(x) J^{-\ell} \neq \tau(x)$, since the commuting algebra of τ consists of only scalar matrices by Schur's Lemma. It is enough to show this for $\tau|_{M(O)}$, since the intertwining of this irreducible restriction already determines J^ℓ . We give details only for the case $r = 1$, since the general case involves no additional ideas. Recall that τ is the lift of a cuspidal representation of $M(k_D)$ to $M(O)$. By results of S. I. Gelfand, quoted in [SZ], the restriction $\tau_{\Omega(k_d)}$ is irreducible, where $\Omega(k_D)$ denotes the Gelfand subgroup of $M(k_D) = \text{GL}_m(k_D)$. Moreover, the irreducible representation of $\Omega(k_D)$ is independent of the choice of cuspidal representation and is simply $\text{Ind}_{U_0(k_d)}^{\Omega(k_d)} \mu$, where $U_0(k_D)$ denotes the upper triangular unipotent subgroup of $M(k_D)$ and μ denotes any generic character of $U_0(k_D)$. It follows that we may choose a basis for the representation space of τ which consists of generic vectors for conjugates of $U_0(k_D)$. The Galois group $\text{Gal}(k_D|k)$ acts as a permutation group on this basis of \mathcal{W}_τ . Since the set of generic vectors for $U_0(k_D)$ consists of a single orbit under the multiplicative group of diagonal matrices $T(k_D)$ acting by conjugation with only

the center acting trivially, and the group $\text{Gal}(k_D|k)$ acts on $T(k_D)$ mod the center with only the identity acting trivially, it follows that

$$x \mapsto \tau(\varpi^{\ell/r} x \varpi^{-\ell/r}) = J^\ell \tau(x) J^{-\ell} = \tau(x)$$

for all $x \in M(O)$ if and only if $\frac{d}{l} \mid \frac{\ell}{r}$, since no smaller power of ϖ permutes the basis for \mathcal{W}_τ trivially. Thus only when $\frac{dr}{l} \mid \ell$ can $\tilde{\mathcal{F}}^\ell$ be a multiple of \bar{e} . We conclude that $\mathcal{F} \mapsto \tilde{\mathcal{F}}$ defines an isomorphism

$$\mathbb{C}[\langle [h] \rangle] / ([h^{dr/l}] - [1]) \xrightarrow{\sim} \mathbb{C}[\langle \tilde{\mathcal{F}} \rangle].$$

The rest of the proof involves checking that the conjugation action of \mathcal{F} on the affine part of the Hecke algebra $\mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ factors through the projection $\mathcal{F} \mapsto \tilde{\mathcal{F}}$ and this we leave to the reader. ■

As a special case of the foregoing, we consider the algebra $A' := M_r(F_{ds})$, where the extension $F_{ds}|F$ is unramified of degree ds . In A'^\times , we have the Iwahori subgroup $J'^\times := \mathfrak{A}'^\times$ and for τ' we take the trivial representation $1 := 1_{J'^\times}$. The support of the Hecke algebra $\mathcal{H}(A'^\times, J'^\times, 1)$ is the group of monomial matrices $\bar{W}_{A'}$ with entries which are powers of ϖ_F . As before, we have the isomorphism

$$\Phi_1: \mathcal{H}(\bar{W}_{A'}, q^{ds}) \longrightarrow \mathcal{H}(A'^\times, J'^\times, 1),$$

where $h' := \begin{bmatrix} 0_{r-1,1} & I_{r-1} \\ \varpi_F & 0_{1,r-1} \end{bmatrix}$, $\mathcal{F}' \in \mathcal{H}(A'^\times, J'^\times, 1)$ has support $J'h'J' = J'h'$ with $\mathcal{F}'(h') = 1$, and $\Phi_1(h') = \mathcal{F}'$. (Note that $\mathcal{H}(A'^\times, J'^\times, 1)$ is an algebra of scalar-valued functions, since “1” is one-dimensional. Put precisely, $\mathcal{H}(A'^\times, J'^\times, 1)$ is the convolution algebra consisting of all compactly supported functions which are constant on J' double cosets.)

Recall that a set of representatives for the \mathfrak{A}_r^\times double cosets lying in the support of the algebra $\mathcal{H}(\tau) := \mathcal{H}(A^\times, \mathfrak{A}_r^\times, \tau)$ may be chosen to be the set of all matrices of the form $\varpi^{lv} p \otimes I_s$, where ϖ is the prime element of D which was fixed earlier, l is the length of the Galois orbit $\text{Gal}(k_D|k) \cdot \sigma$, $v = (v_1, \dots, v_r) \in \mathbb{Z}^r$, $\varpi^{lv} = \text{diag}(\varpi^{lv_1}, \dots, \varpi^{lv_r})$, and p is an $r \times r$ permutation matrix. We note that the extended Coxeter system $\tilde{W} = (W, S) \rtimes \langle h \rangle$ is naturally isomorphic to the extended Coxeter system $\tilde{W}' := (W', S') \rtimes \langle h' \rangle$ via the map

$$(a.7) \quad \varpi^{lv} p \otimes I_s \mapsto \varpi_F^v p,$$

v and p as above. It is important to observe that the mapping (a.7) sends $\varpi_F I_m \in A^\times$ to $\varpi_F^{d/l} I_r \in A'^\times$. We have the isomorphisms:

$$(a.8) \quad \mathcal{H}(\tau) \xleftarrow{\Phi_{\mathcal{F}}} \mathcal{H}(\tilde{W}, q^{ds}) \cong \mathcal{H}(\tilde{W}', q^{ds}) \xrightarrow{\Phi_1} \mathcal{H}(A'^\times, J'^\times, 1).$$

Using (a.7) we may regard these isomorphisms as support-preserving.

Next, passing to quotients and using (a.7) again we have

$$(a.9) \quad \tilde{\mathcal{H}}(\tilde{W}, q^{ds}) \cong \tilde{\mathcal{H}}(\tilde{W}', q^{ds}),$$

where $\tilde{\mathcal{H}}(\tilde{W}', q^{ds}) := \mathcal{H}(\tilde{W}', q^{ds})/([\varpi_F^{d/l}] - [1])$. Instead of (a.5) we now use the projection mapping

$$(a.10) \quad \mathcal{H}(A'^{\times}, J'^{\times}, 1) \longrightarrow \mathcal{H}(A'^{\times}, J'^{\times} \langle \varpi_F^{d/l} \rangle, 1),$$

sending $f' \in \mathcal{H}(A'^{\times}, J'^{\times}, 1)$ to $\tilde{f}'(x) := \sum_{i \in \mathbb{Z}} f'(\varpi_F^{-id/l} x) \in \mathcal{H}(A'^{\times}, J'^{\times} \langle \varpi_F^{d/l} \rangle, 1)$. The projection mapping $f' \mapsto \tilde{f}'$ is clearly an algebra homomorphism. From (a.8), (a.10), and Lemma A.6 we deduce the diagram of isomorphisms

$$(a.11) \quad \mathcal{H}(\tau_{\omega}) \xleftarrow{\Phi_{\tilde{\mathcal{H}}}} \tilde{\mathcal{H}}(\tilde{W}, q^{ds}) \cong \mathcal{H}(\tilde{W}', q^{ds}) \xrightarrow{\Phi_1} \mathcal{H}(A'^{\times}, J'^{\times} \langle \varpi_F^{d/l} \rangle, 1),$$

in which $\mathcal{H}(\tau_{\omega}) := \mathcal{H}(A^{\times}, \mathfrak{A}_r^{\times} \langle \varpi_F \rangle, \tau_{\omega})$. Since the maps $\Phi_{\mathcal{H}}$ and Φ_1 of (a.8) are unitary according to Bushnell/Kutzko's definition (see above Proposition A.3), (a.8) preserves the prehilbert structures on $\mathcal{H}(\tau)$ and $\mathcal{H}(A'^{\times}, J'^{\times}, 1)$, respectively. To introduce prehilbert structures on the algebras of (a.11) it is therefore enough to treat any single quotient algebra and then transport the structure to the remaining algebras, using the equivalence of the norms in (a.8). It is enough to treat the case of $\mathcal{H}(\tilde{W}, q^{ds})$. For $w \in \tilde{W}$, the image under the projection $[w] \mapsto \overline{[w]}$ is a function with support $\bar{w} \in \tilde{W}/\langle \varpi_F \rangle$. If $\overline{[w_1]} = \overline{[w_2]} \in \tilde{\mathcal{H}}(\tilde{W}, q^{ds})$, then it is clear that $\|w_1\| = \|w_2\|$. Therefore, we may use any set of representatives for $\tilde{W}/\langle \varpi_F \rangle$ to define the norms of the projections to an orthogonal basis for $\tilde{\mathcal{H}}(\tilde{W}, q^{ds})$, i.e., we set $\|\overline{[w]}\| = \|w\|$. Thus we may conclude that (a.11) induces isomorphisms between the completions of the various constituents which are compatible with module structures with respect to the algebras. We denote the completions of an algebra and its projection, respectively, by \mathcal{H}^2 and $\tilde{\mathcal{H}}^2$.

We consider the categories $\mathcal{M} := \mathcal{M}(A^{\times}, \varpi_F, \omega)$ and $\mathcal{M}' := \mathcal{M}(A'^{\times}, \varpi_F^{d/l}, 1)$ of smooth representations, where, in the first case, ϖ_F acts via $\omega(\varpi_F)$ and, in the second, $\varpi_F^{d/l}$ acts trivially. These categories are closed under forming subquotients. Therefore, the pairs $(\mathfrak{A}_r^{\times} \langle \varpi_F \rangle, \tau_{\omega})$ and $(J'^{\times}, \langle \varpi_F^{d/l} \rangle, 1)$ are types corresponding to subcategories of \mathcal{M} and \mathcal{M}' , respectively.

Simplifying our notation, we denote our pairs by (K, ρ) and the corresponding Hecke algebras by $\mathcal{H}(\rho)$. Let \mathcal{M}_{ρ} denote the subcategory of \mathcal{M} or \mathcal{M}' which consists of all representations which are generated by their ρ -isotypic components. We have the equivalences

$$(a.12) \quad \mathcal{M}_{\rho} \ni \Pi \longmapsto \text{Hom}_K(\rho, \Pi) \in \mathcal{H}(\rho)^{\text{opp}}\text{-modules.}$$

We may also interpret $\mathcal{M}, \mathcal{M}'$ as the category of \mathcal{H} -modules, where \mathcal{H} denotes the algebra of functions on A^{\times}, A'^{\times} respectively, such that $f(\varpi_F g) = \omega(\varpi_F) f(g)$ and $f'(\varpi_F^{d/l} g') = f'(g')$, respectively, and where the support is relatively compact. We embed \mathcal{H} in each case into the Hilbert space completion \mathcal{H}^2 , where the condition

of relative compactness is replaced by square integrability on $A^\times / \langle \varpi_F \rangle, A'^\times / \langle \varpi_F^{d/l} \rangle$ respectively. Now we refer to the arguments of [BK1, (7.7.3)–(7.7.5)] and apply them in the present context. We replace the notations $\mathcal{H}(G, \omega), \mathcal{H}^2(G, \omega), \mathcal{H}(G, \lambda\omega),$ and $\mathcal{H}^2(G, \lambda\omega)$ of [BK1] with the notations $\mathcal{H}, \mathcal{H}^2, \mathcal{H}(\rho),$ and $\mathcal{H}^2(\rho)$ (see (a.11), (a.12)), where (K, ρ) are the types we are interested in. Then we may obtain the following formulation from [BK1, (7.7.5)]. In the following we write $\mathcal{M}_\rho^{\text{finite}} \subset \mathcal{M}_\rho$ for the full subcategory consisting of representations with finite composition series.

Proposition A.7 *Under the equivalence (a.12) of categories a representation $\Pi \in \mathcal{M}_\rho^{\text{finite}}$ is square integrable if and only if the $\mathcal{H}(\rho)^{\text{opp}}$ -module $\text{Hom}_K(\rho, \Pi)$ can be embedded into $\mathcal{H}^2(\rho)$.*

We wish to formulate a consequence of this assertion. Let $(K, \rho) := (\mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega)$ and $(K', \rho') := (\mathcal{J}'^\times \langle \varpi_F^{d/l} \rangle, 1)$. Let $\mathcal{M}_\rho \subset \mathcal{M}$ and $\mathcal{M}'_{\rho'} \subset \mathcal{M}'$ be the corresponding subcategories.

Corollary A.8 *The isomorphisms of (a.11) induce a discrete-series preserving equivalence of the categories $\mathcal{M}_\rho^{\text{finite}} \rightarrow \mathcal{M}'_{\rho'}{}^{\text{finite}}$.*

Proof Let $\Pi \in \mathcal{M}_\rho^{\text{finite}}$ and consider the $\mathcal{H}(\rho)$ -module $\text{Hom}_K(\rho, \Pi)$. Using (a.11) we may consider $\text{Hom}_K(\rho, \Pi)$ as an $\mathcal{H}(\rho')$ -module. Since (K', ρ') is a type, we also obtain a representation $\Pi' \in \mathcal{M}'_{\rho'}{}^{\text{finite}}$ such that

$$(a.13) \quad \text{Hom}_{K'}(\rho', \Pi') = \text{Hom}_K(\rho, \Pi)$$

as $\mathcal{H}(\rho')$ -modules. By Proposition A.7, Π is discrete series if and only if $\text{Hom}_K(\rho, \Pi)$ embeds into $\mathcal{H}^2(\rho)$. Since (a.11) implies that the isomorphism $\mathcal{H}(\rho) \cong \mathcal{H}(\rho')$ extends to a unitary isomorphism of $\mathcal{H}^2(\rho) \cong \mathcal{H}^2(\rho')$, it follows that $\text{Hom}_K(\rho, \Pi)$ embeds in $\mathcal{H}^2(\rho)$ if and only if the isomorphic module $\text{Hom}_{K'}(\rho', \Pi')$ embeds in $\mathcal{H}^2(\rho')$. This means that $\Pi \in \mathcal{M}_\rho^{\text{finite}}$ is discrete series if and only if the same is true of $\Pi' \in \mathcal{M}'_{\rho'}{}^{\text{finite}}$. ■

Proposition A.9 *If $\Pi \in \mathcal{M}_\rho^{\text{finite}}$ is an irreducible discrete series module, then $\dim_{\mathbb{C}}(\text{Hom}_K(\rho, \Pi)) = 1$, i.e., ρ occurs in Π with multiplicity one.*

Proof If Π is irreducible and discrete series, then the same holds for the corresponding representation $\Pi' \in \mathcal{M}'_{\rho'}{}^{\text{finite}}$. But $A'^\times = \text{GL}_r(F_{d_s})$ and for this split group Bernstein/Zelevinsky have proved that the discrete series which occur in $\text{Irr}(\Omega)$, where Ω is an arbitrary Bernstein component, occur as at most a single unitary inertial class of representations [Ze, 9.3]. Therefore, the only discrete series which occur in the unramified principal series of $\text{GL}_r(F_{d_s})$, i.e., in the set of irreducible discrete series

representations associated to the Bernstein component of $GL_r(F_{ds})$ for which (K', ρ') is a type, have to be unramified twists of the Steinberg representation of $GL_r(F_{ds})$. For the Steinberg representation St of an arbitrary p -adic reductive group G it is known that the corresponding Iwahori bi-invariant matrix coefficient gives a scalar-valued homomorphism of the algebra $\mathcal{H}(G, J, 1)$ (see [Bo, Section 5], where a more general result is given though only for semi-simple groups). This implies that 1_J occurs simply in St_J and obviously also the same will remain true for any unramified twist of St . This implies that $\text{Hom}_{K'}(\rho', \Pi')$ has \mathbb{C} -dimension one as an $\mathcal{H}(\rho')$ -module. From (a.13) we obtain the proposition. ■

Corollary A.10 *The number of irreducible discrete series representations Π of A^\times which contain $(\mathfrak{A}_r^\times \langle \varpi_F \rangle, \tau_\omega)$ is dr/l .*

Proof It is enough to count the number of irreducible discrete series representations of A'^\times which contain $(J'^\times \langle \varpi_F^{d/l} \rangle, 1)$. As we have seen in the proof of Proposition A.9, each of these representations is an unramified twist of the Steinberg representation St of $GL_r(F_{ds})$. Since the central character of St is trivial, it follows that the central character of $St \otimes \tilde{\chi}$, where χ is an unramified character and $\tilde{\chi} := \chi \circ \text{Nrd}_{A'|F_{ds}}$, is χ^r . In order to have a central character which is trivial on $\langle \varpi_F^{d/l} \rangle$ we therefore have to choose χ of F^\times such that χ^r is trivial on $\langle \varpi_F^{d/l} \rangle$. We see that there are dr/l such possibilities and this implies the result. ■

References

- [Ba] A. Badulescu, *Correspondance entre GL_n et ses formes intérieures en caractéristique positive*. Thèse présentée pour obtenir le grade de Docteur en Sciences, Université de Paris-Sud, Centre d'Orsay, 1999.
- [BD] J. Bernstein (rédigé par P. Deligne), *Le "centre" de Bernstein*. In: Représentations des groupes réductifs sur un corps local, collection dirigée par Jean Dieudonné, Hermann, Paris, 1984, 1–32.
- [BR] J. Bernstein and C. Rumelhart, *Lectures on Representations of Reductive p -adic Groups*. Manuscript, 1996.
- [Bo] A. Borel, *Admissible Representations of a Semi-Simple Group over a Local Field with Vectors Fixed under an Iwahori Subgroup*. *Invent. Math.* **35**(1976), 233–259.
- [BK1] C. Bushnell and P. Kutzko, *The Admissible Dual of $GL(N)$ via Compact Open Subgroups*. *Annals of Math. Stud.* **129**, Princeton University Press, Princeton, NJ, 1993.
- [BK2] ———, *Smooth Representations of Reductive p -adic Groups: Structure Theory via Types*. *Proc. London Math. Soc.* **77**(1998), 582–634.
- [DKV] P. Deligne, D. Kazhdan and M.-F. Vignéras, *Représentations des algèbres centrales simples p -adiques*. In: Représentations des groupes réductifs sur un corps local, collection dirigée par Jean Dieudonné, Hermann, Paris, 1984, 33–117.
- [DM] F. Digne and J. Michel, *Representations of Finite Groups of Lie Type*. London Mathematical Society Student Texts **21**, Cambridge University Press, Cambridge, 1991.
- [GSZ] M. Grabitz, A. Silberger, and E.-W. Zink, *Level Zero Types and Hecke Algebras for Local Central Simple Algebras*. *J. Number Theory* **91**(2001), 92–125.
- [Gr] J. Green, *The characters of the finite general linear groups*. *Trans. Amer. Math. Soc.* **80**(1955), 402–447.
- [HC] Harish-Chandra, *A Submersion Principle and Its Applications*. In: Harish-Chandra Collected Papers, vol. **IV**, Springer, New York, 1984, 439–446.
- [Ro] J. Rogawski, *Representations of $GL(n)$ and division algebras over a p -adic field*. *Duke Math. J.* **50**(1983), 161–169.

- [SZ] A. Silberger and E.-W. Zink, *The Characters of the Generalized Steinberg Representations of Finite General Linear Groups on the Regular Elliptic Set*. Trans. Amer. Math. Soc. **352**(2000), 3339–3356.
- [Ze] A. Zelevinsky, *Induced Representations of Reductive p -adic Groups II*. Ann. Sci. École Norm. Sup. (4) **13**(1980), 165–210.

Cleveland State University
Cleveland, Ohio 44115
USA
email: silberger@math.csuohio.edu

Humboldt-Universität
Institut für Mathematik
Rudower Chaussee 25
10099 Berlin
Germany
email: zink@mathematik.hu-berlin.de