



Counting points by height in semigroup orbits

Jason P. Bell , Wade Hindes , and Xiao Zhong

Abstract. We improve known estimates for the number of points of bounded height in semigroup orbits of polarized dynamical systems. In particular, we give exact asymptotics for generic semigroups acting on the projective line. The main new ingredient is the Wiener-Ikehara Tauberian theorem, which we use to count functions in semigroups of bounded degree.

1 Introduction

A general principle in arithmetic dynamics is that the arithmetic behavior of orbits of rational points of algebraic varieties under self-maps can often be understood in terms of dynamical properties of the associated dynamical systems. In particular, dynamical properties of algebraic dynamical systems often put strong constraints on the shapes of orbits of rational points. It is then natural to study the extent to which measures of complexity of orbits of rational points are reflected by the corresponding measures for complexity of dynamical systems. Typically, one uses some notion of asymptotic behavior of heights in orbits as a measure of the complexity of an orbit and uses some notion of degree for iterates to measure the complexity of a dynamical system. The Kawaguchi-Silverman conjecture [9], which asserts that the dynamical degree should determine the height growth for Zariski dense orbits of $\overline{\mathbb{Q}}$ -points, is perhaps the most striking prediction concerning the connection between these seemingly unrelated notions of complexity. Their conjecture has since been verified in a number of cases [11, 12, 13, 14, 15] but remains open.

In this paper, we build on earlier work of the second-named author [6] and investigate the growth of heights under semigroup orbits. More precisely, we consider the setting when one has a projective variety V and a finite set $S = \{\phi_1, \dots, \phi_r\}$ of regular self-maps defined over a number field K that are polarizable with respect to a common ample invertible sheaf; that is, there is an ample invertible sheaf \mathcal{L} and integers $d_1, \dots, d_r \geq 2$ such that $\phi_i^*(\mathcal{L}) \cong \mathcal{L}^{\otimes d_i}$ for $i = 1, \dots, r$. We note that this polarizable condition holds for every set of regular maps that are not automorphisms when $V = \mathbb{P}^N$ with $\mathcal{L} = \mathcal{O}(1)$. Now let M_S denote the semigroup of endomorphisms of V generated by S under composition. In particular, when S is polarized, one has a natural notion of degree for each $f \in M_S$ by declaring that the degree of f , denoted by $\deg_{\mathcal{L}}(f)$, is the unique integer d such that $f^*\mathcal{L} \cong \mathcal{L}^{\otimes d}$. Likewise, we may choose

Received by the editors July 24, 2023; revised April 8, 2024; accepted July 8, 2024.

AMS subject classification: 37P15, 37P05, 37P55.

Keywords: arithmetic dynamics, semigroup dynamics, heights.



a height function $h_{\mathcal{L}} : V \rightarrow \mathbb{R}$ associated to \mathcal{L} , and to any point $P \in V(\overline{\mathbb{Q}})$ study the orbit growth of P , which is the function

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\}.$$

Then the general principle stated above suggests that the asymptotics of the orbit growth should be heavily constrained by the asymptotic behavior of the corresponding degree growth function

$$\#\{f \in M_S : \text{deg}_{\mathcal{L}}(f) \leq X\}.$$

It is relatively straightforward to give upper bounds for the orbit growth in terms of the degree growth. Indeed, Kawaguchi and Silverman [9] showed that the analogous phenomenon holds for dominant rational self-maps. However, it is very difficult in general to give strong lower bounds for the orbit growth in terms of the degree growth. The reason for this is that distinct maps in M_S can send a point P to a common point Q , and this sort of “collapse” can, in some cases, be extreme. For example, if P is a common fixed point of all elements of S , then the orbit growth function is $O(1)$, while the degree growth function can be exponential in X .

In an earlier work, the second-named author [6] considered orbit growth and degree growth for semigroups of polarizable morphisms. In this paper, we extend those results by getting optimal asymptotics for degree growth and getting optimal asymptotics for orbit growth for certain classes of semigroups of morphisms of the projective line. However, similar to the case of iterating a single function, we must avoid preperiodic points, defined below.

Definition 1.1 Let $S = \{\phi_1, \dots, \phi_r\}$ be a set of endomorphisms on a variety V . We say that a point $P \in V$ is *preperiodic* for M_S if there exist $f, g \in M_S$, with g not the identity, such that $g \circ f(P) = f(P)$.

Our first main result is the following. In what follows, given real functions F and G , we write $F \ll G$ if there is a constant C such that $F(x) \leq C \cdot G(x)$ holds for all x sufficiently large. Likewise, we write $F \asymp G$ to mean both $F \ll G$ and $G \ll F$. Finally, we write $F \sim G$ to mean $\lim_{x \rightarrow \infty} F(x)/G(x) = 1$.

Theorem 1.2 Let V be a projective variety, let $S := \{\phi_1, \dots, \phi_r\}$ be a polarized set of endomorphisms on V , and say $\phi_i^* \mathcal{L} \cong \mathcal{L}^{\otimes d_i}$ for some $d_i > 1$. Moreover, assume that $r \geq 2$, and let ρ be the unique real number satisfying

$$\frac{1}{d_1^\rho} + \dots + \frac{1}{d_r^\rho} = 1.$$

Then, the following statements hold for all non-preperiodic $P \in V$:

- (1) In all cases, $\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \ll X^\rho$.
- (2) If M_S contains a free, finitely generated sub-semigroup, then there is a positive constant $\rho' \leq \rho$ such that

$$X^{\rho'} \ll \#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \ll X^\rho.$$

- (3) If M_S is free, then $\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \asymp X^\rho$.

(4) If M_S is free and the d_1, \dots, d_r are not all pairwise multiplicatively dependent, then there is a positive constant c such that

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \sim cX^{\rho}.$$

Here in all cases, the implicit constants depend on P and S .

Remark 1.3 In particular, if the individual sets of preperiodic points $\text{Prep}(\phi)$ of each $\phi \in S$ are not all identical, then we obtain bounds of the form

$$X^{\rho'} \ll \#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \ll X^{\rho}$$

for all non-preperiodic $P \in V$ by [1, Theorem 1.3].

Remark 1.4 In fact, an exact asymptotic as in statement (4) of Theorem 1.2 is not possible when the degrees of the maps in S are all powers of a single integer. For a concrete example, consider the set of polynomials $S = \{2x^8, 3x^8\}$ and the point $P = 1$. Note that $\rho = 1/3$ in this case. Now for $X > 0$, define the function

$$\Theta(X) = \#\{f \in M_S : h(f(P)) \leq X\} / X^{1/3},$$

where $h(\cdot)$ is the usual Weil height on \mathbb{P}^1 . Then, one can show that

$$0 < \liminf \Theta(X) < \limsup \Theta(X) < \infty$$

in this case.

However, since every point in the total *semigroup orbit* of P ,

$$\text{Orb}_S(P) := \{f(P) : f \in M_S\},$$

of height at most X is determined by at least one function $f \in M_S$ satisfying $h_{\mathcal{L}}(f(P)) \leq X$, we easily obtain a general upper bound for the number of points of bounded height in non-preperiodic orbits. However, with some additional work, we can prove that such an upper bound holds for all orbits.

Theorem 1.5 Let V be a projective variety, and let $S := \{\phi_1, \dots, \phi_r\}$ be a polarized set of endomorphisms on V . Moreover, assume that $r \geq 2$ and that ρ and \mathcal{L} are as in Theorem 1.2. Then,

$$\#\{Q \in \text{Orb}_S(P) : h_{\mathcal{L}}(Q) \leq X\} \ll X^{\rho}$$

holds for all points $P \in V$.

However, we prove an even stronger result for generic sets of maps on $V = \mathbb{P}^1$ and establish an asymptotic for the number of *points* in an orbit of bounded height, instead of just an asymptotic for the number of *functions* yielding a bounded height relation. In what follows, given a positive integer d , we let Rat_d be the space of rational functions in one variable of degree d ; see [21, §4.3] for details.

Theorem 1.6 Let d_1, \dots, d_r be integers all at least four. Moreover, assume that $r \geq 2$, and let ρ be the unique positive real number satisfying $\sum d_i^{-\rho} = 1$. Then, there is a proper Zariski open subset $U \subseteq \text{Rat}_{d_1} \times \dots \times \text{Rat}_{d_r}$ such that

$$\#\{Q \in \text{Orb}_S(P) : h(Q) \leq X\} \asymp X^{\rho}$$

for all $S \in U$ and all P such that $\text{Orb}_S(P)$ is infinite. Moreover, if in addition the d_1, \dots, d_r are not all pairwise multiplicatively dependent, then

$$\#\{Q \in \text{Orb}_S(P) : h(Q) \leq X\} \sim cX^\rho$$

for some positive constant c , which depends on P and S .

Remark 1.7 The constants in Theorem 1.6 are not explicit since they incorporate height bounds for rational points on curves. However, the subset U is explicit and consists of all critically separate and critically simple sets of maps (cf. [6, Theorem 1.5]); see section 3 for details.

Remark 1.8 Similar hypotheses were recently used in [7] to give strong lower bounds for the number of points in semigroup orbits modulo primes. Moreover, these results are analogous to known results for abelian varieties [17]. With this analogy in mind, the number of points of height at most X on an abelian variety A over a number field K is asymptotic to $X^{\rho/2}$, where ρ is the rank of the Mordell-Weil group $A(K)$; see [19, p. 54]. Thus, the present paper, as well as the papers [6, 7], suggest that the analogy

$$\left(\begin{array}{l} \text{arithmetic of points} \\ \text{of an abelian variety} \end{array} \right) \iff \left(\begin{array}{l} \text{arithmetic of points in orbits} \\ \text{of a dynamical system} \end{array} \right)$$

described in [2] and [22, §6.5] may be more accurate when the dynamical system on the right-hand side is generated by at least two noncommuting maps, rather than using orbits coming from iteration of a single map.

1.1 Notation

Throughout this paper, we use the following notation. If S is a set of self-maps, we let M_S denote the semigroup generated by S under composition. If M_S is free on the set S , we let S_n (respectively, $S_{\leq n}$) denote the set of n -fold compositions of elements of S (respectively, elements that are compositions of at most n elements of S), and for $f \in M_S$, we let $|f|$ denote the unique natural number n such that f is an n -fold composition of elements of S .

2 Counting functions in polarized systems

2.1 Functions of bounded degree

Let V be a projective variety, and let $S = \{\phi_1, \dots, \phi_r\}$ be a set of endomorphisms on V . Moreover, we assume that S is *polarized* with respect to some ample line bundle \mathcal{L} on V , meaning that there are $d_1, \dots, d_r > 1$ satisfying $\phi_i^* \mathcal{L} \cong \mathcal{L}^{\otimes d_i}$. In particular, we may define the \mathcal{L} -degree (or simply degree when \mathcal{L} is understood) of a function f in the semigroup generated by S as follows:

$$\deg_{\mathcal{L}}(f) := \prod_{j=1}^n d_{i_j} \quad \text{for } f = \phi_{i_1} \circ \dots \circ \phi_{i_n} \in M_S.$$

In particular, $f^* \mathcal{L} \cong \mathcal{L}^{\otimes \deg_{\mathcal{L}}(f)}$ follows from standard results in algebraic geometry. However, if we fix a height function associated to \mathcal{L} on V and a point $P \in V$, then the

height of $f(P)$ tends to grow according to $\deg_{\mathcal{L}}(f)$ (a fact used by Call and Silverman to define canonical heights). Therefore, as a first step in counting points of bounded height in orbits, we count functions of bounded degree. In fact, we will prove in this section that if M_S is free, then there are positive constants ρ , c_1 , and c_2 (depending only on the initial degrees d_1, \dots, d_r) such that

$$(2.0.1) \quad c_1 X^\rho \leq \#\{f \in M_S : \deg_{\mathcal{L}}(f) \leq X\} \leq c_2 X^\rho$$

for all X sufficiently large. Here, the constant ρ is the unique positive real number satisfying

$$\frac{1}{d_1^\rho} + \dots + \frac{1}{d_r^\rho} = 1.$$

In fact, we can give more exact asymptotics. However, the form of these asymptotics turns out to depend on whether or not the degrees of the maps in S are all powers of a single integer, and so we put off stating these more precise results until later.

To begin, we collect some basic facts about counting words in free semigroups of bounded, *multiplicative* weight. This problem is very similar to a classical arithmetic problem of counting bounded multiplicative compositions of integers [3, 10], though not exactly the same, since we want to allow factors of equal size (precisely to allow maps of equal degree in our main dynamical theorems). In particular, due to the lack of an exact reference, we include the relevant statements and proofs here. Let us fix some notation. Let M be a multiplicative semigroup generated by a finite set $S = \{\alpha_1, \dots, \alpha_r\}$. We say that M is a *weighted semigroup* with respect to a vector $\mathbf{d} = (d_1, \dots, d_r)$ of positive real weights if there is a well-defined multiplicative weight function $|\cdot|_{\mathbf{d}} : M \rightarrow \mathbb{R}$ given by

$$(2.0.2) \quad |\omega|_{\mathbf{d}} := \prod_{j=1}^m d_{i_j}, \quad \omega = \alpha_{i_1} \cdots \alpha_{i_m} \in M.$$

In the case when M is the free semigroup on S , then every set of positive weights endows M with the structure of a weighted semigroup. In general, however, to be weighted, one requires that all representations of an element of M as a composition of elements of S have the same weight.

Then, given a real number X , we would like to obtain an asymptotic formula for the number of words of weight at most X ,

$$(2.0.3) \quad \#\{\omega \in M : |\omega|_{\mathbf{d}} \leq X\}.$$

In particular, we have the following general estimate, which assumes only that $r \geq 2$ and that $\min\{d_i\} > 1$.

Remark 2.1 Strictly speaking, the result below is not necessary for proving our main theorems in this paper; it is superseded by the asymptotics given in Theorem 2.6. Nevertheless, we include this result since it may be useful for future work. For one, the more precise asymptotics to come assume integral weights, and non-integral weights do appear elsewhere in dynamics [8, 20]. Hence, the result below may still be useful in a more general setting.

Proposition 2.2 *Let M be a semigroup generated by a set $S = \{\alpha_1, \dots, \alpha_r\}$, and suppose that M is weighted with respect to the vector $\mathbf{d} = (d_1, \dots, d_r)$ of positive real numbers. Assume that $r \geq 2$ and that $\min\{d_1, \dots, d_r\} > 1$, and let ρ be the unique real number satisfying $d_1^{-\rho} + \dots + d_r^{-\rho} = 1$. Then, there is a positive constant c_2 such that*

$$\#\{\omega \in M : |\omega|_{\mathbf{d}} \leq X\} \leq c_2 X^\rho$$

holds for all X sufficiently large. Moreover, if M is free on the set S , then there is a positive constant c_1 such that

$$c_1 X^\rho \leq \#\{\omega \in M : |\omega|_{\mathbf{d}} \leq X\}.$$

Proof We first consider the case when M is free, and we adjoin an identity to make M a monoid with identity element of weight 1. Let $w(x)$ be the number of elements of M of weight at most x . Then,

$$w(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1, \\ 1 + \sum_{i=1}^r w\left(\frac{x}{d_i}\right) & \text{if } 1 \leq x < \infty. \end{cases}$$

Indeed, there are no words of weight less than 1, and any nonempty word ω of weight at most x is uniquely of the form $\omega = \alpha_{i_1} \omega'$ for some word ω' of weight at most x/d_{i_1} (and there are exactly $w(x/d_{i_1})$ such ω'). Now let $W(x) := w(x) + \frac{1}{r-1}$, and note that it suffices to show that $W(x) \asymp x^\rho$, to conclude the same for $w(x)$; the reason we consider W instead of w is that it satisfies a slightly simpler recursion:

$$(2.2.1) \quad W(x) = \begin{cases} 1/(r-1) & \text{if } 0 \leq x < 1, \\ \sum_{i=1}^r W\left(\frac{x}{d_i}\right) & \text{if } 1 \leq x < \infty, \end{cases}$$

which follows easily from w 's functional equation. Now assume that the d 's are arranged in nondecreasing order, $d_1 \leq \dots \leq d_r$, and let $c_1 = W(1)/d_r^\rho$ and $c_2 = W(d_r)$. We will show that

$$(2.2.2) \quad c_1 x^\rho \leq W(x) \leq c_2 x^\rho \quad \text{for all } 1 \leq x \leq d_r \cdot d_1^n$$

and all $n \geq 0$ by induction on n . In particular, since $d_1 > 1$, it follows that $c_1 x^\rho \leq W(x) \leq c_2 x^\rho$ for all $x \geq 1$. Note first that if $1 \leq x \leq d_r$, then

$$\begin{aligned} c_1 x^\rho &= \frac{W(1)}{d_r^\rho} x^\rho \leq \frac{W(1)}{d_r^\rho} d_r^\rho = W(1) \leq W(x) \\ &\leq W(d_r) \leq W(d_r) \cdot x^\rho = c_2 x^\rho \end{aligned}$$

since $W(x)$ and x^ρ are an increasing function. Hence, (2.2.2) is true for $n = 0$. Now to the induction step: suppose that (2.2.2) holds and that $1 \leq x \leq d_r \cdot d_1^{n+1}$. In particular, since we have established (2.2.2) for all $1 \leq x \leq d_r$, we may assume without loss that $d_r \leq x \leq d_r \cdot d_1^{n+1}$. But then,

$$1 \leq \frac{d_r}{d_i} \leq \frac{x}{d_i} \leq d_r \cdot \frac{d_1}{d_i} \cdot d_1^n \leq d_r \cdot d_1^n$$

for all $1 \leq i \leq r$. In particular, $1 \leq x/d_i \leq d_r \cdot d_1^n$, and so we have that $c_1(x/d_i)^\rho \leq W(x/d_i) \leq c_2(x/d_i)^\rho$ by the induction hypothesis. Hence,

$$\begin{aligned} c_1x^\rho &= c_1x^\rho \left(\sum_{i=1}^r \frac{1}{d_i^\rho} \right) = \sum_{i=1}^r c_1 \left(\frac{x}{d_i} \right)^\rho \leq \sum_{i=1}^r W \left(\frac{x}{d_i} \right) \\ &\leq \sum_{i=1}^r c_2 \left(\frac{x}{d_i} \right)^\rho = c_2x^\rho \left(\sum_{i=1}^r \frac{1}{d_i^\rho} \right) = c_2x^\rho. \end{aligned}$$

However, (2.2.1) implies that $W(x) = \sum_{i=1}^r W(x/d_i)$, and we deduce that $c_1x^\rho \leq W(x) \leq c_2x^\rho$ for all $x \geq 1$, as claimed.

We note that when M is not free, the estimates giving the upper bound still hold since for general semigroups, we still have the inequality

$$W(x) \leq \sum_{i=1}^r W(x/d_i),$$

and so we can use the same proof verbatim to obtain the upper bound in the non-free case. ■

In particular, we immediately deduce the bounds in (2.0.1) for polarized sets of endomorphisms from Proposition 2.2 and the fact that the degrees of endomorphisms are multiplicative with respect to composition. However, it is possible to obtain more exact asymptotics for the quantity in (2.0.3) in the case of integer weights by analyzing the analytic properties of an associated generating function. With this in mind, we assume for the remainder of this section that the vector \mathbf{d} consists of positive integer weights. Then for any $n \in \mathbb{Z}$, we let

$$(2.2.3) \quad a_{\mathbf{d},n} := \#\{\omega \in F_S : |\omega|_{\mathbf{d}} = n\}$$

be the number of words of weight equal to n and let

$$(2.2.4) \quad \mathcal{D}_{\mathbf{d}}(s) := \sum_{n \geq 0} a_{\mathbf{d},n} n^{-s}$$

be the associated Dirichlet series generating function. In particular, we will improve upon Proposition 2.2 in this case by applying known Tauberian theorems to (2.2.4). However, there is a catch: we must make two separate cases depending on whether the generating weights are all powers of a single integer or not. With this in mind, we make the following definition.

Definition 2.3 A vector $\mathbf{d} = (d_1, \dots, d_r)$ of positive integers is called *cyclic* if there exists a single positive integer d and a collection of positive integers n_1, \dots, n_r such that $d_j = d^{n_j}$ for all $j \geq 1$. Moreover, if this is not the case, then we call \mathbf{d} *acyclic*. If (d_1, \dots, d_r) is cyclic, then there is a unique positive integer d such that $d_i = d^{a_i}$ for some positive integers a_1, \dots, a_r with $\gcd(a_1, \dots, a_r) = 1$, and we call this integer d the *base* of \mathbf{d} .

We use the following elementary fact from arithmetic.

Lemma 2.4 Let d_1, \dots, d_r be positive integers all at least 2. If the ratio $\log(d_i)/\log(d_j) \in \mathbb{Q}$ for all i, j , then the vector $\mathbf{d} = (d_1, \dots, d_r)$ is cyclic.

Remark 2.5 When d, e are positive integers with $\log(d)/\log(e) \in \mathbb{Q}$, we say that d and e are *multiplicatively dependent*, and when this does not hold, we say that d and e are *multiplicatively independent*. Then, Lemma 2.4 gives that $\mathbf{d} = (d_1, \dots, d_r)$ is cyclic if and only if the d 's are all pairwise multiplicatively dependent.

With these notions in place, we now establish an exact asymptotic for the number of words of bounded multiplicative weight in free semigroups.

Theorem 2.6 Let $S = \{\alpha_1, \dots, \alpha_r\}$ be an alphabet, let F_S be the free semigroup generated by S , and let $\mathbf{d} = (d_1, \dots, d_r)$ be a vector of positive integers all at least 2. Moreover, assume that $r \geq 2$, and let $\rho_{\mathbf{d}} > 0$ be the unique real number satisfying $G_{\mathbf{d}}(\rho_{\mathbf{d}}) = 1$, where

$$G_{\mathbf{d}}(s) = d_1^{-s} + d_2^{-s} + \dots + d_r^{-s}.$$

Then, the following statements hold:

- (1) If \mathbf{d} is acyclic, then

$$\mathcal{D}_{\mathbf{d}}(s) = \frac{1}{1 - G_{\mathbf{d}}(s)}$$

converges for all $\Re(s) \geq \rho_{\mathbf{d}}$ except for a simple pole at $s = \rho_{\mathbf{d}}$. Hence, the Wiener-Ikehara Tauberian theorem [16, §8.3] implies that

$$\#\{\omega \in F_S : |\omega|_{\mathbf{d}} \leq X\} = \sum_{n \leq X} a_{\mathbf{d},n} = \frac{-1}{\rho_{\mathbf{d}} G'(\rho_{\mathbf{d}})} X^{\rho_{\mathbf{d}}} + o(X^{\rho_{\mathbf{d}}})$$

as X tends to infinity.

- (2) If \mathbf{d} is cyclic of base d , then there is a positive constant C such that

$$\#\{\omega \in F_S : |\omega|_{\mathbf{d}} \leq X\} = C \cdot d^{\lfloor \log_d(X) \rfloor \rho_{\mathbf{d}}} + o(d^{\lfloor \log_d(X) \rfloor \rho_{\mathbf{d}}})$$

as X tends to infinity.

Before we can establish Theorem 2.6, we need an elementary fact, which is a simple consequence of the strong triangle inequality.

Lemma 2.7 Let $z_1, \dots, z_r \in \mathbb{C}$ not all zero be such that

$$z_1 + \dots + z_n = |z_1| + \dots + |z_n|.$$

Then, $z_i = |z_i|$ for all i (i.e., $\arg(z_i) = 2k_i\pi$ for some $k_i \in \mathbb{Z}$).

Proof of Theorem 2.6 For all $m \geq 0$, let $F_S(m)$ be the set of sequences of length m in F_S . Then, it follows from the definition of weight in (2.0.2) that

$$G_{\mathbf{d}}(s)^m = \sum_{\omega \in F_S(m)} |\omega|_{\mathbf{d}}^{-s} = \sum_{n \geq 1} \sum_{|\omega|_{\mathbf{d}}=n} n^{-s} = \sum_{n \geq 1} a_{\mathbf{d},n,m} n^{-s},$$

where $a_{\mathbf{d},n,m} = \#\{\omega \in F_S(m) : |\omega|_{\mathbf{d}} = n\}$. In particular, we see that

$$\begin{aligned} \frac{1}{1 - G_{\mathbf{d}}(s)} &= \sum_{m \geq 0} G_{\mathbf{d}}(s)^m = \sum_{m \geq 0} \sum_{n \geq 1} (a_{\mathbf{d},n,m}) n^{-s} \\ &= \sum_{n \geq 1} \sum_{m \geq 0} (a_{\mathbf{d},n,m}) n^{-s} = \sum_{n \geq 1} a_{\mathbf{d},n} n^{-s} = \mathcal{D}_{\mathbf{d}}(s) \end{aligned}$$

holds formally. As for the convergence of this expression, note first that since $d_i \geq 2$ for all i , $G_{\mathbf{d}}(s)$ is a strictly decreasing continuous function on the real line. Moreover, $G_{\mathbf{d}}(0) = r \geq 2$ and $\lim_{x \rightarrow \infty} G_{\mathbf{d}}(x) = 0$, and so there is a unique, simple, real solution $x = \rho_{\mathbf{d}} > 0$ to the equation $G_{\mathbf{d}}(x) = 1$. However, for all $\Re(s) > \rho_{\mathbf{d}}$, we have that

$$\begin{aligned} |G_{\mathbf{d}}(s)| &= |d_1^{-s} + d_2^{-s} + \dots + d_r^{-s}| \\ &\leq |d_1^{-s}| + |d_2^{-s}| + \dots + |d_r^{-s}| \\ &= d_1^{-\Re(s)} + d_2^{-\Re(s)} + \dots + d_r^{-\Re(s)} \\ &< d_1^{-\rho_{\mathbf{d}}} + d_2^{-\rho_{\mathbf{d}}} + \dots + d_r^{-\rho_{\mathbf{d}}} \\ &= 1. \end{aligned}$$

In particular, the meromorphic function $\mathcal{D}_{\mathbf{d}}(s) = (1 - G_{\mathbf{d}}(s))^{-1}$ converges in $\Re(s) > \rho_{\mathbf{d}}$ and has a simple pole at $s = \rho_{\mathbf{d}}$. However, this is, in fact, the only pole on the line $\Re(s) = \rho_{\mathbf{d}}$. To see this, suppose that $G_{\mathbf{d}}(s) = 1$ for some $\Re(s) = \rho_{\mathbf{d}}$, and write $s = \rho_{\mathbf{d}} + ci$ for some $c \in \mathbb{R}$. Then setting $z_i = d_i^{-s}$, we deduce from Lemma 2.7 that

$$\log(d_i)c = 2\pi k_i$$

for some $k_i \in \mathbb{Z}$. In particular, if $c \neq 0$, then $\log(d_i)/\log(d_j) \in \mathbb{Q}$ for all i, j , and Lemma 2.4 implies that \mathbf{d} is cyclic. Therefore, if \mathbf{d} is acyclic, then $\mathcal{D}_{\mathbf{d}}(s) = (1 - G_{\mathbf{d}}(s))^{-1}$ converges whenever $\Re(s)$ is at least $\rho_{\mathbf{d}}$, except for the simple pole at $s = \rho_{\mathbf{d}}$. This completes the proof of statement (1).

However, the cyclic case (say with base d) can be handled with somewhat simpler analysis. Write $d_i = d^{a_i}$ for some positive integers a_i with $\gcd(a_1, \dots, a_r) = 1$. Since F_S is free on $S = \{\alpha_1, \dots, \alpha_r\}$, there are exactly $\binom{n_1 + \dots + n_r}{n_1, \dots, n_r}$ words with n_i occurrences of α_i for all $i = 1, \dots, r$. Moreover, the weight of each such element is exactly $d^{\sum a_i n_i}$. In particular, the number of elements in F_S of weight $\leq d^L$ is the sum of the coefficients of x^i for $i = 0, \dots, L$ in the polynomial

$$\sum_{n \geq 0} (x^{a_1} + \dots + x^{a_r})^n.$$

In particular, we see that for a positive integer L , the number of elements in F_S of weight $\leq d^L$ is the coefficient of x^L in the formal power series

$$F(x) := (1 - x)^{-1} \cdot (1 - x^{a_1} - \dots - x^{a_r})^{-1}.$$

Observe that $G(x) := 1 - x^{a_1} - \dots - x^{a_r}$ is strictly decreasing on $(0, \infty)$, and since the function is 0 when $x = 0$ and is negative when $x = 1$, it has a unique zero θ in $(0, 1)$. Since the derivative of G is negative on $(0, \infty)$, this is a simple zero. Observe further

that $x = \theta$ is the unique pole of $F(x)$ in the closed disc of radius θ . To see this, observe that if β is another pole in this disc, then we must have $G(\beta) = 0$. Then,

$$|G(\beta)| \geq 1 - |\beta|^{a_1} - \dots - |\beta|^{a_r} \geq 1 - \theta^{a_1} - \dots - \theta^{a_r} = 0,$$

with equality only if $|\beta| = \theta$. Thus, it suffices to consider β of the form $\theta e^{2\pi i \mu}$. Then, $G(\beta) = 1 - \theta^{a_1} \exp(2\pi i a_1 \mu) - \dots - \theta^{a_r} \exp(2\pi i a_r \mu)$, and since $\sum \theta^{a_j} = 1$, this can only be zero if $\exp(2\pi i a_1 \mu) = \dots = \exp(2\pi i a_r \mu) = 1$. Since the a_j have gcd 1, we then see this forces μ to be an integer, and so $\beta = \theta$ as required. Using partial fractions and the fact that $F(x)$ has a simple pole at $x = \theta$ and no other poles in the closed disc of radius θ , we see that

$$F(x) = C/(1 - x/\theta) + F_0(x)$$

for some positive constant C and some rational function $F_0(x)$ whose radius of convergence is strictly larger than θ . In particular, the coefficient of x^L in $F(x)$ is asymptotic to $C\theta^L$, and so we have that

$$\#\{\omega \in F_S : |\omega|_{\mathbf{d}} \leq d^L\} \sim C\theta^L.$$

Since θ is the unique positive solution to $\sum \theta^{a_i} = 1$, we see that if ρ is such that $d^\rho = \theta$, then ρ is the unique positive solution to $\sum 1/d_i^\rho = 1$, and so $\rho = \rho_{\mathbf{d}}$, and we have

$$\#\{\omega \in F_S : |\omega|_{\mathbf{d}} \leq d^L\} \sim C(d^L)^{\rho_{\mathbf{d}}}.$$

Now for X large, there is a unique L such that $d^L \leq X < d^{L+1}$. Hence, in general, we have that

$$\#\{\omega \in F_S : |\omega|_{\mathbf{d}} \leq X\} \sim Cd^{\lfloor \log_d(X) \rfloor \cdot \rho_{\mathbf{d}}},$$

a fact equivalent to statement (2). ■

In particular, we obtain an improved asymptotic for the number of functions of bounded degree in free semigroups generated by polarized sets.

Corollary 2.8 *Let V be a projective variety, let $S := \{\phi_1, \dots, \phi_r\}$ be a polarized set of endomorphisms on V , and say $\phi_i^* \mathcal{L} \cong \mathcal{L}^{\otimes d_i}$ for some integers $d_i \geq 2$. Moreover, assume that $r \geq 2$, that M_S is free, and that ρ is the real number satisfying $d_1^{-\rho} + \dots + d_r^{-\rho} = 1$. Then the following statements hold.*

(1) *If $\mathbf{d} = (d_1, \dots, d_r)$ is acyclic, there is a positive constant c such that*

$$\#\{f \in M_S : \deg_{\mathcal{L}}(f) \leq X\} \sim cX^\rho.$$

(2) *If $\mathbf{d} = (d_1, \dots, d_r)$ is cyclic of base d , there is a positive constant C such that*

$$\#\{f \in M_S : \deg_{\mathcal{L}}(f) \leq X\} \sim C \cdot d^{\lfloor \log_d(X) \rfloor \cdot \rho}.$$

Remark 2.9 We note that our results on degrees in this section work over an arbitrary ground field. However, for the remainder of this paper (when we instead count by heights), we will restrict ourselves to number fields.

2.2 From degrees to heights

Now that we have a handle on the growth of degrees in polarized semigroups, we use this to study the growth of heights in orbits. To do this, we fix some notation. As before, let $S = \{\phi_1, \dots, \phi_r\}$ be a polarized set of endomorphisms on a projective variety V with $\phi_i^* \mathcal{L} \cong \mathcal{L}^{\otimes d_i}$ for some $d_i > 1$ and some ample line bundle \mathcal{L} . Moreover, we may choose a height function $h_{\mathcal{L}} : V \rightarrow \mathbb{R}_{\geq 0}$ associated to \mathcal{L} , well-defined up to $O(1)$. Then the basic tool that makes the conversion between degrees and heights possible is the following fact: for each ϕ_i there is a constant C_i such that

$$(2.9.1) \quad \left| h_{\mathcal{L}}(\phi_i(P)) - d_i h_{\mathcal{L}}(P) \right| \leq C_i$$

holds for all $P \in V$; see, for instance, [21, Theorem 7.29]. In particular, one may iterate the bound above and use a telescoping argument (due to Tate) to obtain a similar statement for semigroups; see [5, Lemma 2.1].

Lemma 2.10 *Let $d_S = \min\{d_i\}$, and let $C_S = \max\{C_i\}$ be as in (2.9.1). Then,*

$$\left| \frac{h_{\mathcal{L}}(f(P))}{\deg_{\mathcal{L}}(f)} - h_{\mathcal{L}}(P) \right| \leq \frac{C_S}{d_S - 1}$$

holds for all $f \in M_S$ and all $P \in V$.

In particular, we may deduce the following result, which says first, that the set of preperiodic points of M_S is a set of bounded height, and second, that if P is not a preperiodic point, then given a positive constant B , there are at most finitely many elements $f \in M_S$ for which $h_{\mathcal{L}}(f(P)) < B$. We note that this does not require freeness of the semigroup M_S but heavily uses the fact that the endomorphisms in S are polarizable.

Lemma 2.11 *There is a positive constant C such that*

- (1) *if P is a preperiodic point for the semigroup M_S , then $h_{\mathcal{L}}(P) \leq C$;*
- (2) *if $h_{\mathcal{L}}(P) > C$, then $h_{\mathcal{L}}(f(P)) > h_{\mathcal{L}}(P)$ for all $f \in M_S$.*

In particular, if P is not preperiodic for M_S and if B is a positive constant, then there is an $N := N(V, S, P, B)$ such that $h_{\mathcal{L}}(f(P)) > B$ whenever there is some $m > N$ such that $f \in M_S$ can be expressed as a length- m composition of elements of S .

Proof Notice that if P is a preperiodic point for M_S and $Q := f(P) = g \circ f(P)$, then

$$\left| \frac{h_{\mathcal{L}}(g(Q))}{\deg_{\mathcal{L}}(g)} - h_{\mathcal{L}}(Q) \right| \leq \frac{C_S}{d_S - 1},$$

and since $g(Q) = Q$ and $\deg_{\mathcal{L}}(g) \geq 2$, we then see that

$$h_{\mathcal{L}}(Q) \leq 2C_S \cdot (d_S - 1)^{-1}.$$

Similarly, we have

$$\left| \frac{h_{\mathcal{L}}(f(P))}{\deg_{\mathcal{L}}(f)} - h_{\mathcal{L}}(P) \right| \leq \frac{C_S}{d_S - 1},$$

and since $f(P) = Q$, we use the inequality on the height of Q to get that $h_{\mathcal{L}}(P) \leq 2 \frac{C_S}{d_S - 1}$. Also, observe that if $h_{\mathcal{L}}(P) \geq 2C_S$, then if $f \in M_S$, then Lemma 2.10 gives that $h_{\mathcal{L}}(f(P))/\deg_{\mathcal{L}}(f) > h_{\mathcal{L}}(P)/2$, and so $h_{\mathcal{L}}(f(P)) > h_{\mathcal{L}}(P)$ for all $f \in M_S$. In particular, setting $C = 2C_S$, we deduce both claims (1) and (2).

For the final claim, assume that P is not preperiodic for M_S and that $B > 0$. Note that without loss of generality, we may assume that $B \geq C$. Now let K be a number field over which P and every map in S is defined. Then, $f(P) \in V(K)$ for all $f \in M_S$, and

$$N := \#\{Q \in V(K) : h_{\mathcal{L}}(P) \leq B\}$$

is finite by Northcott’s theorem. We claim that we can take $N(V, S, P, B)$ equal to this N . To see this, suppose that $f = \theta_m \circ \theta_{m-1} \circ \dots \circ \theta_1$ for some $\theta_i \in S$ with $m > N$ and that $h_{\mathcal{L}}(f(P)) \leq B$. For $1 \leq i \leq m$, we let $f_i := \theta_i \circ \theta_{i-1} \circ \dots \circ \theta_1$. Note first that if $h_{\mathcal{L}}(f_i(P)) > B \geq C$ for some i , then $h_{\mathcal{L}}(f(P)) > B$ also by property (2). Therefore, we may assume that $h_{\mathcal{L}}(f_i(P)) \leq B$ for all $i \leq m$. However, $m > N$, so that $f_i(P) = f_j(P)$ for some $i > j$ by the pigeon-hole principle and the definition of N . But then,

$$g_{ij} \circ f_j(P) = f_i(P) = f_j(P)$$

for $g_{ij} = \theta_i \circ \dots \circ \theta_{j+1}$. Since $\theta_1, \dots, \theta_m$ are polarizable, we see that g_{ij} is not the identity map, and this now contradicts our assumption that P is not preperiodic. Therefore, $h_{\mathcal{L}}(f(P)) > B$, as claimed. ■

Lastly, before proving Theorem 1.2, we need the following result, which provides a key new ingredient for improving the estimates in [6].

Lemma 2.12 *Suppose that M_S is free and that P is not a preperiodic point of V . Then, there is a positive constant $\beta = \beta(S, \mathcal{L}, P)$ such that*

$$\lim_{n \rightarrow \infty} \sum_{g \in S_n} h_{\mathcal{L}}(g(P))^{-\rho} = \beta.$$

Proof For n , let $\beta_n = \beta_n(P) = \sum_{g \in S_n} h_{\mathcal{L}}(g(P))^{-\rho}$. Since M_S is free, S_{n+1} is the disjoint union of $\phi_1 \circ S_n, \dots, \phi_r \circ S_n$. Then,

$$|h_{\mathcal{L}}(\phi_i(f(P))) - d_i h_{\mathcal{L}}(f(P))| \leq C_S,$$

and so

$$d_i h_{\mathcal{L}}(f(P)) - C_S \leq h_{\mathcal{L}}(\phi_i(f(P))) \leq d_i h_{\mathcal{L}}(f(P)) + C_S.$$

In particular, the mean value theorem gives the inequality

$$|h_{\mathcal{L}}(\phi_i(f(P)))^{-\rho} - (d_i h_{\mathcal{L}}(f(P)))^{-\rho}| \leq C_S \rho |d_i \cdot h_{\mathcal{L}}(f(P)) - C_S|^{-\rho-1}.$$

Assume first that $h_{\mathcal{L}}(P) > C := 2C_S$. (We note that this C is the constant used in the proof of Lemma 2.11.) Then, since $\sum_{i=1}^r 1/d_i^{\rho} = 1$, we have

$$\begin{aligned} |\beta_{n+1} - \beta_n| &= \left| \sum_{i=1}^r \sum_{f \in S_n} h_{\mathcal{L}}(\phi_i \circ f(P))^{-\rho} - (d_i h_{\mathcal{L}}(f(P)))^{-\rho} \right| \\ (2.12.1) \quad &\leq \sum_{i=1}^r \sum_{f \in S_n} C_S \rho |d_i \cdot h_{\mathcal{L}}(f(P)) - C_S|^{-\rho-1}. \end{aligned}$$

By Lemma 2.10, we have

$$h_{\mathcal{L}}(f(P)) \geq \deg_{\mathcal{L}}(f)h_{\mathcal{L}}(P) - \deg_{\mathcal{L}}(f) \frac{C_S}{d_S - 1} > \deg_{\mathcal{L}}(f) \left(2C_S - \frac{C_S}{d_S - 1} \right),$$

and so $h_{\mathcal{L}}(f(P)) > C_S \deg_{\mathcal{L}}(f)$. It follows that

$$d_i h_{\mathcal{L}}(P) - C_S > C_S (d_i \deg_{\mathcal{L}}(f) - 1),$$

and so using Equation (2.12.1), we see

$$\begin{aligned} |\beta_{n+1} - \beta_n| &\leq \sum_{i=1}^r \sum_{f \in \mathcal{S}_n} \frac{C_S \rho}{(C_S (d_i \deg_{\mathcal{L}}(f) - 1))^{\rho+1}} \\ &\leq C_S^{-\rho} 2^{\rho+1} \rho \sum_{i=1}^r \sum_{f \in \mathcal{S}_n} 1 / (d_i \deg_{\mathcal{L}}(f))^{\rho+1} \\ &= C_S^{-\rho} 2^{\rho+1} \rho \sum_{j_1 + \dots + j_r = n+1} \binom{n+1}{j_1, \dots, j_r} d_1^{-j_1(\rho+1)} \dots d_r^{-j_r(\rho+1)} \\ &= C_S^{-\rho} 2^{\rho+1} \rho \cdot (1/d_1^{\rho+1} + \dots + 1/d_r^{\rho+1})^{n+1}. \end{aligned}$$

In particular, since $C' := \sum 1/d_i^{\rho+1} < 1$ and since $\sum_j (C')^j$ converges, we see that $\{\beta_n(P)\}$ is a Cauchy sequence converging to some $\beta(P)$, as claimed. This completes the proof when $h_{\mathcal{L}}(P) \geq 2C_S$. For the general case, assume only that P is not preperiodic for S . Then Lemma 2.11 implies that there exists N such that $h_{\mathcal{L}}(f(P)) > 2C_S$ for all $|f| \geq N$. However, for $n > N$, we have that

$$\sum_{g \in \mathcal{S}_n} h_{\mathcal{L}}(g(P))^{-\rho} = \sum_{f \in \mathcal{S}_N} \sum_{F \in \mathcal{S}_{n-N}} h_{\mathcal{L}}(F(f(P)))^{-\rho} = \sum_{f \in \mathcal{S}_N} \beta_{n-N}(f(P))$$

since M_S is free. Moreover, for each $f \in M_S$, the sequence $\{\beta_{n-N}(f(P))\}$ converges by our argument above. Therefore, $\{\beta_n(P)\}$, a finite sum of convergent sequences, must converge also. ■

In particular, we may use Lemma 2.10, Lemma 2.12, and the degree growth estimates in Section 2 to bound the number of functions in a polarized semigroup yielding a bounded height relation; see Theorem 1.2 from the Introduction.

Proof of Theorem 1.2 Assume that $r = \#S \geq 2$, let ρ be the unique positive number satisfying $d_1^{-\rho} + \dots + d_r^{-\rho} = 1$, and let $b_S = C_S / (d_S - 1)$; here, d_S and C_S are the constants in Lemma 2.10. We first prove (1) in the case when $P \in V$ satisfies $h_{\mathcal{L}}(P) > b_S$ and then show how the general non-preperiodic case follows from this one. In particular, if $h_{\mathcal{L}}(P) > b_S$, then Lemma 2.10 implies

$$\begin{aligned} \left\{ f \in M_S : \deg_{\mathcal{L}}(f) \leq \frac{X}{h_{\mathcal{L}}(P) + b_S} \right\} &\subseteq \left\{ f \in M_S : h_{\mathcal{L}}(f(P)) \leq X \right\} \\ &\subseteq \left\{ f \in M_S : \deg_{\mathcal{L}}(f) \leq \frac{X}{h_{\mathcal{L}}(P) - b_S} \right\}. \end{aligned}$$

In particular, Proposition 2.2, using our degree function to assign weights to elements of our semigroup, gives that there is a constant c such that

$$\begin{aligned} \#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} &\leq \#\left\{f \in M_S : \deg_{\mathcal{L}}(f) \leq \frac{X}{h_{\mathcal{L}}(P) - b_S}\right\} \\ &\leq c\left(\frac{X}{h_{\mathcal{L}}(P) - b_S}\right)^p \end{aligned}$$

holds for all X sufficiently large.

Hence, we obtain statement (1) in the case when $h_{\mathcal{L}}(P) > b_S$, as claimed. In general, if P is not preperiodic, then by Lemma 2.11, there are only finitely many elements f of our semigroup for which $h_{\mathcal{L}}(f(P)) \leq b_S$. In particular, there is some N such that if $g = \phi_{i_1} \circ \dots \circ \phi_{i_N} \in M_S$, then either g is equivalent in M_S to a composition of elements of S of length $< N$ or $h_{\mathcal{L}}(g(P)) > b_S$. Let g_1, \dots, g_r denote the length- N compositions of elements of S that are not equivalent in M_S to a shorter composition of elements of S .

Then, since all but finitely many elements of M_S have a right compositional factor from $\{g_1, \dots, g_r\}$, we have

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \leq \sum_{i=1}^r \#\{f \in M_S : h_{\mathcal{L}}(f(g_i(P))) \leq X\} + O(1).$$

Since $h_{\mathcal{L}}(g(P)) > b_S$, we see that each of the summands on the right side is $\ll X^p$, and so we deduce (1) in general.

We next prove (3) and (4) in the case when $h_{\mathcal{L}}(P) > b_S$, and so we now assume that M_S is free. Then, Proposition 2.2 and Lemma 2.10 imply that there is a positive constant c' such that

$$\begin{aligned} \#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} &\geq \#\left\{f \in M_S : \deg_{\mathcal{L}}(f) \leq \frac{X}{h_{\mathcal{L}}(P) + b_S}\right\} \\ &\geq c'\left(\frac{X}{h_{\mathcal{L}}(P) + b_S}\right)^p \end{aligned}$$

holds for all X sufficiently large. Combining this inequality with (1), we see that $\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \asymp X^p$ for all sufficiently large X , as claimed in statement (3).

To prove (4) in the case when $h_{\mathcal{L}}(P) > b_S$, suppose that M_S is free and that the vector of degrees $\mathbf{d} = (d_1, \dots, d_r)$ is acyclic. Moreover, let $\varepsilon > 0$, and let

$$\beta := \lim_{n \rightarrow \infty} \sum_{g \in S_n} h_{\mathcal{L}}(g(P))^{-p}$$

be the constant from Lemma 2.12. Then, Lemma 2.11 gives that there are only finitely many elements of M_S with $h_{\mathcal{L}}(g(P)) \leq b_S/\varepsilon$. Thus, there is a natural number n such that

$$\left| \sum_{g \in S_n} h_{\mathcal{L}}(g(P))^{-p} - \beta \right| < \varepsilon$$

and

$$h_{\mathcal{L}}(g(P)) - b_S > (1 - \varepsilon)h_{\mathcal{L}}(g(P))$$

for all $g \in S_n$. Then, statement (1) of Corollary 2.8 implies that for all $g \in S_n$ and all large X , we have that

$$\begin{aligned} & \#\{f \in M_S : h_{\mathcal{L}}(f \circ g(P)) \leq X\} \\ & \leq \#\left\{f \in M_S : \text{deg}_{\mathcal{L}}(f) \leq \frac{X}{h_{\mathcal{L}}(g(P)) - b_S}\right\} \\ & \leq c(1 + \varepsilon)\left(\frac{X}{h_{\mathcal{L}}(g(P)) - b_S}\right)^{\rho} \\ & \leq c(1 + \varepsilon)(1 - \varepsilon)^{-\rho} X^{\rho} / h_{\mathcal{L}}(g(P))^{\rho}. \end{aligned}$$

However, for large X , we have that

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} = \#S_{<n} + \sum_{g \in S_n} \#\{f \in M_S : h_{\mathcal{L}}(f \circ g(P)) \leq X\},$$

and so we get that

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \leq \#S_{<n} + c(1 + \varepsilon)(1 - \varepsilon)^{\rho} X^{\rho} (\beta + \varepsilon)$$

for X sufficiently large. Moreover, by a completely analogous argument, we get a lower bound of

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} \geq \#S_{<n} + c(1 - \varepsilon)(1 + \varepsilon)^{\rho} X^{\rho} (\beta - \varepsilon)$$

for all large X . Then, since n is fixed and ε is arbitrarily small, we see that

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} = c\beta X^{\rho}(1 + o(1)),$$

as claimed in statement (4).

Next, we assume only that P is a non-preperiodic for M_S and prove (3) and (4) in this case. In particular, we still assume that M_S is free. Then, Lemma 2.11 implies that there exists N such that $h_{\mathcal{L}}(g(P)) > b_S$ for all $|g| \geq N$. In particular, for all large X , we have that

$$\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\} = \#S_{\leq N} + \sum_{g \in S_N} \#\{f \in M_S : h_{\mathcal{L}}(f(g(P))) \leq X\}.$$

However, by replacing P with $g(P)$ in our arguments above, we see that each of the finitely many terms $\#\{f \in M_S : h_{\mathcal{L}}(f(g(P))) \leq X\}$ for $g \in S_N$ exhibit growth according to statements (3) and (4) of Theorem 1.2, assuming the corresponding hypotheses. In particular, the total count $\#\{f \in M_S : h_{\mathcal{L}}(f(P)) \leq X\}$ exhibits the desired growth also.

Finally, we observe that (2) quickly follows from (3). To see this, if M_S contains a free semigroup with finite basis $S' \subset M_S$, let ρ' be the unique positive number

satisfying $\sum_{g \in S'} 1/\deg(g)^{\rho'} = 1$. Then, (3) gives that there is a positive constant c'' such that

$$\#\{f \in M_{S'} : h_{\mathcal{L}}(f(P)) \leq X\} \geq c'' \left(\frac{X}{h_{\mathcal{L}}(P) + b_S} \right)^{\rho'}$$

holds for all X sufficiently large. In particular, we deduce statement (2) as claimed; moreover, clearly $\rho' \leq \rho$ since $X^{\rho'} \ll X^{\rho}$ by combining the bounds in statements (1) and (2) with the fact that $M_{S'} \subseteq M_S$. ■

Lastly, we note that our upper bound on the number of functions $f \in M_S$ such that $h_{\mathcal{L}}(f(P)) \leq X$ for non-preperiodic basepoints P leads to a general upper bound for the number of points of bounded height in orbits; see Theorem 1.5 from the Introduction. The last step is the following decomposition lemma.

Lemma 2.13 *Let $S = \{\phi_1, \dots, \phi_r\}$ be a polarizable set of endomorphisms on a variety V , let $P \in V$ be such that $\text{Orb}_S(P)$ is infinite, and let $B > 0$. Then, there is a finite subset $F \subseteq \text{Orb}_S(P)$ and distinct points Q_1, \dots, Q_n in $\text{Orb}_S(P)$ such that the following statements hold:*

- (1) *Each Q_i satisfies $h_{\mathcal{L}}(Q_i) > B$.*
- (2) *$\text{Orb}_S(P) = F \cup \text{Orb}_S(Q_1) \cup \dots \cup \text{Orb}_S(Q_n)$.*

In particular, we may assume that Q_1, \dots, Q_n are not preperiodic for S .

Proof Let C be the constant in Lemma 2.11. As a reminder, $C = 2C_S$, where C_S is the constant from Lemma 2.10. Moreover, let $d = \max\{d_i\}$, where d_i is as in (2.9.1). Now let $P \in V$ have infinite orbit and $B > 0$. Without loss, we may assume $B \geq C$. Note that the desired conclusion follows easily if $h_{\mathcal{L}}(P) > B$ by taking $n = 1$, $P = Q_1$, and $F = \emptyset$. Therefore, we may assume that $h_{\mathcal{L}}(P) \leq B$. Now define the set

$$T := \{Q \in \text{Orb}_S(P) : B < h_{\mathcal{L}}(Q) \leq 2dB\}.$$

In particular, $T = Q_1, \dots, Q_n$ is a finite (possibly empty) set by Northcott’s theorem (seen by working over a fixed number field K over which P and every map in S is defined). Likewise, $F := \{Q \in \text{Orb}_S(P) : h_{\mathcal{L}}(Q) \leq B\}$ is a finite set, and we will show that F and the points Q_i have the desired properties. To see this, suppose that $Q = \theta_m \circ \dots \circ \theta_1(P)$ for some $\theta_i \in S$ satisfies $h_{\mathcal{L}}(Q) > B$ (such Q must exist since $\text{Orb}_S(P)$ is infinite), and define

$$m_0 = \min \{s : h_{\mathcal{L}}(\theta_s \circ \dots \circ \theta_1(P)) > B\}.$$

Note that m_0 exists since m is in the defining set above. Moreover, $m_0 \geq 1$ since $h_{\mathcal{L}}(P) \leq B$ by assumption. We claim that $Q' := \theta_{m_0} \circ \dots \circ \theta_1(P) \in T$ (in particular, T is nonempty): if not, then necessarily $h_{\mathcal{L}}(Q') > 2dB$. However, we then see that

$$\begin{aligned} 2dB < h_{\mathcal{L}}(Q') &= h_{\mathcal{L}}(\theta_{m_0}(\theta_{m_0-1} \circ \dots \circ \theta_1(P))) \\ &\leq \deg(\theta_{m_0})(h_{\mathcal{L}}(\theta_{m_0-1} \circ \dots \circ \theta_1(P)) + C_S) \leq 2dB, \end{aligned}$$

a contradiction; here, we use Lemma 2.10 and the minimality of m_0 . In particular, we have shown that for any $Q \in \text{Orb}_S(P)$ of height bigger than B , there is a point

$Q_i \in T$ such that $Q \in \text{Orb}_S(Q_i)$. Hence, we deduce that $\text{Orb}_S(P) = F \cup \text{Orb}_S(Q_1) \cup \dots \cup \text{Orb}_S(Q_n)$, as claimed. Finally, we note that since each $h_{\mathcal{L}}(Q_i) > B$ and we have taken $B \geq 2C_S$, which is the constant C in the proof of Lemma 2.11, we have that the Q_i are not preperiodic by Lemma 2.11. ■

In particular, since every point in $\text{Orb}_S(P)$ of height at most X is determined by at least one function $f \in M_S$ satisfying $h_{\mathcal{L}}(f(P)) \leq X$ and (outside of a finite set) all orbits are covered by non-preperiodic orbits, we see that

$$\#\{Q \in \text{Orb}_S(P) : h_{\mathcal{L}}(Q) \leq X\} \ll X^p$$

holds for all $P \in V$ by combining Theorem 1.2 part (1) with Lemma 2.13. Hence, we have Theorem 1.5 from the Introduction.

3 Orbit counts in \mathbb{P}^1

3.1 Generic bounds

Now that we have some handle on the growth of functions with various boundedness properties in a semigroup, we turn to the asymptotic growth of points in orbits of bounded height. To make this transition, we need some control on the functions that agree at a particular value. This is possible when $V = \mathbb{P}^1$ precisely because we have some control on the set of rational (or integral) points on curves. To state this refinement in dimension 1, it will be useful to have the following general definition.

Definition 3.1 Let S be a set of endomorphisms of a variety V that are polarizable with respect to a common ample invertible sheaf \mathcal{L} . Then, we say that M_S has the *cancellation property* if for each $P \in V$, there is a positive constant $B = B(P)$ such that if $f(Q_1) = g(Q_2)$ for some $f, g \in M_S$ and some $Q_1, Q_2 \in \text{Orb}_S(P)$ with $h_{\mathcal{L}}(Q_i) > B$, then we have compositional factorizations $f = \phi \circ f_0$ and $g = \phi \circ g_0$ for some $\phi \in S$ and $f_0, g_0 \in M_S \cup \{\text{id}\}$ with $f_0(Q_1) = g_0(Q_2)$.

In particular, if M_S has the cancellation property above, then an asymptotic on the number of functions yielding a bounded height relation leads to an asymptotic for the number of points of bounded height in orbits.

Proposition 3.2 Let $S = \{\phi_1, \dots, \phi_r\}$ be a polarizable set of endomorphisms of a variety V generating a free semigroup with the cancellation property, and suppose that $P \in V$ is such that $\text{Orb}_S(P)$ is infinite. Then, there exists a $B > 0$ and points $Q_1, \dots, Q_t \in \text{Orb}_S(P)$ such that $h_{\mathcal{L}}(Q_i) > B$ and

$$(3.2.1) \quad \#\{Q \in \text{Orb}_S(P) : h_{\mathcal{L}}(Q) \leq X\} \sim \sum_{i=1}^t \#\{f \in M_S : h_{\mathcal{L}}(f(Q_i)) \leq X\}.$$

In particular, we may choose non-preperiodic points $Q_1, \dots, Q_t \in \text{Orb}_S(P)$ such that (3.2.1) holds.

Proof Let B_0 be the constant from Lemma 2.11, let B_1 be the constant from Definition 3.1, and let $B = \max(B_1, B_0)$. Then, Lemma 2.13 implies that there exists a finite set F and points Q_1, \dots, Q_n satisfying $h_{\mathcal{L}}(Q_i) > B$ such that

$$(3.2.2) \quad \text{Orb}_S(P) = F \cup \bigcup_{i=1}^n \text{Orb}_S(Q_i).$$

We will show that the union over the $\text{Orb}_S(Q_i)$ may be written as a disjoint union. To see this, suppose that $f(Q_i) = g(Q_j)$ for some $f, g \in M_S$. If $|f| = |g|$, then repeated application of the cancellation property implies that $Q_i = Q_j$. However, if $|f| \neq |g|$, say without loss $|f| > |g|$, then repeated application of the cancellation property implies that $f_0(Q_i) = Q_j$ for some $f_0 \in M_S$. Hence, $\text{Orb}_S(Q_j) \subseteq \text{Orb}_S(Q_i)$ in this case. Thus, after reordering if necessary, we may choose $t \leq n$ such that

$$(3.2.3) \quad \text{Orb}_S(P) = F \cup \bigsqcup_{i=1}^t \text{Orb}_S(Q_i).$$

However, for each Q_i , a similar argument implies that the evaluation map $f \rightarrow f(Q_i)$ is injective: if $f(Q_i) = g(Q_i)$ for some $|f| = |g|$, then repeated application of the cancellation property implies that $f = g$. Likewise, if $|f| > |g|$, then repeated application of the cancellation property implies that $f_0(Q_i) = Q_i$ for some non-identity map $f_0 \in M_S$. But then Q_i is preperiodic, contradicting Lemma 2.11 and the fact that $h_{\mathcal{L}}(Q_i) > B$. Therefore, for each Q_i , the evaluation map $f \rightarrow f(Q_i)$ is injective, and thus,

$$(3.2.4) \quad \#\{Q \in \text{Orb}_S(Q_i) : h_{\mathcal{L}}(Q) \leq X\} = \#\{f \in M_S : h_{\mathcal{L}}(f(Q_i)) \leq X\}.$$

Hence, combining (3.2.3) and (3.2.4), we obtain (3.2.1) as claimed. ■

We next note that many sets of rational maps on $V = \mathbb{P}^1$ generate semigroups with the cancellation property above. To make this precise, recall that $w \in \mathbb{P}^1(\overline{\mathbb{Q}})$ is called a *critical value* of $\phi \in \overline{\mathbb{Q}}(x)$ if $\phi^{-1}(w)$ contains fewer than $\deg(\phi)$ elements. Likewise, we call a critical value w of ϕ *simple* if $\phi^{-1}(w)$ contains exactly $\deg(\phi) - 1$ points. In particular, we have the corresponding notions for sets:

Definition 3.3 Let $S = \{\phi_1, \dots, \phi_r\}$ be a set of rational maps on \mathbb{P}^1 , and let \mathcal{C}_{ϕ_i} denote the set of critical values of ϕ_i . Then, S is called *critically separate* if $\mathcal{C}_{\phi_i} \cap \mathcal{C}_{\phi_j} = \emptyset$ for all $i \neq j$. Moreover, S is called *critically simple* if every critical value of every $\phi \in S$ is simple.

In particular, critically separate and critically simple sets of maps have the cancellation property; see also Proposition 4.1 and Lemma 4.8 in [6].

Proposition 3.4 Let $S = \{\phi_1, \dots, \phi_r\}$ be a critically separate and critically simple set of rational maps on \mathbb{P}^1 all of degree at least four. The following statements hold:

- (1) M_S is a free semigroup.
- (2) For all non-preperiodic P , there is a constant $t_{S,P}$ such that

$$\#\{f \in M_S : f(P) = Q\} \leq t_{S,P}$$

holds for all $Q \in \text{Orb}_S(P)$.

- (3) The semigroup M_S has the cancellation property.

Proof The first two statements are proved in [6, §4]. Therefore, it only remains to prove (3). Let K be a number field over which P and all maps in S are defined. By the main results in [18] (see also [6, Proposition 4.6]), we have that the curves $C_k := \{(x, y) : \frac{\phi_k(x) - \phi_k(y)}{x - y} = 0\}$ for $k = 1, \dots, r$ and $C_{i,j} := \{(x, y) : \phi_i(x) = \phi_j(y)\}$ for $i \neq j$ have only finitely many K -points, and we let Σ denote the finite set of K -points of the union of these curves.

Let

$$B_0 = \max_{(P_1, P_2) \in \Sigma} \{ \max\{h_{\mathcal{L}}(P_1), h_{\mathcal{L}}(P_2)\} \},$$

let B_1 be the constant from Lemma 2.11, and let $B = \max\{B_0, B_1\}$. Now suppose that $f, g \in M_S$ and $Q_1, Q_2 \in \mathbb{P}^1$ satisfy

$$\min(h_{\mathcal{L}}(Q_1), h_{\mathcal{L}}(Q_2)) > B$$

and $f(Q_1) = g(Q_2)$. We will show that B satisfies the conditions needed in Definition 3.1. To see this, write $f = \phi_i \circ f_0$ and $g = \phi_j \circ g_0$ for $f_0, g_0 \in M_S \cup \{\text{id}\}$ and $1 \leq i, j \leq r$. There are now two cases to consider. The first is when $i \neq j$. In this case, $(f_0(Q_1), g_0(Q_2)) \in C_{i,j}$ and hence must be in Σ , and so

$$\max(h_{\mathcal{L}}(f_0(Q_1)), h_{\mathcal{L}}(g_0(Q_2))) \leq B_0 \leq B.$$

However, since $\min(h_{\mathcal{L}}(Q_1), h_{\mathcal{L}}(Q_2)) > B \geq B_1$, it follows from Lemma 2.11 that

$$\min(h_{\mathcal{L}}(f_0(Q_1)), h_{\mathcal{L}}(g_0(Q_2))) > B,$$

a contradiction.

The remaining case is when $i = j$ and $f_0(Q_1) \neq g_0(Q_2)$. In this case, we have that $(f_0(Q_1), g_0(Q_2)) \in C_i$ and hence is in Σ , and the result again follows as in the preceding case. ■

We can now improve the main result in [6].

Theorem 3.5 *Let $S = \{\phi_1, \dots, \phi_r\}$ be a critically separate and critically simple set of rational maps on \mathbb{P}^1 all of degree at least four. Moreover, assume that $r \geq 2$, and let ρ be the positive real number satisfying $\sum \deg(\phi_i)^{-\rho} = 1$. Then,*

$$\#\{Q \in \text{Orb}_S(P) : h(Q) \leq X\} \asymp X^\rho$$

for all $P \in \mathbb{P}^1(\overline{\mathbb{Q}})$ such that $\text{Orb}_S(P)$ is infinite. Moreover, if the degrees of the maps in S are acyclic, then

$$\#\{Q \in \text{Orb}_S(P) : h(Q) \leq X\} \sim cX^\rho$$

for some positive constant c , depending on P and S .

Proof Assume that P is a point in \mathbb{P}^1 that satisfies that $\text{Orb}_S(P)$ is infinite. Then, Proposition 3.4 implies that M_S is free and has the cancellation property. Hence, Proposition 3.2 implies that there exists non-preperiodic points $Q_1, \dots, Q_t \in \text{Orb}_S(P)$ such that

$$(3.5.1) \quad \#\{Q \in \text{Orb}_S(P) : h(Q) \leq X\} \sim \sum_{i=1}^t \#\{f \in M_S : h(f(Q_i)) \leq X\}.$$

From here, the result now follows directly from Theorem 1.2. For instance, if the degrees of the maps in S are acyclic, then statement (4) of Theorem 1.2 implies that for each Q_i there are positive constants c_{S,Q_i} such that

$$(3.5.2) \quad \#\{f \in M_S : h(f(Q_i)) \leq X\} \sim c_{S,Q_i} X^\rho.$$

Here, we use that the Q_i are not preperiodic. In particular, combining (3.5.1) and (3.5.2) yields the desired conclusion. Likewise, the general case follows from (3.5.1) and statement (3) of Theorem 1.2. ■

In particular, if we fix integers d_1, \dots, d_r all at least four, then the bound for orbits in Theorem 3.5 holds on a Zariski open subset of $\text{Rat}_{d_1} \times \dots \times \text{Rat}_{d_r}$.

Proof of Theorem 1.6 For each $d_i \geq 4$, the proof of [18, Theorem 1.4] on page 211 implies that there is a proper open subset $U_i \subset \text{Rat}_{d_i}$ such that each $\phi \in U_i$ is critically simple. Likewise, the proof of [18, Theorem 1.2] on pages 208–209 implies that for each pair (d_i, d_j) , there are proper open subsets $V_{ij} \subset \text{Rat}_{d_i}$ and $W_{ij} \subset \text{Rat}_{d_j}$ such that $\{\phi, \psi\}$ is a critically separate set of maps for all $\phi \in V_{ij}$ and $\psi \in W_{ij}$. Now define the open set

$$U := O_1 \times \dots \times O_r \subset \text{Rat}_{d_1} \times \dots \times \text{Rat}_{d_r}, \quad \text{where } O_i = \bigcap_{j=1}^r (U_i \cap V_{ij} \cap W_{ji}).$$

In particular, if $(\phi_1, \dots, \phi_r) \in U$, then it follows that $S = \{\phi_1, \dots, \phi_r\}$ is a critically simple and critically separate set of maps whose orbits satisfy the desired height bounds by Theorem 3.5. ■

Although Theorems 3.5 and 1.6 do not apply directly to polynomials, similar statements can be made in this case assuming the disjointness and simplicity of affine critical values and that the degrees of the polynomials in S are at least 5; see, for instance, [4, §3]. However, such conditions are stronger than those in the following result; compare to [6, Theorem 1.6].

Theorem 3.6 Let $S = \{\phi_1, \dots, \phi_r\}$ be a set of polynomials defined over a number field K with distinct and acyclic degrees all at least 2. Moreover, let ρ be the real number satisfying $\sum_{\phi \in S} \deg(\phi)^{-\rho} = 1$, and assume that the following conditions hold:

- (1) Each $\phi_i \in S$ is not of the form $R \circ E \circ L$ for some polynomial $R \in \overline{\mathbb{Q}}[x]$, some cyclic or Chebyshev polynomial E with $\deg(E) \geq 2$, and some linear polynomial $L \in \overline{\mathbb{Q}}[x]$;
- (2) For all $i \neq j$, we have that $\phi_j \neq \phi_i \circ F$ for any $F \in \overline{\mathbb{Q}}[x]$.

Then, M_S is a free semigroup, and for all points $P \in \mathbb{P}^1(K)$ such that $\text{Orb}_S(P)$ is infinite, we have that

$$\#\{Q \in \text{Orb}_S(P) : h(P) \leq X\} \sim cX^\rho$$

for some positive constant c depending on S and P .

Proof The semigroup M_S is free by [6, Theorem 1.6]. For the rest, take $\mathcal{O} \subset K$ to be a ring of \mathcal{S} -integers in K containing P and all the coefficients of polynomials in S . Then, the curves $C_\phi := \left\{ (x, y) : \frac{\phi(x) - \phi(y)}{(x-y)} = 0 \right\}$ and $C_{\phi, \psi} := \{(x, y) : \phi(x) = \psi(y)\}$

have finitely many \mathcal{O} -points by [6, Proposition 4.5] for all distinct $\phi, \psi \in S$; that is, the set

$$\Sigma := \{(x, y) \in \mathcal{O} \times \mathcal{O} : (x, y) \in C_\phi \text{ or } (x, y) \in C_{\phi, \psi} \text{ for some } \phi, \psi \in S\}$$

is finite. From here, an identical proof to that of statement (3) of Proposition 3.4 implies that M_S has the cancellation property; specifically, if

$$B_0 = \max_{(x, y) \in \Sigma} \{ \max\{h(x), h(y)\} \},$$

if B_1 is the constant from Lemma 2.11, and if $B = \max\{B_0, B_1\}$, then $B = B(P)$ satisfies the conditions of Definition 3.1. In particular, the desired asymptotics now follow directly from Proposition 3.2 and Theorem 1.2. \blacksquare

Acknowledgements We thank the anonymous referee for many helpful comments and suggestions that greatly improved this paper.

References

- [1] J. P. Bell, K. Huang, W. Peng and T. J. Tucker, *A tits alternative for rational functions*. J. Eur. Math. Soc., to appear. Preprint, [arXiv:2103.09994](https://arxiv.org/abs/2103.09994).
- [2] R. Benedetto, P. Ingram, R. Jones, M. Manes, J. H. Silverman and T. J. Tucker, *Current trends and open problems in arithmetic dynamics*. Bull. Amer. Math. Soc. (N.S.) 56(2019), no. 4, 611–685.
- [3] P. Erdős, *On some asymptotic formulas in the theory of the “factorisation numerorum.”* Ann. of Math. (2) 42(1941), 989–993.
- [4] H. K. Ha and C. C. Yang, *On the functional equation $P(f) = Q(g)$* . In Value distribution theory and related topics, Adv. Complex Anal. Appl., Vol. 3, Kluwer Acad. Publ., Boston, MA, 2004, 201–207.
- [5] V. Olsiewski Healey and W. Hindes, *Stochastic canonical heights*. J. Number Theory 201(2019), 228–256.
- [6] W. Hindes, *Counting points of bounded height in monoid orbits*. Math. Z. 301(2022), no. 4, 3395–3416.
- [7] W. Hindes and J. H. Silverman, *The size of semigroup orbits modulo primes*. Pacific J. Math. 325(2023), no. 2, 281–297.
- [8] S. Kawaguchi, *Canonical heights, invariant currents, and dynamical eigensystems of morphisms for line bundles*. J. Reine Angew. Math. 597(2006), 135–173.
- [9] S. Kawaguchi and J. H. Silverman, *On the dynamical and arithmetic degrees of rational self-maps of algebraic varieties*. J. Reine Angew. Math. 713(2016), 21–48.
- [10] A. Knopfmacher and M. E. Mays, *A survey of factorization counting functions*. Int. J. Number Theory 1(2005), no. 4, 563–581.
- [11] J. Lesieutre and M. Satriano, *Canonical heights on hyper-Kähler varieties and the Kawaguchi-Silverman conjecture*. Int. Math. Res. Not. IMRN 10(2021), 7677–7714.
- [12] J.-L. Lin, *On the arithmetic dynamics of monomial maps*. Ergodic Theory Dynam. Systems 39(2019), no. 12, 3388–3406.
- [13] Y. Matsuzawa, *Kawaguchi-Silverman conjecture for endomorphisms on several classes of varieties*. Adv. Math. 366(2020), 107086, 26.
- [14] Y. Matsuzawa and K. Sano, *Arithmetic and dynamical degrees of self-morphisms of semi-abelian varieties*. Ergodic Theory Dynam. Systems 40(2020), no. (6), 1655–1672.
- [15] Y. Matsuzawa and S. Yoshikawa, *Kawaguchi-Silverman conjecture for endomorphisms on rationally connected varieties admitting an int-amplified endomorphism*. Math. Ann. 382(2022), no. 3–4, 1681–1704.
- [16] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory. I. Classical theory*, Cambridge Studies in Advanced Mathematics, Vol. 97, Cambridge University Press, Cambridge, 2007.
- [17] M. Ram Murty, M. Rosen and J. H. Silverman, *Variations on a theme of Romanoff*. Internat. J. Math. 7 (1996), no. 3, 373–391.

- [18] F. Pakovich, *Algebraic curves $P(x) - Q(y) = 0$ and functional equations*. Complex Var. Elliptic Equ. 56 (2011), no. 1–4, 199–213.
- [19] J.-P. Serre, M. Brown, and M. Waldschmidt, *Lectures on the Mordell-Weil theorem*, Springer, 1989.
- [20] J. H. Silverman, *Rational points on $K3$ surfaces: A new canonical height*. Invent. Math. 105(1991), no. 2, 347–373.
- [21] J. H. Silverman, *The arithmetic of dynamical systems*, Graduate Texts in Mathematics, Vol. 241, Springer, New York, 2007.
- [22] J. H. Silverman, *Moduli spaces and arithmetic dynamics*, CRM Monograph Series, Vol. 30, American Mathematical Society, Providence, RI, 2012.

Department of Mathematics, University of Waterloo, Waterloo, ON, N2L 3G1, Canada
e-mail: jpbell@uwaterloo.ca

Department of Mathematics, Texas State University, 601 University Dr., San Marcos, TX 78666 USA
e-mail: wmh33@txstate.edu

Department of Mathematics, University of Waterloo, Waterloo, ON, N2L 3G1, Canada
e-mail: xiao.zhong@uwaterloo.ca