

Modulatory effect of α -linolenic acid-rich garden cress (*Lepidium sativum* L.) seed oil on inflammatory mediators in adult albino rats

Bastihalli Tukaramrao Diwakar¹, Belur Ramaswamy Lokesh² and Kamatham Akhilender Naidu^{1*}

¹Department of Biochemistry and Nutrition, Central Food Technological Research Institute, Council of Scientific and Industrial Research, Mysore 570 020, India

²Lipid Science and Traditional Foods, Central Food Technological Research Institute, Council of Scientific and Industrial Research, Mysore 570 020, India

(Received 9 September 2010 – Revised 20 January 2011 – Accepted 21 January 2011 – First published online 27 April 2011)

Abstract

Vegetable oils containing α -linolenic acid (ALA; 18:3n-3) have been shown to modulate the functions of immunocompetent cells. The aim of the present study was to understand the modulatory effect of ALA-rich garden cress (*Lepidium sativum* L.) seed oil (GCO) on lipid composition, spleen lymphocyte (SL) proliferation and inflammatory mediator production by peritoneal macrophages (PM Φ) in rats. Female Wistar rats were fed diets containing either GCO (2.5, 5.0 and 10%, w/w) or sunflower oil (SFO, 10% w/w) for 8 weeks. *Ex vivo* proliferation of SL was measured by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay. IL-2 and TNF- α in SL and PM Φ were analysed by ELISA. Inflammatory mediators such as NO, leukotriene B₄ (LTB₄) and H₂O₂ were measured in mitogen-activated PM Φ . GCO significantly increased the levels of ALA, EPA and DHA, but reduced linoleic acid and arachidonic acid in SL and PM Φ lipids. GCO (10%) significantly decreased the concanavalin A (Con-A)- and phytohaemagglutinin-mediated proliferation of SL by 54 and 38%, respectively, in comparison with SFO. A marginal decrease in IL-2 and TNF- α was observed in Con-A-stimulated SL and lipopolysaccharide-stimulated PM Φ . LTB₄ levels in Ca ionophore-stimulated PM Φ were reduced by 40% in GCO-fed rats. NO release in response to various stimuli was significantly decreased in PM Φ of GCO-fed rats. The present study is the first report on the modulatory efficacy of GCO on immunomodulators in rats. GCO modulated inflammatory mediators such as NO and LTB₄, and thus may play a role in alleviating inflammatory conditions favourably.

Key words: Garden cress seed oil: Peritoneal macrophages: Spleen lymphocytes: Nitric oxide: Leukotriene B₄: TNF- α

Fatty acids in membrane phospholipids determine the structural and physiological functions of the cell membrane⁽¹⁾. Since dietary fats can modify the fatty acid composition of membrane phospholipids, they can perturb membrane-mediated functions of immunocompetent cells⁽²⁾. Long-chain (LC) PUFA serve as precursors for inflammatory lipid mediators. Arachidonic acid (AA; 20:4), an n-6 LCPUFA, gives rise to pro-inflammatory mediators, namely leukotriene B₄ (LTB₄) and PGE₂, whereas EPA (20:5) and DHA, the n-3 LCPUFA, produce less inflammatory mediators such as leukotriene B₅ and PGE₃, resolvins and protectins^(2,3). n-3 LCPUFA have been reported to alter the lipid composition of the T-cell membrane⁽⁴⁾ and the signalling of IL-2 receptors⁽⁵⁾. n-3 LCPUFA have also been reported to decrease the expression of genes involved in eicosanoid synthesis, scavenger receptor activity, adipogenesis, NF- κ B and hypoxia signalling in human

blood mononuclear cells⁽⁶⁾. Therefore, n-3 LCPUFA may have a therapeutic value in acute and chronic inflammation and also in disorders involving impaired immune function.

Mammals can convert α -linolenic acid (ALA; 18:3n-3) to EPA and DHA; however, there are concerns regarding the efficiency of its conversion to n-3 LCPUFA^(7,8). It has been reported that a decrease in linoleic acid (LA; 18:2n-6) and an increase in ALA content in the diet can alter the levels of AA and EPA in the tissues of hamsters⁽⁹⁾ and rats^(10,11). An ALA-rich diet has been reported to increase the levels of ALA and EPA in the phospholipid fraction of erythrocytes⁽¹²⁾, suppress the *ex vivo* proliferation of lymphocytes⁽¹³⁾ and inhibit TNF- α and IL-1 β production in human mononuclear cells⁽¹⁴⁾. ALA has been reported to modulate the T-helper 1/T-helper 2 balance when ingested at normal dietary concentrations⁽¹⁵⁾. These studies have demonstrated that the efficacy

Abbreviations: 5-LO, 5-lipoxygenase; AA, arachidonic acid; ALA, α -linolenic acid; Con-A, concanavalin A; GCO, garden cress seed oil; LA, linoleic acid; LC, long chain; LPS, lipopolysaccharide; LTB₄, leukotriene B₄; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PHA, phytohaemagglutinin; PMA, phorbol 12-myristate-13-acetate; PM Φ , peritoneal macrophage; RPMI, Roswell Park Memorial Institute; SFO, sunflower oil; SL, spleen lymphocyte.

*Corresponding author: Dr K. A. Naidu, fax +91 821 2517233, email akhilnaidu@gmail.com

of ALA to modulate the functions of immunocompetent cells may be of importance in altering the pathophysiological effects of inflammatory diseases.

Garden cress seed oil (GCO) is a good source of PUFA. It contains 32% of ALA and 12% of LA. GCO has a fairly balanced ratio of SFA:MUFA:PUFA (1:2.6:3) compared with flaxseed and perilla seed oil. GCO is a relatively stable oil due to the presence of a high concentration of natural antioxidants such as tocopherols (1330 mg/kg) and carotenoids (5.32 mg/kg)⁽¹⁶⁾. The major part of ALA in GCO is esterified at the sn-2 carbon of TAG⁽¹⁶⁾, hence it can be easily absorbed by the intestine. We have reported a significant increase in EPA and DHA levels in the serum, liver, heart and brain with a concomitant decrease in LA and AA in rats fed with a GCO diet⁽¹⁰⁾. The aim of the present study was to assess the modulatory effect of GCO on some key functions of immunocompetent cells such as proliferation and release of inflammatory mediators *ex vivo*.

Materials and methods

Materials

Garden cress seeds were purchased from a local market in Mysore. The seeds were identified and authenticated at the Department of Horticultural Sciences, University of Agriculture Sciences, Bangalore. Seeds were air-dried and flaked in a roller flaker (Kvarnmaskiner, Malmö, Sweden). Flaked garden cress seeds were pressed in a hydraulic press (B Sen Barry and Company, New Delhi, India), at a pressure of 98.1 kPa for 10 min, to expel the oil. Oil was collected, flushed with N₂ and stored at -20°C⁽¹⁶⁾.

All chemicals used were of analytical grade. Fatty acid methyl ester standards, B trifluoride, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), dimethyl sulphoxide, Ca ionophore (A23187), concanavalin A (Con-A), phorbol 12-myristate-13-acetate (PMA), phytohaemagglutinin (PHA) and lipopolysaccharide (LPS) were procured from Sigma Chemical Company (St Louis, MO, USA). Roswell Park Memorial Institute (RPMI)-1640 medium and antibiotics were procured from Thermo Scientific Company (Rockford, IL, USA). Rat IL-2 ELISA kit was purchased from R&D Systems, Inc. (Minneapolis, MN, USA). TNF- α was purchased from Koma Biotech (Seoul, South Korea).

Animals and diets

Weaned female Wistar rats (OUTB-Wistar, IND-cft (2c)) weighing 55–62 g, bred in the animal house facility at the Central Food Technological Research Institute, Mysore, India, were used in the present study. The experimental protocol adopted in the present study was approved by the Institute's animal ethical committee. Animals were housed individually in stainless-steel cages, in a room, where the temperature was maintained at 25 \pm 2°C. Animals were divided into four groups, with twelve animals in each group. They were fed with a semi-synthetic diet containing sunflower oil (SFO)10 (SFO 10 g/100 g), GCO2.5 (GCO2.5 g + SFO 7.5 g/100 g),

GCO5 (GCO 5 g + SFO 5 g/100 g) and GCO10 (GCO 10 g/100 g), respectively. The composition of the isoenergetic, semi-synthetic diet and its fatty acid profile are presented in Tables 1 and 2, respectively. Diets were prepared every week and stored at 4°C. Animals were given a fresh diet *ad libitum* for 8 weeks. Animals had free access to their respective diets and water at all times throughout the study. After 8 weeks of feeding, rats were fasted overnight and were euthanised under diethyl ether anaesthesia. Blood was drawn by cardiac puncture, and serum was separated by centrifugation at 600 g for 10 min (after allowing the blood to clot at room temperature for 2 h). Serum from animals of each group was pooled, heat inactivated at 54°C for 30 min, filtered through a 0.2 μ m filter and used at a 2.5% level in culture media for primary cell culture experiments.

Spleen lymphocyte preparation

Spleens were aseptically removed, dilacerated and passed through a sterile cell strainer to form a cell suspension. The cell suspension was centrifuged for 10 min at 150 g, and the resulting pellet was suspended in RPMI-1640 medium. Spleen lymphocyte (SL) were collected by centrifugation at 400 g for 30 min on Ficol Histopaque gradient, washed and re-suspended in complete media (RPMI-1640 containing 25 mM-HEPES supplemented with 2.5% autologous serum, penicillin (60 mg/l), streptomycin (100 mg/l), 2 mM-L-glutamine and 10 μ M-2-mercaptoethanol). The number of viable cells was counted by the trypan blue exclusion method using a haemocytometer, and cell number was adjusted to 1.5 \times 10⁶ cells/ml.

Proliferation assay of rat spleen lymphocytes

Proliferation of SL was evaluated by the MTT assay⁽¹⁷⁾. SL were cultured at a cell density of 1.5 \times 10⁵ cells/well in complete medium (RPMI-1640 containing 25 mM-HEPES supplemented with 2.5% autologous serum, penicillin (60 mg/l), streptomycin

Table 1. Composition of the diets

Component (g/kg)	Diet			
	SFO10	GCO2.5	GCO5	GCO10
Casein	200.0	200.0	200.0	200.0
Maize starch	500.0	500.0	500.0	500.0
Sucrose	100.0	100.0	100.0	100.0
Cellulose	50.0	50.0	50.0	50.0
Mineral mix*	35.0	35.0	35.0	35.0
Methionine	2.0	2.0	2.0	2.0
Choline chloride	3.0	3.0	3.0	3.0
Vitamin mix†	10.0	10.0	10.0	10.0
SFO	100.0	75.0	50.0	0.0
GCO	0.0	25.0	50.0	100.0

SFO10, sunflower oil (SFO) 10 g/100 g; GCO2.5, garden cress seed oil (GCO) 2.5 g + SFO 7.5 g/100 g; GCO5, GCO 5 g + SFO 5 g/100 g; GCO10, GCO 10 g/100 g.

* Bernhart Tommarelli salt mixture.

† AIN-76A containing per kg: 1.8 g vitamin A, 0.125 g vitamin D₂, 22 g DL- α -tocopherol, 45 g ascorbic acid, 5 g inositol, 2.25 g menadione, 5 g *p*-aminobenzoic acid, 4.25 g niacin, 1 g riboflavin, 1 g pyridoxine hydrochloride, 1 g thiamine hydrochloride, 3 g calcium pantothenate, 0.02 g biotin, 0.09 g folic acid, 0.00135 g vitamin B₁₂ and 833.46 g sucrose carrier.

Table 2. Fatty acid composition of the experimental diets*
(Mean values with their standard errors, *n* 4)

Fatty acids (% total fatty acids)	SFO10		GCO2.5		GCO5		GCO10	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
SFA								
16:0	5.7 ^a	0.08	6.6 ^b	0.12	8.9 ^c	0.1	11.1 ^d	0.09
18:0	2.3 ^{b,c}	0.07	2.6 ^c	0.10	2.2 ^b	0.13	1.6 ^a	0.13
20:0	0.00	0.00	0.62 ^a	0.04	1.6 ^b	0.05	2.5 ^c	0.10
MUFA								
18:1 <i>n</i> -9	38.4 ^d	1.1	36.2 ^c	0.57	31.5 ^b	0.49	22.1 ^a	0.36
20:1	0.00	0.00	3.0 ^a	0.32	6.0 ^b	0.06	11.1 ^c	0.31
22:1	0.00	0.00	1.4 ^a	0.12	2.2 ^b	0.04	4.3 ^c	0.05
PUFA								
18:2 <i>n</i> -6	53.6 ^d	0.33	40.0 ^c	0.63	32.2 ^b	0.95	9.9 ^a	0.21
18:3 <i>n</i> -3	0.45 ^a	0.09	8.8 ^b	0.22	16.7 ^c	0.41	32.2 ^d	0.63
<i>n</i> -6: <i>n</i> -3 ratio	105.9 ^b	6.7	4.6 ^a	0.15	1.9 ^a	0.09	0.31 ^a	0.08

SFO10, sunflower oil (SFO) 10 g/100 g; GCO2.5, garden cress seed oil (GCO) 2.5 g + SFO 7.5 g/100 g; GCO5, GCO 5 g + SFO 5 g/100 g; GCO10, GCO 10 g/100 g.

^{a,b,c,d}Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$; ANOVA, $a < b < c < d$).

* For details of the diets, see Table 1.

(100 mg/l), 2 mM-L-glutamine and 10 μ M-2-mercaptoethanol) in ninety-six-well flat-bottomed culture plates (Nunc, Roskilde, Denmark) in the presence or absence of 5 μ g/ml of T-cell mitogens (Con-A and PHA) and a B-cell mitogen (LPS) with triplicate wells for each sample. Cells were incubated in a CO₂ incubator at 37°C for 72 h. At the 69th hour, 20 μ l of MTT (5 mg/ml) prepared in PBS (pH 7.4) were added and incubated for 3 h. The plates were spun at 500 *g*, the culture medium was discarded, the solubilising reagent dimethyl sulfoxide (100 μ l) was added and then the absorbance was measured at 550 nm.

IL-2 measurement

SL were cultured at a cell density of 2.5×10^5 cells/well in complete medium, as described earlier, and stimulated with 5 μ g of Con-A or PHA for 48 h. Culture medium was aspirated and stored at -80°C . Subsequently, the culture medium was thawed, vortexed and centrifuged (4°C) at 500 *g* for 5 min. An aliquot (50 μ l) of the culture supernatant was analysed quantitatively for IL-2 levels using a commercially available ELISA kit (Quantakine, IL-2; R&D Systems, Inc.). Results are expressed as pg of IL-2/ 2.5×10^5 cells, and data are presented as means with their standard errors (*n* 4).

Isolation and activation of peritoneal macrophages

The GCO and SFO diets were fed to rats for 8 weeks. After 8 weeks, rats were injected intraperitoneally with 1% sodium caseinate in PBS to elicit the migration of macrophages to the peritoneal cavity. After 4 d, rats were anaesthetised with diethyl ether, and peritoneal exudate cells were collected aseptically by irrigating the peritoneal cavity with chilled RPMI-1640 medium. Peritoneal exudates were centrifuged, washed and resuspended in complete RPMI-1640 medium. Cells were seeded at a density of 1×10^5 cells/well in ninety-six-well culture plates (Nunc) and incubated for 2 h at 37°C in a CO₂ incubator. Non-adherent cells were removed

by washing (three times) with serum-free medium, and adherent cells were cultured overnight in complete medium. More than 95% of the adherent cell population was found to be peritoneal macrophages (PM Φ) as determined by morphology and non-specific esterase staining.

Stimulation of peritoneal macrophages for nitric oxide production

PM Φ (1×10^5 cells/well) were cultured for 12 h at 37°C in a CO₂ incubator, and treated with either Con-A (10 μ g/ml), PHA (1 μ g/ml), LPS (10 μ g/ml), A23187 (1 μ g/ml) or PMA (500 ng/ml). Samples were drawn after 6, 12 and 24 h incubation and centrifuged, and the supernatant (100 μ l) was transferred to ninety-six-well assay plates. Nitrite concentration in the cell-free culture supernatant was measured by a spectrophotometric assay based on the Griess reaction as described elsewhere⁽¹⁸⁾. Briefly, 100 μ l of the culture supernatant were mixed with an equal volume of Griess reagent (one part of 0.1% (w/v) naphthylethylenediamine dihydrochloride in distilled water and one part of 1% (w/v) sulphanilamide in 5% (v/v) H₃PO₄) at room temperature. After 15 min, the absorbance was measured using an ELISA plate reader at 540 nm. Nitrite content (μ mol/ 10^5 cells) was quantified using a sodium nitrite standard curve. Determinations were carried out in triplicate (*n* 6).

Measurement of TNF- α

PM Φ (1×10^5 cells/well) in RPMI-1640 medium were treated with LPS (10 μ g/ml) in ninety-six-well culture plates (Nunc). After 10 h incubation at 37°C in a CO₂ incubator, the cell-free culture supernatant was removed and stored at -80°C . An aliquot (50 μ l) of the culture supernatant was analysed quantitatively for TNF- α using a commercially available ELISA kit (Koma Biotec). Results are expressed as pg of TNF- α / 1×10^5 cells. Data are expressed as means with their standard errors (*n* 4).

Measurement of hydrogen peroxide release

H_2O_2 released in the culture medium by PM Φ was determined according to the method of Yaqoob & Calder⁽¹⁸⁾. PM Φ (1×10^5 cells/well) were incubated in 100 μ l Hanks balanced salt solution containing 0.1% (w/v) dextrose, 0.1% phenol red and horseradish peroxidase (40 μ g/ml). PMA (10 U/ml) was added and incubated for 2 h at 37°C in a CO₂ incubator. The reaction was stopped by the addition of 10 μ l of 0.1 M-NaOH, and the absorbance was measured at 610 nm. Results are expressed as nm of $H_2O_2/1 \times 10^5$ PM Φ . H_2O_2 concentration was determined by the extinction coefficient $E = 43.6$ cm/m.

In vitro lysosomal phosphatase enzyme activity

Cellular lysosomal phosphatase enzyme activity was determined according to the method described by Manosroi *et al.*⁽¹⁹⁾. Briefly, PM Φ (1×10^5 cells/well) were solubilised with 20 μ l of 0.1% Triton X-100, 100 μ l of 10 mM-*p*-nitrophenyl phosphate solution and 50 μ l of 0.1 M-citrate buffer (pH 5.0), and incubated for 30 min at 37°C. The reaction was terminated by adding 150 μ l of 0.2 M-borate buffer (pH 9.8), and the absorbance was measured at 405 nm. Results are expressed as a percentage of lysosomal phosphatase activity.

$$\text{Lysosomal enzyme activity} = \frac{\text{OD}_{\text{sample}} - \text{OD}_{\text{control}}}{\text{OD}_{\text{control}}} \times 100,$$

where OD is optical density.

Measurement of leukotriene B₄

LTB₄ secretion by PM Φ in response to LPS stimulation was measured by the HPLC method⁽²⁰⁾. PM Φ (3×10^6) were incubated with or without Ca ionophore (A23187) at a concentration of 5 μ g/ml in a twenty-four-well culture plate (Nunc) for 90 min at 37°C in a CO₂ incubator. The reaction was stopped by adding 30 μ l of 1 M-HCl on ice. The supernatants were centrifuged at 2000 g for 20 min at 4°C and loaded onto Sep-Pack C₁₈ cartridges (Waters Millipore Corporation, Milliford, MA, USA). Leukotrienes were eluted with ethyl acetate. The ethyl acetate fraction was evaporated to dryness, resuspended in methanol and stored at -80°C. Identification and quantification of LTB₄ was carried out by the HPLC method. A Shimadzu LC-10A HPLC system with a UV detector and a Hypersil reverse-phase C18 column (150 \times 4.6 mm; 5 μ m) was used. An isocratic elution program was employed using a mobile phase containing methanol-ammonium acetate (70:30, v/v). The flow rate was 0.5 ml/min, and the chromatogram was monitored at 280 nm. LTB₄ content was quantified based on the calibration curve of standard LTB₄ (Sigma Chemicals Company). Results are reported as ng of LTB₄/3 $\times 10^6$ cells. Determinations were done in duplicate (n 6).

Measurement of peritoneal macrophage 5-lipoxygenase activity

PM Φ were suspended in PBS and sonicated for 20–30 s at 20 kHz to release the cytosolic 5-lipoxygenase (5-LO)

enzyme. This solution was centrifuged at 2000 g for 30 min at 4°C. The supernatant was used as the source of the enzyme. Protein was estimated by Lowry's method⁽²¹⁾ using bovine serum albumin as standard. Assay of 5-LO was performed according to the method of Raghavenra *et al.*⁽²⁰⁾. The standard reaction mixture for the 5-LO assay contained 100 mM-phosphate buffer (pH 7.4), 50 mM-dithiothreitol, 200 mM-ATP, 300 mM-CaCl₂, 150 mM-AA and 5.0 mg protein. The enzymatic reaction was carried out at 28 \pm 2°C. Enzyme activity was measured as 5-hydroperoxyeicosatetraenoic acid (5-HETE) formed at 234 nm using a Shimadzu spectrophotometer. The molar extinction coefficient of 25/mm cm was used to calculate the specific activity of 5-LO. Enzyme activity was expressed as μ M-5-HETE/min per mg protein. Determinations were done in duplicate (n 6).

Lipid extraction and fatty acid analysis

Lipids were extracted from SL and PM Φ with a chloroform-methanol mixture (2:1, v/v). Fatty acid analysis was carried out according to the method of Morisson & Smith⁽²²⁾. Extracted lipid samples were saponified for 1 h with 1 ml of 0.7 M-methanolic KOH at 60°C, followed by neutralisation with 1 ml of methanolic HCl (0.7 M). The resulting NEFA were extracted with hexane and evaporated to dryness. The fatty acids were methylated using boron trifluoride (14% in methanol) and 0.2 ml benzene. The fatty acid methyl esters were extracted with hexane, washed with water and evaporated to dryness. Fatty acid analysis was performed using a gas-liquid chromatograph (Shimadzu, GC-14B; Shimadzu Corporation, Kyoto, Japan) fitted with a fused silica capillary column (BP 21, 30 m in length, 0.30 mm in inner diameter). The gas-liquid chromatograph was equipped with a flame ionisation detector and a Clarity Lite 420 integrator. The column temperature was set at 220°C, the injector temperature at 230°C and the detector temperature at 240°C. N₂ was used as the carrier gas, and the flow rate was 1 ml/min. Individual fatty acids in the sample were identified by comparison with retention times of standard fatty acid methyl esters. Fatty acid determinations were carried out in duplicate (n 6).

Statistical analysis

Results were analysed using SPSS-10 software (SPSS, Inc., Chicago, IL, USA). The significance of mean differences between the groups was determined by one-way ANOVA, followed by Duncan's multiple-range test as a *post hoc* test for multiple comparisons whenever a significant *F* value emerged. Significance was set at $P < 0.05$.

Results

Fatty acid composition of lymphocytes and peritoneal macrophages

The fatty acid composition of lipids extracted from diets is presented in Table 2. The *n*-6:*n*-3 ratio was greatly affected by the addition of GCO in diets. The *n*-6:*n*-3 ratio was 114.5 in the

Table 3. Fatty acid composition of spleen lymphocyte (SL) and peritoneal macrophage (PM Φ) lipids isolated from rats fed with the experimental diets* (Mean values with their standard errors, *n* 6)

Fatty acids (% total fatty acids)	SFO10				GCO2.5				GCO5				GCO10				
	SL		PM Φ		SL		PM Φ		SL		PM Φ		SL		PM Φ		
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	
SFA																	
16:0	27.4	2.2	31.2	1.6	26.4	1.3	30.0	1.4	25.7	0.93	30.6	1.6	28.3	0.82	31.1	1.0	
18:0	15.7	1.2	20.0	1.5	13.1	0.48	19.0	1.6	14.0	1.2	19.5	2.0	13.1	1.4	18.7	1.3	
20:0	0.0 ^a	0.0	0.0	0.0	0.40 ^a	0.06	0.0	0.0	0.18 ^{a,b}	0.12	0.0	0.0	0.83 ^b	0.30	0.0	0.0	
Total SFA	43.1	2.4	51.2	2.6	39.9	1.3	49.0	2.3	39.8	1.8	50.1	3.1	42.3	1.4	49.8	2.0	
MUFA																	
18:1 <i>n</i> -9	14.0 ^a	1.6	16.1	1.0	16.3 ^a	1.5	18.0	1.4	14.4 ^a	1.1	16.3	1.6	20.6 ^b	0.85	19.6	0.91	
20:1 <i>n</i> -9	0.0 ^a	0.0	0.27 ^a	0.09	1.7 ^b	0.14	0.57 ^a	0.18	2.1 ^b	0.31	1.2 ^b	0.22	2.4 ^b	0.36	1.4 ^b	0.29	
Total MUFA	14.0 ^a	1.3	16.4 ^a	1.1	18.1 ^b	1.4	18.6 ^{a,b}	1.5	16.5 ^{a,b}	1.3	17.5 ^{a,b}	1.7	23.6 ^c	1.0	21.5 ^b	1.0	
PUFA																	
18:2 <i>n</i> -6	15.3 ^b	1.8	9.7 ^b	0.32	14.7 ^b	0.96	9.9 ^b	1.3	13.7 ^b	0.73	8.2 ^{a,b}	1.1	9.4 ^a	0.91	7.0 ^a	0.26	
20:4 <i>n</i> -6	17.1 ^c	0.93	16.8 ^b	1.6	14.3 ^{b,c}	1.0	15.1 ^{a,b}	2.4	13.1 ^{a,b}	0.58	13.5 ^{a,b}	1.5	11.1 ^a	1.1	10.2 ^a	1.9	
Total <i>n</i> -6 PUFA	32.4 ^c	1.0	26.6 ^b	1.5	29.0 ^b	0.8	23.3 ^{a,b}	3.1	26.8 ^b	0.9	21.6 ^{a,b}	1.6	20.5 ^a	1.4	17.2 ^a	1.8	
18:3 <i>n</i> -3	0.0 ^a	0.0	0.0 ^a	0.0	0.43 ^{a,b}	0.05	0.30 ^{a,b}	0.09	1.0 ^b	0.12	0.60 ^b	0.15	3.2 ^c	0.49	0.97 ^c	0.2	
20:5 <i>n</i> -3	0.0 ^a	0.0	0.0 ^a	0.0	0.56 ^b	0.05	0.51 ^{a,b}	0.14	0.56 ^b	0.02	0.77 ^{b,c}	0.27	1.7 ^c	0.14	1.1 ^c	0.2	
22:6 <i>n</i> -3	0.0 ^a	0.0	0.0 ^a	0.0	0.68 ^b	0.08	0.0 ^a	0.0	0.97 ^b	0.09	0.60 ^b	0.07	1.7 ^c	0.24	1.6 ^c	0.17	
Total <i>n</i> -3 PUFA	0.0 ^a	0.0	0.00 ^a	0.00	1.7 ^b	0.10	0.80 ^a	0.2	2.5 ^b	0.15	2.0 ^b	0.3	6.6 ^c	0.55	3.6 ^c	0.45	
Total PUFA	32.4	1.0	26.6	1.5	30.6	0.84	24.1	3.2	29.3	0.94	23.6	1.8	27.1	1.2	20.8	1.9	
<i>n</i> -6: <i>n</i> -3 ratio	–	–	–	–	17.6 ^c	0.90	33.0 ^b	4.2	10.9 ^b	0.6	12.7 ^a	2.4	3.2 ^a	0.4	5.1 ^a	0.86	

SFO10, sunflower oil (SFO) 10 g/100 g; GCO2.5, garden cress seed oil (GCO) 2.5 g + SFO 7.5 g/100 g; GCO5, GCO 5 g + SFO 5 g/100 g; GCO10, GCO 10 g/100 g.

^{a,b,c} Mean values within a row with unlike superscript letters were significantly different: (*P* < 0.05; ANOVA, *a* < *b* < *c*).

* For details of the diets see Table 1 and for procedures see Materials and methods.

SFO10 diet. The ratio decreased to 4.6, 1.9 and 0.3, respectively, in the GCO2.5, GCO5 and GCO10 diets.

The fatty acid composition of lipids extracted from SL and PM Φ is presented in Table 3. SFA (16:0 + 18:0) content in SL and PM Φ did not change among the dietary groups. However, an increase of 51.4 and 27.9% was observed in MUFA levels in both SL and PM Φ , respectively, in the GCO10 group compared with the SFO10 group. ALA, EPA and DHA contents increased by 3.2, 1.72, 1.7% in SL lipids and 0.97, 1.08, 1.6% in PM Φ lipids, respectively, in the GCO groups compared with the SFO10 group. LA and AA content was decreased by 28.05 and 39.5% in PM Φ , and 38.35 and 35.37% in SL, respectively, in the GCO10 group compared with the SFO10 group (Table 3).

Lymphocyte proliferation and IL-2 production

The effect of the experimental diets on SL proliferation is summarised in Fig. 1. SL proliferation induced by Con-A and PHA was significantly inhibited in the GCO groups. GCO at 2.5% caused a 40 and 36% reduction in the proliferation of SL induced by Con-A and PHA, respectively. An insignificant decrease was observed in the proliferation of SL induced by LPS in the GCO10 group (1.06 (SEM 0.09)) compared with the SFO10 group (1.95 (SEM 0.42)). A marginal decrease in IL-2 levels was observed in SL stimulated with Con-A in the GCO10 group (21.32 (SEM 1.9) pg/ 2.5×10^5 cells) compared with the SFO10 group (22.5 (SEM 1.0) pg/ 2.5×10^5 cells).

Nitric oxide release by peritoneal macrophages

A stable end product, nitrite, was determined to measure NO release in PM Φ . Nitrite content at different time intervals (6, 12 and 24 h) after post-stimulation of PM Φ with Con-A, LPS,

A23187, PHA and PMA is presented in Fig. 2. In the GCO groups, a significant decrease in nitrite production was observed at 12 h post-treatment with Con-A, PHA and A23187 (Fig. 2(a)–(c), respectively) compared with the SFO10 group. However, a significant decrease in nitrite content was observed at 6, 12 and 24 h in PHA-treated PM Φ , in all GCO groups. A non-significant downward trend was observed in nitrite production in LPS- and PMA-stimulated PM Φ (Fig. 2(d) and (e), respectively) in GCO-fed rats.

TNF- α and leukotriene B₄ release by peritoneal macrophages

Concentration of TNF- α released by PM Φ after 10 h post-stimulation of LPS did not differ much between the GCO10 (125 (SEM 6.9) pg/ 1×10^5 cells) and SFO10 (145 (SEM 3.0) pg/ 1×10^5 cells) groups. Concentration of LTB₄ released by PM Φ in response to A23187 stimulation was measured using reverse-phase HPLC and is depicted in Fig. 3. LTB₄ content was reduced significantly in the GCO5 and GCO10 groups compared with the SFO10 group.

Leukotriene B₄ release by peritoneal macrophages

Concentration of TNF- α released by PM Φ after 10 h of post-stimulation of LPS did not differ much between the GCO and SFO groups. Concentration of LTB₄ released by PM Φ in response to A23187 stimulation was measured using reverse-phase HPLC and is depicted in Fig. 3. LTB₄ content was reduced significantly in the GCO5 (17.0 (SEM 0.9) ng/ 3×10^6 cells) and GCO10 groups (12.0 (SEM 1.0) ng/ 3×10^6 cells) compared with the SFO10 group (20.0 (SEM 1.2) ng/ 3×10^6 cells).




Fig. 1. Effect of dietary lipids on the proliferation of spleen lymphocytes stimulated with concanavalin-A (Con-A, 5 μ g/ml) or phytohaemagglutinin (PHA, 5 μ g/ml). Rats were fed with the sunflower oil (SFO)- and garden cress seed oil (GCO)-supplemented diets for 8 weeks as described in Materials and methods. Spleen lymphocytes (1.5×10^5 cells/well) were isolated and cultured for 72 h in Roswell Park Memorial Institute-1640 complete medium and 2.5% autologous serum in the presence or absence of mitogens. The amount of formazan formed by the cells was measured spectrophotometrically at 550 nm. Values are means, with their standard errors represented by vertical bars (n 6). * Mean values were significantly different compared with the SFO 10 g/100 g group ($P < 0.05$). A, absorbance.




Fig. 2. Effect of dietary lipids on nitric oxide produced by rat peritoneal macrophages (PMΦ). PMΦ from rats fed with the experimental diets for 8 weeks were stimulated *ex vivo* with (a) concanavalin-A (10 µg/ml), (b) phytohaemagglutinin (1 µg/ml), (c) calcium ionophore (1 µg/ml), (d) lipopolysaccharide (10 µg/ml) and (100 mg/l) (e) phorbol 12-myristate 13-acetate (0.5 µg/ml). Culture supernatants were removed after 6, 12 and 24 h post-treatment of mitogens. Concentration of nitrite in the medium was measured by the Griess method. Values are means, with their standard errors represented by vertical bars (*n* 6). *Mean values were significantly different compared with the 10 g/100 g sunflower oil group (SFO10); *P* < 0.05. □, SFO10; ■, garden cress seed oil (GCO)2.5; ▒, GCO5; ■, GCO10.

Hydrogen peroxide release by stimulated peritoneal macrophages

No significant difference was observed in H₂O₂ levels released by PMΦ stimulated with PMA in the GCO groups.

Acid phosphatase and 5-lipoxygenase activity in peritoneal macrophages

Acid phosphatase and 5-LO activities in PMΦ were unaltered among the different GCO groups.

Discussion

PMΦ and SL are key cells of the immune system involved in host defence, inflammation and immune response. Since the action of these cells is mediated by membrane-bound components, an alteration in the membrane lipid composition could modulate immune function. It is known that dietary lipids can affect the fatty acid composition of immunocompetent cells. In the present study, we demonstrated that partial or complete replacement of LA (18:2*n*-6) with ALA (18:3*n*-3) in diets through GCO can alter the fatty acid composition and modulate inflammatory mediators produced by PMΦ and SL.

Immunocompetent cells (PMΦ and SL) originate in the bone marrow. Lymphocytes mature in primary lymphoid tissues and stay in peripheral lymphoid organs to keep a vigil on invading microbes. Rapidly dividing promonocytes are progenitor cells of blood monocytes, which give rise to PMΦ. When stimulated, blood monocytes divide rapidly and move to the peritoneal cavity as PMΦ⁽²³⁾. The type and amount of fatty acids in

the blood fatty acid pool during blastogenesis determine the fatty acid profile of PMΦ. Earlier studies have shown an increase in the ALA content of PMΦ and SL in rats fed with flaxseed and perilla oil. These oils contain 55–56% of ALA, 13 and 18% of LA and a *n*-6:*n*-3 ratio of 0.24 and 0.32, respectively^(11,14,24). Lymphocytes are cells that specifically recognise and respond to foreign antigens. In response to stimuli, T-lymphocytes secrete cytokines and undergo cell division⁽²⁵⁾. Proliferation of lymphocytes leads to an increase in the number of antigen-specific lymphocytes and the activation of B-lymphocytes, natural killer cells and macrophages. Therefore, proliferation is a key process in the regulation, amplification and memory of the cell-mediated immune response⁽²⁵⁾. In the present study, a significant decrease was observed in the *ex vivo* proliferation of lymphocytes in response to Con-A or PHA stimuli in the GCO groups. Substitution of SFO with 50% GCO in the diet significantly decreased the proliferation of SL. A further increase in GCO in the diet had no additive effect on the proliferation of lymphocytes. A similar effect has been observed when a small amount of SFO was replaced by linseed oil (*n*-6:*n*-3 PUFA ratio of 14:75) in the diet of rats⁽¹¹⁾. An ALA-rich diet has been reported to decrease the *ex vivo* proliferation of blood lymphocytes in response to mitogenic stimuli in poultry, rats and human subjects^(13,26–29). The exact mechanisms by which *n*-3 PUFA modulate lymphocyte proliferation are far from clear, but certain mechanisms have been suggested. A decreased expression in IL-2 in activated lymphocytes by *n*-3 fatty acids has been suggested for the inhibition of lymphocyte proliferation. However, in the present study, we observed no significant change in the IL-2




Fig. 3. Effect of dietary lipids on leukotriene B₄ production by peritoneal macrophages (PMΦ, 3×10^6 cells/ml) stimulated with calcium ionophore (A23187, 5 μ g/ml). Rats were fed with the sunflower oil (SFO)- and garden cress seed oil (GCO)-supplemented diets as described in Materials and methods. PMΦ were isolated and incubated for 90 min in Roswell Park Memorial Institute-1640 complete medium with 2.5% autologous serum in the presence or absence of A23187 (5 μ g/ml). Leukotriene B₄ in the culture supernatant was analysed by reverse-phase HPLC as described in Materials and methods. Values are means, with their standard errors represented by vertical bars (n 6). * Mean values were significantly different compared with the SFO group (10 g/100 g) ($P < 0.05$).

of lymphocytes in GCO-fed rats. Earlier studies with fish oil (EPA and DHA) have reported a decrease in IL-2 levels^(30,31) and or no effect on IL-2 production in lymphocytes of rats and human subjects^(29,32–35). However, a decreased expression in IL-2 receptors has been suggested to play a role in the suppression of lymphocyte proliferation⁽³⁶⁾. LCPUFA in fish oil have been reported to differentially modulate T-cell rafts and soluble membrane phospholipids, causing a displacement of acylated proteins from rafts^(4,37). When T-cells are activated, protein kinase C θ , phospholipase C- γ , linker for the activation of T-cells are translocated to lipid rafts, which induce the proliferation of lymphocytes^(38,39). The conditions that modify the integrity of T-cell rafts can disrupt the early steps of T-cell activation⁽⁴⁰⁾. Therefore, the incorporation of n -3 PUFA into the membrane microdomain may influence signal transduction and modulate T-cell activation⁽⁴⁾.

PMΦ constitute a major group of phagocytic leucocytes, which are crucial to immune surveillance against invading pathogens. Activated PMΦ release an array of mediators including pro-inflammatory cytokines (TNF- α , IL-1 and IL-6), growth factors, bioactive eicosanoids (LTB₄, PGE₂ and thromboxane A₂), hydrolytic enzymes (peroxidase and protease), reactive oxygen intermediates and NO. Each of the above secretory molecules is implicated in the pathogenesis of chronic inflammatory conditions and considered as pharmacological targets of inflammatory diseases. In the present study, we assessed the effect of dietary GCO on inflammatory mediators released by PMΦ. NO is considered as one of the principal effector molecules in PMΦ-mediated cytotoxicity⁽⁴¹⁾. The amount of NO released by PMΦ isolated from the GCO group of rats was significantly reduced in comparison with NO released by PMΦ of the SFO group. The percentage of reduction was found to be 57, 62 and 56%, respectively, after 6, 12 and 24 h post-stimulation of PHA. NO release was also significantly reduced by 37 and 44%, respectively, at

12 h post-stimulation with Con-A and A23187 in the GCO10 group compared with the SFO10 group. Previous studies have shown decreased^(42,43), increased or no change in the production of NO by the PMΦ of animals fed with fish oil^(18,30,44,45) and linseed oil⁽⁴⁶⁾. Dietary LCPUFA have been reported to decrease the expression of inducible NO synthase in cancer cells and LPS-stimulated RAW 264.7 macrophages^(47,48). Ren & Chung⁽⁴⁹⁾ reported that ALA markedly inhibits LPS-induced NO production and inducible NO synthase gene expressions by blocking NF- κ B activation and also the phosphorylation of mitogen-activated protein kinases in RAW 264.7 macrophages. In the present study, macrophages were cultured with 2.5% of autologous serum from the respective SFO- and GCO-fed rats. In a previous study, serum from GCO2.5-, GCO5- and GCO10-fed rats contained 2.2, 4.5 and 8.3% of ALA, 0.8, 1.2 and 5.4% of EPA and 1.72, 2.9 and 2.7% of DHA, respectively⁽¹⁰⁾. Thus, the exogenous pool of n -3 PUFA added to PMΦ might play an important role in the reduction of NO observed in the present study. Furthermore, it has been reported that γ -tocopherol can also inhibit inducible NO synthase in activated macrophages⁽⁵⁰⁾. Since GCO contains high amount of γ -tocopherol (87.74 mg/100 g)⁽¹⁶⁾, we hypothesise that the decrease in NO production in *ex vivo*-stimulated PMΦ can be attributed to the additive effect of ALA and γ -tocopherol present in GCO. However, further studies are needed to elucidate the mechanism of modulation of inducible NO synthase and NO production by physiological concentrations of ALA and γ -tocopherol.

The SFO (18:2 n -6)-rich diet increased the levels of AA in PMΦ lipids. A concomitant increase in LTB₄ was observed in A23187-stimulated PMΦ. On the contrary, the GCO10 diet reduced AA and LTB₄ levels by 38 and 40%, respectively, in PMΦ lipids. Leukotrienes are synthesised by the 5-LO pathway, and their activity was unaltered in the PMΦ of rats fed with the SFO and GCO diets. Therefore, the decrease in LTB₄ observed in the present study can be attributed to the

decrease in AA levels in the cellular lipids of PM Φ in GCO-fed rats. Since LTB₄ is a potent chemotactic agent, recruits immune cells at the site of injury and orchestrates inflammatory events leading to tissue damage, modulating its levels by *n*-3 fatty acids may be beneficial in chronic inflammatory conditions.

Phagocytosis by polymorphonuclear leucocytes or PM Φ results in increased oxygen consumption and production of reactive oxygen intermediates, which contribute to the microbicidal action of these immunocompetent cells⁽⁵¹⁾. Furthermore, lysosomal acid phosphatase activity is often considered as a marker and correlates with the bactericidal action of macrophages⁽⁵²⁾. Therefore, we investigated the effect of SFO or GCO on H₂O₂ release and *in vitro* lysosomal phosphatase enzyme activity in PMA-stimulated PM Φ . The *ex vivo* release of H₂O₂ and lysosomal phosphatase activity were not affected in the PM Φ of rats fed with SFO or GCO. This observation shows that GCO may not modulate the bactericidal activity of macrophages.

Conclusion

We report that dietary supplementation of GCO increased ALA, EPA and DHA levels in the membrane lipids of immunocompetent cells. GCO significantly suppressed the *ex vivo* proliferation of SL in response to stimulation by T-cell mitogens with little or no effect on IL-2 secretion. GCO suppressed the release of inflammatory mediators such as LTB₄, NO and, to a lesser extent, TNF- α in activated PM Φ . Our data add further evidence that dietary supplementation of ALA alters the lipid composition of immune cells and modulates certain inflammatory mediators in normal rats. However, further studies are needed to understand the long-term effect of ALA in the modulation of inflammatory mediators in ulcerative colitis, arthritis, asthma and allergic disorders.

Acknowledgements

The authors thank Dr V. Prakash, Director, Central Food Technological Research Institute, and Dr P. V. Salimath, Head of the Department of Biochemistry and Nutrition, for their encouragement and support. K. A. N. gratefully acknowledges the financial assistance in the form of a project from the Indian Council of Medical Research, New Delhi. Financial support to B. T. D. through a Senior Research Fellowship from the Council for Scientific and Industrial Research, New Delhi, India, is also gratefully acknowledged. There are no conflicts of interest whatsoever among the authors. B. T. D. was responsible for conducting the experiments. K. A. N. supervised the experiments and prepared the manuscript. B. R. L. helped in the interpretation of the results and preparation of the manuscript.

References

- Clandinin MT, Cheema S, Field CJ, *et al.* (1991) Dietary fat: exogenous determination of membrane structure and cell function. *FASEB J* **5**, 2761–2769.
- Calder PC (1998) Dietary fatty acids and lymphocyte functions. *Proc Nutr Soc* **57**, 487–502.
- Simopoulos AP (2002) The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed Pharmacother* **56**, 365–379.
- Fan YY, McMurray DN, Ly LH, *et al.* (2003) Dietary (*n*-3) polyunsaturated fatty acids remodel mouse T-cell lipid rafts. *J Nutr* **133**, 1913–1920.
- Li Q, Wang M, Tan L, *et al.* (2005) Docosahexaenoic acid changes lipid composition and interleukin-2 receptor signaling in membrane rafts. *J Lipid Res* **46**, 1904–1913.
- Bouwens M, Rest O, Dellschaft, *et al.* (2009) Fish-oil supplementation induces antiinflammatory gene expression profiles in human blood mononuclear cells. *Am J Clin Nutr* **90**, 415–424.
- Burdge GC, Jones AE & Wooton SA (2002) Eicosapentaenoic and docosapentaenoic acids are the principal products of α -linolenic acid metabolism in young men. *Br J Nutr* **88**, 355–363.
- Pawlosky RJ, Hibbeln JR, Novotny JA, *et al.* (2001) Physiological compartmental analysis of α -linolenic acid metabolism in adult humans. *J Lipid Res* **42**, 1257–1265.
- Morisea A, Combeb N, Bouébé C, *et al.* (2004) Dose effect of α -linolenic acid on PUFA conversion, bioavailability and storage in the hamster. *Lipids* **39**, 325–334.
- Diwakar BT, Dutta PK, Lokesh BR, *et al.* (2008) Bioavailability and metabolism of *n*-3 fatty acid rich garden cress (*Lepidium sativum*) seed oil in albino rats. *Prostaglandins Leukot Essent Fatty Acids* **78**, 123–130.
- Jeffery NM, Sanderson P, Sherrington EJ, *et al.* (1996) The ratio of *n*-6 to *n*-3 polyunsaturated fatty acids in the rat diet alters serum lipid levels and lymphocyte functions. *Lipids* **31**, 737–745.
- Cobljnj GB, Murphy EJ, Othman R, *et al.* (2009) Flaxseed oil and fish-oil capsule consumption alters human red blood cell *n*-3 fatty acid composition: a multiple-dosing trial comparing 2 sources of *n*-3 fatty acid. *Am J Clin Nutr* **88**, 801–809.
- Kelley DS, Branch LB, Love JE, *et al.* (1991) Dietary (α -linolenic acid and immunocompetence in humans. *Am J Clin Nutr* **53**, 40–46.
- Caughey GE, Mantzioris E, Gibson RA, *et al.* (1996) Interleukin 1 production of diets enriched in *n*-3 fatty acids from vegetable oil or fish oil. *Am J Clin Nutr* **63**, 116–122.
- Mizota T, Kambara CF, Matsuya N, *et al.* (2009) Effect of dietary fatty acid composition on Th1/Th2 polarization in lymphocytes. *J Parent Enteral Nutr* **33**, 390–396.
- Diwakar BT, Dutta PK, Lokesh BR, *et al.* (2010) Physicochemical properties of ω -3 fatty acid rich garden cress (*Lepidium sativum* L.) seed oil. *J Am Oil Chem Soc* **87**, 539–548.
- Sladowski D, Steer SJ, Clothier RH, *et al.* (1993) An improved MTT assay. *J Immunol Methods* **157**, 203–207.
- Yaqoob P & Calder P (1995) Effects of dietary lipid manipulation upon inflammatory mediator production by murine macrophages. *Cell Immunol* **163**, 120–128.
- Manosroi A, Saraphanchotiwitthaya A & Manosroi J (2003) Immunomodulatory activities of *Clausena excavate* Burm. f. wood extracts. *J Ethnopharmacol* **89**, 155–160.
- Raghavenra H, Diwakra BT, Lokesh BR, *et al.* (2006) Eugenol – the active principle from cloves inhibits 5-lipoxygenase activity and leukotriene-C₄ in human PMNL cells. *Prostaglandins Leukot Essent Fatty Acids* **74**, 23–27.
- Lowry OH, Rosebrough NJ, Farr A, *et al.* (1951) Protein measurement with Folin phenol reagent. *J Biol Chem* **143**, 265–271.

22. Morisson WR & Smith LM (1964) Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride methanol. *J Lipid Res* **5**, 600–608.
23. Furth RV & Colin ZA (1968) The origin and kinetics of mononuclear phagocytes. *J Exp Med* **128**, 415–435.
24. Kaku S, Yunoki SI, Ohkura KI, *et al.* (2001) Interactions of dietary fats and proteins on fatty acid composition of immune cells and LTB₄ production by peritoneal exudate cells of rats. *Biosci Biotechnol Biochem* **65**, 315–321.
25. Abbas AK, Lichtman AH & Pober JS (1994) *Cellular and Molecular Immunology*. Philadelphia, PA: WB Saunders.
26. Wang YW, Field CJ & Sim JS (2000) Dietary polyunsaturated fatty acids alter lymphocyte subset proportion and proliferation, serum immunoglobulin G concentration, and immune tissue development in chicks. *Poult Sci* **79**, 1741–1748.
27. Jeffery NM, Newsholme EA & Calder PC (1997) The level of polyunsaturated fatty acids and the *n*-6 to *n*-3 polyunsaturated fatty acid ratio in the rat diet both affect serum lipid levels and lymphocyte functions. *Prostaglandins Leukot Essent Fatty Acids* **57**, 149–160.
28. Calder PC, Costa-Rosa LF & Curi R (1994) Effects of feeding lipids of different fatty acid compositions upon rat lymphocyte proliferation. *Life Sci* **56**, 455–463.
29. Thies F, Nebe-von-Caron G, Powell JR, *et al.* (2001) Dietary supplementation with γ -linolenic acid or fish oil decreases T lymphocyte proliferation in healthy older humans. *J Nutr* **131**, 1918–1927.
30. Turek JJ, Schoenlein IA, Clark LK, *et al.* (1994) Dietary polyunsaturated fatty acids effects on immune cells of the porcine lung. *J Leukoc Biol* **56**, 599–604.
31. Wallace FA, Miles EA, Evans C, *et al.* (2001) Dietary fatty acids influence the production of Th1- but not Th2-type cytokines. *J Leukoc Biol* **69**, 449–457.
32. Meydani SN, Lichtenstein AH, Cornwall S, *et al.* (1993) Immunologic effects of national cholesterol education panel step-2 diets with and without fish-derived *n*-3 fatty acid enrichment. *J Clin Invest* **92**, 105–113.
33. Yaqoob P, Pala HS, Cortina-Borja M, *et al.* (2000) Encapsulated fish oil enriched in α -tocopherol alters plasma phospholipid and mononuclear cell fatty acid compositions but not mononuclear cell functions. *Eur J Clin Invest* **30**, 260–274.
34. Endres S, Meydani SN, Ghorbani R, *et al.* (1993) Dietary supplementation with *n*-3 fatty acids suppresses interleukin-2 production and mononuclear cell proliferation. *J Leukoc Biol* **54**, 599–603.
35. Babu US, Wiesenfeld PW, Collins TFX, *et al.* (2003) Impact of high flaxseed diet on mitogen-induced proliferation, IL-2 production, cell subsets and fatty acid composition of spleen cells from pregnant and F1 generation Sprague–Dawley rats. *Food Chem Toxicol* **41**, 905–915.
36. Jolly CA, Mac Murray DN & Chapkin RS (1998) Effect of dietary *n*-3 fatty acids on interleukin-2 and interleukin-2 receptor α expression in activated murine lymphocytes. *Prostaglandins Leukot Essent Fatty Acids* **58**, 289–293.
37. Stulnig TM, Huber J, Leitinger N, *et al.* (2001) Polyunsaturated eicosapentaenoic acid displaces proteins from membrane rafts by altering raft lipid composition. *J Biol Chem* **276**, 37335–37340.
38. Ebinu JO, Stang SL, Teixeira C, *et al.* (2000) Ras GRP links T cell receptor signaling to Ras. *Blood* **95**, 3199–3203.
39. Bi K & Altman A (2001) Membrane lipid microdomains and the role of PKC ζ in T cell activation. *Semin Immunol* **13**, 139–146.
40. Xavier R, Brennan T, Li Q, *et al.* (1998) Membrane compartmentalization is required for efficient T cell activation. *Immunity* **8**, 723–732.
41. Godge MP (1998) How cytotoxic is nitric oxide? *Exp Nephrol* **6**, 12–16.
42. Boutard V, Fouquery B, Philippe C, *et al.* (1994) Fish oil supplementation and essential fatty acid deficiency reduce nitric oxide synthesis by rat macrophages. *Kidney Int* **46**, 1200–1286.
43. Joe B & Lokesh BR (1994) Role of capsaicin, curcumin and dietary *n*-3 fatty acids in lowering the generation of reactive oxygen species in rat peritoneal macrophages. *Biochim Biophys Acta* **1224**, 255–263.
44. Renier G, Skamene E, de Sanctis J, *et al.* (1993) Dietary *n*-3 polyunsaturated fatty acids prevent the development of atherosclerotic lesions in mice: modulation of macrophage secretory activities. *Arterioscler Thromb* **13**, 1515–1524.
45. Chaet MS, Garcia VF, Arya G, *et al.* (1994) Dietary fish oil enhances macrophage production of nitric oxide. *J Surg Res* **57**, 65–68.
46. Hubbard NE, Chapkin RS & Erickson KL (1994) Effect of dietary linseed oil on tumoricidal activity and eicosanoid production in murine macrophages. *Lipids* **29**, 651–655.
47. Narayanan BA, Narayanan KN, Simi B, *et al.* (2003) Modulation of inducible nitric oxide synthase and related proinflammatory genes by the omega-3 fatty acid docosahexaenoic acid in human colon cancer cells. *Cancer Res* **63**, 972–979.
48. Aldridge C, Razzak A, Babcock TA, *et al.* (2008) LPS-stimulated RAW 264.7 macrophage inducible nitric oxide synthase (iNOS) and nitric oxide production is decreased by an omega-3 fatty acid lipid emulsion. *J Surg Res* **149**, 296–302.
49. Ren J & Chung SH (2007) Anti-inflammatory effect of (γ -linolenic acid and its mode of action through the inhibition of nitric oxide production and inducible nitric oxide synthase gene expression via NF- κ B and mitogen-activated protein kinase pathways. *J Agr Food Chem* **55**, 5073–5080.
50. Jiang Q, Schwab IE, Courtemanche C, *et al.* (2000) γ -Tocopherol and its major metabolite, in contrast to α -tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc Natl Acad Sci U S A* **97**, 11494–11499.
51. Allen RC (1994) Role of oxygen in phagocyte microbicidal action. *Environ Health Perspect* **102**, Suppl. 10, 201–208.
52. Schnyder J & Baggiolini M (1978) Secretion of lysosomal hydrolases by stimulated and nonstimulated macrophages. *J Exp Med* **148**, 435–450.