

THE JOURNAL OF AGRICULTURAL SCIENCE

EDITED BY

G. D. H. BELL, C.B.E., PH.D., F.R.S., Plant Breeding Institute, Cambridge

K. L. BLAXTER, PH.D., N.D.A. (HONS.), D.SC., F.R.S.E., F.R.S.,
Rowett Research Institute, Bucksburn, Aberdeen

G. W. COOKE, PH.D., F.R.I.C., F.R.S., Rothamsted Experimental Station, Harpenden

JOHN HAMMOND, Jr., M.A., School of Agriculture, Cambridge

PROF. SIR J. B. HUTCHINSON, C.M.G., SC.D., F.R.S., School of Agriculture,
Cambridge

H. L. PENMAN, O.B.E., PH.D., F.R.S., Rothamsted Experimental Station, Harpenden

W. J. RIDGMAN, M.A., School of Agriculture, Cambridge

H. H. ROGERS, B.SC., DIP.AG.SCI., Plant Breeding Institute, Cambridge

PROF. E. W. RUSSELL, C.M.G., PH.D., F.INST.P., Department of Soil Science,
University of Reading

F. YATES, C.B.E., SC.D., F.R.S., Rothamsted Experimental Station, Harpenden



CAMBRIDGE UNIVERSITY PRESS

BENTLEY HOUSE, 200 EUSTON ROAD, LONDON, N.W. 1

AMERICAN BRANCH: 32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Published bi-monthly

Annual subscription £9 net (U.S.A. \$29.00)

Single parts £2 net (U.S.A. \$6.00)

Second class postage paid at New York, N.Y.

INSTRUCTIONS TO AUTHORS

Failure to comply with the Instructions to Authors may delay publication

Papers intended for publication should be submitted to Dr G. D. H. Bell, C.B.E., F.R.S., Plant Breeding Institute, Cambridge, CB2 2LQ or to one of the Associate Editors. It must be understood that, if accepted by the Editorial Board, the paper will not be published elsewhere in the same form.

MANUSCRIPTS. Papers, written in English, are accepted from any country and should be typed in double-spacing on one side of the paper with a margin at least 4 cm wide on the left-hand side. Authors should instruct typists on the style required. A top copy and one carbon copy should be submitted.

SHORT NOTES may be accepted provided they are based on adequate experimental evidence; special provision is made for their publication with the least possible delay. MSS should not exceed 1500 words in length or their equivalent. For tabulated matter allow 25 words per line of the table (including headings). For line illustrations allow 225 words per quarter of a page.

TABLES must be self-explanatory. They should be typed on separate sheets, numbered consecutively and carry an appropriate title. When possible, tables should be arranged so that they can be printed in the normal orientation of the text and without rules.

LINE DRAWINGS (with photocopies) should be 25 cm wide and drawn in black waterproof ink on Bristol board, graph paper with blue lines or tracing paper. Legends should be typed on a separate copy and numbering inserted lightly and clearly in soft pencil on the drawing. Tables and figures should not reproduce the same data.

The approximate position of tables and figures should be noted in the text.

PLATES should make a definite contribution to the value of the paper and the number submitted should be kept to a minimum. They should be good quality, unmounted, glossy prints and be lightly numbered in pencil on the reverse side. If several, or coloured, plates are submitted the author may be asked to contribute to the cost of reproducing them.

TITLE. The title must be specific and suitable for indexing by the mechanical methods now being employed. The full name and address of the institution in which the research has been carried out should be stated. Change of address may be given as a footnote. A short title, not exceeding 50 characters, should be provided for the running headlines.

STYLE. Experimental details and results should be recorded in the past tense and there should be no unnecessary repetition or loose phrases. Manuscripts are likely to be returned for modification if the presentation is not clear and precise.

LAYOUT. Authors are recommended to study '*General Notes on the Preparation of Scientific Papers*' (Royal Society, London, 2nd edn, 1965). The Editorial Board do not insist on a rigid format but it is usually convenient to divide the paper into sections, e.g. Introduction, Materials and Methods, Results, and Discussion. An excess of headings and sub-headings should be avoided.

Authors are advised to note the following points: a detailed review of literature is not necessary; relevant details should be given of the plant or animal material, the experimental design and chemical or other techniques employed; mean results with their relevant standard errors should be presented rather than detailed data; the statistical methods used should be clearly stated; the discussion should relate the author's experiments to other work on the subject and give the author's conclusions. Footnotes should be avoided.

SUMMARIES of papers are placed at the beginning of the text and authors should submit MSS with the summaries so placed. The summary should be factual and suitable for use in abstracting journals; paragraphs should not be numbered.

REFERENCES. The bibliography must be given in the form—Surname of authors, initials, year of publication (in parentheses), title of paper, name of journal (abbreviated according to the *World List of Scientific Periodicals*, 4th edn, Butterworths, London), volume and pages of reference. References should be in alphabetical order. In the text a reference should be quoted by the author's name and date (in parentheses). Where there are more than two authors, the initial reference in the text should include the names of all authors, but subsequent citations should be in the form—first author followed by *et al.* Authors should check that all references in the text appear at the end of the paper and vice versa, and that names and dates correspond in the two places.

PROOF CORRECTION. Standard proof correction marks (British Standard 1219) should be made as legibly as possible in ink, not pencil. Directions to the printer which are not to be set up in type should be encircled. Captions to illustrations and all references should be checked. Queries marked by the printer should be answered. Proofs are provided in order that authors can check the correctness of the type-setting—excessive alterations may be charged to the author.

OFFPRINTS. Contributors will receive 25 copies of their papers free and can order others when they receive the proofs.

Aldrin

makes the earth a healthy place for crops to grow in

Aldrin is a highly effective Shell insecticide.

It works *underground* to destroy the full range of soil insect pests, which attack the vital root system of your crop. For the major groups of soil pests, wireworms, cutworms, white grubs and mole-crickets, it is recognised as the most efficient insecticide there is. It can be applied to any type of soil in the world, and is harmless to beneficial organisms in the soil.


Used for row treatment, as an overall application, as a root dip, or as a seed-treatment, aldrin is especially suited for use with maize and other cereals, sugar cane, sugar beet, potatoes and tobacco.

Aldrin has excellent residual action. It is stable in the presence of other agricultural chemicals, does not affect germination, and has no taint. Used as directed, aldrin is harmless to both farmers and consumers. It is available as the technical material and high

concentrate dusts for further formulation, and also in forms suitable for the grower: emulsifiable concentrates, wettable powders, field strength dusts and granules.

For details of approved uses and advice on application of aldrin consult your Shell company or Shell chemicals distributor.

Shell Chemicals



JOURNAL OF REPRODUCTION AND FERTILITY

The Journal publishes original papers, abstracts of Proceedings, and occasionally reviews and bibliographies, dealing with the morphology, physiology, biochemistry and pathology of reproduction in man and animals, and with the biological, medical and veterinary problems of fertility and lactation. Nine numbers are issued to give three volumes a year.


Papers appearing during 1969 include:

- D. P. BOSHIER. A histological and histochemical examination of implantation and early placentome formation in sheep.
- J. MATOUSEK. Effects on spermatogenesis in guinea-pigs, rabbits and sheep after their immunization with sexual organ fluids of bulls.
- T. D. QUINLIVAN and T. J. ROBINSON. Numbers of spermatozoa in the genital tract after artificial insemination of progestagen-treated ewes.
- L. A. JOHNSON, R. J. GERRITS and E. P. YOUNG. Quantitative analysis of porcine spermatozoa and seminal plasma phospholipids as affected by frequency of ejaculation.
- A. N. BRUERE, R. B. MARSHALL and D. P. J. WARD. Testicular hypoplasia and XXY sex chromosome complement in two rams: the ovine counterpart of Klinefelter's syndrome in man.
- J. C. BOURSNELL and P. A. BRIGGS. Boar seminal plasma proteins. 2. Electrophoretic identification of the haemagglutinin.
- R. H. SCHULTZ, H. B. BURCALOW, M. L. FAHNING, E. F. GRAHAM and A. F. WEBER. A karyometric study of epithelial cells lining the glands of the bovine endometrium.
- D. WIERZBOWSKI and E. WIERZCHOS. Cannulation of the vas deferens in the boar.
- B. HOWARTH, Jr. Fertility in the ram following exposure to elevated ambient temperature and humidity.
- L. E. APPELGREN. Histochemical demonstration of drug interference with progesterone synthesis.
- F. L. CHERMS. Influence of the histocompatibility system on reproductive efficiency of turkeys.
- B. M. AUSTERN and A. M. GAWIENOWSKI. The isolation of retinal from the bovine corpus luteum.
- J. M. O'DONNELL. Intracellular levels of sodium and potassium in bull spermatozoa in relation to cell metabolism.
- I. S. WHEATLEY and H. M. RADFORD. Luteinizing hormone secretion during the oestrous cycle of the ewe as determined by radio-immunoassay.

Editorial correspondence should be sent to the Editor, Journal of Reproduction and Fertility, 8 Jesus Lane, Cambridge, England, and business correspondence to the publishers.

The subscription rate for 1969 is £7 per volume or £21 per year (three volumes). U.S.A. and Canada \$23.50 per volume or \$70.00 per year.

BLACKWELL SCIENTIFIC PUBLICATIONS
5 ALFRED STREET OXFORD



Stainless Steel

MICRO HAMMER MILL

pulverizes almost anything—whether hay, cotton, wool or rock samples.
Only two minutes needed to pulverize most samples. Even a few grammes can be ground;
 maximum charge 120 c.c. Dust-proof closure. Hardened stainless steel
 grinding mechanism. Slide-in screens for particle size adjustment.
 Variable speed motor for better grinding control. Price, complete as shown; £84.
For full details please quote AS/63/M.H.



GLEN CRESTON

The Red House, 37 The Broadway, Stanmore, Middlesex.
 Telephone: 01-954 4218


Experimental Agriculture

Volume 5, Number 2. April 1969

- P. C. OWEN The growth of four rice varieties as affected by temperature and photoperiod with uniform daily periods of daylight.
- A. R. BROMFIELD Uptake of phosphorus and other nutrients by maize in western Nigeria.
- P. C. OWEN Comparative potential yields of rice varieties in the Australian tropics.
- O. L. OKE Sulphur nutrition of legumes.
- K. C. ANANTH Timing and frequency of spraying for control of coffee leaf rust in southern India.
- H. K. PANDE and PANJAB SINGH Effects of moisture and nitrogen on growth, yield and mineral content of rice.
- R. W. SMITH Fertilizer responses by coconuts (*Cocos nucifera*) on two contrasting Jamaican soils.
- ABDALLA AHMED ABDALLA Effects of planting date and spacing on yields of onion seed under hot arid conditions in the Sudan.
- D. A. HOLLAND Component analysis: an aid to the interpretation of data.
- BOOK REVIEWS.

35s. net (\$5.00). Annual subscription £5 net (\$16.50) for four issues.

CAMBRIDGE UNIVERSITY PRESS



COLUMBIA

COLUMBIA

THE RISE AND FALL OF T. D. LYSENKO

Zhores A. Medvedev

Translated by I. Michael Lerner

This book tells the story of Soviet genetics in the period 1937-1964. In these years T. D. Lysenko, an illiterate and fanatical charlatan, was allowed absolute dictatorship and control over research in biology and practical agriculture in the Soviet Union. His domination in these subjects not only stifled the development of science but also had far-reaching and destructive influence on Soviet economy. Geneticists and agriculturists in the West have long speculated about the machinery of the Lysenko 'take-over' in biology and about the situation that made it possible. This book provides an answer. \$10.00

COLUMBIA UNIVERSITY PRESS, 440 West 110th Street, New York, N.Y. 10025
Address for orders: 136 South Broadway, Irvington, N.Y. 10533



ANIMAL PRODUCTION

JOURNAL OF THE BRITISH SOCIETY OF ANIMAL PRODUCTION

Contents of Volume 10, Part 3, August 1968 include the following

- CUTHBERTSON and PEASE. The inter-relationships of various measurements, visual assessments and dissection results of pigs of 200 lb. live weight.
- WATSON and BROADBENT. Inherited variation in carcass composition of Suffolk x Welsh lambs.
- SLEE. Body temperature and vasomotor responses in Scottish Blackface and Tasmanian Merino sheep subjected to slow cooling.
- MARSHALL, WATSON and SLATTERY. Prediction of eye-muscle area from introscope measurements on carcasses of pigs of bacon weight.
- TROELSEN and CAMPBELL. Voluntary consumption of forage by sheep and its relation to the size and shape of particles in the digestive tract.
- ROBINSON and FORBES. The effect of protein intake during gestation on ewe and lamb performance.
- DAVIES. The effect of cereal and protein source on the energy intake and nitrogen balance of fattening lambs given all-concentrate diets.
- GRIFFITHS. Observations on neonatal changes of body temperature in Blackface lambs.
- LEVY, HOLZER and VOLCANI. The effect of age and live weight on feed conversion and yield of saleable meat of intact Israeli Friesian male calves.
- HINKS and ANDERSON, BECH. Phenotypic aspects of growth and carcass quality in veal calves.
- MORRISON, HINTZ and GRIVENS. A note on effect of exercise on behaviour and performance of confined swine.

The Journal is published quarterly in February, May, August and November. The annual subscription rate is 90s. (U.S.A. \$14.00). Single copy 25s. (U.S.A. \$4.00). Orders and subscriptions may be sent to any bookseller or direct to:

OLIVER AND BOYD LTD.

Tweeddale Court, 14 High Street, Edinburgh 1

ANIMAL PRODUCTION

JOURNAL OF THE BRITISH SOCIETY OF ANIMAL PRODUCTION

Contents of Volume 11, Part 1, February 1969 include the following:

- OWEN, DAVIES and RIDGMAN. The effects of varying the quantity and distribution of liquid feed in lambs reared artificially.
- LEAVER, CAMPLING and HOLMES. The effect of level of feeding on the digestibility of diets for sheep and cattle.
- MILES, WALTERS and EVANS. Dry-matter intake and live-weight gain of cattle and sheep offered different grass varieties with and without clover.
- SUZUKI, SHOZO, FUJITA, HIROSHI and SHINDE, YOZO. Change in the rate of eating during a meal and the effect of the interval between meals on the rate at which cows eat roughages.
- HINKS and ANDERSON, BECH. Genetic aspects of growth and carcass quality in veal calves.
- KING and GAJIC. The repeatability of maternal performance in inbred, outbred and linecross Large White sows.
- KING. The distribution of sheep B-lactoglobulins.
- SPOONER and OLIVER. Albumin polymorphism in British cattle.
- SYKES and SLEE. Cold exposure of Southdown and Welsh Mountain sheep. 1. Effects of breed, place of nutrition and acclimatization to cold upon resistance to body cooling.
- SYKES and SLEE. Cold exposure of Southdown and Welsh Mountain sheep. 2. Effects of breed, place of nutrition and previous acclimatization to cold upon skin temperature, heart rate, shivering and respiration rate.
- SYKES, FIELD and SLEE. Cold exposure of Southdown and Welsh Mountain sheep. 3. Changes in plasma calcium, phosphorus, magnesium, sodium and potassium levels.
- WINFIELD, BROWN and LUCAS. Sheltering behaviour at lambing by Welsh Mountain ewes.
- TOMOV. A note on commercial crossing of Bulgarian fine-wool ewes with rams of English short-wool breeds.
- ROLLINS, JULIAN and CARROLL. A note on the body compositions of a double muscled female and a normal female from a linebred Aberdeen Angus herd.
- THE BRITISH SOCIETY OF ANIMAL PRODUCTION. MAULE, J. P. 25 years in retrospect.
- PROCEEDINGS OF THE 48TH MEETING OF THE BRITISH SOCIETY OF ANIMAL PRODUCTION, ABERDEEN, 9-12 SEPTEMBER 1968.

Contents of Volume 11, Part 2, May 1969 include the following:

- LODGE. The effects of pattern of feed distribution during the reproductive cycle on the performance of sows.
- MCCULLOUGH. A study of factors affecting the voluntary intake of food by cattle.
- BROADBENT, BALL and DODSWORTH. The effect of plane of nutrition during calthood on the subsequent performance of Hereford × Ayrshire steers.
- LEAVER, CAMPLING and HOLMES. The influence of flexible and rigid grazing management and of supplementary feed on output per hectare and per cow.
- ANDREWS, KAY and ØRSKOV. The effect of different dietary energy concentrations of the voluntary intake and growth of intensively-fed lambs.
- ØRSKOV, ANDREWS and GILL. Effect of replacing rolled barley with swedes or potatoes on the intake and rumen volatile fatty acid composition of lambs.
- PRESCOTT, EL-SHOBOKSHY and ARMSTRONG. Ammonium salts of fatty acids for milk production.
- HUTTON, PRESCOTT, SEELEY and ARMSTRONG. Ammonium salts of fatty acids for milk production.
- HULET, FOOTE and PRICE. Factors affecting growth of ovine foetuses during early gestation.
- ELSLEY and others. The effect of level of feed intake in pregnancy and lactation upon the productivity of sows.
- MACDOUGALL, CUTHBERTSON and SMITH. The assessment of pig meat paleness by reflectance photometry.
- AGAR, RAWAT and ROY. Haemoglobin variants in Indian sheep.
- LAND, DICKINSON and READ. The influence of seasonal variation on fertility and early body growth in sheep.

The Journal is published quarterly in February, May, August and November. The annual subscription rate is 90s. (U.S.A. \$14.00). Single copy 25s. (U.S.A. \$4.00). Orders and subscriptions may be sent to any bookseller or direct to:

OLIVER AND BOYD LTD.

Tweeddale Court, 14 High Street, Edinburgh 1

Azodrin

does its job inside out-outside in

... and that means double trouble for cotton pests, because Shell's powerful new organo-phosphorus insecticide Azodrin works *two* ways.

By contact, immediately destroying insects on the plant surface. And systemically, by penetrating foliage rapidly where it kills sucking and chewing pests as they try to eat.

This sophisticated dual action makes Azodrin by far the most wide-ranging insecticide yet devised giving *continuing* control over *more* pests *more* effectively.

Azodrin has already been employed with outstanding success in the United States and

Central America, and is now proving world-wide to be one of the cotton grower's most powerful allies.

Suddenly pests that took a bite out of profits are losing their appetite.

For further information or supplies of Azodrin contact your Shell company or Shell chemicals distributor. Shell works today for your more profitable tomorrow.

Shell Chemicals

