

Conclusions. The results show that the virtual diagnosis network based on a telemedicine platform can enhance significantly the community hospital diagnostic services, maximizing professional time and productivity, increasing access and equity, and reducing costs. However, before carrying out its countrywide implementation, a contextualization with the regional epidemiological profile must be performed.

VP29 Designing A Mobile Clinical Decision Support System For Dementia

Noemí Robles (nrobles@uoc.edu), Carme Carrion i Ribas and Marta Aymerich

Introduction. E-health offers the opportunity of supporting the management of several diseases, but most of these tools are far from being based on scientific evidence and demonstrating their effectiveness and efficacy. The PSICODEM Project aims to develop a mobile personalized clinical decision support system (CDSS) based on evidence for contributing to e-health interventions addressed to the management of dementia that require not only a pharmacological approach but also psychosocial interventions for improving patients' quality of life and reducing emotional, cognitive and behavioral symptoms. The present communication focuses on the identification of the evidence on which the CDSS algorithm will be developed.

Methods. Three systematic reviews were carried out in order to identify the existing scientific evidence published in relation to the effectiveness of behavioral, emotional and cognitive therapies addressing dementia (January 2009 to December 2017). The main databases were consulted (PubMed, Cochrane Library, PsychoInfo) and only randomized control trials (RCT) were considered. Articles were reviewed by two independent reviewers. The quality of the selected publications was assessed according to the SIGN criteria.

Results. Forty-seven RCTs were selected for cognitive therapies, thirty-two for emotional ones and fifteen for behavioral interventions. Those therapies with more support of evidence were skills training for cognitive therapies and reminiscence interventions for emotional interventions; however, in behavioral interventions a variety of therapeutically approaches were found. Wide differences were found between studies in terms of types and levels of dementia, forms of intervention (number, length and frequency of sessions) and outcome measures.

Conclusions. In-depth analysis of evidence will allow the identification of those interventions more appropriate for each patient according to their symptoms and level of dementia. According to this evidence, the mobile CDSS algorithm will be developed. Additionally, these findings point out the gaps in psychosocial intervention research.

VP30 Evaluation Of CINAHL In Six Systematic Reviews On Maternal Care

Inga Overesch (inga.overesch@iqwig.de), Dorothea Sow, Elke Hausner and Nina Peterwerth

Introduction. Information retrieval for systematic reviews (SRs) should include sensitive searches in several bibliographic databases. In addition to standard databases (i.e., MEDLINE, Embase and CENTRAL), researchers might consider subject-specific ones. In the fields of nursing and midwifery, a SR would typically include CINAHL as a subject-specific database. The aim of this study was to analyze the number and relevance of references retrieved from CINAHL in six SRs on maternal care.

Methods. We conducted a retrospective analysis of six SRs (e.g., benefit of intrapartum ultrasound or one-to-one care during labor). The study type was limited to randomized controlled trials (RCTs) in all but three SRs. In all cases, MEDLINE, Embase, CENTRAL and CINAHL were searched for primary studies. Further information sources (e.g., study registries and reference lists of SRs) were also considered. The proportion of the additional number of hits and studies included from CINAHL as well as the corresponding number of participants were calculated.

Results. Overall, the reviewers screened 12,013 references from bibliographic databases and identified forty relevant studies. CINAHL contained 2,643 (22 percent) of the references. In five out of six SRs, no additional studies were identified in CINAHL. In the remaining SR on birthing positions, the reviewers included thirteen RCTs of which one was a feasibility study with 68 participants indexed only in CINAHL. This corresponds to 0.9 percent of the women participating in all thirteen RCTs ($n = 7,861$). However, this study was cited in a journal article on a subsequent RCT that was identified and included via MEDLINE and ClinicalTrials.gov.

Conclusions. It is not necessary to search CINAHL in SRs on maternal care if standard databases and further information sources are considered. An additional study from CINAHL was included in one out of six SRs, a small feasibility study that could have been identified without CINAHL via a subsequent RCT.

VP31 Searching Non-English Literature For HTA Reports May Be Unnecessary

Lisa Schell (lisa.schell@iqwig.de), Elke Hausner, Lina Rodenhäuser, Oliver Assall, Anke Schulz, Wiebke Sieben, Kerstin van der Leck and Stefan Sauerland

Introduction. Currently, the Institute for Quality and Efficiency in Health Care (IQWiG) does not restrict literature searches by language. Given limited resources, it is unclear whether the effort put into screening and translating studies published in non-English and non-German (nEnG) languages yields much new information when compared to including only English and German literature. Therefore, we aimed to analyze the impact of nEnG literature on the conclusion of IQWiG's health technology assessments (HTAs).

Methods. We checked for seventy-two IQWiG HTAs (all non-drug intervention HTAs published until August 2018 and three additional HTAs on drugs) whether they included nEnG studies. For all HTAs including at least one nEnG study, we analyzed whether the statistical significance would have changed for any endpoint without the respective nEnG study(ies). If no