


ERRATUM

## Influence of trap colour, type, deployment height, and a host volatile on monitoring *Orchestes fagi* (Coleoptera: Curculionidae) in Nova Scotia, Canada—ERRATUM

Joel T.L. Goodwin<sup>1,2\*</sup> , Simon P. Pawlowski<sup>1</sup>, Peter D. Mayo<sup>3</sup>, Peter J. Silk<sup>3</sup>, Jon D. Sweeney<sup>3</sup>, and N. Kirk Hillier<sup>1</sup>

<https://doi.org/10.4039/tce.2019.63>. Published by Cambridge University Press, 19 December 2019

<sup>1</sup>Department of Biology, Acadia University, 33 Westwood Avenue, Wolfville, Nova Scotia, B4P 2R6, Canada <sup>2</sup>Faculty of Forestry, University of Toronto, 33 Wilcox Street, Toronto, ON, M5S 3E8, Canada; and <sup>3</sup>Natural Resources Canada, Canadian Forest Service-Atlantic Forestry Centre, 1350 Regent Street, Fredericton, New Brunswick, E3C 2G6, Canada

Owing to a printer's error, the affiliations in the original online version of this article were incorrectly listed. They have been corrected here and in both the print and revised online version of the article.

We regret the error.

### Reference

Goodwin, J.T.L., Pawlowski, S.P., Mayo, P.D., Silk, P.J., Sweeney, J.D., and Hillier, N.K. 2019. Influence of trap colour, type, deployment height, and a host volatile on monitoring *Orchestes fagi* (Coleoptera: Curculionidae) in Nova Scotia, Canada. The Canadian Entomologist, 152: 98–109.