

- (f) Examine groups of drugs by side-effect profile. For example, susceptibility to metabolic side-effects such as maternal weight gain, gestational hypertension and gestational diabetes, and low-potency FGA/typicals should be grouped with -pines; thus, for example the 'typical' chlorpromazine grouped with 'atypical' olanzapine and clozapine.
- (g) Compare outcomes between siblings and generations to account for non-measured family-wide effects.
- (h) Examine evidence of longer-term developmental effects.

**Kathryn M. Abel**, MA, MBBS, MRCP, MRCPsych, PhD, Centre for Women's Health, 3rd Floor Jean McFarlane Building, University of Manchester, Manchester M13 9PL, UK. Email: kathryn.abel@manchester.ac.uk

First received 11 Jul 2012, final revision 24 Oct 2012, accepted 31 Oct 2012

## Acknowledgements

With thanks to Dr Robert Bodén, Professors Cecilia Magnusson, Christina Dalman and Peter Brocklehurst for useful discussions on the use of Scandinavian and other population samples in pharmaco-epidemiology.

## References

- McKenna K, Koren G, Tetelbaum M, Wilton L, Shakir S, Diav-Citrin O, et al. Pregnancy outcome of women using atypical antipsychotic drugs: a prospective comparative study. *J Clin Psychiatry* 2005; **66**: 444–9.
- Webb RT, Howard L, Abel KM. Antipsychotic drugs for non-affective psychosis during pregnancy and postpartum. *Cochrane Database Syst Rev* 2004; **2**: CD004411.
- National Institute for Health and Clinical Excellence. *Antenatal and Postnatal Mental Health. Clinical Guideline 45*. NICE, 2007.
- Barnes TR. Evidence-based guidelines for the pharmacological treatment of schizophrenia: recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2011; **25**: 567–620.
- Webb RT, Pickles AR, King-Hele SA, Appleby L, Mortensen PB, Abel KM. Parental mental illness and fatal birth defects in a national birth cohort. *Psychol Med* 2008; **38**: 1495–503.
- Altshuler LL, Cohen L, Szuba MP, Burt VK, Gitlin M, Mintz J. Pharmacological management of psychiatric illness during pregnancy: dilemmas and guidelines. *Am J Psychiatry* 1996; **153**: 592–606.
- Reis M, Kallen B. Maternal use of antipsychotics in early pregnancy and delivery outcome. *J Clin Psychopharmacol* 2008; **28**: 279–88.
- Gentile S. Antipsychotic therapy during early and late pregnancy. A systematic review. *Schizophr Bull* 2010; **36**: 518–44.
- Bodén R, Lundgren M, Brandt L, Reutfors J, Kieler H. Antipsychotics during pregnancy: relation to fetal and maternal metabolic effects. *Arch Gen Psychiatry* 2012; **69**: 715–21.
- Lin HC, Chen IJ, Chen YH, Lee HC, Wu FJ. Maternal schizophrenia and pregnancy outcome: does the use of antipsychotics make a difference? *Schizophr Res* 2010; **116**: 55–60.
- Abel KM, Morgan VA. Mental illness, women, mothers, and their children. In *Textbook in Psychiatric Epidemiology (3rd edn)* (eds MT Tsang, M Tohen, P Jones): 483–514. Wiley-Blackwell, 2011.
- Bodén R, Lundgren M, Brandt L, Reutfors J, Andersen M, Kieler H. Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. *BMJ* 2012; **345**: e7085.
- The Health and Social Care Information Centre. *Prescription Cost Analysis: England 2010*. The Health and Social Care Information Centre, 2011.
- Newport DJ, Calamaras MR, DeVane CL, Donovan J, Beach AJ, Winn S, et al. Atypical antipsychotic administration during late pregnancy: placental passage and obstetrical outcomes. *Am J Psychiatry* 2007; **164**: 1214–20.

Poems  
by  
doctors

## Ode to Anxiety

Ankur Sharma

Whine of mangled nerves  
Like convoluting steel rails  
Conducting trains of worry  
Deafeningly.

Blur of contracting sight  
Like curtains falling  
Leaden lids, shutting down  
Hazily.

Anxiety, you are potent  
Like cyanide  
Choking on contact  
Gaspingly.

Leave me, leave me  
Like life from a corpse  
Stop this galloping pulse  
Hurriedly.

©Ankur Sharma.

The British Journal of Psychiatry (2013)  
202, 323. doi: 10.1192/bjp.bp.111.105791