

Introduction: Amnesia and palimpsests occurring and recurring in alcohol addicts due to alcohol intoxication (Ebrietas alcoholica) are accompanied by hazardous memory failures, gradual mental degradation and psychoorganic syndrome, which evidences urgent clinical, therapeutic and therapeutic issue in addictology, psychiatry, forensic medicine, sociology, medical psychology, etc. At EPA initiative (2019), research interest in non-invasive brain stimulation tools and methods for such populations was activated.

Objectives: Development of a patentable method of treatment in addictology using pyracetam and nicotinic acid transcerebral electrophoresis (TCE).

Methods: Valid clinical-diagnostic, laboratory, biochemical, electrophysiological, psychological (scaling, testing), statistical methods for identification of alcohol dependence complicated by amnesic disorders.

Results: The method of treatment of alcohol dependence complicated by amnesic disorders (Patent 141785 UA) provides complex pharmacological and drug-free therapy. Antiamnesic drugs are administered by TCE bilaterally; pyracetam 20% solution to the left orbit through active negative electrodes, and nicotinic acid 0.1% solution to the right orbit (positive electrode in the occipital fossa), current of 2-4 mA, 20-30 minutes exposure. The procedure was performed daily with a TCE device, for a 10-day course of treatment along with psychotherapeutic potentiation. TCE provides the ionic implementation of pharmacological agents in the brain and their physiological electrical stimulation.

Conclusions: In a representative clinical trial, using statistical methods and generated bank of patient-specific observations, significant potentiating effects of combined drug-free, non-invasive transcerebral electrical stimulation and electrophoretic implementation of pyracetam and nicotinic acid were demonstrated.

Keywords: pharmacotherapy; Alcohol addiction; Amnesic disorders

EPP1325

Quality of life of alcohol dependence patients who have been having acute psychotic disorder

V. Kuzminov

Department Of Emergency Psychiatry And Narcology, SI Institute of Neurology, Psychiatry and Narcology NAMS of Ukraine, Kharkiv, Ukraine

doi: 10.1192/j.eurpsy.2021.1503

Introduction: Severe acute psychosis significantly alters patient's quality of life in patients with alcohol dependence. The aim of the investigation were examination value quality of the life patients with alcohol dependence who have recently suffered of acute psychotic disorder. The factor influencing the quality of life is the psychoorganic syndrome after acute psychosis.

Objectives: 120 patients with alcohol dependence who had recent history of acute psychosis were examined.

Methods: Psychopathological.

Results: The psychorganic syndromes at these patients were investigated. The Index quality of the life in these patients was assessed due to type of the psychorganic syndromes. The dynamics of the Index quality of the life at patients with psychorganic syndrome during the treatment were described. The subjective assessment of their condition in patients with hard psychorganic syndrome was dissociated from the assessment of doctors and relatives. The explaining the characteristics of the consequences of the transferred psychotic disorder to the patients turned out to be important for overcoming anosognosia. The Index quality of the life in these

patients was assessed repeatedly at the same time, there was a significant decrease in the difference in the assessment of the quality of life by patients with relatives.

Conclusions: The importance of value quality of the life from the point of the patient, relatives of the patient and physician was underlined. The assessment of Index quality of the life is important important to explain the peculiarities of the postpsychotic state to the patients and their relatives in order to develop rehabilitation programs and carrying out psychotherapeutic activities.

Keywords: Alcohol dependence; acute psychotic disorder; quality of life

EPP1326

Impact of addictive behaviors on productivity at work among employees working on an onshore oil field

N. Rmadi¹, N. Kotti¹, R. Masmoudi^{2*}, F. Dhouib¹, K. Jmal Hammami¹, M. Larbi Masmoudi¹, J. Masmoudi² and M. Hajjeji¹

¹Department Of Occupational Medicine, HEDI CHAKER hospital, SFAX, Tunisia and ²Psychiatrie "a" Department, Hedi Chaker Hospital University -Sfax - Tunisia, sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1504

Introduction: Addictive behaviors on the workplace are a real public health problem because of its consequences not only on workers but also on productivity at work.

Objectives: To explore the relationship between addictive behaviors and productivity at work among employees of a Tunisian oil rig.

Methods: A cross-sectional study was conducted in the first half of 2018. The assessment of work productivity was done using the validated WPAI-GH questionnaire. Smoking dependence was assessed via the Fagerström score and alcohol abuse by the FACE questionnaire.

Results: It was 94 employees working in an onshore oil field with an average age of 41.1 years. Average job seniority was 14.3 years. Active smoking was noted in 34.7% of cases. Alcohol consumption was noted in 19.1% of cases. In the 7 days preceding the survey, the average percentage of absenteeism was $3.64 \pm 21.7\%$ and the presenteeism was $17.66 \pm 25.58\%$. The average decline in productivity was $14.8 \pm 43.7\%$ and the average decline in daily activities was $20.21 \pm 31.45\%$. These parameters were not correlated with smoking and alcoholism.

Conclusions: Addictive behaviors in the workplace still a denied reality. Increasing awareness and clarifying expectations can be a good first step in order to ameliorate employee functioning and decrease productivity problems.

Keywords: addictive behaviors; onshore oil workers

EPP1329

Cannabis linked to improved sleep quality: A preliminary study

J. Cebrian^{1*} and G. Gonzalez-Cuevas²

¹Psychology, European University of Madrid, Madrid, Spain and

²Biomedical And Pharmaceutical Sciences, Idaho State University, Meridian, United States of America

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1505