

INDEX

	PAGE
Banaschewski, B. & Bhutani, K. R. Boolean algebras in a localic topos	43
Ben-Israel, A. Generalized inverses of matrices: a perspective of the work of Penrose	407
Bernau, S. J. Additivity and superadditivity in L_p -spaces	133
Bhutani, K. R. & Banaschewski, B. Boolean algebras in a localic topos	43
Brownawell, W. D. & Masser, D. W. Vanishing sums in function fields	427
Bruce, J. W. Envelopes and characteristics	475
Cassels, J. W. S. Mordell's finite basis theorem revisited	31
Chetwynd, A. G. & Hilton, A. J. W. Star multigraphs with three vertices of maximum degree	303
Cordier, J.-M. & Porter, T. Vogt's theorem on categories of homotopy coherent diagrams	65
Elias, J. On the analytic equivalence of curves	57
Evertse, J. H. & Silverman, J. H. Uniform bounds for the number of solutions to $Y^n = f(x)$	237
Falconer, K. J. Random fractals	559
Flapan, E. A prime strongly positive amphicheiral knot which is not slice	533
Friedlander, E. M. & Mislin, G. Locally finite approximation of Lie groups II	505
Gersten, S. M. & Howie, J. Counterexamples to a conjecture about spherical diagrams	539
Ghahramani, F. & Medgalchi, A. R. Compact multipliers on weighted hypergroup algebras II	145
Gibbon, G. Intersection properties of functions on cardinals	265
Godwin, H. J. Explicit expressions for Sturm sequences	225
Gopalsamy, K. On the global attractivity in a generalized delay-logistic differential equation	183
Hall, R. R. On some theorems of Hurwitz and Sheil-Small	365
Hildebrand, A. Multiplicative functions at consecutive integers	229
Hilton, A. J. W. & Chetwynd, A. G. Star multigraphs with three vertices of maximum degree	303
Howie, J. & Gersten, S. M. Counterexamples to a conjecture about spherical diagrams	539
Isbell, J. General function spaces, products and continuous lattices	193
Janezko, S. Generating families for images of Lagrangian submanifolds and open swallowtails	91
Janson, S. Random trees in a graph and trees in a random graph	319
Jefferies, B. Egorov measurability and generator cores	137
Johnson, F. E. A. A class of non-Kählerian manifolds	519
Jupp, P. E. Characterization of matrix probability distributions by mean residual lifetime	583
Kassem, S. E. Wave source potentials for two superposed fluids, each of finite depth	595
Katz, E., Morris, S. A. & Nickolas, P. Free subgroups of free abelian topological groups	347
Landweber, P. S. A survey of bordism and cobordism	207
Leinen, F. & Phillips, R. E. Existentially closed central extensions of locally finite p -groups	281
Lickorish, W. B. R. A relationship between link polynomials	109
Lines, D. On even-dimensional fibred knots obtained by plumbing	117
Lipson, A. S. An evaluation of a link polynomial	361
Maddox, J. J. Sequence spaces defined by a modulus	161

Masser, D. W. & Brownawell, W. D. Vanishing sums in function fields	427
Medgalchi, A. R. & Ghahramani, F. Compact multipliers on weighted hypergroup algebras II	145
Millett, K. & Rolfsen, D. A theorem of Borsuk–Ulam type for Seifert-fibred 3-manifolds	423
Mislin, G. & Friedlander, E. M. Locally finite approximation of Lie groups II	505
Morris, S. A., Katz, E. & Nickolas, P. Free subgroups of free abelian topological groups	347
Nestoridis, V. Averages of holomorphic mappings	371
Nickolas, P., Morris, S. A. & Katz, E. Free subgroups of free abelian topological groups	347
Palka, Z. Some remarks about extreme degrees in a random graph	167
Pascual-Gainz, A. P. On the simple object associated to a diagram in a closed model category	459
Patterson, S. J. Tori in metaplectic covers of GL_2 and applications to a formula of Loxton-Matthews	249
Phillips, R. E. & Leinen, F. Existentially closed central extensions of locally finite p -groups	281
Pinch, R. G. E. Elliptic curves with good reduction away from 2: II	435
Porter, D. The reduction of a pair of singular integral equations	175
Porter, T. & Cordier, J.-M. Vogt’s theorem on categories of homotopy coherent diagrams	65
Rankin, R. A. Fourier coefficients of cusp forms	5
Rhoades, B. E. Matrix summability of Fourier series based on inclusion theorems	545
Rolfsen, D. & Millett, K. A theorem of Borsuk–Ulam type for Seifert-fibred 3-manifolds	523
Silverman, J. H. & Evertse, J. H. Uniform bounds for the number of solutions to $Y^n = f(x)$	237
Sofi, M. A. Some criteria for nuclearity	151
Szücs, A. Multiple points of singular maps	331
Taylor, S. J. The measure theory of random fractals	383
Toland, J. F. A necessary geometric condition for the existence of certain homoclinic orbits	591
Trace, B. A note concerning Seifert manifolds for 2-knots	113
Wu, Y.-Q. On the Arf invariant of links	355
Yiu, P. Y. H. Quadratic forms between spheres and the non-existence of sums of squares formulae	493

THE PREPARATION OF MANUSCRIPTS

The attention of authors is particularly directed to the following requests.

1. Papers should be typed, double-spaced, on one side of white paper (of which A4, 210 by 297 mm, is a suitable size). The pages must be numbered. Margins of 30 mm should be left at the side, top and bottom of each page. Two clear copies should be sent.

A cover page should give the title, the author's name and institution, with the address at which mail is to be sent.

The title, while brief, must be informative (e.g. *A new proof of the prime-number theorem*, whereas *Some applications of a theorem of G. H. Hardy* would be useless).

The first paragraph or two should form a summary of the main theme of the paper, providing an abstract intelligible to mathematicians.

For a typescript to be accepted for publication, it must accord with the standard requirements of publishers, and be presented in a form in which the author's intentions regarding symbols etc. are clear to a printer (who is not a mathematician).

The following notes are intended to help the author in preparing the typescript. New authors may well enlist the help of senior colleagues, both as to the substance of their work and the details of setting it out correctly and attractively.

2. Notation

Notation should be chosen carefully so that mathematical operations are expressed with all possible neatness, to enlighten the task of the compositor and reduce the chance of error.

For instance n_k (n sub k) is common usage, but avoid if possible using c sub n sub k . Fractions are generally best expressed by a solidus. Complicated exponentials like

$$\exp \{z^2 \sin \theta / (1 + y^2)\}$$

should be shown in this and no other way.

In the manuscript, italics, small capitals and capitals are specified by single, double and triple underlinings. Bold-faced type is shown by wavy underlining; wavy will be printed wavy.

It helps if displayed equations or statements which will be quoted later are numbered in order on the right of their line. They can then be referred to by, for example, 'from (7)'

The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as o , O , o , O , 0 ; x , X , \times ; ϕ , Φ , \emptyset ; l , 1 ; ϵ , \in ; κ , k .

Greek letters can be denoted by Gk in the margin.

If an author wishes to mark the end of the proof of a theorem, the sign \square may be used.

Footnotes should be avoided.

3. Diagrams

It is extremely helpful if diagrams are drawn in Indian ink on white card, faintly blue or green-lined graph paper, or tracing cloth or paper. Symbols, legends and captions should be given on a transparent overlay. Each text figure must be numbered as Figure 1, Figure 2, ... and its intended position clearly indicated in the manuscript:

Figure 1 here

The author's name in pencil must be on all separate sheets of diagrams.

A figure is expensive to reproduce and should be included only when the subject matter demands it, or when it greatly clarifies the exposition.

The Society recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and it is therefore willing to have such diagrams re-drawn, provided that they are clear.

4. Tables

Tables should be numbered (above the table) and set out on separate sheets. Indicate the position of each in the text as for figures:

Table 3 here

5. References

References should be collected at the end of the paper numbered in alphabetical order of the author's names. Titles of journals should be abbreviated as in *Mathematical Reviews*. The following examples show the preferred style for references to a paper in a journal, a paper in a proceedings volume, a book and an unpublished dissertation:

- [1] J. F. ADAMS. On the non-existence of elements of Hopf invariant one. *Ann of Math.* (2) 72 (1960), 20–104.
- [2] M. P. FOURMAN and D. S. SCOTT. Sheaves and logic. In *Applications of Sheaves*, Lecture Notes in Math. vol. 753 (Springer-Verlag, 1979), pp. 302–401.
- [3] P. T. JOHNSTONE. *Stone Spaces*. Cambridge Studies in Advanced Math. no. 3 (Cambridge University Press, 1982).
- [4] F. W. LAWVERE. Functional semantics of algebraic theories. Ph.D. thesis, Columbia University (1963).

*Mathematical Proceedings of
the Cambridge Philosophical Society*

MPCPCO 100 (Pt 3) 383–605 (1986) 0305–0041 November 1986

CONTENTS

	PAGE
TAYLOR, S. JAMES. The measure theory of random fractals	383
BEN-ISRAEL, A. Generalized inverses of matrices: a perspective of the work of Penrose	407
BROWNAWELL, W. D. & MASSER, D. W. Vanishing sums in function fields	427
PINCH, R. G. E. Elliptic curves with good reduction away from 2: II	435
PASCUAL-GAINZA, PERE. On the simple object associated to a diagram in a closed model category	459
BRUCE, J. W. Envelopes and characteristics	475
YIU, PAUL Y. H. Quadratic forms between spheres and the non-existence of sums of squares formulae	493
FRIEDLANDER, ERIC M. & MISLIN, GUIDO. Locally finite approximation of Lie groups. III	505
JOHNSON, F. E. A. A class of non-Kählerian manifolds	519
MILLETT, KENNETH & ROLFSEN, DALE. A theorem of Borsuk–Ulam type for Seifert-fibred 3-manifolds	523
FLAPAN, ERICA. A prime strongly positive amphicheiral knot which is not slice	533
GERSTEN, S. M. & HOWIE, JAMES. Counterexamples to a conjecture about spherical diagrams	539
RHOADES, B. E. Matrix summability of Fourier series based on inclusion theorems	545
FALCONER, K. J. Random fractals	559
JUPP, P. E. Characterization of matrix probability distributions by mean residual lifetime	583
TOLAND, J. F. A necessary geometric condition for the existence of certain homoclinic orbits	591
KASSEM, S. E. Wave source potentials for two superposed fluids, each of finite depth	595
PROCEEDINGS 1985–86	601

© The Cambridge Philosophical Society 1986

CAMBRIDGE UNIVERSITY PRESS

THE PITT BUILDING, TRUMPINGTON STREET, CB2 1RP

32 EAST 57TH STREET, NEW YORK, NY 10022, USA

10 STAMFORD ROAD, OAKLEIGH, MELBOURNE 3166, AUSTRALIA

Price £19.00 net (USA and Canada US \$44.00)

*Subscription price £50.50 per volume (£101.00 per annum) net post free
(US \$118.50 per volume (US \$237.00 per annum) in USA and Canada)*

Printed in Great Britain by the University Press, Cambridge