



COMPOSITIO MATHEMATICA

Modular Koszul duality

Simon Riche, Wolfgang Soergel and Geordie Williamson

Compositio Math. **150** (2014), 273–332.

[doi:10.1112/S0010437X13007483](https://doi.org/10.1112/S0010437X13007483)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY



Modular Koszul duality

Simon Riche, Wolfgang Soergel and Geordie Williamson

ABSTRACT

We prove an analogue of Koszul duality for category \mathcal{O} of a reductive group G in positive characteristic ℓ larger than 1 plus the number of roots of G . However, there are no Koszul rings, and we do not prove an analogue of the Kazhdan–Lusztig conjectures in this context. The main technical result is the formality of the dg-algebra of extensions of parity sheaves on the flag variety if the characteristic of the coefficients is at least the number of roots of G plus 2.

1. Introduction

1.1 The scaffolding of Koszul duality

Given a \mathbb{Z} -graded ring E let us consider the abelian category $\text{Mod}^{\mathbb{Z}}\text{-}E$ of all \mathbb{Z} -graded right E -modules. There are two ways to ‘forget a part of the grading’ on its derived category. That is, there are two triangulated functors.

$$\mathcal{D}(\text{Mod}\text{-}E) \xleftarrow{v} \mathcal{D}(\text{Mod}^{\mathbb{Z}}\text{-}E) \xrightarrow{\bar{v}} \text{dgDer}\text{-}E.$$

On the left is simply the derived functor v of forgetting the grading $\text{Mod}^{\mathbb{Z}}\text{-}E \rightarrow \text{Mod}\text{-}E$. On the right is the derived category of the differential graded ring $(E, d=0)$, which is obtained as the localization of the homotopy category of differential graded right modules at quasi-isomorphisms (see for example [BL94] for a thorough discussion). The right-hand functor \bar{v} sends a complex of graded modules, thought of as a bigraded abelian group $(M^{i,j})$ (with $(-)\cdot a : M^{i,j} \rightarrow M^{i,j+|a|}$ for all $a \in E$ homogeneous of degree $|a|$ and differential $d : M^{i,j} \rightarrow M^{i+1,j}$) to the differential graded E -module $\bar{v}M$ with $(\bar{v}M)^n := \bigoplus_{i+j=n} M^{i,j}$ and obvious differential. It is this construction which provides the basic homological scaffolding of Koszul duality.

In order to obtain Koszul duality in the sense of [BGS96] we equip the above picture with some finiteness conditions. Let \mathbb{k} be a field and let E be a finite-dimensional \mathbb{Z} -graded \mathbb{k} -algebra of finite global dimension. Consider the categories $\text{Modf}\text{-}E$ (respectively $\text{Modf}^{\mathbb{Z}}\text{-}E$) of finite-dimensional (respectively finite-dimensional \mathbb{Z} -graded) right E -modules. Let $\text{dgDerf}\text{-}E \subset \text{dgDer}\text{-}E$ denote the full triangulated subcategory with objects finite-dimensional differential graded right E -modules (see § 5.4). In this setting Koszul duality for category \mathcal{O} means the existence of a finite-dimensional \mathbb{Z} -graded \mathbb{C} -algebra E as above together with vertical

Received 5 December 2012, accepted in final form 13 May 2013, published online 13 December 2013.

2010 Mathematics Subject Classification 14M15 (primary), 20G05, 16S37 (secondary).

Keywords: Koszul duality, category \mathcal{O} , formality, parity sheaves, perverse sheaves.

S.R. was supported by ANR Grants No. ANR-09-JCJC-0102-01 and No. ANR-10-BLAN-0110.

W.S. acknowledges the support of the DFG in the framework of SPP 1388.

This journal is © Foundation Compositio Mathematica 2013.

Moreover, for all $x \in W$ there exists a finite-dimensional graded E -module \widetilde{M}_x which specializes to the standard object M_x in $\mathcal{O}_0(\mathbb{F})$ on the left and to $i_{x*}\mathbb{F}_{BxB/B}[\ell(x)]$ on the right. In formulas,

$$M_x \leftarrow \widetilde{M}_x \mapsto i_{x*}\mathbb{F}_{BxB/B}[\ell(x)].$$

Let us emphasize, however, the difference to the non-modular case: in the modular case it is not clear whether the \mathbb{Z} -graded algebra E can be chosen such that it vanishes in negative degrees and is semi-simple in degree 0 (unless we make some strong assumptions, see §§ 5.7 and 7.3). Hence we are in no position to discuss the Koszulity of E in general.

We believe that the subquotient around the Steinberg weight has no special meaning in itself. However, this subquotient does see some of the multiplicities of simple modules in Weyl modules. One of the goals of this paper was to try to develop an accessible model which might help us approach Lusztig’s modular conjecture. (Indeed, Lusztig’s conjecture for multiplicities around the Steinberg weight is equivalent to the existence of a positive grading as discussed above.)

A large part of the proof of this modular Koszul duality can already be found in [Soe00]. More precisely, in [Soe00] the second author considered the full subcategory $\mathcal{K} \subset \mathcal{D}_{(B)}^b(G/B, \mathbb{F})$ consisting of all direct images of constant sheaves on Bott–Samelson varieties, together with their direct sums, summands and shifts. It was then shown that for all $y \in W$ there exists a unique indecomposable object \mathcal{E}_y in \mathcal{K} whose support is contained in $\overline{ByB/B}$ and whose restriction to ByB/B is isomorphic to $\mathbb{F}_{ByB/B}[\ell(y)]$. In modern language \mathcal{E}_y is a parity sheaf, and in [JMW09] their existence and uniqueness is shown by purely geometric arguments, without restriction on the characteristic. Finally, one of the main results of [Soe00] is an equivalence of categories

$$\text{Modf-}E \xrightarrow{\sim} \mathcal{O}_0(\mathbb{F})$$

where $E = \text{Ext}^\bullet(\mathcal{E}, \mathcal{E})$ is the algebra of self-extensions of $\mathcal{E} := \bigoplus_{y \in W} \mathcal{E}_y$. More precisely, E^i consists of all morphisms $\mathcal{E} \rightarrow \mathcal{E}[i]$ in the derived category $\mathcal{D}_{(B)}^b(G/B, \mathbb{F})$. Hence, all that remains in order to construct a diagram as in (1.2.2) is the construction of an equivalence of triangulated \mathbb{F} -categories

$$\text{dgDerf-}E \xrightarrow{\sim} \mathcal{D}_{(B)}^b(G/B, \mathbb{F}).$$

Remark 1.2.3. In [Fie11], Fiebig uses moment graphs, the combinatorics of [AJS94], and what in [FW10] was explained to be parity sheaves to develop an approach to the full Lusztig’s modular conjecture, which has led in particular to an explicit bound above which the conjecture holds [Fie12].

1.3 Some homological algebra

We briefly recall some general constructions of homological algebra, which allow us to establish such an equivalence. Let \mathcal{A} be an abelian category. We call a set of complexes \mathcal{C} ‘end-acyclic’ if, for all $T, T' \in \mathcal{C}$ and $n \in \mathbb{Z}$, the natural map gives an isomorphism

$$\text{Hom}_{\mathcal{K}(\mathcal{A})}(T, T'[n]) \xrightarrow{\sim} \text{Hom}_{\mathcal{D}(\mathcal{A})}(T, T'[n])$$

between morphisms in the homotopy and derived categories. Given a finite end-acyclic family T_1, \dots, T_m of complexes we can consider the complex $T = \bigoplus_i T_i$. Its endomorphism complex $E := \underline{\text{End}}_{\mathcal{A}}^\bullet(T)$ has a natural structure of a dg-ring with idempotents $1_i \in E$ given by the projection to each factor. Then the functor $\underline{\text{Hom}}_{\mathcal{A}}^\bullet(T, -)$ induces an equivalence of triangulated categories between the full triangulated subcategory

$$\langle T_1, \dots, T_m \rangle_{\Delta} \subset \mathcal{D}(\mathcal{A})$$

generated by the objects T_i and the full subcategory

$$\langle 1_1 E, \dots, 1_m E \rangle_{\Delta} \subset \text{dgDer} - E$$

generated by the right dg-modules $e_i E$. An important stepping stone in proving this claim is provided by the full triangulated subcategory

$$\langle T_1, \dots, T_m \rangle_{\Delta} \subset \mathcal{K}(A)$$

which by assumption is equivalent to the first triangulated subcategory above. It is on this category that the functor $\underline{\text{Hom}}_{\mathbf{A}}^{\bullet}(T, -)$ first really makes sense.

Furthermore, it is known that for any quasi-isomorphism $D \xrightarrow{\text{qis}} E$ of dg-rings (that is, a homomorphism of dg-rings which induces an isomorphism on cohomology) the restriction induces an equivalence of triangulated categories

$$\text{dgDer} - E \xrightarrow{\sim} \text{dgDer} - D.$$

Possible sources for statements of this type are [Kel94, Ric89].

1.4 Construction of the equivalence

We apply the above to the abelian category of all sheaves of \mathbb{F} -vector spaces. It is straightforward to see that the parity sheaves generate the triangulated subcategory of Bruhat constructible sheaves. In formulas,

$$\langle \mathcal{E}_y \mid y \in W \rangle_{\Delta} = \mathcal{D}_{(B)}^b(G/B, \mathbb{F}).$$

Let us choose a bounded below injective resolution \mathcal{E}_y^{\bullet} of each parity sheaf \mathcal{E}_y and set $\mathcal{E}^{\bullet} := \bigoplus_{y \in W} \mathcal{E}_y^{\bullet}$. If we consider the endomorphism dg-ring $E^{\bullet} := \underline{\text{End}}^{\bullet}(\mathcal{E}^{\bullet})$ together with the obvious idempotents 1_y then the constructions of § 1.3 give an equivalence of triangulated categories

$$\mathcal{D}_{(B)}^b(G/B, \mathbb{F}) \xrightarrow{\sim} \langle 1_y E^{\bullet} \mid y \in W \rangle_{\Delta} \subset \text{dgDer} - E^{\bullet}.$$

By definition the cohomology of the dg-ring E^{\bullet} is the graded ring $E = \text{Ext}^{\bullet}(\mathcal{E}, \mathcal{E})$ from above. Let e_y ($y \in W$) be the obvious idempotents in E . Suppose that we can find another dg-ring D and quasi-isomorphisms $E \xleftarrow{\text{qis}} D \xrightarrow{\text{qis}} E^{\bullet}$ together with homogeneous idempotents in D which are sent to the appropriate idempotents in E^{\bullet} and E . Then we obtain a further equivalence of derived categories

$$\mathcal{D}_{(B)}^b(G/B, \mathbb{F}) \xrightarrow{\sim} \langle e_y E \mid y \in W \rangle_{\Delta} \subset \text{dgDer} - E.$$

Hence, having found D together with these idempotents, all that remains is to compare finiteness conditions and deduce that the above can also be described using $\text{dgDerf} - E$.

In order to find our dg-ring D and the desired quasi-isomorphisms we adapt the techniques of [DGMS75] to the context of modular étale sheaves. Here, the Frobenius action plays the part that Hodge theory plays in [DGMS75]. In particular, we do not actually work on a complex flag variety, but rather on the flag variety over a finite field \mathbb{F}_p of characteristic different from ℓ . Also, in order to construct our dg-ring E^{\bullet} with compatible Frobenius action we instead work with complexes of perverse sheaves (and projective resolutions). These technical demands account in part for the length of this paper. We explain in Remark 7.1.4 how to deduce Theorem 1.2.1 from its étale counterpart.

1.5 Formality in characteristic zero

Except in special situations it is a difficult problem to establish whether or not a given dg-algebra is formal (i.e. quasi-isomorphic to its cohomology). However, there is a trick,

originally due to Deligne, which gives a general method of establishing formality: any bigraded dg-algebra $A = \bigoplus A^{i,j}$ with $d(A^{i,j}) \subset A^{i+1,j}$ and cohomology concentrated on the diagonal $\{i = j\}$ is formal. The proof is easy: it suffices to shear the bigrading by setting $A^{(i,j)} := A^{i-j,j}$ to arrive at a bigraded dg-ring with cohomology concentrated in the cohomological degree $i = 0$, which is well known to be formal.

In particular, this trick can be applied to establish the formality of the extension algebra of Bruhat constructible intersection cohomology sheaves with coefficients in \mathbb{C} on the flag variety, the idea being that if E denotes this extension algebra then E can be equipped with an additional ‘weight’ grading. The pointwise purity of intersection cohomology complexes means that this grading is diagonal. Hence, if the dg-algebra E^\bullet computing E can also be equipped with a ‘weight’ grading then the above trick can be used to conclude that E^\bullet is formal.

There are at least two settings where this approach can be carried out. The first one is mixed étale sheaves on the flag variety over the algebraic closure of a finite field \mathbb{F}_q , and the second one is mixed Hodge modules on the complex flag variety. In both settings a key role is played by the ‘realization’ equivalence

$$\mathcal{D}^b(\text{Perv}_{(B)}(G/B, \mathbb{C})) \xrightarrow{\sim} \mathcal{D}_{(B)}^b(G/B, \mathbb{C})$$

between $\mathcal{D}_{(B)}^b(G/B)$ and the corresponding bounded derived category of perverse sheaves. It is much easier to work in the abelian category of Bruhat constructible perverse sheaves than the abelian category of all sheaves on G/B . For example, $\text{Perv}_{(B)}(G/B, \mathbb{C})$ has enough projective objects and one even has an inductive algorithm for their construction.

In both settings $\text{Perv}_{(B)}(G/B, \mathbb{C})$ has an ‘enhancement’ (given by perverse sheaves on the flag variety over the finite field \mathbb{F}_q in the étale case, and by mixed Hodge modules in the complex case). Moreover, one can construct resolutions of the intersection cohomology complexes in this enhanced category, which become projective resolutions when one forgets the enhancement. It is the existence of these enhanced resolutions that allows one to equip the dg-algebra E^\bullet with an extra grading and deduce formality. (For a lucid explanation of this argument in the case of mixed Hodge modules the reader is referred to [Sch11].)

1.6 Formality in characteristic ℓ

In this paper, we adapt the above arguments to coefficients in a finite field \mathbb{F} of positive characteristic ℓ . In order to have a suitable enhancement we work on a flag variety defined over a finite field. (As we have already remarked, the case of the flag variety over \mathbb{C} can be deduced from this.) More precisely, fix a finite field \mathbb{F}_q of characteristic different from ℓ and let G_\circ denote a split semi-simple group over \mathbb{F}_q with Borel subgroup $B_\circ \subset G_\circ$ and flag variety G_\circ/B_\circ . We follow the convention of [BBD82]: a subscript ‘ \circ ’ denotes an object (variety or sheaf) defined over \mathbb{F}_q , and suppression of the subscript denotes the extension of scalars to the algebraic closure. Recall that our goal is to obtain an algebraic description of $\mathcal{D}_{(B)}^b(G/B, \mathbb{F})$ in terms of the extension algebra of parity sheaves.

Here the first natural question arises: in the formality statement, why do we have to replace intersection cohomology complexes by parity sheaves? With coefficients in \mathbb{F} there seems to be no good notion of purity, and intersection cohomology complexes can be badly behaved. (For example, their stalks may not satisfy parity vanishing as they do in characteristic 0.) Instead, one can consider indecomposable direct summands of ‘Bott–Samelson sheaves’: those sheaves obtained as direct images from Bott–Samelson resolutions. These are the parity sheaves and it




FIGURE 1. Picture of the ‘bigrading’ on E^\bullet (the stars indicate possible non-zero terms in $E = H^\bullet(E^\bullet)$).

is straightforward to see that their Ext-algebra satisfies a weak form of purity (Theorem 5.1.5): it vanishes in odd degree and all eigenvalues of Frobenius in degree $2n$ are equal to the image of q^{-n} in \mathbb{F}_ℓ .

Analogously to the case of coefficients of characteristic zero one can construct resolutions of Bott–Samelson sheaves in $\text{Perv}_{(B_\circ)}(G_\circ/B_\circ, \mathbb{F})$ which become projective resolutions when pulled back to G/B . Taking endomorphisms we obtain a dg-algebra E^\bullet of \mathbb{F} -vector spaces with compatible automorphism induced by the Frobenius. It turns out that all eigenvalues of Frobenius on E^\bullet belong to the subgroup $\Sigma \subset \mathbb{F}^\times$ generated by the image of q . Here we encounter a problem: with coefficients in characteristic zero the decomposition into generalized Frobenius eigenspaces gives a $\mathbb{Z} \times \mathbb{Z}$ -grading, whereas with coefficients in \mathbb{F}_ℓ we only obtain a $\mathbb{Z} \times \Sigma$ -grading. We have tried to depict the difference between these two situations in Figure 1. In order to apply the trick to deduce formality we would need to know, for example, that any fixed Frobenius eigenspace of E^\bullet (the columns in Figure 1) has cohomology concentrated in only one degree. It is possible to conclude in this way, but it leads to unpleasant bounds on ℓ in terms of the dimension of G/B .

1.7 Formality over \mathbb{O}

Instead, we fix a finite extension \mathbb{O} of the ℓ -adic integers \mathbb{Z}_ℓ with residue field \mathbb{F} and work with coefficients in \mathbb{O} . It turns out that all essential parts of the construction of E^\bullet can be lifted to \mathbb{O} : projective perverse sheaves over \mathbb{F} admit lifts to \mathbb{O} , morphism spaces between these sheaves are free \mathbb{O} -modules and the Bott–Samelson sheaves admit resolutions via perverse \mathbb{O} -sheaves on G_\circ/B_\circ which become projective resolutions when pulled back to G/B . Taking endomorphisms we obtain a dg-algebra $E_\mathbb{O}^\bullet$ which is free as an \mathbb{O} -module and has a compatible Frobenius automorphism ϕ , all of whose eigenvalues are integral powers of q . However, an \mathbb{O} -module with an endomorphism, even if it is free of finite rank over \mathbb{O} , need not decompose as the direct sum of its generalized eigenspaces (see §3.1). In order to conclude we need to prove that $E_\mathbb{O}^\bullet$ is ‘ q -decomposable’: that is, that $E_\mathbb{O}^\bullet$ is isomorphic to the direct sum of its generalized ϕ -eigenspaces with eigenvalues powers of q .

To explain the proof of the q -decomposability we need to explain in more detail how the resolutions of Bott–Samelson sheaves are constructed. We first obtain a purely algebraic description of the category of perverse \mathbb{O} -sheaves on G_{\circ}/B_{\circ} by constructing a perverse sheaf \mathcal{P}° on G_{\circ}/B_{\circ} such that its pullback \mathcal{P}° to G/B is a projective generator of $\text{Perv}_{(B)}(G/B, \mathbb{O})$. Hence, if we write $A = \text{End}(\mathcal{P}^{\circ})$ we have an equivalence

$$\text{Perv}_{(B)}(G/B, \mathbb{O}) \xrightarrow{\sim} \text{Modf-}A.$$

Moreover, because \mathcal{P}° is obtained by pullback from G_{\circ}/B_{\circ} , A is equipped with an automorphism ϕ and we have an equivalence

$$\text{Perv}_{(B_{\circ})}(G_{\circ}/B_{\circ}, \mathbb{O}) \xrightarrow{\sim} \text{Modf-}(A, \phi),$$

where $\text{Modf-}(A, \phi)$ denotes the category of pairs (M, ϕ_M) where M is a finitely generated right A -module and ϕ_M is an automorphism of M compatible with ϕ (see §3.1), and $\text{Perv}_{(B_{\circ})}(G_{\circ}/B_{\circ}, \mathbb{O})$ denotes the category of perverse sheaves on G_{\circ}/B_{\circ} whose pullback belongs to $\text{Perv}_{(B)}(G/B, \mathbb{O})$. Under these equivalences the pullback functor from G_{\circ}/B_{\circ} to G/B is given by forgetting the endomorphism ϕ_M of M . These considerations allow us to conclude that the realization functor

$$\mathcal{D}^b(\text{Perv}_{(B_{\circ})}(G_{\circ}/B_{\circ}, \mathbb{O})) \rightarrow \mathcal{D}^b_{(B_{\circ})}(G_{\circ}/B_{\circ}, \mathbb{O})$$

is an equivalence (see Proposition 3.4.1). We also prove similar results for partial flag varieties $G_{\circ}/P_{s,\circ}$, where $B_{\circ} \subset P_{s,\circ} \subset G_{\circ}$ is a minimal parabolic subgroup corresponding to a simple reflection $s \in W$: there exists an \mathbb{O} -algebra A^s with automorphism ϕ^s such that the analogues for G/P_s of the above results hold. Moreover, for any such s we have a morphism of algebras $A \rightarrow A^s$ and the direct and inverse image functors $\pi_{s!}$ and $\pi_s^!$ (where $\pi_s : G/B \rightarrow G/P_s$ denotes the projection) can be described algebraically in terms of derived induction and restriction along this morphism (Proposition 5.2.6).

Finally, the Bott–Samelson sheaves may be obtained by repeatedly applying the functors $\pi_s^! \pi_{s!}$ to the skyscraper sheaf on the base point $B/B \in G/B$. The above results allow us to translate the problem of constructing resolutions of the Bott–Samelson sheaves into a purely algebraic construction involving the algebras A and A^s . In particular, this allows us to conclude that the dg-algebra $E_{\mathbb{O}}^{\bullet}$ is q -decomposable if A and A^s are (Proposition 5.3.6). We prove that A and A^s are q -decomposable if the order of q in \mathbb{F}^{\times} is at least the number of roots plus one by examining the eigenvalues of Frobenius that may occur during the inductive construction of projective perverse sheaves. This in turn is a consequence of explicit bounds on the weights in the cohomology of an intersection of a Bruhat cell with an opposite Bruhat cell in G/B (see §§4.2–4.4).

1.8 Organization of the paper

This paper contains three parts.

Part I. The first two sections after the introduction are devoted to proving basic properties of the category of perverse sheaves with coefficients in a finite extension \mathbb{O} of \mathbb{Z}_{ℓ} on a sufficiently nice stratified variety. The corresponding statements for coefficients in a field are all well known.

Section 2. We consider projective perverse sheaves when our variety is defined over an algebraically closed field.

Section 3. We consider varieties defined over a finite field (with coefficients in \mathbb{O} or its residue field).

Part II. In the next two sections, we specialize our results to perverse sheaves on the flag variety of a reductive group.

Section 4. We prove results on the action of Frobenius on morphisms between standard objects, and derive consequences for the structure of projective perverse sheaves.

Section 5. We prove our formality theorem.

Part III. The last two sections are concerned with modular Koszul duality.

Section 6. We recall the results of [Soe00] concerning the modular category \mathcal{O} .

Section 7. We prove Theorem 1.2.1.

This article is the result of our efforts to join the two partially wrong and quite incomplete preprints [Soe11] and [RW11] into one readable article.

1.9 Notation

If D is a ring (respectively \mathbb{Z} -graded ring) we let $\text{Mod-}D$ (respectively $\text{Mod}^{\mathbb{Z}}-D$) be the category of right D -modules (respectively \mathbb{Z} -graded right D -modules). We also denote by $\text{Modf-}D$ (respectively $\text{Modf}^{\mathbb{Z}}-D$) the subcategory of finitely generated modules. We denote by $\text{Projf-}D$ the category of finitely generated projective right D -modules. We write $\text{Hom}_{-D}(-, -)$ for $\text{Hom}_{\text{Mod-}D}(-, -)$. We denote by $\langle 1 \rangle : \text{Mod}^{\mathbb{Z}}-D \rightarrow \text{Mod}^{\mathbb{Z}}-D$ the autoequivalence which sends a graded module $M = \bigoplus_i M^i$ to the graded module whose i th component is M^{i-1} . We denote by $\langle n \rangle$ the n th power of $\langle 1 \rangle$. With this convention, the functor \bar{v} satisfies

$$\bar{v}(M\langle 1 \rangle) = \bar{v}(M)[-1].$$

Similarly, we denote by $D\text{-Modf}$ the category of finitely generated left modules over the ring D .

If D is a dg-ring, we denote by $\text{dgHo-}D$ the homotopy category of right D -dg-modules, and by $\text{dgDer-}D$ the associated derived category.

In the whole paper, we fix a prime number ℓ . We also fix a finite extension \mathbb{K} of \mathbb{Q}_{ℓ} , and denote its ring of integers by \mathbb{O} . It is a finite extension of \mathbb{Z}_{ℓ} . We fix a uniformizer π , and set $\mathbb{F} = \mathbb{O}/(\pi)$; it is a finite field of characteristic ℓ . We denote by $\bar{\mathbb{K}}$ an algebraic closure of \mathbb{K} .

PART I. GENERALITIES ON \mathbb{O} -PERVERSE SHEAVES

2. Projective perverse sheaves

Let us fix an integer p which is either 0 or a prime number different from ℓ . In this section, X is a variety (i.e. a separated reduced scheme of finite type) over an algebraically closed field of characteristic p , endowed with a finite stratification by locally closed subvarieties

$$X = \bigsqcup_{s \in \mathcal{S}} X_s,$$

each isomorphic to an affine space. For each $s \in \mathcal{S}$ we denote by $i_s : X_s \hookrightarrow X$ the inclusion.

Note that the constructions in this section work similarly if we assume X is a complex algebraic variety, and work with the classical topology instead of the étale topology.

2.1 Definitions and first results

Consider the constructible derived categories $\mathcal{D}_c^b(X, \mathbb{O})$ and $\mathcal{D}_c^b(X, \mathbb{F})$, see e.g. [BBD82, § 2.2] or [Jut09, § 2] for definitions. For any $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$ let $\mathcal{L}_{s,m}$ denote the constant local system on X_s with stalk $\mathbb{O}/(\pi^m)$. (Here, by convention, $\mathbb{O}/(\pi^{\infty}) = \mathbb{O}$.) For simplicity we sometimes write \mathcal{L}_s

for $\mathcal{L}_{s,\infty}$. We write $\mathcal{L}_{s,\mathbb{F}}$ for the constant \mathbb{F} -local system of rank one on X_s . We assume that the following condition is satisfied:

$$\text{for all } s, t \in \mathcal{S} \text{ and } n \in \mathbb{Z}, \mathcal{H}^n(i_t^* i_{s*} \mathcal{L}_s) \text{ is constant.} \tag{2.1.1}$$

(If $p > 0$ then the strata X_s are not simply-connected unless X_s is a point, and so this condition is *a priori* stronger than requiring that $\mathcal{H}^n(i_t^* i_{s*} \mathcal{L}_s)$ be locally constant.) This implies the analogous condition for $\mathcal{L}_{s,m}$ and $\mathcal{L}_{s,\mathbb{F}}$. (See [Yun09, § 2.1] for a discussion of this condition; note that in our case an extension of constant sheaves on a stratum is constant.)

For $\mathbb{E} = \mathbb{O}$ or \mathbb{F} we denote by $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$ the full subcategory of $\mathcal{D}_c^b(X, \mathbb{E})$ consisting of objects \mathcal{F} such that for any $s \in \mathcal{S}$ and $n \in \mathbb{Z}$, $\mathcal{H}^n(i_s^* \mathcal{F})$ is a constant local system of finitely generated \mathbb{E} -modules. Our assumption (2.1.1) guarantees that if $i : Y \hookrightarrow X$ is a locally closed inclusion of a union of strata, the functors i_* , $i!$, i^* , $i^!$ restrict to functors between $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$ and $\mathcal{D}_{\mathcal{S}}^b(Y, \mathbb{E})$ (where for simplicity we write \mathcal{S} also for the restriction of the stratification \mathcal{S} to Y), so that $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$ can be endowed with the perverse t-structure whose heart we denote by

$$\text{Perv}_{\mathcal{S}}(X, \mathbb{E}) \subset \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E}).$$

Recall (see e.g. [BBD82, Jut09]) that if we take coefficients in \mathbb{O} we cannot speak of *the* subcategory of perverse sheaves on X with respect to this stratification. Instead, one has two ‘natural’ perverse t-structures p and p^+ with hearts $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ and $\text{Perv}_{\mathcal{S}}^+(X, \mathbb{O})$ respectively; Verdier duality exchanges these two hearts. (The reader unfamiliar with these facts is encouraged to think about the case of $X = \text{pt.}$) In this paper we will only ever need to consider the perversity p (except in Lemma 2.1.2). Note that the category $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is noetherian but not artinian.

We have a modular reduction functor

$$\mathbb{F} := \mathbb{F} \overset{L}{\otimes}_{\mathbb{O}} (-) : \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{O}) \rightarrow \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{F}).$$

By definition, an object \mathcal{M} of $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{O})$ is a collection $(\mathcal{M}_k)_{k \geq 1}$ (where each \mathcal{M}_k is an object of the derived category of étale sheaves of $\mathbb{O}/(\pi^k)$ -modules satisfying certain properties) together with isomorphisms $\mathcal{M}_{k+1} \overset{L}{\otimes}_{\mathbb{O}/(\pi^{k+1})} \mathbb{O}/(\pi^k) \cong \mathcal{M}_k$; then we have $\mathbb{F}(\mathcal{M}) = \mathcal{M}_1$. By definition, this functor commutes with all direct and inverse image functors, see e.g. [KW01, Appendix A] and references therein.

The functor \mathbb{F} does not preserve the subcategory of perverse sheaves (again, this is already false for a point). But it is right exact and if $\mathcal{F} \in \text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ then ${}^p\mathcal{H}^i(\mathbb{F}\mathcal{F}) = 0$ if $i \notin \{0, -1\}$.

Below we will need the following result.

LEMMA 2.1.2. *The functors i_{s*} and $i_{s!}$ are exact for the perverse t-structure, for coefficients \mathbb{O} or \mathbb{F} .*

*Proof.*¹ The morphism i_s is affine. Hence, in the case of coefficients \mathbb{F} the result is proved in [BBD82, Corollaire 4.1.3]. Let us consider now the case of coefficients \mathbb{O} . As Verdier duality exchanges the t-structures p and p^+ , it is sufficient to prove that the functor i_{s*} is exact for both t-structures p and p^+ . As the subcategories ${}^p\mathcal{D}^{\geq 0}$ and ${}^{p^+}\mathcal{D}^{\geq 0}$ are defined in terms of functors $i_i^!$, it is easy to check that this functor is left exact for both t-structures. The same arguments as for \mathbb{F} prove that i_{s*} is also right exact for the t-structure p . Finally, it is explained in [BBD82, § 3.3.4] that an \mathbb{O} -linear triangulated functor from $\mathcal{D}_{\mathcal{S}}^b(X_s)$ to $\mathcal{D}_{\mathcal{S}}^b(X)$ is right exact for the t-structure p if and only if it is for the t-structure p^+ , which finishes the proof. \square

¹ We thank Daniel Juteau and Carl Mautner for explaining this proof to us.

Remark 2.1.3. Similar arguments show more generally that if f is an affine morphism and for coefficients \mathbb{O} , f_* is right exact and $f_!$ is left exact for both t-structures p and p^+ . See [BBD82, Théorème 4.1.1, Corollaire 4.1.2] for a proof of the case of coefficients in a field (\mathbb{F} or \mathbb{K}).

If $Y \subset X$ is a locally closed union of strata, and if \mathbb{E} is \mathbb{O} or \mathbb{F} , we denote by

$$\mathrm{IC}(Y, -) : \mathrm{Perv}_{\mathcal{S}}(Y, \mathbb{E}) \rightarrow \mathrm{Perv}_{\mathcal{S}}(X, \mathbb{E})$$

the intermediate extension functor. This functor is fully faithful [Jut09, Proposition 2.29] and preserves injections and surjections [Jut09, Proposition 2.27], but is not exact.² If \mathcal{L} is a local system on Y we will sometimes abuse notation and write $\mathrm{IC}(Y, \mathcal{L})$ for $\mathrm{IC}(Y, \mathcal{L}[\dim Y])$.

LEMMA 2.1.4. *Any object in $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is a successive extension of $\mathrm{IC}(X_s, \mathcal{L}_{s,m})$ for some $s \in \mathcal{S}$ and $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$.*

Proof. We prove this result by induction on the number of strata, the case when X consists of a single stratum being obvious (since in this case the category $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is equivalent to $\mathrm{Mod}\text{-}\mathbb{O}$).

Let X_s be an open stratum, and let $Y = X \setminus X_s$ (a closed union of strata). Let $i : Y \hookrightarrow X$ denote the inclusion. Because the functor $i_* : \mathrm{Perv}_{\mathcal{S} \setminus \{s\}}(Y, \mathbb{O}) \rightarrow \mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is fully faithful we may assume by induction that the lemma is true for objects supported on Y . Now let \mathcal{F} be an arbitrary object of $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$. The morphism $\mathcal{F} \rightarrow i_*^p \mathcal{H}^0(i^* \mathcal{F})$ induced by adjunction is surjective [BBD82, Proposition 1.4.17(ii)] and hence we have an exact sequence of perverse sheaves

$$\mathrm{Ker} \hookrightarrow \mathcal{F} \twoheadrightarrow i_*^p \mathcal{H}^0(i^* \mathcal{F}).$$

Hence by induction it is enough to prove the result for Ker . One sees easily that $i_*^p \mathcal{H}^0(i^* \mathrm{Ker}) = 0$ and hence Ker has no non-zero quotient supported on Y . Now, we have a dual exact sequence (again by [BBD82, Proposition 1.4.17(ii)])

$$i_*^p \mathcal{H}^0(i^! \mathrm{Ker}) \hookrightarrow \mathrm{Ker} \twoheadrightarrow \mathrm{Coker}$$

and hence it is sufficient to prove the result for Coker . Similarly to above one sees that Coker has no non-trivial subobject or quotient supported on Y . By definition, $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is obtained by recollement from $\mathrm{Perv}_{\mathcal{S} \setminus \{s\}}(Y, \mathbb{O})$ and $\mathrm{Perv}_{\{s\}}(X_s, \mathbb{O})$, and hence we can apply [BBD82, Corollaire 1.4.25] to conclude that $\mathrm{Coker} \cong \mathrm{IC}(X_s, i_s^* \mathrm{Coker})$. By definition of $\mathrm{Perv}_{\{s\}}(X_s, \mathbb{O})$, $i_s^* \mathrm{Coker}$ is a constant local system, hence isomorphic to a direct sum of local systems of the form $\mathcal{L}_{s,m}$. The lemma now follows. \square

Remark 2.1.5. One can deduce from this lemma (using induction on the number of strata) that the category $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is generated (under extensions) by a finite number of objects, namely the collection of objects $\mathrm{IC}(X_s, \mathcal{L}_{s,1})$ and $\mathrm{IC}(X_s, \mathcal{L}_{s,\infty})$ for $s \in \mathcal{S}$.

Recall that an additive category \mathbf{A} satisfies the Krull–Schmidt property if every object in \mathbf{A} is isomorphic to a finite direct sum of indecomposable objects, and $\mathrm{End}_{\mathbf{A}}(M)$ is local for any indecomposable object M . In such a category, the decomposition as a direct sum of indecomposable objects is unique up to isomorphism and permutation of factors.

LEMMA 2.1.6. *The category $\mathrm{Perv}_{\mathcal{S}}(X, \mathbb{O})$ satisfies the Krull–Schmidt property.*

² In [Jut09], it is assumed on p. 1196 that the categories of perverse sheaves are noetherian and artinian; however this assumption is not used in the proofs of Propositions 2.27 and 2.29.

Proof. The endomorphism ring of any object of $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is a finitely generated \mathbb{O} -module, and hence is semi-perfect by [Lam01, Example 23.3]. Now it is easily checked (using e.g. [Lam01, Theorem 23.6]) that if \mathbf{A} is an abelian category in which every object has a semi-perfect endomorphism ring, then \mathbf{A} satisfies the Krull–Schmidt property. \square

In [BGS96, § 3.3] the authors study some properties of the category of perverse sheaves on a complex variety stratified by affine spaces, with complex coefficients. However their methods adapt directly to our category $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$. When necessary we will use these properties and give only the reference for the analogous result in [BGS96], without further details.

2.2 Standard and costandard objects over \mathbb{O} and \mathbb{F}

For any $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$, set

$$\Delta_{s,m} = i_{s!}\mathcal{L}_{s,m}[\dim X_s], \quad \nabla_{s,m} = i_{s*}\mathcal{L}_{s,m}[\dim X_s], \quad \text{IC}_{s,m} = \text{IC}(X_s, \mathcal{L}_{s,m}).$$

These objects are all in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ (see Lemma 2.1.2). We often abbreviate $\Delta_s = \Delta_{s,\infty}$, $\nabla_s = \nabla_{s,\infty}$, $\text{IC}_s = \text{IC}_{s,\infty}$.

Let $\mathcal{L}_{s,\mathbb{F}}$ denote the constant local system on s with stalk \mathbb{F} . For the analogous objects over \mathbb{F} , we use the notation

$$\Delta_{s,\mathbb{F}} = i_{s!}\mathcal{L}_{s,\mathbb{F}}[\dim X_s], \quad \nabla_{s,\mathbb{F}} = i_{s*}\mathcal{L}_{s,\mathbb{F}}[\dim X_s], \quad \text{IC}_{s,\mathbb{F}} = \text{IC}(X_s, \mathcal{L}_{s,\mathbb{F}})$$

in $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$. We have $\mathbb{F}(\Delta_s) \cong \Delta_{s,\mathbb{F}}$, $\mathbb{F}(\nabla_s) \cong \nabla_{s,\mathbb{F}}$.

LEMMA 2.2.1. *Let \mathcal{F} be in $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{O})$. If $\mathbb{F}(\mathcal{F}) = 0$, then $\mathcal{F} = 0$.*

Proof. If $\mathcal{F} \neq 0$, there exists $s \in \mathcal{S}$ such that $i_s^*\mathcal{F} \neq 0$. Then it is easy to check that $\mathbb{F}(i_s^*\mathcal{F}) = i_s^*(\mathbb{F}(\mathcal{F})) \neq 0$, hence $\mathbb{F}(\mathcal{F}) \neq 0$. \square

We will say that an object \mathcal{F} of $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ has a Δ -filtration (respectively a ∇ -filtration) if it admits a filtration in the abelian category $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ with subquotients of the form Δ_t (respectively ∇_t) for $t \in \mathcal{S}$. Note that here we only allow \mathbb{O} -free (co-)standard objects. Similarly, we will say that an object \mathcal{F} of $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$ has a $\Delta_{\mathbb{F}}$ -filtration (respectively a $\nabla_{\mathbb{F}}$ -filtration) if it admits a filtration in the abelian category $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$ with subquotients of the form $\Delta_{t,\mathbb{F}}$ (respectively $\nabla_{t,\mathbb{F}}$) for $t \in \mathcal{S}$.

LEMMA 2.2.2. *Let \mathcal{F} be in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$. Assume that $\mathbb{F}(\mathcal{F})$ is in $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$, and has a $\Delta_{\mathbb{F}}$ -filtration. Then \mathcal{F} has a Δ -filtration.*

Proof. We proceed by induction on the number of strata. We can assume that the support of M is X , and let X_s be an open stratum. Let $Y = X \setminus X_s$, and let $i : Y \hookrightarrow X$ be the (closed) inclusion.

Consider the restriction $i_s^*\mathcal{F}$. It is a shift of a constant local system on X_s . We have $\mathbb{F}(i_s^*\mathcal{F}) = i_s^*(\mathbb{F}(\mathcal{F}))$, hence by assumption $\mathbb{F}(i_s^*\mathcal{F})$ is also a shift (by the same integer) of a local system on X_s . This implies that $i_s^*\mathcal{F}$ has no torsion (otherwise $\mathbb{F}(i_s^*\mathcal{F})$ would be non-zero in degree $-\dim(X_s) - 1$), hence that $i_{s!}i_s^*\mathcal{F}$ is a direct sum of finitely many copies of Δ_s . Consider the exact triangle

$$i_{s!}i_s^*\mathcal{F} \rightarrow \mathcal{F} \rightarrow i_*i^*\mathcal{F} \xrightarrow{[1]} . \tag{2.2.3}$$

Its modular reduction is the similar triangle for $\mathbb{F}(\mathcal{F})$

$$i_{s!}i_s^*\mathbb{F}(\mathcal{F}) \rightarrow \mathbb{F}(\mathcal{F}) \rightarrow i_*i^*\mathbb{F}(\mathcal{F}) \xrightarrow{[1]} .$$

As $\mathbb{F}(\mathcal{F})$ has a filtration with subquotients of the form $\Delta_{t,\mathbb{F}}$, this triangle is an exact sequence of perverse sheaves. In particular, $\mathbb{F}(i_*i^*\mathcal{F})$ is a perverse sheaf.

We claim that $i_*i^*\mathcal{F}$ is a perverse sheaf. Indeed, using (2.2.3) it can have non-zero perverse cohomology sheaves only in degrees -1 and 0 . Consider the truncation triangle

$${}^p\mathcal{H}^{-1}(i_*i^*\mathcal{F})[1] \rightarrow i_*i^*\mathcal{F} \rightarrow {}^p\mathcal{H}^0(i_*i^*\mathcal{F}) \xrightarrow{[1]},$$

and its modular reduction

$$\mathbb{F}({}^p\mathcal{H}^{-1}(i_*i^*\mathcal{F}))[1] \rightarrow \mathbb{F}(i_*i^*\mathcal{F}) \rightarrow \mathbb{F}({}^p\mathcal{H}^0(i_*i^*\mathcal{F})) \xrightarrow{[1]}.$$

We have seen above that the middle term is a perverse sheaf, while the left-hand side (respectively the right-hand side) is concentrated in perverse degrees -2 and -1 (respectively -1 and 0). It easily follows that $\mathbb{F}({}^p\mathcal{H}^{-1}(i_*i^*\mathcal{F})) = 0$, hence ${}^p\mathcal{H}^{-1}(i_*i^*\mathcal{F}) = 0$ by Lemma 2.2.1, which proves the claim.

By this claim, triangle (2.2.3) is an exact sequence of perverse sheaves. Moreover, $i_*i^*\mathcal{F}$ again satisfies the conditions of the lemma. Hence, by induction, it has a filtration with subquotients of the form Δ_t . The result follows. \square

2.3 Projective objects: existence and some consequences

PROPOSITION 2.3.1. *For any $s \in \mathcal{S}$ there exists a projective object \mathcal{P} in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ which admits a Δ -filtration, and a surjection $\mathcal{P} \twoheadrightarrow \text{IC}_s$.*

Proof. We prove the proposition by induction on the number of strata. First, the result is clearly true when X consists of a single stratum.

Now, let X_s be an open stratum, and set $Y = X \setminus X_s$. We will abbreviate $\mathbb{D} = \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{O})$, $\mathbb{P} = \text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ and $\mathbb{P}_Y = \text{Perv}_{\mathcal{S}}(Y, \mathbb{O})$. Given $\mathcal{F} \in \mathbb{P}$, we have that $\mathcal{F} \in \mathbb{P}_Y$ if and only if $i_s^!\mathcal{F} (= i_s^*\mathcal{F}) = 0$. Also, note that if $\mathcal{F}, \mathcal{G} \in \mathbb{P}_Y$ then

$$\text{Ext}_{\mathbb{P}_Y}^1(\mathcal{F}, \mathcal{G}) = \text{Ext}_{\mathbb{P}}^1(\mathcal{F}, \mathcal{G}).$$

(If one thinks about both groups as classifying extensions, then any such extension has to be supported on Y .) Similarly,

$$\text{Ext}_{\mathbb{P}}^1(\mathcal{F}, \mathcal{G}) \cong \text{Ext}_{\mathbb{D}}^1(\mathcal{F}, \mathcal{G})$$

if $\mathcal{F}, \mathcal{G} \in \mathbb{P}$. We will use these equalities without further details below.

By adjunction, for any \mathcal{F} in \mathbb{P} we have

$$\text{Ext}_{\mathbb{P}}^1(\Delta_s, \mathcal{F}) = \text{Hom}_{\mathbb{D}}(\Delta_s, \mathcal{F}[1]) = \text{Hom}_{\mathcal{D}_{\{s\}}^b(X_s, \mathbb{O})}(\mathcal{L}_s, i_s^!\mathcal{F}[1]) = 0,$$

which implies that $\Delta_s \in \mathbb{P}$ is projective (and Δ -filtered). Hence we have found our first surjection $\Delta_s \twoheadrightarrow \text{IC}_s$.

To construct the other projectives we will adopt the following strategy (essentially copying [BGS96]): by induction we may assume that for each $t \in \mathcal{S} \setminus \{s\}$ we can find a surjection $\mathcal{P} \twoheadrightarrow \text{IC}_t$, with \mathcal{P} projective and Δ -filtered in \mathbb{P}_Y ; we then see what needs to be done to enlarge \mathcal{P} , so that it is projective in \mathbb{P} .

So, fix $t \in \mathcal{S} \setminus \{s\}$ and assume that \mathcal{P}_Y is a Δ -filtered projective object in \mathbb{P}_Y which surjects to IC_t . Certainly we have

$$\text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \mathcal{F}) = 0 \quad \text{for any } \mathcal{F} \in \mathbb{P}_Y. \tag{2.3.2}$$

Now, consider $E = \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \Delta_s)$ and let E_{free} be a finitely generated free \mathbb{O} -module endowed with a surjection to E . We denote by E_{free}^* the dual \mathbb{O} -module. The sequence of canonical

morphisms

$$\mathbb{O} \rightarrow E_{\text{free}}^* \otimes_{\mathbb{O}} E_{\text{free}} \rightarrow E_{\text{free}}^* \otimes_{\mathbb{O}} \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \Delta_s) \cong \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s)$$

gives us a distinguished extension

$$E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s \hookrightarrow \mathcal{P} \rightarrow \mathcal{P}_Y$$

in \mathbb{P} . Composing $\mathcal{P} \rightarrow \mathcal{P}_Y$ with $\mathcal{P}_Y \rightarrow \text{IC}_t$ yields a surjection of \mathcal{P} onto IC_t , and \mathcal{P} is Δ -filtered. We claim that \mathcal{P} is projective, which will conclude the proof. To prove this claim we need four preliminary steps.

Step 1. $\text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \mathcal{F}) = 0$ for $\mathcal{F} \in \mathcal{P}_Y$. This is clear from (2.3.2), the long exact sequence

$$\cdots \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \mathcal{F}) \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \mathcal{F}) \rightarrow \text{Ext}_{\mathbb{P}}^1(E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s, \mathcal{F}) \rightarrow \cdots$$

and the fact (already used above) that $\Delta_s \in \mathbb{P}$ is projective.

Step 2. $\text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \Delta_s) = 0$. Consider the long exact sequence

$$\begin{aligned} \cdots \rightarrow \text{Hom}_{\mathbb{P}}(E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s, \Delta_s) &\rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \Delta_s) \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \Delta_s) \\ &\rightarrow \text{Ext}_{\mathbb{P}}^1(E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s, \Delta_s) \rightarrow \cdots \end{aligned}$$

Because $\text{Ext}_{\mathbb{P}}^1(\Delta_s, \Delta_s) = 0$, it is enough to show that the first map above is surjective. However under the canonical isomorphisms

$$\text{Hom}_{\mathbb{P}}(E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_s, \Delta_s) \cong E_{\text{free}} \otimes_{\mathbb{O}} \text{Hom}_{\mathbb{P}}(\Delta_s, \Delta_s) \cong E_{\text{free}}$$

this map corresponds to the map $E_{\text{free}} \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}_Y, \Delta_s)$, which is surjective by construction.

Step 3. $\text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \mathcal{F}) = 0$ for all $\mathcal{F} \in \mathbb{P}$. By Lemma 2.1.4 it is enough to show that for all $t \in \mathcal{S}$, $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$ we have $\text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \text{IC}_{t,m}) = 0$. The short exact sequence

$$\text{IC}_{u,m} \hookrightarrow \nabla_{u,m} \rightarrow \text{Coker}$$

leads to a long exact sequence

$$\cdots \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \text{Coker}) \rightarrow \text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \text{IC}_{u,m}) \rightarrow \text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \nabla_{u,m}) \rightarrow \cdots$$

Now $\text{Coker} \in \mathcal{P}_Y$ and hence $\text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \text{Coker}) = 0$ by Step 1. Also, \mathcal{P} is Δ -filtered, and hence $\text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \nabla_{u,m}) = 0$. It follows that $\text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \text{IC}_{u,m}) = 0$ as claimed.

Step 4. $\text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \text{IC}_{s,m}) = 0$ for all m . The short exact sequence

$$\text{Ker} \hookrightarrow \Delta_s \rightarrow \text{IC}_{s,m}$$

leads to a long exact sequence:

$$\cdots \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \Delta_s) \rightarrow \text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \text{IC}_{s,m}) \rightarrow \text{Ext}_{\mathbb{D}}^2(\mathcal{P}, \text{Ker}) \rightarrow \cdots$$

Steps 2 and 3 show that the first and third terms are zero, which proves the claim.

Finally, we have shown in Steps 1 and 4 that $\text{Ext}_{\mathbb{P}}^1(\mathcal{P}, \text{IC}_{t,m}) = 0$ for all $t \in \mathcal{S}$ and $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. Hence \mathcal{P} is projective by Lemma 2.1.4. \square

COROLLARY 2.3.3. *There are enough projective objects in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$. More precisely, for any \mathcal{M} in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ there exists a projective object \mathcal{P} in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ which admits a Δ -flag and a surjection $\mathcal{P} \rightarrow \mathcal{M}$.*

Proof. By Lemma 2.1.4, it is enough to prove the result for $\mathcal{M} = \text{IC}_{s,m}$ ($s \in \mathcal{S}$, $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$). The case $m = \infty$ is treated in Proposition 2.3.1; the general case follows from the fact that the natural morphism $\text{IC}_{s,\infty} \rightarrow \text{IC}_{s,m}$ is surjective, since intermediate extension preserves surjections (see § 2.1). \square

COROLLARY 2.3.4. (i) *The realization functor*

$$\mathcal{D}^b\text{Perv}_{\mathcal{S}}(X, \mathbb{O}) \rightarrow \mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})$$

(see [Bei87]) *is an equivalence of categories.*

(ii) *The category $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ has finite projective dimension, i.e. there exists $N > 0$ such that every object of $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ admits a projective resolution of length at most N .*

Proof. (i) The idea of this proof is taken from [BGS96, Corollary 3.3.2]. As $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ generates the category $\mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})$, it is enough to prove that the realization functor is fully faithful, or equivalently that it induces an isomorphism

$$\text{Hom}_{\mathcal{D}^b\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}^i(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} \text{Hom}_{\mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})}^i(\mathcal{F}, \mathcal{G}) \tag{2.3.5}$$

for all $i \in \mathbb{Z}$ and all \mathcal{F}, \mathcal{G} in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$.

First, we claim that both sides in (2.3.5) vanish if \mathcal{F} is projective with a Δ -filtration and $i \geq 1$ (and for any \mathcal{G}). This claim is obvious by definition for the left-hand side. We prove the claim for the right-hand side by induction on i . If $i = 1$, then the morphism (2.3.5) is an isomorphism by [BBD82, Remarque 3.1.17], which proves the vanishing of the right-hand side. Now let $i \geq 2$, and assume the claim is known for $i - 1$. By Lemma 2.1.4 it suffices to treat the case $\mathcal{G} = \text{IC}_{s,m}$ for $s \in \mathcal{S}$ and $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. Consider the exact sequence

$$\text{IC}_{s,m} \hookrightarrow \nabla_{s,m} \rightarrow \text{Coker}.$$

Applying $\text{Hom}(\mathcal{F}, -)$ we obtain an exact sequence

$$\text{Hom}_{\mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})}^{i-1}(\mathcal{F}, \text{Coker}) \rightarrow \text{Hom}_{\mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})}^i(\mathcal{F}, \text{IC}_{s,m}) \rightarrow \text{Hom}_{\mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})}^i(\mathcal{F}, \nabla_{s,m}).$$

The left term vanishes by induction. The right term vanishes because \mathcal{F} has a Δ -filtration. Hence the middle term also vanishes, which finishes the proof of the claim.

Now we deduce that (2.3.5) is an isomorphism in general. First, both sides vanish if $i < 0$, and the realization functor induces an isomorphism for $i = 0$ by definition and for $i = 1$ by [BBD82, Remarque 3.1.17]. We treat the case $i \geq 2$ by induction. So, assume that $i \geq 2$ and that the result is known for $i - 1$. Let \mathcal{P} be a projective object with a Δ -filtration surjecting onto \mathcal{F} (see Corollary 2.3.3), and consider an exact sequence

$$\text{Ker} \hookrightarrow \mathcal{P} \rightarrow \mathcal{F}.$$

We have a commutative diagram with exact rows

$$\begin{array}{ccccccc} \text{Hom}_{\mathcal{D}^b\mathcal{P}}^{i-1}(\mathcal{P}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}^b\mathcal{P}}^{i-1}(\text{Ker}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}^b\mathcal{P}}^i(\mathcal{F}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}^b\mathcal{P}}^i(\mathcal{P}, \mathcal{G}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{Hom}_{\mathcal{D}}^{i-1}(\mathcal{P}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}}^{i-1}(\text{Ker}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}}^i(\mathcal{F}, \mathcal{G}) & \longrightarrow & \text{Hom}_{\mathcal{D}}^i(\mathcal{P}, \mathcal{G}) \end{array}$$

where $\mathcal{P} = \text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ and $\mathcal{D} = \mathcal{D}^b_{\mathcal{S}}(X, \mathbb{O})$. By our claim above, the terms in the right and the left column vanish. And by induction the second vertical arrow is an isomorphism. It follows that the third vertical arrow is also an isomorphism.

(ii) The idea for this proof is taken from [BGS96, Corollary 3.2.2]. By Remark 2.1.5 it is enough to prove that any object IC_s or $IC_{s,1}$ ($s \in \mathcal{S}$) has a finite projective resolution. Then, by induction on $\dim(X_s)$, it is enough to prove the same claim for objects Δ_s and $\Delta_{s,1}$ ($s \in \mathcal{S}$). There is an exact sequence

$$\Delta_s \hookrightarrow \Delta_s \twoheadrightarrow \Delta_{s,1}$$

since the functor $i_{s!}$ is exact (see Lemma 2.1.2). Hence we only have to prove the claim for objects Δ_s , $s \in \mathcal{S}$. Then we use a decreasing induction on $\dim(X_s)$. If this dimension is maximal, then Δ_s is projective. The induction step follows from the property that there exists a projective \mathcal{P}_s surjecting onto Δ_s such that the kernel has a Δ -filtration with subquotients Δ_t with $\dim(X_t) > \dim(X_s)$ (see the proof of Proposition 2.3.1). \square

Thanks to Corollary 2.3.4, we do not have to mention whether we consider Ext-groups in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ or in $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{O})$.

2.4 Projective objects: properties

Recall that the projective objects in the finite length category $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$ can be described using [BGS96, Theorem 3.2.1]. In particular, there are enough projective objects in this category, and any such object has a $\Delta_{\mathbb{F}}$ -filtration. Moreover, by [BGS96, Corollary 3.3.2], the realization functor

$$\mathcal{D}^b\text{Perv}_{\mathcal{S}}(X, \mathbb{F}) \rightarrow \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{F})$$

is an equivalence of categories. For $s \in \mathcal{S}$, we denote by $\mathcal{P}_{s,\mathbb{F}}$ the projective cover of $IC_{s,\mathbb{F}}$ in $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$.

PROPOSITION 2.4.1. *Let \mathcal{P} be a projective object in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$.*

- (i) *The object $\mathbb{F}(\mathcal{P})$ is a perverse sheaf, which is projective in $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$.*
- (ii) *For any $s \in \mathcal{S}$, the \mathbb{O} -module $\text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}, \Delta_s)$ is free, and the natural morphism*

$$\mathbb{F} \otimes_{\mathbb{O}} \text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}, \Delta_s) \rightarrow \text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{F})}(\mathbb{F}(\mathcal{P}), \Delta_{s,\mathbb{F}})$$

is an isomorphism.

- (iii) *The object \mathcal{P} admits a Δ -filtration.*
- (iv) *For any \mathcal{Q} projective in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$, the \mathbb{O} -module*

$$\text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}, \mathcal{Q})$$

is free. Moreover, the natural morphism

$$\mathbb{F} \otimes_{\mathbb{O}} \text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}, \mathcal{Q}) \rightarrow \text{Hom}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{F})}(\mathbb{F}(\mathcal{P}), \mathbb{F}(\mathcal{Q}))$$

is an isomorphism.

Proof. We prove (i) and (ii) simultaneously. By Corollary 2.3.3, there exists a projective object \mathcal{Q} of $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ which admits a Δ -filtration and a surjection $\mathcal{Q} \twoheadrightarrow \mathcal{P}$. By projectivity, \mathcal{P} is a direct factor of \mathcal{Q} . As \mathcal{Q} admits a Δ -filtration, $\mathbb{F}(\mathcal{Q})$ is a perverse sheaf; it follows that $\mathbb{F}(\mathcal{P})$ is also perverse.

Now we observe that we have

$$\mathbb{F}(R \text{Hom}(\mathcal{P}, \Delta_s)) \cong R \text{Hom}(\mathbb{F}(\mathcal{P}), \Delta_{s,\mathbb{F}}).$$

The left-hand side is concentrated in degrees -1 and 0 by Corollary 2.3.4(i), and the right-hand side is in non-negative degrees since $\mathbb{F}(\mathcal{P})$ is perverse. It follows that both complexes are

concentrated in degree 0, and (ii) follows. Then one easily deduces the projectivity of $\mathbb{F}(\mathcal{P})$ from the fact that $\text{Ext}^{>0}(\mathbb{F}(\mathcal{P}), \Delta_{s,\mathbb{F}}) = 0$ for any $s \in \mathcal{S}$.

By (i) and the remarks before the corollary, $\mathbb{F}(\mathcal{P})$ is a perverse sheaf and it admits a $\Delta_{\mathbb{F}}$ -filtration; (iii) follows using Lemma 2.2.2.

Finally, (iv) follows easily from (i), (ii), (iii) and the five lemma. □

COROLLARY 2.4.2. *For any $s \in \mathcal{S}$, there exists, up to isomorphism, a unique indecomposable projective object \mathcal{P}_s in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ such that $\mathbb{F}(\mathcal{P}_s) \cong \mathcal{P}_{s,\mathbb{F}}$. Moreover, these objects are pairwise non-isomorphic, and any projective object in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ is a direct sum of objects \mathcal{P}_s , $s \in \mathcal{S}$.*

Proof. By Proposition 2.3.1 and its proof, there exists an indecomposable projective \mathcal{P}_s surjecting onto Δ_s . By Proposition 2.4.1(i), $\mathbb{F}(\mathcal{P}_s)$ is a projective perverse sheaf. It also surjects onto $\Delta_{s,\mathbb{F}}$, and hence onto $\text{IC}_{s,\mathbb{F}}$. Next, recall that $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$ and $\text{Perv}_{\mathcal{S}}(X, \mathbb{F})$ both satisfy the Krull–Schmidt property (see Lemma 2.1.6 for \mathbb{O}), and hence an object is indecomposable if and only if its endomorphism ring is local. Then indecomposability of $\mathbb{F}(\mathcal{P}_s)$ follows from Proposition 2.4.1(iv), using the fact that one can lift idempotents from $\mathbb{F} \otimes_{\mathbb{O}} \text{End}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}_s)$ to $\text{End}_{\text{Perv}_{\mathcal{S}}(X, \mathbb{O})}(\mathcal{P}_s)$ (see [Fei82, Theorem 12.3]). As $\mathbb{F}(\mathcal{P}_s)$ surjects onto $\text{IC}_{s,\mathbb{F}}$, we deduce that $\mathbb{F}(\mathcal{P}_s) \cong \mathcal{P}_{s,\mathbb{F}}$.

Let us show uniqueness. Let \mathcal{Q}_s be another indecomposable projective such that $\mathbb{F}(\mathcal{Q}_s) \cong \mathcal{P}_{s,\mathbb{F}}$. By Proposition 2.4.1, the identity morphism of $\mathcal{P}_{s,\mathbb{F}}$ can be lifted to a morphism $f : \mathcal{P}_s \rightarrow \mathcal{Q}_s$. By Lemma 2.2.1, f is an isomorphism.

The fact that the projectives \mathcal{P}_s are pairwise non-isomorphic is obvious, since so are the projectives $\mathcal{P}_{s,\mathbb{F}}$. Finally, let \mathcal{P} be any projective object in $\text{Perv}_{\mathcal{S}}(X, \mathbb{O})$. Then $\mathbb{F}(\mathcal{P})$ is a projective perverse sheaf by Proposition 2.4.1(i), hence there exist integers n_s ($s \in \mathcal{S}$) and an isomorphism

$$\mathbb{F}(\mathcal{P}) \cong \bigoplus_{s \in \mathcal{S}} \mathcal{P}_{s,\mathbb{F}}^{\oplus n_s}.$$

By Proposition 2.4.1(iv) this isomorphism can be lifted to a morphism

$$\mathcal{P} \rightarrow \bigoplus_{s \in \mathcal{S}} \mathcal{P}_s^{\oplus n_s},$$

which is an isomorphism again by Lemma 2.2.1. □

3. Perverse sheaves on a variety over a finite field

3.1 Generalities on \mathbb{O} -modules with an endomorphism

Given an endomorphism ϕ of an \mathbb{O} -module M and $\lambda \in \mathbb{O}$ let us put

$$M_\lambda := \{m \in M \mid (\phi - \lambda \cdot \text{id})^n(m) = 0 \text{ for } n \gg 0\}.$$

We call M_λ the generalized eigenspace of M with eigenvalue λ .

PROPOSITION 3.1.1. *Let M be a free finite-rank \mathbb{O} -module with an endomorphism ϕ . Suppose that $\chi(\phi) = 0$ for some $\chi \in \mathbb{O}[X]$ such that $\chi = \prod_{\lambda \in \Lambda} (X - \lambda)^{n_\lambda}$ for some finite set $\Lambda \subset \mathbb{O}$ and some positive integers n_λ . Assume moreover that if $\lambda, \mu \in \Lambda$ have the same image in \mathbb{F} then $\lambda = \mu$. Then M decomposes into the direct sum of its generalized eigenspaces, i.e. the natural map*

$$\bigoplus_{\lambda \in \mathbb{O}} M_\lambda \rightarrow M \tag{3.1.2}$$

is an isomorphism.

Proof. First observe that $M_\lambda = \{0\}$ unless $\lambda \in \Lambda$, so that the sum in the left-hand side of (3.1.2) reduces to $\bigoplus_{\lambda \in \Lambda} M_\lambda$. Then, consider M as a module over $\mathbb{O}[X]/\langle \chi \rangle$. The natural morphism

$$P_\chi : \mathbb{O}[X]/\langle \chi \rangle \rightarrow \prod_{\lambda \in \Lambda} \mathbb{O}[X]/\langle (X - \lambda)^{n_\lambda} \rangle$$

is a morphism between two free \mathbb{O} -modules of the same finite rank, and its reduction modulo π is an isomorphism by our assumption on Λ and the Chinese Remainder Theorem. We deduce that P_χ is already an isomorphism. Pulling back the obvious idempotents from the right-hand side gives us idempotents $e_\lambda \in \mathbb{O}[X]/\langle \chi \rangle$ ($\lambda \in \Lambda$), and one can easily check that the assignment $m \mapsto (e_\lambda \cdot m)_{\lambda \in \Lambda}$ is an inverse to (3.1.2). \square

Given a free finite-rank \mathbb{O} -module M with an endomorphism ϕ let us say that M is decomposable under ϕ if the natural map is an isomorphism

$$\bigoplus_{\lambda \in \mathbb{O}} M_\lambda \xrightarrow{\sim} M.$$

(Using extension of scalars to \mathbb{K} , one can check that this map is always injective.)

LEMMA 3.1.3. *Let M be a free finite-rank \mathbb{O} -module which is decomposable under an endomorphism ϕ . Then given a ϕ -stable submodule $N \subset M$ such that M/N is free over \mathbb{O} as well, both N and M/N are decomposable under the induced endomorphisms.*

Proof. What we have to show is that if $m = \sum_\lambda m_\lambda$ is the generalized eigenspace decomposition of $m \in N$, then all m_λ already belong to N . Linear algebra tells us that all m_λ belong to $\mathbb{K} \otimes_{\mathbb{O}} N \subset \mathbb{K} \otimes_{\mathbb{O}} M$. It follows that for every λ there exists $n > 0$ such that $\pi^n m_\lambda \in N$. Therefore, the image of m_λ in M/N is killed by some power of π , and hence it is zero since M/N has no torsion, i.e. $m_\lambda \in N$. \square

Remark 3.1.4. (i) The assumption that M/N has no torsion cannot be removed. For example, assume that $\mathbb{O} = \mathbb{Z}_\ell$. Then $M = \mathbb{Z}_\ell^2$ is clearly decomposable under ϕ acting by the matrix

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 + \ell \end{pmatrix},$$

and its submodule $N := \mathbb{Z}_\ell(1, 1) \oplus \mathbb{Z}_\ell(0, \ell)$ is stable under ϕ . However, N is not decomposable under ϕ , as will be explained in (ii).

(ii) Observe that in the setting of the lemma, it can happen that both N and M/N are decomposable, but M is not. For example, consider $M = \mathbb{Z}_\ell^2$, with ϕ acting as

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 + \ell \end{pmatrix},$$

and $N = \mathbb{Z}_\ell(0, 1)$; M is not decomposable under ϕ , since on its reduction modulo π our ϕ is just a Jordan block. Note that this M is, as a \mathbb{Z}_ℓ -module with endomorphism, isomorphic to our N from (i).

Next let A be an \mathbb{O} -algebra which is finitely generated as an \mathbb{O} -module, endowed with an \mathbb{O} -algebra automorphism ϕ . We switch here to automorphisms as opposed to endomorphisms to avoid the introduction of additional terminology later. We denote by

$$\text{Modf}-(A, \phi)$$

the abelian category of pairs (M, ϕ_M) where M is a finitely generated right A -module and $\phi_M : M \xrightarrow{\sim} M$ is an automorphism which satisfies

$$\phi_M(m \cdot a) = \phi_M(m) \cdot \phi(a)$$

for $a \in A, m \in M$. Morphisms in $\mathbf{Modf}-(A, \phi)$ between objects (M, ϕ_M) and (N, ϕ_N) are morphisms of A -modules $f : M \rightarrow N$ such that $f \circ \phi_M = \phi_N \circ f$. Objects of $\mathbf{Modf}-(A, \phi)$ will sometimes be called (A, ϕ) -modules. By our assumptions on A , objects of $\mathbf{Modf}-(A, \phi)$ are always finitely generated as \mathbb{O} -modules. We will often omit the automorphism ϕ_M from notation. An important particular case is when $A = \mathbb{O}$ and $\phi = \text{id}$; in this case we call our objects (\mathbb{O}, ϕ) -modules.

LEMMA 3.1.5. *Let (A, ϕ) be an \mathbb{O} -algebra which is finitely generated as an \mathbb{O} -module, together with an automorphism ϕ . For any (M, ϕ_M) in $\mathbf{Modf}-(A, \phi)$, there exist an object (P, ϕ_P) in $\mathbf{Modf}-(A, \phi)$ such that P is A -free, and a surjection $(P, \phi_P) \twoheadrightarrow (M, \phi_M)$ in $\mathbf{Modf}-(A, \phi)$.*

Proof. First, consider (M, ϕ_M) as an (\mathbb{O}, ϕ) -module. Then there exist (N, ϕ_N) in $\mathbf{Modf}-(\mathbb{O}, \phi)$ such that N is \mathbb{O} -free, and a surjection $(N, \phi_N) \twoheadrightarrow (M, \phi_M)$. Indeed, let e_1, \dots, e_n be a family of vectors in M whose images in $\mathbb{F} \otimes_{\mathbb{O}} M$ form a basis. By Nakayama’s lemma, this family generates M as an \mathbb{O} -module. Choose a matrix $D \in \mathcal{M}_n(\mathbb{O})$ such that $\phi_M(e_i) = \sum_j a_{ji} e_j$; this matrix is automatically invertible since its image in $\mathcal{M}_n(\mathbb{F})$ is invertible. Then one can take $N = \mathbb{O}^n$, with ϕ_N acting with matrix D , and the natural surjection $N \twoheadrightarrow M$. Finally, set $P := N \otimes_{\mathbb{O}} A$, endowed with the automorphism $\phi_P := \phi_N \otimes \phi$. □

Now let us fix a unit $q \in \mathbb{O}^\times$ which is not a root of unity. Suppose M is an object in $\mathbf{Modf}-(\mathbb{O}, \phi)$, i.e. a finitely generated \mathbb{O} -module with an automorphism ϕ . Given some finite subset $I \subset \mathbb{Z}$, we say that M has q -weights obtained from I , if

$$\prod_{i \in I} (\phi - q^i \cdot \text{id})$$

acts nilpotently on M . In addition, let us call M q -decomposable if it is decomposable under ϕ and has q -weights obtained from some finite subset of \mathbb{Z} . Observe that if we have an exact sequence

$$N \hookrightarrow M \twoheadrightarrow P$$

and if N , respectively P , has q -weights obtained from I , respectively J , then M has q -weights obtained from $I \cup J$. Similarly, if M has q -weights obtained from I and N has q -weights obtained from J , then $(M \otimes_{\mathbb{O}} N, \phi_M \otimes \phi_N)$ has q -weights obtained from $I + J = \{i + j \mid i \in I, j \in J\}$.

Remark 3.1.6. If (M, ϕ) is free of finite rank over \mathbb{O} and has q -weights obtained from I , and if moreover the map $i \mapsto \bar{q}^i$ is an injection $I \hookrightarrow \mathbb{F}$ (where \bar{q} is the image of q in \mathbb{F}), then M is q -decomposable by Proposition 3.1.1. This criterion will be crucial for our proofs.

LEMMA 3.1.7. *Let M be in $\mathbf{Modf}-(\mathbb{O}, \phi)$, and assume M has q -weights obtained from I . Then there exists an object M' in $\mathbf{Modf}-(\mathbb{O}, \phi)$, which is \mathbb{O} -free and has q -weights obtained from I , together with a surjection $M' \twoheadrightarrow M$.*

Proof. We can consider M as a finitely generated module over the \mathbb{O} -free algebra

$$R := \mathbb{O}[x] / \left\langle \prod_{i \in I} (x - q^i \text{id})^n \right\rangle$$

for some positive integer n . Then take M' to be a free finite-rank R -module surjecting to M , with ϕ acting as multiplication by x . (Observe that multiplication by x is an automorphism since its reduction modulo π is invertible by our assumption $q \in \mathbb{O}^\times$.) \square

Now let (A, ϕ) be an \mathbb{O} -algebra, which is a free finite-rank \mathbb{O} -module, endowed with an \mathbb{O} -algebra automorphism ϕ , such that A is q -decomposable. We denote by

$$\mathbf{Modf}_{\text{dec}}-(A, \phi)$$

the full subcategory of $\mathbf{Modf}-(A, \phi)$ whose objects are free of finite rank over \mathbb{O} and q -decomposable as (\mathbb{O}, ϕ) -modules.

LEMMA 3.1.8. *Assume that (A, ϕ) is \mathbb{O} -free of finite rank and q -decomposable as above. For any M in $\mathbf{Modf}_{\text{dec}}-(A, \phi)$ there exists an object Q in $\mathbf{Modf}_{\text{dec}}-(A, \phi)$ which is projective as an A -module along with a surjection $Q \rightarrow M$.*

Proof. Take $Q := M \otimes_{\mathbb{O}} A$, with the automorphism $\phi_M \otimes \phi$. \square

3.2 Categories with an autoequivalence

Let \mathbf{A} be an abelian category endowed with an autoequivalence $\Phi : \mathbf{A} \xrightarrow{\sim} \mathbf{A}$. Let $\mathbf{A}[\Phi]$ be the category defined as follows:

- objects are pairs (M, s_M) where M is an object of \mathbf{A} and $s_M : M \xrightarrow{\sim} \Phi(M)$ is an isomorphism;
- morphisms between (M, s_M) and (N, s_N) are morphisms $f : M \rightarrow N$ in \mathbf{A} such that the following diagram commutes.

$$\begin{array}{ccc} M & \xrightarrow{s_M} & \Phi M \\ f \downarrow & & \downarrow \Phi f \\ N & \xrightarrow{s_N} & \Phi N \end{array}$$

We might think of \mathbf{A} as an abelian category with an action of the group \mathbb{Z} and of $\mathbf{A}[\phi]$ as the category of some sort of \mathbb{Z} -equivariant objects of \mathbf{A} . Certainly $\mathbf{A}[\Phi]$ is again abelian.

Example 3.2.1. With the notation of § 3.1, the category $\mathbf{Modf}-(A, \phi)$ is the category $\mathbf{A}[\Phi]$ where $\mathbf{A} = \mathbf{Modf}-A$ and Φ sends M to the right A -module, whose underlying \mathbb{O} -module is M , and whose A -action is the morphism $M \otimes_{\mathbb{O}} A \rightarrow M$ given by $m \otimes a \mapsto m \cdot \phi(a)$.

LEMMA 3.2.2. *Let (\mathbf{A}, Φ) be an abelian category with an autoequivalence, and let $(P, t) \in \mathbf{A}[\Phi]$ be an object with $P \in \mathbf{A}$ projective. Then for any $(M, s) \in \mathbf{A}[\Phi]$ we have*

$$\text{Hom}_{\mathcal{D}^b(\mathbf{A}[\Phi])}^i((P, t), (M, s)) = 0 \quad \text{for } i \geq 2.$$

Proof. Let $i \geq 2$. By definition, a morphism from (P, t) to $(M, s)[i]$ in $\mathcal{D}^b(\mathbf{A}[\Phi])$ is a diagram of morphisms of complexes

$$(M, s)[i] \leftarrow C^\bullet \xrightarrow{\text{qis}} (P, t)$$

where C^\bullet is a bounded complex of objects of $\mathbf{A}[\Phi]$. Clearly, one can assume that this complex is concentrated in non-positive degrees. To prove the lemma it suffices to prove that there exist a complex D^\bullet and a quasi-isomorphism $f : D^\bullet \xrightarrow{\text{qis}} C^\bullet$ whose composition with our morphism $C^\bullet \rightarrow (M, s)[i]$ is zero. To establish this it is enough to show that there exists a complex D^\bullet concentrated in degrees -1 and 0 and a quasi-isomorphism $f : D^\bullet \xrightarrow{\text{qis}} C^\bullet$.

Write the complex C^\bullet as

$$C^\bullet = (\dots \rightarrow (C^{-2}, u^{-2}) \xrightarrow{d^{-2}} (C^{-1}, u^{-1}) \xrightarrow{d^{-1}} (C^0, u^0) \rightarrow 0 \rightarrow \dots).$$

Let $B^{-1} := C^{-1}/\ker(d^{-1})$, and let $p^{-1} : C^{-1} \rightarrow B^{-1}$ be the projection. We consider the object (B^{-1}, v^{-1}) in $\mathbf{A}[\Phi]$, where $v^{-1} : B^{-1} \xrightarrow{\sim} \Phi(B^{-1})$ is the isomorphism induced by u^{-1} . As C^\bullet maps quasi-isomorphically to (P, t) , we have an exact sequence

$$(B^{-1}, v^{-1}) \hookrightarrow (C^0, u^0) \twoheadrightarrow (P, t).$$

As P is projective, the image of this exact sequence in \mathbf{A} splits, hence we have an isomorphism $(C^0, u^0) \cong (B^{-1} \oplus P, w^0)$ in $\mathbf{A}[\Phi]$, where

$$w^0 = \begin{pmatrix} v^{-1} & a \\ 0 & t \end{pmatrix} : B^{-1} \oplus P \xrightarrow{\sim} \Phi(B^{-1}) \oplus \Phi(P)$$

for some morphism $a : P \rightarrow \Phi(B^{-1})$ in \mathbf{A} . Let us fix such an isomorphism. As P is projective, there exists a morphism $b : P \rightarrow \Phi(C^{-1})$ in \mathbf{A} whose composition with the projection $\Phi(p^{-1}) : \Phi(C^{-1}) \rightarrow \Phi(B^{-1})$ is a . Then define $D^0 := C^{-1} \oplus P$, and consider the object (D^0, x^0) of $\mathbf{A}[\Phi]$, where

$$x^0 := \begin{pmatrix} u^{-1} & b \\ 0 & t \end{pmatrix} : C^{-1} \oplus P \xrightarrow{\sim} \Phi(C^{-1}) \oplus \Phi(P).$$

Also, let $(D^{-1}, x^{-1}) := (C^{-1}, u^{-1})$. Then (D^0, x^0) is an object of $\mathbf{A}[\Phi]$, and there exists a natural injection $(D^{-1}, x^{-1}) \hookrightarrow (D^0, x^0)$ whose cokernel is (P, t) . Moreover, we have a quasi-isomorphism

$$\begin{array}{ccccccc} \dots & \longrightarrow & 0 & \longrightarrow & (D^{-1}, x^{-1}) & \longrightarrow & (D^0, x^0) \longrightarrow 0 \longrightarrow \dots \\ & & \downarrow & & \parallel & & \downarrow c \\ \dots & \longrightarrow & (C^{-2}, u^{-2}) & \longrightarrow & (C^{-1}, u^{-1}) & \longrightarrow & (C^0, u^0) \longrightarrow 0 \longrightarrow \dots \end{array}$$

where

$$c = \begin{pmatrix} p^{-1} & 0 \\ 0 & \text{id}_P \end{pmatrix} : C^{-1} \oplus P \rightarrow B^{-1} \oplus P.$$

This finishes the proof. □

3.3 Perverse sheaves on a variety over a finite field

In this subsection we write \mathbb{E} for either \mathbb{O} or \mathbb{F} .

Let X_\circ be a variety over a finite field \mathbb{F}_q , endowed with a finite stratification

$$X_\circ = \bigsqcup_{s \in \mathcal{S}} X_{s,\circ}$$

by affine spaces. Also, let $X := X_\circ \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{\mathbb{F}}_q)$, endowed with the induced stratification (which we also denote by \mathcal{S}). We assume that this stratification satisfies condition (2.1.1). Let

$$\text{For} : \mathcal{D}_c^b(X_\circ, \mathbb{E}) \rightarrow \mathcal{D}_c^b(X, \mathbb{E})$$

be the inverse image functor under the natural projection $X \rightarrow X_\circ$. We denote by $\mathcal{D}_{\mathcal{S}}^b(X_\circ, \mathbb{E})$ the full subcategory of $\mathcal{D}_c^b(X_\circ, \mathbb{E})$ consisting of objects \mathcal{F} such that $\text{For}(\mathcal{F})$ is in $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$. As for X , $\mathcal{D}_{\mathcal{S}}^b(X_\circ, \mathbb{E})$ is endowed with the perverse t-structure whose heart we denote by $\text{Perv}_{\mathcal{S}}(X_\circ, \mathbb{E})$ (for the perversity p if $\mathbb{E} = \mathbb{O}$). Note that if \mathcal{M}_\circ is in $\mathcal{D}_{\mathcal{S}}^b(X_\circ, \mathbb{E})$ then \mathcal{M}_\circ is in $\text{Perv}_{\mathcal{S}}(X_\circ, \mathbb{E})$ if and only if $\text{For}(\mathcal{M}_\circ)$ is in $\text{Perv}_{\mathcal{S}}(X, \mathbb{E})$. As in [BBD82], objects of $\text{Perv}_{\mathcal{S}}(X_\circ, \mathbb{E})$ will always be

denoted with a subscript ‘ \circ ’. Their image under the functor For will usually be denoted by the same symbol, with the subscript removed.

Let $\text{pt}_\circ := \text{Spec } \mathbb{F}_q$ and $f : X_\circ \rightarrow \text{pt}_\circ$ denote the projection. Let $\mathbb{E}(m) \in \mathcal{D}_c^b(\text{pt}_\circ, \mathbb{E})$ denote the m th Tate sheaf (see [Del74, § 2.13]). Given any object $\mathcal{F}_\circ \in \mathcal{D}_c^b(X_\circ, \mathbb{E})$ we define its m th Tate twist by $\mathcal{F}_\circ(m) := \mathcal{F}_\circ \otimes_{\mathbb{E}}^L f^* \mathbb{E}(m)$.

Before we go into the technical details, let us discuss some generalities concerning Galois actions on étale cohomology. Given a scheme X_\circ over a field k and a field extension K/k , certainly the Galois group will act on the scheme $X = K \times_k X_\circ$, and thus will also act on its étale cohomology. More generally, given an étale sheaf \mathcal{F}_\circ on X_\circ and its pullback \mathcal{F} to X , the Galois group will act on the étale cohomology of X with coefficients in \mathcal{F} . Formally, if $\omega : X \rightarrow X_\circ$ is the extension of scalars, for any γ in the Galois group and letting $\gamma = \gamma \times \text{id}$, we have $\omega \circ \gamma = \omega$ and from there get isomorphisms $\gamma^* \omega^* \mathcal{F}_\circ \xrightarrow{\sim} \omega^* \mathcal{F}_\circ$ alias $\gamma^* \mathcal{F} \xrightarrow{\sim} \mathcal{F}$, which make \mathcal{F} what one might call a Galois equivariant sheaf on X . The action of γ on $H^n(X; \mathcal{F})$ is then defined as the composition $H^n(X; \mathcal{F}) \rightarrow H^n(X; \gamma^* \mathcal{F}) \rightarrow H^n(X; \mathcal{F})$ of the pullback by γ on cohomology with the map induced by our isomorphism $\gamma^* \mathcal{F} \xrightarrow{\sim} \mathcal{F}$. Even more generally, if \mathcal{G}_\circ is another étale sheaf or complex of sheaves on X_\circ , we will get a Galois action on the Ext-groups between \mathcal{F} and \mathcal{G} by transport of structure, and we could go on like that.

We will use these Galois actions in the case that the field extension K/k consists of our finite field \mathbb{F}_q with an algebraic closure $\overline{\mathbb{F}}_q$. In this case, additional features try to create confusion. Namely, given a prime p , on any commutative ring A over $\mathbb{Z}/p\mathbb{Z}$, even on any scheme S over $\mathbb{Z}/p\mathbb{Z}$ there is a canonical endomorphism, the absolute Frobenius $\text{Fr}_{\text{ab}} : S \rightarrow S$, given on rings as the map $\text{Fr}_{\text{ab}} : A \rightarrow A$ with $a \mapsto a^p$. This is a natural transformation from the identity functor on the category of schemes over $\mathbb{Z}/p\mathbb{Z}$ to itself. In particular, if we have two schemes S, T over $\mathbb{Z}/p\mathbb{Z}$, on their product we have $\text{Fr}_{\text{ab}} = (\text{Fr}_{\text{ab}} \times \text{id}) \circ (\text{id} \times \text{Fr}_{\text{ab}})$. From the remarkable fact that for any étale morphism $\pi : T \rightarrow S$ the diagram

$$\begin{array}{ccc} T & \xrightarrow{\text{Fr}_{\text{ab}}} & T \\ \pi \downarrow & & \downarrow \pi \\ S & \xrightarrow{\text{Fr}_{\text{ab}}} & S \end{array}$$

is Cartesian, we get for any étale sheaf \mathcal{F} on S a natural isomorphism $\text{Fr}_{\text{ab}}^* \mathcal{F} \xrightarrow{\sim} \mathcal{F}$, and the map induced by this on the étale cohomology of \mathcal{F} is just the identity.

Now, in our case of a scheme X_\circ over \mathbb{F}_q with $q = p^r$, on the scheme $X = \overline{\mathbb{F}}_q \times_{\mathbb{F}_q} X_\circ$ we have as endomorphisms the arithmetic Frobenius $\text{Fr}_a = \text{Fr}_{\text{ab}}^r \times \text{id}$ and the geometric Frobenius $\text{Fr} = \text{Fr}_g = \text{id} \times \text{Fr}_{\text{ab}}^r = \text{id} \times \text{Fr}_\circ$. Clearly $\text{Fr}_g \circ \text{Fr}_a = \text{Fr}_a \circ \text{Fr}_g = \text{Fr}_{\text{ab}}^r$, thus our natural isomorphism leads to a natural isomorphism $\text{Fr}_a^*(\text{Fr}^* \mathcal{F}) \xrightarrow{\sim} \mathcal{F}$ for any étale sheaf \mathcal{F} on X . On the other hand, it also leads to a natural isomorphism $\mathcal{F}_\circ \xrightarrow{\sim} \text{Fr}_\circ^* \mathcal{F}_\circ$ and by pulling back we obtain a natural isomorphism $\mathcal{F} \xrightarrow{\sim} \text{Fr}^* \mathcal{F}$ for any étale sheaf \mathcal{F}_\circ on X_\circ . To get back from this isomorphism our Galois equivariance $\gamma^* \mathcal{F} \xrightarrow{\sim} \mathcal{F}$ alias $\text{Fr}_a^* \mathcal{F} \xrightarrow{\sim} \mathcal{F}$ from above, just apply Fr_a^* to it and compose on the left with the natural isomorphism $\text{Fr}_a^*(\text{Fr}^* \mathcal{F}) \xrightarrow{\sim} \mathcal{F}$ discussed before. This is explained in [SGA4½, p. 84]. In the following, we always work directly with the Frobenius action coming from the natural isomorphism $\mathcal{F} \xrightarrow{\sim} \text{Fr}^* \mathcal{F}$, which might be conceptually a less direct approach, but permits us to connect much more directly to the existing literature.

So let $\text{Fr} : X \rightarrow X$ be the Frobenius endomorphism, obtained from the Frobenius $\text{Fr}_\circ : X_\circ \rightarrow X_\circ$ by base change. For any \mathcal{M}_\circ in $\mathcal{D}_{\mathcal{S}}^b(X_\circ, \mathbb{E})$ there exists a canonical isomorphism $\mathcal{M}_\circ \xrightarrow{\sim} \text{Fr}_\circ^* \mathcal{M}_\circ$, hence also a canonical isomorphism $\psi_{\mathcal{M}} : \mathcal{M} \xrightarrow{\sim} \text{Fr}^* \mathcal{M}$, where $\mathcal{M} = \text{For}(\mathcal{M}_\circ)$.

For details, see [SGA5, Exposé XIV, § 2.1] or [Mil80, § VI.13]. We will need the following well-known result.

LEMMA 3.3.1. *The functor*

$$\text{Fr}^* : \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E}) \rightarrow \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$$

is an equivalence of categories, which restricts to an equivalence $\text{Perv}_{\mathcal{S}}(X, \mathbb{E}) \xrightarrow{\sim} \text{Perv}_{\mathcal{S}}(X, \mathbb{E})$.

Proof. The category $\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$ is generated by the essential image of the functor

$$\text{For} : \mathcal{D}_{\mathcal{S}}^b(X_{\circ}, \mathbb{E}) \rightarrow \mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})$$

(e.g. by standard or costandard objects). Hence, it is enough to prove that for any $\mathcal{M}_{\circ}, \mathcal{N}_{\circ}$ in $\mathcal{D}_{\mathcal{S}}^b(X_{\circ}, \mathbb{E})$, with $\mathcal{M} := \text{For}(\mathcal{M}_{\circ}), \mathcal{N} := \text{For}(\mathcal{N}_{\circ})$, the morphism

$$\text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\mathcal{M}, \mathcal{N}) \rightarrow \text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\text{Fr}^* \mathcal{M}, \text{Fr}^* \mathcal{N})$$

is an isomorphism. To prove this fact it is enough to prove that the adjunction morphism $\mathcal{N} \rightarrow \text{Fr}_* \text{Fr}^* \mathcal{N}$ is an isomorphism. However, this morphism is obtained by extension of scalars from the adjunction morphism $\mathcal{N}_{\circ} \rightarrow \text{Fr}_{\circ*} \text{Fr}_{\circ}^* \mathcal{N}_{\circ}$, which is an isomorphism, since Fr_{\circ} is a universal homeomorphism, see [SGA5, Exposé XIV, § 1, Proposition 2(a)]. \square

Recall that, by [BBD82, Proposition 5.1.2], the functor which sends \mathcal{M}_{\circ} to the pair $(\mathcal{M}, \psi_{\mathcal{M}})$, where $\mathcal{M} = \text{For}(\mathcal{M}_{\circ})$ and $\psi_{\mathcal{M}}$ is defined above, induces an equivalence of abelian categories

$$\text{Perv}_{\mathcal{S}}(X_{\circ}, \mathbb{E}) \xrightarrow{\sim} \text{Perv}_{\mathcal{S}}(X, \mathbb{E})[\text{Fr}^*]. \tag{3.3.2}$$

Using this equivalence, one can make sense of the tensor product $V \otimes_{\mathbb{E}} \mathcal{M}_{\circ}$ of an object \mathcal{M}_{\circ} of $\text{Perv}_{\mathcal{S}}(X_{\circ}, \mathbb{E})$ with an \mathbb{E} -free (\mathbb{E}, ϕ) -module (V, ϕ_V) . Indeed, using (3.3.2) one can see (V, ϕ_V) as an object of $\text{Perv}(\text{pt}_{\circ}, \mathbb{E})$, and we define $V \otimes_{\mathbb{E}} \mathcal{M}_{\circ}$ as the tensor product of the pullback of this object to X_{\circ} with \mathcal{M}_{\circ} .

Example 3.3.3. When $X_{\circ} = \text{pt}_{\circ} = \text{Spec}(\mathbb{F}_q)$, we have $\text{Fr} = \text{id}$, and $\text{Perv}(\text{pt}, \mathbb{E}) \cong \text{Modf-}\mathbb{E}$. The image of $\mathbb{E}(m) \in \text{Perv}(\text{pt}_{\circ}, \mathbb{E})$ under (3.3.2) is a rank 1 free right \mathbb{E} -module endowed with the automorphism given by multiplication by q^m .

LEMMA 3.3.4. *There exists an object \mathcal{P}_{\circ} in $\text{Perv}_{\mathcal{S}}(X_{\circ}, \mathbb{E})$ such that $\mathcal{P} = \text{For}(\mathcal{P}_{\circ})$ is a projective generator of $\text{Perv}_{\mathcal{S}}(X, \mathbb{E})$.*

Proof. We write the details only for $\mathbb{E} = \mathbb{O}$; the case $\mathbb{E} = \mathbb{F}$ is similar. By the proof of Corollary 2.3.3, it is enough to prove that the objects \mathcal{P} constructed in Proposition 2.3.1 can be chosen in such a way that they are in the essential image of the functor For or, equivalently, using equivalence (3.3.2), in such a way that there exists an isomorphism $\mathcal{P} \xrightarrow{\sim} \text{Fr}^* \mathcal{P}$. Now, using the notation of this proof, the \mathbb{O} -module E has a natural automorphism ϕ induced by Fr^* ; it is enough to observe that E_{free} can be endowed with an automorphism ϕ_{free} such that the quotient $E_{\text{free}} \rightarrow E$ is compatible with ϕ and ϕ_{free} , which follows from Lemma 3.1.5 (for $A = \mathbb{O}$). (See the proof of Proposition 4.4.1 below for details on this construction.) \square

Let $\mathcal{M}_{\circ}, \mathcal{N}_{\circ}$ be in $\mathcal{D}_{\mathcal{S}}^b(X_{\circ}, \mathbb{E})$, with $\mathcal{M} := \text{For}(\mathcal{M}_{\circ}), \mathcal{N} := \text{For}(\mathcal{N}_{\circ})$. Composing the isomorphism

$$\begin{cases} \text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\mathcal{M}, \mathcal{N}) \rightarrow \text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\text{Fr}^* \mathcal{M}, \text{Fr}^* \mathcal{N}) \\ f \mapsto \psi_{\mathcal{N}} \circ f \circ \psi_{\mathcal{M}}^{-1} \end{cases}$$

with the inverse of the isomorphism

$$\text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\mathcal{M}, \mathcal{N}) \rightarrow \text{Hom}_{\mathcal{D}_{\mathcal{S}}^b(X, \mathbb{E})}(\text{Fr}^* \mathcal{M}, \text{Fr}^* \mathcal{N})$$

induced by Fr^* (see Lemma 3.3.1) we obtain an automorphism $\phi_{\mathcal{M}, \mathcal{N}}$ of $\text{Hom}_{\mathcal{D}_{\mathcal{F}}^b(X, \mathbb{E})}(\mathcal{M}, \mathcal{N})$. Hence, $\text{Hom}_{\mathcal{D}_{\mathcal{F}}^b(X, \mathbb{E})}(\mathcal{M}, \mathcal{N})$ is naturally an object of $\text{Modf}-(\mathbb{E}, \phi)$, which we will denote by $\text{Hom}_{X, \mathbb{E}}(\mathcal{M}, \mathcal{N})$ for simplicity.

Remark 3.3.5. Using the same arguments as in Lemma 3.3.1, for any variety X_{\circ} over $\text{Spec}(\mathbb{F}_q)$ (not necessarily satisfying our assumptions) we obtain an automorphism (induced by the Frobenius morphism) of the \mathbb{E} -module $\mathbf{H}_{\circ}^{\bullet}(X; \mathbb{E}) = \text{Hom}^{\bullet}(\mathbb{E}_{\text{pt}}, f_! \mathbb{E}_X)$. (Here, $f : X_{\circ} \rightarrow \text{pt}_{\circ}$ is the projection.) For example, with our conventions the automorphism of $\mathbf{H}_{\circ}^2(\mathbb{A}^1; \mathbb{E}) \cong \mathbf{H}_{\circ}^2(\mathbb{A}^1 \setminus \{0\}; \mathbb{E})$ is multiplication by q^{-1} , and the automorphism of $\mathbf{H}_{\circ}^1(\mathbb{A}^1 \setminus \{0\}; \mathbb{E})$ is the identity.

Let us fix an object \mathcal{P}_{\circ} as in Lemma 3.3.4, and let

$$A := \text{End}_{\text{Perv}_{\mathcal{F}}(X, \mathbb{E})}(\mathcal{P}_{\circ}).$$

Since \mathcal{P} is a projective generator of $\text{Perv}_{\mathcal{F}}(X, \mathbb{E})$ we have an equivalence of categories

$$\mathbf{F}_{\mathcal{P}} := \text{Hom}(\mathcal{P}, -) : \text{Perv}_{\mathcal{F}}(X, \mathbb{E}) \xrightarrow{\sim} \text{Modf}-A \tag{3.3.6}$$

(see [Bas68, Exercise on p. 55]). Moreover, as $\mathcal{P} = \text{For}(\mathcal{P}_{\circ})$, we have an automorphism $\phi := \phi_{\mathcal{P}, \mathcal{P}}$ of A , so that we can consider the category $\text{Modf}-(A, \phi)$ as in § 3.1.

LEMMA 3.3.7. *The functor which sends \mathcal{M}_{\circ} to $(\mathbf{F}_{\mathcal{P}}(\mathcal{M}), \phi_{\mathcal{P}, \mathcal{M}})$ (where as usual $\mathcal{M} = \text{For}(\mathcal{M}_{\circ})$) induces an equivalence of abelian categories*

$$\text{Perv}_{\mathcal{F}}(X_{\circ}, \mathbb{E}) \xrightarrow{\sim} \text{Modf}-(A, \phi). \tag{3.3.8}$$

Moreover, the diagram

$$\begin{array}{ccc} \text{Perv}_{\mathcal{F}}(X_{\circ}, \mathbb{E}) & \xrightarrow[\sim]{(3.3.8)} & \text{Modf}-(A, \phi) \\ \text{For} \downarrow & & \downarrow \\ \text{Perv}_{\mathcal{F}}(X, \mathbb{E}) & \xrightarrow[\sim]{(3.3.6)} & \text{Modf}-A \end{array}$$

commutes, where the right vertical functor sends (M, ϕ_M) to M .

Proof. Consider the equivalences

$$\text{Perv}_{\mathcal{F}}(X_{\circ}, \mathbb{E}) \stackrel{(3.3.2)}{\cong} \text{Perv}_{\mathcal{F}}(X, \mathbb{E})[\text{Fr}^*] \cong (\text{Modf}-A)[\mathbf{F}_{\mathcal{P}} \circ \text{Fr}^* \circ (\mathbf{F}_{\mathcal{P}})^{-1}]$$

where the second equivalence is induced by $\mathbf{F}_{\mathcal{P}}$. We claim that the functors Φ (defined as in Example 3.2.1) and $\mathbf{F}_{\mathcal{P}} \circ \text{Fr}^* \circ (\mathbf{F}_{\mathcal{P}})^{-1}$ are isomorphic, or equivalently that the functors $\Phi \circ \mathbf{F}_{\mathcal{P}}$ and $\mathbf{F}_{\mathcal{P}} \circ \text{Fr}^*$ are isomorphic; indeed, an isomorphism is provided by the composition of isomorphisms

$$\Phi \circ \mathbf{F}_{\mathcal{P}}(\mathcal{M}) = \text{Hom}(\mathcal{P}, \mathcal{M}) \xrightarrow{\text{Fr}^*} \text{Hom}(\text{Fr}^* \mathcal{P}, \text{Fr}^* \mathcal{M}) \xrightarrow{(-) \circ \psi_{\mathcal{P}}} \text{Hom}(\mathcal{P}, \text{Fr}^* \mathcal{M}) = \mathbf{F}_{\mathcal{P}} \circ \text{Fr}^*(\mathcal{M})$$

for \mathcal{M} in $\text{Perv}_{\mathcal{F}}(X, \mathbb{E})$. Equivalence (3.3.8) follows. We leave it to the reader to check that this equivalence can be described by the formula in the lemma, and that the diagram commutes. \square

3.4 Derived categories of perverse sheaves

In this subsection we keep the same assumptions and notation as in § 3.3. The subcategories $\text{Perv}_{\mathcal{F}}(X_{\circ}, \mathbb{E}) \subset \mathcal{D}_{\mathcal{F}}^b(X_{\circ}, \mathbb{E})$, $\text{Perv}_{\mathcal{F}}(X, \mathbb{E}) \subset \mathcal{D}_{\mathcal{F}}^b(X, \mathbb{E})$ are hearts of t-structures, and the triangulated categories $\mathcal{D}_{\mathcal{F}}^b(X_{\circ}, \mathbb{E})$ and $\mathcal{D}_{\mathcal{F}}^b(X, \mathbb{E})$ have natural filtered analogues. Hence, by [Bei87] one has realization functors

$$\text{real}_{\circ} : \mathcal{D}^b \text{Perv}_{\mathcal{F}}(X_{\circ}, \mathbb{E}) \rightarrow \mathcal{D}_{\mathcal{F}}^b(X_{\circ}, \mathbb{E}), \quad \text{real} : \mathcal{D}^b \text{Perv}_{\mathcal{F}}(X, \mathbb{E}) \rightarrow \mathcal{D}_{\mathcal{F}}^b(X, \mathbb{E}).$$

Moreover, these functors fit in the commutative diagram

$$\begin{CD} \mathcal{D}^b\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E}) @>\text{real}_o>> \mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E}) \\ @VVV @VVV \\ \mathcal{D}^b\text{Perv}_{\mathcal{F}}(X, \mathbb{E}) @>\text{real}>> \mathcal{D}^b_{\mathcal{F}}(X, \mathbb{E}) \end{CD}$$

where vertical functors are induced by For (see [Bei87, Lemma A.7.1]). It is known that the functor real is an equivalence of categories, see Corollary 2.3.4 if $\mathbb{E} = \mathbb{O}$ and [BGS96, Corollary 3.3.2].

PROPOSITION 3.4.1. *The functor real_o is an equivalence of categories.*

Proof. The category $\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E})$ generates $\mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E})$. Hence, it is enough to prove that for any $\mathcal{M}_o, \mathcal{N}_o$ in $\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E})$ and any $i \in \mathbb{Z}$ the natural morphism

$$\text{Hom}^i_{\mathcal{D}^b\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E})}(\mathcal{M}_o, \mathcal{N}_o) \rightarrow \text{Hom}^i_{\mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E})}(\mathcal{M}_o, \mathcal{N}_o)$$

is an isomorphism. We prove this fact by induction on i . In fact, both sides vanish if $i < 0$; if $i = 0$ then the morphism is an isomorphism by definition; and the morphism is an isomorphism also if $i = 1$ by [BBD82, Remarque 3.1.7(ii)].

Now, assume that $i \geq 2$, and that the result is known for $i - 1$ (and any $\mathcal{M}_o, \mathcal{N}_o$). Let \mathcal{P}_o be an object of $\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E})$ surjecting onto \mathcal{M}_o and such that $\text{For}(\mathcal{P}_o)$ is projective. (Such an object exists by Lemmas 3.3.7 and 3.1.5, or an obvious analogue if $\mathbb{E} = \mathbb{F}$.) Consider an exact sequence

$$\ker_o \hookrightarrow \mathcal{P}_o \twoheadrightarrow \mathcal{M}_o,$$

and the following associated commutative diagram with exact rows.

$$\begin{CD} \text{Hom}^{i-1}_{\mathcal{D}^b\mathcal{P}_o}(\mathcal{P}_o, \mathcal{N}_o) @>>> \text{Hom}^{i-1}_{\mathcal{D}^b\mathcal{P}_o}(\ker_o, \mathcal{N}_o) @>>> \text{Hom}^i_{\mathcal{D}^b\mathcal{P}_o}(\mathcal{M}_o, \mathcal{N}_o) @>>> \text{Hom}^i_{\mathcal{D}^b\mathcal{P}_o}(\mathcal{P}_o, \mathcal{N}_o) \\ @VVV @VVV @VVV @VVV \\ \text{Hom}^{i-1}_{\mathcal{D}^b_o}(\mathcal{P}_o, \mathcal{N}_o) @>>> \text{Hom}^{i-1}_{\mathcal{D}^b_o}(\ker_o, \mathcal{N}_o) @>>> \text{Hom}^i_{\mathcal{D}^b_o}(\mathcal{M}_o, \mathcal{N}_o) @>>> \text{Hom}^i_{\mathcal{D}^b_o}(\mathcal{P}_o, \mathcal{N}_o) \end{CD}$$

Here, $\mathcal{P}_o := \text{Perv}_{\mathcal{F}}(X_o, \mathbb{E})$ and $\mathcal{D}^b_o := \mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E})$. The first two vertical morphisms in this diagram are isomorphisms by induction. We claim that both \mathbb{E} -modules in the right column vanish, which will conclude the proof (by the five lemma). Indeed, the top right term is 0 by Lemmas 3.3.7 and 3.2.2 (since $i \geq 2$). The lower right term is also 0 by the exact sequences [BBD82, (5.1.2.5)] and the fact that

$$\text{Hom}^k_{\mathcal{D}^b_{\mathcal{F}}(X, \mathbb{E})}(\text{For}(\mathcal{P}_o), \text{For}(\mathcal{N}_o)) = 0$$

for $k \geq 1$. (The latter vanishing follows from the facts that real is an equivalence and that $\text{For}(\mathcal{P}_o)$ is a projective perverse sheaf.) \square

Remark 3.4.2. Let Φ be the equivalence considered in Example 3.2.1. We denote similarly the equivalence induced on derived categories, so that every object M_o in $\mathcal{D}^b\text{Modf}(A, \phi)$, with image M in $\mathcal{D}^b\text{Modf}A$, comes equipped with an isomorphism $\phi_M : M \xrightarrow{\sim} \Phi(M)$. If M_o, N_o are in $\mathcal{D}^b\text{Modf}(A, \phi)$ then there is a canonical automorphism of $\text{Hom}_{\mathcal{D}^b\text{Modf}A}(M, N)$ which sends a morphism f to $\Phi^{-1}(\phi_N \circ f \circ \phi_M^{-1})$. One can check that if $\mathcal{M}_o, \mathcal{N}_o$ are in $\mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E})$ and if M_o, N_o are their images under the equivalence

$$\mathcal{D}^b_{\mathcal{F}}(X_o, \mathbb{E}) \xrightarrow{\text{real}_o^{-1}} \mathcal{D}^b\text{Perv}_{\mathcal{F}}(X_o, \mathbb{E}) \xrightarrow{(3.3.8)} \mathcal{D}^b\text{Modf}(A, \phi),$$

then this automorphism of $\text{Hom}(M, N)$ coincides with the automorphism $\phi_{\mathcal{M}, \mathcal{N}}$ of $\text{Hom}(\mathcal{M}, \mathcal{N})$ defined in §3.3 under the isomorphism $\text{Hom}(\mathcal{M}, \mathcal{N}) \cong \text{Hom}(M, N)$ induced by the equivalence $F_{\mathcal{P}} \circ \text{real}^{-1}$.

PART II. FORMALITY OF THE CONSTRUCTIBLE DERIVED CATEGORY OF THE FLAG VARIETY

4. Bounding weights

As in §3.3, we use \mathbb{E} to denote either \mathbb{F} or \mathbb{O} . We fix a finite field \mathbb{F}_q (where q is a prime power which is invertible in \mathbb{F}) and an algebraic closure $\overline{\mathbb{F}}_q$ of \mathbb{F}_q .

4.1 Notation

We let G_{\circ} be a split connected reductive algebraic group over \mathbb{F}_q , $B_{\circ} \subset G_{\circ}$ be a Borel subgroup, and $T_{\circ} \subset B_{\circ}$ be a maximal torus. We let W be the Weyl group of G_{\circ} with respect to T_{\circ} . The main player in the rest of this paper will be the flag variety

$$X_{\circ} := G_{\circ}/B_{\circ},$$

endowed with the Bruhat stratification

$$X_{\circ} = \bigsqcup_{w \in W} X_{w, \circ}, \quad X_{w, \circ} := B_{\circ}wB_{\circ}/B_{\circ}.$$

For any subset $I \subset W$, we denote by $X_{I, \circ}$ the union of the $X_{v, \circ}$ for $v \in I$. Following the notation in the previous sections, we denote by $i_w : X_{w, \circ} \hookrightarrow X_{\circ}$ the inclusion, and we omit the index ‘ \circ ’ when considering the varieties obtained by extension of scalars from \mathbb{F}_q to $\overline{\mathbb{F}}_q$. The category of \mathbb{E} -perverse sheaves on X_{\circ} (respectively X) for the Bruhat stratification will be denoted by $\text{Perv}_{(B_{\circ})}(X_{\circ}, \mathbb{E})$ (respectively $\text{Perv}_{(B)}(X, \mathbb{E})$), and similarly for derived categories. We use similar notation for B_{\circ} -stable subvarieties of X_{\circ} .

Let $B_{\circ}^{-} \subset G_{\circ}$ be the Borel subgroup opposite to B_{\circ} with respect to T_{\circ} . We will also consider the opposite Bruhat cells

$$X_{w, \circ}^{-} := B_{\circ}^{-}wB_{\circ}/B_{\circ} \subset X_{\circ}$$

($w \in W$).

The choice of $B_{\circ} \subset G_{\circ}$ determines a subset $S \subset W$ of simple reflections and of positive roots, which we choose such that the Lie algebra of the unipotent radical of B_{\circ} is spanned by positive root spaces. If $s \in S$, we denote by $P_{s, \circ}$ the associated minimal standard parabolic subgroup. We let $X_{\circ}^s := G_{\circ}/P_{s, \circ}$. This variety has a natural stratification by Bruhat cells,

$$X_{\circ}^s = \bigsqcup_{w \in W^s} X_{w, \circ}^s, \quad X_{w, \circ}^s := B_{\circ}wP_{s, \circ}/P_{s, \circ},$$

where $W^s := \{w \in W \mid ws > w\}$. As for X_{\circ} , we denote by $\text{Perv}_{(B_{\circ})}(X_{\circ}^s, \mathbb{E})$ (respectively $\text{Perv}_{(B)}(X^s, \mathbb{E})$) the category of \mathbb{E} -perverse sheaves on X_{\circ}^s (respectively X^s) for the Bruhat stratification.

We denote by ℓ the length function on W with respect to S and by w_0 the longest element of W .

We denote by $\pi_s : X_{\circ} \rightarrow X_{\circ}^s$ the natural projection. We have functors

$$\text{Perv}_{(B_{\circ})}(X_{\circ}, \mathbb{E}) \begin{array}{c} \xrightarrow{\pi_s^{[!]} := {}^p\mathcal{H}^0(\pi_{s!}(-)[1])} \\ \xleftarrow{\pi_s^{[!]} := \pi_s^![-1]} \end{array} \text{Perv}_{(B_{\circ})}(X_{\circ}^s, \mathbb{E}),$$

and similarly for the categories $\text{Perv}_{(B)}(X, \mathbb{E})$ and $\text{Perv}_{(B)}(X^s, \mathbb{E})$. The functor $\pi_s^{[1]}$ is exact, and is right adjoint to $\pi_s^{[0]}$. It follows that $\pi_s^{[0]}$ sends projective objects in $\text{Perv}_{(B)}(X, \mathbb{E})$ to projective objects in $\text{Perv}_{(B)}(X^s, \mathbb{E})$.

4.2 Morphisms between standard perverse sheaves

For any $w \in W$, following the notation of § 2.2 we set $\Delta_{w,\circ} := i_{w!}\mathbb{E}_{X_{w,\circ}}[\ell(w)] \in \text{Perv}_{(B_\circ)}(X_\circ, \mathbb{E})$, and $\Delta_w := \text{For}(\Delta_{w,\circ})$. Recall (see § 3.3 and Remark 3.3.5) that for $u, v \in W$ and $i \in \mathbb{Z}$ the \mathbb{E} -modules $\text{Hom}_{X, \mathbb{E}}^i(\Delta_u, \Delta_v)$ and $\text{H}_c^i(X_v \cap X_u^-, \mathbb{E})$ have a natural automorphism induced by the Frobenius morphism.

PROPOSITION 4.2.1. *For any $u, v \in W$ there exists an isomorphism of graded (\mathbb{E}, ϕ) -modules*


$$\text{Hom}_{X, \mathbb{E}}^\bullet(\Delta_u, \Delta_v) \cong \text{H}_c^{\bullet+\ell(v)-\ell(u)}(X_v \cap X_u^-; \mathbb{E}).$$

Proof. If $u \not\leq v$, then both modules are 0, hence there is nothing to prove. Assume from now on that $u \leq v$.

Fix $n \in \mathbb{Z}$. First, adjunction provides an isomorphism of (\mathbb{E}, ϕ) -modules

$$\text{Hom}_{X, \mathbb{E}}^n(\Delta_u, \Delta_v) \cong \text{H}^{n+\ell(v)-\ell(u)}(X_u, i_u^! i_v! \mathbb{E}_{X_v}).$$


Consider the following diagram.



We obtain isomorphisms

$$i_u^! i_v! \mathbb{E}_{X_v} \cong i_{u,v}^! i_v^! j_v^! \mathbb{E}_{X_v} \cong i_{u,v}^! j_v^! \mathbb{E}_{X_v}.$$

Now, consider the following diagram with cartesian square



where j and j' are the open inclusions considered in [KLS0, § 1.4], j'_v is induced by j_v , and k (respectively p) is the obvious inclusion (respectively projection). Then we have

$$i_{u,v}^! j_v^! \mathbb{E}_{X_v} \cong k^! j'^! j_v^! \mathbb{E}_{X_v} \cong k^! j^* j_v^! \mathbb{E}_{X_v} \cong k^! j_v^! \mathbb{E}_{(X_v \cap X_u^-) \times X_u},$$

where the second isomorphism follows from the fact that j is an open embedding, and the third one from base change.

The adjunction $(k_!, k^!)$ provides a morphism of functors $k_! k^! \rightarrow \text{id}$. Composing with $p_!$ gives a morphism of functors $k^! \rightarrow p_!$. By [Soe89, Proposition 1], this morphism induces an isomorphism

$$k^! j_v^! \mathbb{E}_{(X_v \cap X_u^-) \times X_u} \xrightarrow{\sim} p_! j_v^! \mathbb{E}_{(X_v \cap X_u^-) \times X_u}.$$

Combining these isomorphisms we obtain

$$\mathrm{Hom}_{X, \mathbb{E}}^n(\Delta_u, \Delta_v) \cong \mathrm{H}^{n+\ell(v)-\ell(u)}(X_u, (p \circ j'_v)! \mathbb{E}_{(X_v \cap X_u^-) \times X_u}).$$

Now, consider the following (tautological) cartesian diagram,

$$\begin{array}{ccc} (X_v \cap X_u^-) \times X_u & \xrightarrow{q} & X_v \cap X_u^- \\ p \circ j'_v \downarrow & \square & \downarrow b \\ X_u & \xrightarrow{a} & \mathrm{pt} \end{array}$$

where q, a, b are the natural projections. Then by the base change theorem we have

$$(p \circ j'_v)! \mathbb{E}_{(X_v \cap X_u^-) \times X_u} \cong (p \circ j'_v)! q^* \mathbb{E}_{X_v \cap X_u^-} \cong a^* b! \mathbb{E}_{X_v \cap X_u^-}.$$

We deduce isomorphisms

$$\begin{aligned} \mathrm{Hom}_{X, \mathbb{E}}^n(\Delta_u, \Delta_v) &\cong \mathrm{H}^{n+\ell(v)-\ell(u)}(X_u, a^* b! \mathbb{E}_{X_v \cap X_u^-}) \\ &\cong \mathrm{H}^{n+\ell(v)-\ell(u)}(\mathrm{pt}, a_* a^* b! \mathbb{E}_{X_v \cap X_u^-}). \end{aligned}$$

Finally, since X_u is an affine space, we observe that the adjunction (a^*, a_*) induces an isomorphism

$$\mathrm{H}^{n+\ell(v)-\ell(u)}(\mathrm{pt}, b! \mathbb{E}_{X_v \cap X_u^-}) \xrightarrow{\sim} \mathrm{H}^{n+\ell(v)-\ell(u)}(\mathrm{pt}, a_* a^* b! \mathbb{E}_{X_v \cap X_u^-}),$$

which finishes the proof. □

4.3 Weights of cohomology of Deodhar varieties

In the next two lemmas we let \mathbb{k} be an arbitrary field, and we use the notation G, B, X , etc. for a split reductive algebraic group, a Borel subgroup, the flag variety, etc. over \mathbb{k} . We will use these results only when $\mathbb{k} = \mathbb{F}_q$.

The following result is well known, and probably due to Deodhar (see [Deo85], or [Cur88] for a more general result). For this reason we call the varieties $X_v \cap X_u^-$ ‘Deodhar varieties’.

LEMMA 4.3.1. *Let $u, v \in W$ and $s \in S$ be such that $u \leq v, vs > v$ and $us > u$.*

(i) *There exists an isomorphism of \mathbb{k} -varieties*

$$X_{vs} \cap X_{us}^- \cong X_v \cap X_u^-.$$

(ii) *There exists a closed subvariety $Z \subset X_{vs} \cap X_u^-$, with complement U , and isomorphisms of \mathbb{k} -varieties*

$$Z \cong (X_v \cap X_{us}^-) \times \mathbb{A}_{\mathbb{k}}^1, \quad U \cong (X_v \cap X_u^-) \times (\mathbb{A}_{\mathbb{k}}^1 \setminus \{0\}).$$

Proof. Set $X_u^{s,-} := B^- u P_s / P_s \subset X^s$. Then we have

$$(\pi_s)^{-1}(X_u^{s,-}) = X_u^- \sqcup X_{us}^-, \quad (\pi_s)^{-1}(X_v^s) = X_{vs} \sqcup X_v.$$

In each of these decompositions, the first term is open and the second term is closed. On the other hand, as the restriction of π_s to X_v (respectively X_{us}^-) is an isomorphism with X_v^s (respectively $X_u^{s,-}$), this morphism induces isomorphisms

$$\begin{aligned} X_v \cap X_u^- &\cong \pi_s(X_v \cap X_u^-), & X_v \cap X_{us}^- &\cong \pi_s(X_v \cap X_{us}^-), \\ X_{vs} \cap X_{us}^- &\cong \pi_s(X_{vs} \cap X_{us}^-), \end{aligned}$$

and

$$\begin{aligned} X_v^s \cap X_u^{s,-} &= \pi_s(X_v \cap X_u^-) \sqcup \pi_s(X_v \cap X_{us}^-) \\ &= \pi_s(X_{vs} \cap X_{us}^-) \sqcup \pi_s(X_v \cap X_{us}^-). \end{aligned}$$

(In each of these decompositions the first term is open and the second one is closed.) It follows that

$$\pi_s(X_v \cap X_u^-) = \pi_s(X_{vs} \cap X_{us}^-), \tag{4.3.2}$$

which proves (i).

Now, set

$$\begin{aligned} Z &:= (X_{vs} \cap X_u^-) \cap (\pi_s)^{-1}(\pi_s(X_v \cap X_{us}^-)), \\ U &:= (X_{vs} \cap X_u^-) \cap (\pi_s)^{-1}(\pi_s(X_v \cap X_u^-)). \end{aligned}$$

To prove (ii) it is enough to prove that $\pi_s|_Z : Z \rightarrow \pi_s(X_v \cap X_{us}^-)$ is a trivial \mathbb{A}_k^1 -fibration, and that $\pi_s|_U : U \rightarrow \pi_s(X_v \cap X_u^-)$ is a trivial $(\mathbb{A}_k^1 \setminus \{0\})$ -fibration. Consider the \mathbb{A}_k^1 -fibration

$$f : (X_{vs} \cap X_u^-) \sqcup (X_{vs} \cap X_{us}^-) \rightarrow X_v^s \cap X_u^{s,-}$$

induced by π_s , which is trivial since it is the restriction of the trivial \mathbb{A}_k^1 -fibration $X_{vs} \rightarrow X_v^s$ to the inverse image of $X_v^s \cap X_u^{s,-}$. We claim that

$$f^{-1}(\pi_s(X_v \cap X_{us}^-)) = Z, \tag{4.3.3}$$

which will prove our claim about Z . Indeed, let $x \in \pi_s(X_v \cap X_{us}^-)$. Then $\pi_s^{-1}(x) \cap X_v = \text{pt}$, and $\pi_s^{-1}(x) \cap X_{us}^- = \text{pt}$, and by assumption these points coincide. It follows that $\pi_s^{-1}(x) \cap X_{vs} = \pi_s^{-1}(x) \cap X_u^-$, hence that $\pi_s^{-1}(x) \cap (X_{vs} \cap X_{us}^-) = \emptyset$. This proves (4.3.3).

On the other hand, consider $f^{-1}(\pi_s(X_v \cap X_u^-))$, and denote by g the restriction of f to this open subvariety. Then the restriction of g to

$$(X_{vs} \cap X_{us}^-) \subset f^{-1}(\pi_s(X_v \cap X_u^-))$$

is an isomorphism, see (4.3.2). Hence the claim about U follows from Lemma 4.3.5 below. \square

Remark 4.3.4. In case (ii), the closed subvariety Z is empty if $us \not\leq v$.

LEMMA 4.3.5. *Let M be a k -variety, and consider the trivial fibration*

$$a : M \times \mathbb{A}_k^1 \rightarrow M.$$

Assume there exists a subvariety $N \subset M \times \mathbb{A}_k^1$ such that the restriction of a to N is an isomorphism $N \xrightarrow{\sim} M$. Then the restriction of a to the complement of N is a trivial $(\mathbb{A}_k^1 \setminus \{0\})$ -fibration.

Proof. Let ξ be the composition $M \xrightarrow{\sim} N \hookrightarrow M \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$. Then the automorphism

$$\begin{cases} M \times \mathbb{A}_k^1 \rightarrow M \times \mathbb{A}_k^1 \\ (m, t) \mapsto (m, t - \xi(m)) \end{cases}$$

identifies the complement of N with $M \times (\mathbb{A}_k^1 \setminus \{0\})$ as a fibration over M . \square

Now we come back to our varieties over \mathbb{F}_q and $\overline{\mathbb{F}}_q$.

PROPOSITION 4.3.6. *Let $u, v \in W$, with $u \leq v$. The cohomology*

$$H_c^\bullet(X_v \cap X_u^-; \mathbb{O})$$

is concentrated in degrees between $\ell(v) - \ell(u)$ and $2(\ell(v) - \ell(u))$. Moreover, for any $n \in \llbracket \ell(v) - \ell(u), 2(\ell(v) - \ell(u)) \rrbracket$, the (\mathbb{O}, ϕ) -module $H_c^n(X_v \cap X_u^-; \mathbb{O})$ has q -weights obtained from $\llbracket -\lfloor \frac{n}{2} \rfloor, -n + \ell(v) - \ell(u) \rrbracket$.

Proof. We prove the claim by induction on the Bruhat order on v . It is obvious if $v = 1$.

Now, assume the result is known for v , and let $s \in S$ be such that $vs > v$. Let $u \in W$ such that $u \leq vs$. If $us < u$ (and then $us \leq v$), by Lemma 4.3.1(i) we have an isomorphism

$$X_{vs, \circ} \cap X_{u, \circ}^- \cong X_{v, \circ} \cap X_{us, \circ}^-.$$

As $\ell(vs) - \ell(u) = \ell(v) - \ell(us)$, the result follows by induction.

Now, assume that $us > u$ (and then $u \leq v$). Then, by Lemma 4.3.1(ii), $X_{vs, \circ} \cap X_{u, \circ}^-$ is the disjoint union of a closed subvariety Z_{\circ} isomorphic to $(X_{v, \circ} \cap X_{us, \circ}^-) \times \mathbb{A}_{\circ}^1$ and an open subvariety U_{\circ} isomorphic to $(X_{v, \circ} \cap X_{u, \circ}^-) \times (\mathbb{A}_{\circ}^1 \setminus \{0\})$. Consider the associated long exact sequence

$$\cdots \rightarrow H_c^n(U; \mathbb{O}) \rightarrow H_c^n(X_{vs} \cap X_u^-; \mathbb{O}) \rightarrow H_c^n(Z; \mathbb{O}) \rightarrow \cdots .$$

Both $H_c^{\bullet}(\mathbb{A}^1; \mathbb{O})$ and $H_c^{\bullet}(\mathbb{A}^1 \setminus \{0\}; \mathbb{O})$ are free \mathbb{O} -modules and so we can apply the Künneth formula to obtain

$$H_c^{\bullet}(Z; \mathbb{O}) \cong H_c^{\bullet}(X_v \cap X_{us}^-; \mathbb{O}) \otimes_{\mathbb{O}} H_c^{\bullet}(\mathbb{A}^1; \mathbb{O}) \cong H_c^{\bullet-2}(X_v \cap X_{us}^-; \mathbb{O}) \otimes_{\mathbb{O}} H_c^2(\mathbb{A}^1; \mathbb{O})$$

and

$$\begin{aligned} H_c^{\bullet}(U; \mathbb{O}) &\cong H_c^{\bullet}(X_v \cap X_u^-; \mathbb{O}) \otimes_{\mathbb{O}} H_c^{\bullet}(\mathbb{A}^1 \setminus \{0\}; \mathbb{O}) \\ &\cong H_c^{\bullet-1}(X_v \cap X_u^-; \mathbb{O}) \otimes_{\mathbb{O}} H_c^1(\mathbb{A}^1 \setminus \{0\}; \mathbb{O}) \oplus H_c^{\bullet-2}(X_v \cap X_u^-; \mathbb{O}) \otimes_{\mathbb{O}} H_c^2(\mathbb{A}^1 \setminus \{0\}; \mathbb{O}). \end{aligned}$$

Our claim follows, using Remark 3.3.5 and induction. □

Remark 4.3.7. The bounds on weights in Proposition 4.3.6 are exactly the same as the bounds provided in a very general setting for cohomology with coefficients in \mathbb{K} by [Del80, Corollaire 3.3.3]. In particular, if one is only interested in $H_c^{\bullet}(X_v \cap X_u^-; \mathbb{K})$, then the information on the structure of $X_v \cap X_u^-$ given by Lemma 4.3.1 does not give improved bounds. However, here we work with integral coefficients, which are not considered in [Del80].

Combining Proposition 4.2.1 and Proposition 4.3.6 we obtain the following result, which will play a crucial role in the rest of this section.

COROLLARY 4.3.8. *Let $u, v \in W$ such that $u \leq v$. Then*

$$\text{Hom}_{X, \mathbb{O}}^n(\Delta_u, \Delta_v)$$

vanishes unless $0 \leq n \leq \ell(v) - \ell(u)$. If $n \in \llbracket 0, \ell(v) - \ell(u) \rrbracket$, this (\mathbb{O}, ϕ) -module has q -weights obtained from $\llbracket -\lfloor (n + \ell(v) - \ell(u))/2 \rfloor, -n \rrbracket$.

4.4 Weights in the Δ -flag of projective covers

PROPOSITION 4.4.1. *For any $u \in W$, there exists an object $\mathcal{P}_{u, \circ}$ in $\text{Perv}_{(B_{\circ})}(X_{\circ}, \mathbb{O})$ such that $\text{For}(\mathcal{P}_{u, \circ})$ is projective in $\text{Perv}_{(B)}(X, \mathbb{O})$, together with an exact sequence*

$$\mathcal{M}_{u, \circ} \hookrightarrow \mathcal{P}_{u, \circ} \twoheadrightarrow \Delta_{u, \circ}$$

such that $\mathcal{M}_{u, \circ}$ has a filtration whose subquotients are of the form

$$E_{v, u} \otimes_{\mathbb{O}} \Delta_{v, \circ}$$

for $v > u$, where $E_{v, u}$ is an (\mathbb{O}, ϕ) -module which is \mathbb{O} -free and has q -weights obtained from $\llbracket 1, \ell(v) - \ell(u) \rrbracket$.

Proof. We prove by induction that for any $I \subset W$ containing u with X_I closed in X , there exists an object $\mathcal{P}_{u,\circ}^I$ in $\text{Perv}_{(B_\circ)}(X_{I,\circ}, \mathbb{O})$ such that $\mathcal{P}_u^I := \text{For}(\mathcal{P}_{u,\circ}^I)$ is projective in $\text{Perv}_{(B)}(X_I, \mathbb{O})$ together with an exact sequence

$$\mathcal{M}_{u,\circ}^I \hookrightarrow \mathcal{P}_{u,\circ}^I \twoheadrightarrow \Delta_{u,\circ}^I$$

such that $\mathcal{M}_{u,\circ}^I$ has a filtration whose subquotients are of the form

$$E_{v,u}^I \otimes_{\mathbb{O}} \Delta_{v,\circ}^I$$

for $v > u$ ($v \in I$), where $E_{v,u}^I$ is an (\mathbb{O}, ϕ) -module which is \mathbb{O} -free and has q -weights obtained from $[[1, \ell(v) - \ell(u)]]$. (Here, $\Delta_{v,\circ}^I$ is the standard perverse sheaf on $X_{I,\circ}$ associated with v .) The case $I = W$ will give the proposition. If $u \in I$ is maximal, then $\mathcal{P}_{u,\circ}^I = \Delta_{u,\circ}^I$ satisfies the requirements.

Now we take $I = I' \cup \{v\}$ with $X_{I'}$ and X_I closed and containing X_u , and we assume that $\mathcal{P}_{u,\circ}^{I'}$ has been constructed. Consider the (\mathbb{O}, ϕ) -module

$$E := \text{Hom}_{X_{I',\circ}}^1(\mathcal{P}_u^{I'}, \Delta_v^I).$$

(Here we consider $\mathcal{P}_u^{I'}$ as a perverse sheaf on $X_{I'}$.) We claim that this module has q -weights obtained from $[[\ell(u) - \ell(v), -1]]$. Indeed, by induction hypothesis $\mathcal{P}_{u,\circ}^{I'}$ has a filtration with subquotients $E_{w,u}^{I'} \otimes_{\mathbb{O}} \Delta_{w,\circ}^{I'}$, where $E_{w,u}^{I'}$ has q -weights obtained from $[[0, \ell(w) - \ell(u)]]$, and moreover, by Corollary 4.3.8, if it is non-zero then $\text{Hom}_{X_{I',\circ}}^1(\Delta_w^{I'}, \Delta_v^I) = \text{Hom}_{X,\circ}^1(\Delta_w, \Delta_v)$ has q -weights obtained from

$$\left[- \left\lfloor \frac{1 + \ell(v) - \ell(w)}{2} \right\rfloor, -1 \right] \subset [[\ell(w) - \ell(v), -1]].$$

Using Lemma 3.1.7, we deduce that there exists an (\mathbb{O}, ϕ) -module E_{free} which is \mathbb{O} -free, with q -weights obtained from $[[\ell(u) - \ell(v), -1]]$ and a surjection of (\mathbb{O}, ϕ) -modules $E_{\text{free}} \twoheadrightarrow E$. By [BBD82, (5.1.2.5)], there exists a natural surjection

$$\text{Ext}_{\text{Perv}_{(B_\circ)}(X_\circ, \mathbb{O})}^1(\mathcal{P}_{u,\circ}^{I'}, E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_{v,\circ}^I) \twoheadrightarrow (\text{Ext}_{\text{Perv}_{(B)}(X, \mathbb{O})}^1(\mathcal{P}_u^{I'}, E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_v^I))^{\phi\text{-inv}}.$$

The element of $\text{Ext}_{\text{Perv}_{(B)}(X, \mathbb{O})}^1(\mathcal{P}_u^{I'}, E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_v^I)$ considered in the proof of Proposition 2.3.1 is ϕ -invariant by construction, and hence it defines an extension of $E_{\text{free}}^* \otimes_{\mathbb{O}} \Delta_{v,\circ}^I$ by $\mathcal{P}_{u,\circ}^{I'}$, which we denote by $\mathcal{P}_{u,\circ}^I$. By the proof of Proposition 2.3.1, $\text{For}(\mathcal{P}_{u,\circ}^I)$ is projective, which proves the induction. \square

For any $u \in W$ we fix an object $\mathcal{P}_{u,\circ}$ as in Proposition 4.4.1, and we set

$$\mathcal{P}_\circ := \bigoplus_{u \in W} \mathcal{P}_{u,\circ}.$$

Note that this object satisfies the condition of Lemma 3.3.4 (see the proof of Corollary 2.3.3).

PROPOSITION 4.4.2. *The (\mathbb{O}, ϕ) -module*

$$\text{End}_{X,\circ}(\mathcal{P})$$

is \mathbb{O} -free, and has q -weights obtained from $[-\ell(w_0), \ell(w_0)]$. In particular, this (\mathbb{O}, ϕ) -module is q -decomposable if the order of q in \mathbb{F} is strictly bigger than $2\ell(w_0)$.

Proof. The fact that this \mathbb{O} -module is free follows from Proposition 2.4.1(iv).

To prove the statement about weights, it is sufficient to prove that for any $u, v \in W$ the q -weights of $\text{Hom}_{X,\circ}(\mathcal{P}_u, \mathcal{P}_v)$ are in $[-\ell(w_0), \ell(w_0)]$. Now recall that $\mathcal{P}_{v,\circ}$ has a filtration with

subquotients of the form

$$E_{w,v} \otimes_{\mathbb{O}} \Delta_{w,\circ}$$

for $w \geq v$, where $E_{w,v}$ is an (\mathbb{O}, ϕ) -module which has q -weights obtained from $\llbracket 0, \ell(w) - \ell(v) \rrbracket$. Hence, $\text{Hom}_{X,\mathbb{O}}(\mathcal{P}_u, \mathcal{P}_v)$ has a filtration (as an (\mathbb{O}, ϕ) -module) with subquotients of the form

$$E_{w,v} \otimes_{\mathbb{O}} \text{Hom}_{X,\mathbb{O}}(\mathcal{P}_u, \Delta_w).$$

Similarly, $\mathcal{P}_{u,\circ}$ has a filtration with subquotients of the form

$$E_{w',u} \otimes_{\mathbb{O}} \Delta_{w',\circ}$$

for $w' \geq u$, where $E_{w',u}$ is an (\mathbb{O}, ϕ) -module which has q -weights obtained from $\llbracket 0, \ell(w') - \ell(u) \rrbracket$. Hence, it is enough to study weights in

$$F_{u,v,w,w'} := E_{w',u}^* \otimes_{\mathbb{O}} \text{Hom}_{X,\mathbb{O}}(\Delta_{w'}, \Delta_w) \otimes_{\mathbb{O}} E_{w,v}.$$

By Corollary 4.3.8, $\text{Hom}_{X,\mathbb{O}}(\Delta_{w'}, \Delta_w)$ has q -weights obtained from $\llbracket \ell(w') - \ell(w)/2, 0 \rrbracket$. Hence, $F_{u,v,w,w'}$ has q -weights obtained from

$$\left\llbracket \ell(u) - \frac{\ell(w') + \ell(w)}{2}, \ell(w) - \ell(v) \right\llbracket \subset \llbracket -\ell(w_0), \ell(w_0) \rrbracket,$$

which finishes the proof of the claim on weights.

Finally, the last claim of the statement follows from Remark 3.1.6. □

4.5 Partial flag varieties

In this subsection we fix a simple reflection s , and consider the variety X^s . For any $v \in W^s$, we denote by Δ_v^s the standard object associated with the stratum X_v^s .

LEMMA 4.5.1. *For $u, v \in W^s$, the (\mathbb{O}, ϕ) -module*

$$\text{Hom}_{X^s,\mathbb{O}}^n(\Delta_u^s, \Delta_v^s)$$

vanishes unless $0 \leq n \leq \ell(v) - \ell(u)$, in which case it has q -weights obtained from the interval $\llbracket -(n + \ell(v) - \ell(u))/2, -n \rrbracket$.

Proof. The restriction of π_s to X_v is an isomorphism onto X_v^s , hence we have $\pi_{s*}\Delta_v = \pi_{s!}\Delta_v = \Delta_v^s$. Using adjunction, we deduce isomorphisms

$$\text{Hom}_{X^s,\mathbb{O}}^n(\Delta_u^s, \Delta_v^s) \cong \text{Hom}_{X^s,\mathbb{O}}^n(\Delta_u^s, \pi_{s*}\Delta_v) \cong \text{Hom}_{X,\mathbb{O}}^n(\pi_s^*\Delta_u^s, \Delta_v).$$

By the base change theorem we have $\pi_s^*\Delta_u^s = i_u^s! \underline{\mathbb{O}}_{X_u \sqcup X_{us}}[\ell(u)]$, where $i_u^s : X_u \sqcup X_{us} \hookrightarrow X$ is the inclusion. We have an exact sequence of sheaves on $X_u \sqcup X_{us}$

$$j_! \underline{\mathbb{O}}_{X_{us}} \hookrightarrow \underline{\mathbb{O}}_{X_u \sqcup X_{us}} \twoheadrightarrow i_* \underline{\mathbb{O}}_{X_u},$$

where i (respectively j) denotes the inclusion of X_u (respectively X_{us}) which induces a distinguished triangle

$$\Delta_{us}[-1] \rightarrow i_u^s! \underline{\mathbb{O}}_{X_u \sqcup X_{us}}[\ell(u)] \rightarrow \Delta_u \xrightarrow{[1]}.$$

Hence we obtain an exact sequence for any $n \geq 0$,

$$\text{Hom}_{X,\mathbb{O}}^n(\Delta_u, \Delta_v) \rightarrow \text{Hom}_{X^s,\mathbb{O}}^n(\Delta_u^s, \Delta_v^s) \rightarrow \text{Hom}_{X,\mathbb{O}}^{n+1}(\Delta_{us}, \Delta_v).$$

Then the result follows from Corollary 4.3.8. □

Using this lemma, the same proof as that of Proposition 4.4.1 gives the following result.

PROPOSITION 4.5.2. For any $u \in W^s$, there exists an object $\mathcal{P}_{u,\circ}^s$ in $\text{Perv}_{(B_\circ)}(X_\circ^s, \mathbb{O})$ such that $\text{For}(\mathcal{P}_{u,\circ}^s)$ is projective in $\text{Perv}_{(B)}(X^s, \mathbb{O})$, together with an exact sequence

$$\mathcal{M}_{u,\circ}^s \hookrightarrow \mathcal{P}_{u,\circ}^s \twoheadrightarrow \Delta_{u,\circ}^s$$

such that $\mathcal{M}_{u,\circ}^s$ has a filtration whose subquotients are of the form

$$E_{v,u}^s \otimes_{\mathbb{O}} \Delta_{v,\circ}^s$$

for $v > u$ ($v \in W^s$), where $E_{v,u}^s$ is an (\mathbb{O}, ϕ) -module which is \mathbb{O} -free and has q -weights obtained from $[[1, \ell(v) - \ell(u)]]$.

As in § 4.4, we choose objects $\mathcal{P}_{u,\circ}^s$ as in Proposition 4.5.2, and we define the object

$$\mathcal{P}_\circ^s := \bigoplus_{u \in W^s} \mathcal{P}_{u,\circ}^s.$$

Again we observe that this object satisfies the conditions of Lemma 3.3.4 (see the proof of Corollary 2.3.3). We let \mathcal{P}_\circ denote the same object as in § 4.4.

LEMMA 4.5.3. The (\mathbb{O}, ϕ) -module

$$\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[[1]]} \mathcal{P}, \mathcal{P}^s)$$

is \mathbb{O} -free, and has q -weights obtained from $[-\ell(w_0) + 1, \ell(w_0)]$. In particular, it is q -decomposable if the order of q in \mathbb{F} is bigger than $2\ell(w_0)$.

Proof. First, the object $\pi_{s[[1]]} \mathcal{P} \in \text{Perv}_{(B)}(X^s, \mathbb{O})$ is projective (see § 4.1), hence the fact that $\text{Hom}(\pi_{s[[1]]} \mathcal{P}, \mathcal{P}^s)$ is \mathbb{O} -free follows from Proposition 2.4.1(iv).

Next, by adjunction we have

$$\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[[1]]} \mathcal{P}, \mathcal{P}^s) \cong \text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \pi_s^{[[1]]} \mathcal{P}^s) \cong \text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \pi_s^! \mathcal{P}^s[-1]).$$

Now there exists an isomorphism of functors $\pi_s^! \cong \pi_s^*[2](1)$ since π_s is smooth of relative dimension 1, hence we obtain an isomorphism of (\mathbb{O}, ϕ) -modules

$$\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[[1]]} \mathcal{P}, \mathcal{P}^s) \cong \text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \pi_s^* \mathcal{P}^s[1])(1).$$

By construction, \mathcal{P}_\circ^s has a filtration with subquotients of the form

$$E_{v,u}^s \otimes_{\mathbb{O}} \Delta_{v,\circ}^s$$

for $v \geq u$ ($v, u \in W^s$), where $E_{v,u}^s$ is an (\mathbb{O}, ϕ) -module which has q -weights obtained from $[[0, \ell(v) - \ell(u)]]$. Hence, $\pi_s^* \mathcal{P}_\circ^s[1]$ has a filtration with subquotients

$$E_{v,u}^s \otimes_{\mathbb{O}} \pi_s^* \Delta_{v,\circ}^s[1],$$

and then $\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[[1]]} \mathcal{P}, \mathcal{P}^s)$ has a filtration with subquotients

$$A_{v,u}^s := E_{v,u}^s \otimes_{\mathbb{O}} \text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \pi_s^* \Delta_v^s[1])(1).$$

As in the proof of Lemma 4.5.1, we have a surjection $\Delta_{vs} \twoheadrightarrow \pi_s^* \Delta_v^s[1]$, which induces a surjection

$$\text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \Delta_{vs})(1) \twoheadrightarrow \text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \pi_s^* \Delta_v^s[1])(1).$$

We have seen in the proof of Proposition 4.4.2 that $\text{Hom}_{X, \mathbb{O}}(\mathcal{P}, \Delta_{vs})$ has q -weights obtained from $[-\ell(w_0), 0]$. It follows that $A_{v,u}^s$ has q -weights obtained from $[-\ell(w_0) + 1, \ell(w_0)]$, finishing the proof of the claim concerning weights.

The last claim of the statement again follows from Remark 3.1.6. □

PROPOSITION 4.5.4. *The (\mathbb{O}, ϕ) -module*

$$\text{End}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P})$$

is \mathbb{O} -free. If the order of q in \mathbb{F} is bigger than $2\ell(w_0)$, then it is q -decomposable.

Proof. As $\pi_{s[\ell]} \mathcal{P}$ is projective (see § 4.1), by Proposition 2.4.1(iv) its endomorphism algebra is \mathbb{O} -free.

By Lemma 4.5.5 below, the morphism

$$\text{End}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P}) \rightarrow \text{End}_{\text{End}(\mathcal{P}^s)}(\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P}, \mathcal{P}^s))$$

induced by the functor $\text{Hom}(-, \mathcal{P}^s)$ is an isomorphism of (\mathbb{O}, ϕ) -modules. By Lemma 4.5.3, under our assumptions $\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P}, \mathcal{P}^s)$ is a q -decomposable, \mathbb{O} -free (\mathbb{O}, ϕ) -module, hence the same holds for the (\mathbb{O}, ϕ) -module

$$V_s := \text{End}_{\mathbb{O}}(\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P}, \mathcal{P}^s)).$$

As $\text{End}_{\text{End}(\mathcal{P}^s)}(\text{Hom}_{X^s, \mathbb{O}}(\pi_{s[\ell]} \mathcal{P}, \mathcal{P}^s))$ is a sub- (\mathbb{O}, ϕ) -module of V_s with torsion-free quotient, by Lemma 3.1.3, it is also q -decomposable, which finishes the proof. \square

LEMMA 4.5.5. *Let R be a right noetherian ring, and Q be a projective generator of $\text{Modf}-R$. Then the contravariant functor*

$$\text{Hom}_{-R}(-, Q) : \text{Modf}-R \rightarrow \text{End}_{-R}(Q)\text{-Mod}$$

is fully faithful on projectives.

Proof. To simplify notation, let $R' := \text{End}_{-R}(Q)$. By [Bas68, Exercise on p. 55], R' is a right noetherian ring and the functor $\text{Hom}_{-R}(Q, -)$ induces an equivalence of categories between $\text{Modf}-R$ and $\text{Modf}-R'$. In particular, for any P in $\text{Projf}-R$ the morphism

$$\text{Hom}_{-R}(P, Q) \rightarrow \text{Hom}_{-R'}(\text{Hom}_{-R}(Q, P), R') \tag{4.5.6}$$

induced by $\text{Hom}_{-R}(Q, -)$ is an isomorphism, and moreover $\text{Hom}_{-R}(Q, P)$ is in $\text{Projf}-R'$. Using isomorphism (4.5.6), the functor $\text{Hom}_{-R}(-, Q) : \text{Projf}-R \rightarrow R'\text{-Mod}$ is isomorphic to the composition

$$\text{Projf}-R \xrightarrow{\text{Hom}_{-R}(Q, -)} \text{Projf}-R' \xrightarrow{\text{Hom}_{-R'}(-, R')} R'\text{-Mod}.$$

As explained above the first functor is fully faithful, and the second one is easily seen to be fully faithful also. \square

5. Formality

As above, in §§ 5.1–5.2 we let \mathbb{E} be either \mathbb{O} or \mathbb{F} .

5.1 Bott–Samelson sheaves and parity sheaves

Recall [JMW09, Definition 2.4] that a complex $\mathcal{F} \in \mathcal{D}_{(B)}^b(X, \mathbb{E})$ is called *parity* if it admits a decomposition $\mathcal{F} \cong \mathcal{F}_0 \oplus \mathcal{F}_1$ such that for all $w \in W$, $? \in \{!, *\}$ and $j \in \mathbb{Z}$ we have:

- (i) $\mathcal{H}^j(i_w^? \mathcal{F}_k) = 0$ if $j \not\equiv k \pmod{2}$;
- (ii) $\mathcal{H}^j(i_w^? \mathcal{F}_k)$ is an \mathbb{E} -free constant local system if $j \equiv k \pmod{2}$.

An indecomposable parity complex is called a *parity sheaf*. Note that, as the category $\mathcal{D}_{(B)}^b(X, \mathbb{E})$ satisfies the Krull–Schmidt property, every parity complex decomposes (uniquely) as a direct sum of parity sheaves (see [JMW09, § 2.1] for details).

We have the following theorem [JMW09, Theorem 2.12].

THEOREM 5.1.1. *For all $w \in W$ there is (up to isomorphism) at most one parity sheaf \mathcal{E}_w supported on \overline{X}_w and extending the constant local system $\underline{\mathbb{E}}_{X_w}[\ell(w)]$ on X_w . Moreover, any parity sheaf is isomorphic to $\mathcal{E}_w[m]$ for some $w \in W$ and $m \in \mathbb{Z}$.*

If $f = (s, t, \dots, r)$ is a sequence of simple reflections, we consider the corresponding Bott–Samelson complex

$$\mathcal{BS}_{f,\circ}^{\mathbb{E}} := \pi_s^! \pi_s! \pi_t^! \pi_t! \cdots \pi_r^! \pi_r! \text{IC}_{e,\circ}^{\mathbb{E}} \in \mathcal{D}_{(B_\circ)}^b(X_\circ, \mathbb{E}),$$

where $e \in W$ is the unit element and $\text{IC}_{e,\circ}^{\mathbb{E}}$ is the corresponding IC-sheaf. As usual, we denote by $\mathcal{BS}_f^{\mathbb{E}}$ the object obtained by base change to $\overline{\mathbb{F}}_q$. The following is an immediate consequence of Proposition 5.1.7 below in the case $\mathbb{E} = \mathbb{O}$; the case $\mathbb{E} = \mathbb{F}$ follows (or can alternatively be proved along the same lines).

PROPOSITION 5.1.2. *For any sequence f of simple reflections, $\mathcal{BS}_f^{\mathbb{E}}$ is parity. If f is a reduced expression for w^{-1} then $\mathcal{BS}_f^{\mathbb{E}}$ is supported on \overline{X}_w and $i_w^* \mathcal{BS}_f^{\mathbb{E}} \cong \underline{\mathbb{E}}_{X_w}[2\ell(w)]$.*

Remark 5.1.3. Theorem 5.1.1 says nothing about the existence of parity sheaves, and for an arbitrary stratified variety existence may be difficult to establish. The above proposition combined with Theorem 5.1.1 gives the existence and uniqueness of parity sheaves on the flag variety (see [Soe00, Theorem 1.2] and [JMW09, Theorem 4.6]).

Given a sequence $f = (s, t, \dots, r)$ as above we set $\overline{f} := st \cdots r \in W$. Let us fix a family $F = \{f_1, \dots, f_n\}$ of sequences of simple reflections such that $W = \{\overline{f}_1, \dots, \overline{f}_n\}$. Then we set

$$\mathcal{BS}_\circ^{\mathbb{E}} := \bigoplus_{f \in F} \mathcal{BS}_{f,\circ}^{\mathbb{E}},$$

and denote by $\mathcal{BS}^{\mathbb{E}}$ the object obtained by base change to $\overline{\mathbb{F}}_q$.

LEMMA 5.1.4. (i) *The category $\mathcal{D}_{(B)}^b(X, \mathbb{E})$ is generated, as a triangulated category, by the objects $\mathcal{BS}_f^{\mathbb{E}}$ for $f \in F$.*

(ii) *The category $\mathcal{D}_{(B)}^b(X^s, \mathbb{E})$ is generated, as a triangulated category, by the objects $\pi_{s!} \mathcal{BS}_f^{\mathbb{E}}$ for $f \in F$.*

Proof. (i) A straightforward induction on the support using standard distinguished triangles shows that the set $\{\Delta_x \mid x \in W\}$ generates $\mathcal{D}_{(B)}^b(X, \mathbb{E})$ as a triangulated category. (If $\mathbb{E} = \mathbb{O}$, one has to observe that for any $m \in \mathbb{Z}_{\geq 1}$ there is a natural exact sequence $\Delta_x \hookrightarrow \Delta_x \twoheadrightarrow \Delta_{x,m}$.) Hence it is enough to show that one can obtain Δ_x as a successive extension of the complexes $\mathcal{BS}_f^{\mathbb{E}}$. However, if $\overline{f} = x^{-1}$ then, by Proposition 5.1.2, $\mathcal{BS}_f^{\mathbb{E}}$ is supported on \overline{X}_x and $i_x^! \mathcal{BS}_f^{\mathbb{E}} \cong i_x^* \mathcal{BS}_f^{\mathbb{E}} \cong \underline{\mathbb{E}}_{X_x}[2\ell(x)]$. Hence we have a distinguished triangle

$$\Delta_x^{\mathbb{E}}[2\ell(x)] \rightarrow \mathcal{BS}_f^{\mathbb{E}} \rightarrow \mathcal{C} \xrightarrow{[1]}$$

with the support of \mathcal{C} contained in $\overline{X}_x \setminus X_x$. By induction, \mathcal{C} belongs to the triangulated subcategory generated by Bott–Samelson complexes, hence so does Δ_x .

(ii) One can easily check that $\mathcal{D}_{(B)}^b(X^s, \mathbb{E})$ is generated, as a triangulated category, by the essential image of $\pi_{s!} : \mathcal{D}_{(B)}^b(X, \mathbb{E}) \rightarrow \mathcal{D}_{(B)}^b(X^s, \mathbb{E})$. Hence the result follows from (i). \square

Consider the graded algebras

$$E_{\mathbb{E}} := \text{Ext}_{\mathcal{D}_{(B)}^b(X, \mathbb{E})}^{\bullet}(\mathcal{BS}^{\mathbb{E}}, \mathcal{BS}^{\mathbb{E}}),$$

$$E_{\mathbb{E}}^s := \text{Ext}_{\mathcal{D}_{(B)}^b(X^s, \mathbb{E})}^{\bullet}(\pi_{s!}\mathcal{BS}^{\mathbb{E}}, \pi_{s!}\mathcal{BS}^{\mathbb{E}})$$

for $s \in S$. These graded algebras will play a major role in the rest of the paper. Note that, as $\mathcal{BS}^{\mathbb{E}} = \text{For}(\mathcal{BS}_{\circ}^{\mathbb{E}})$ and $\pi_{s!}\mathcal{BS}^{\mathbb{E}} = \text{For}(\pi_{s!}\mathcal{BS}_{\circ}^{\mathbb{E}})$ these algebras are endowed with a natural automorphism ϕ (see §3.3). We denote by e_f the idempotent in $E_{\mathbb{E}}$, respectively $E_{\mathbb{E}}^s$, given by projection to $\mathcal{BS}_f^{\mathbb{E}}$, respectively $\pi_{s!}\mathcal{BS}_f^{\mathbb{E}}$. Note that the functor $\pi_{s!}$ induces a ring morphism $E_{\mathbb{E}} \rightarrow E_{\mathbb{E}}^s$ sending e_f to e_f , which justifies our notation.

From now on we concentrate on the case $\mathbb{E} = \mathbb{O}$. The aim of the rest of this section is to prove the following theorem.

THEOREM 5.1.5. *The (\mathbb{O}, ϕ) -algebras $E_{\mathbb{O}}$ and $E_{\mathbb{O}}^s$ are \mathbb{O} -free and vanish in odd degree. Moreover, for any $m \in \mathbb{Z}$, the (\mathbb{O}, ϕ) -modules $E_{\mathbb{O}}^{2m}$ and $(E_{\mathbb{O}}^s)^{2m}$ have q -weights obtained from $\{-m\}$.*

Recall (see Example 3.3.3) that under the equivalence $\text{Perv}(\text{pt}_{\circ}, \mathbb{O}) \cong \text{Mod}-(\mathbb{O}, \phi)$ (see Lemma 3.3.7 with $\mathcal{P}_{\circ} = \underline{\mathbb{O}}_{\text{pt}_{\circ}}$), the perverse sheaf $\underline{\mathbb{O}}_{\text{pt}_{\circ}}(m)$ corresponds to the \mathbb{O} -module \mathbb{O} endowed with the automorphism $a \mapsto q^m a$. By abuse of notation, we also denote this (\mathbb{O}, ϕ) -module by $\mathbb{O}(m)$.

Using the observation that by adjunction for $f, g \in F$ we have

$$\text{Ext}_{\mathcal{D}_{(B)}^b(X^s, \mathbb{O})}^n(\pi_{s!}\mathcal{BS}_f^{\mathbb{O}}, \pi_{s!}\mathcal{BS}_g^{\mathbb{O}}) \cong \text{Ext}_{\mathcal{D}_{(B)}^b(X, \mathbb{O})}^n(\mathcal{BS}_f^{\mathbb{O}}, \pi_s^!\pi_{s!}\mathcal{BS}_g^{\mathbb{O}}) \cong \text{Ext}_{\mathcal{D}_{(B)}^b(X, \mathbb{O})}^n(\mathcal{BS}_f^{\mathbb{O}}, \mathcal{BS}_{(s,g)}^{\mathbb{O}}),$$

Theorem 5.1.5 is a consequence of the following result.

PROPOSITION 5.1.6. *If f and g are sequences of simple reflections, then the (\mathbb{O}, ϕ) -module $\text{Ext}_{X, \mathbb{O}}^n(\mathcal{BS}_f^{\mathbb{O}}, \mathcal{BS}_g^{\mathbb{O}})$ is zero for odd n and is an extension of copies of $\mathbb{O}(-m)$ for $n = 2m$ even.*

Proof. By adjunction and the isomorphisms $\pi_{s!} \cong \pi_{s*}, \pi_s^! \cong \pi_s^*[2](1)$ we can assume that g is the empty sequence. Then the result follows from the case $v = e$ of Proposition 5.1.7(i) below. \square

PROPOSITION 5.1.7. *Let f be a sequence of simple reflections.*

(i) *For any $v \in W$, $\mathcal{H}^j(i_v^*\mathcal{BS}_{f, \circ}^{\mathbb{O}}) = 0$ for odd j and $\mathcal{H}^j(i_v^*\mathcal{BS}_{f, \circ}^{\mathbb{O}})$ is an extension of copies of $\underline{\mathbb{O}}_{X_{v, \circ}}(-k)$ if $j = 2k$ is even.*

(ii) *If f is a reduced expression for w^{-1} then $\mathcal{BS}_{f, \circ}^{\mathbb{O}}$ is supported on $\overline{X_{w, \circ}}$ and $i_w^*\mathcal{BS}_{f, \circ}^{\mathbb{O}} \cong \underline{\mathbb{O}}_{X_{w, \circ}}[2\ell(w)](\ell(w))$.*

Proof. (i) It is enough to show that if s is a simple reflection and \mathcal{S} is a complex satisfying the conclusions of (i) then so is $\pi_s^!\pi_{s!}\mathcal{S}$. To this end fix $y \in W^s$. Consider the locally closed subvariety Z_{\circ} of X_{\circ} defined via the following Cartesian diagram.

$$\begin{array}{ccc} Z_{\circ} & \xhookrightarrow{u} & X_{\circ} \\ \pi_s \downarrow & \square & \downarrow \pi_s \\ X_{y, \circ}^s & \xhookrightarrow{v} & X_{\circ}^s \end{array}$$

Then Z_{\circ} is the disjoint union of the closed subvariety $X_{y, \circ}$ and its open complement $X_{ys, \circ}$. Denote by i and j the corresponding inclusions

$$X_{ys, \circ} \xrightarrow{j} Z_{\circ} \xleftarrow{i} X_{y, \circ}.$$

On Z_\circ we have a distinguished triangle

$$j_!j^!u^*\mathcal{S} \rightarrow u^*\mathcal{S} \rightarrow i_*i^*u^*\mathcal{S} \xrightarrow{[1]}.$$

Applying $\pi_{s*} = \pi_{s!}$ to this triangle and keeping in mind the base change isomorphism $\pi_{s!}u^* \cong v^*\pi_{s!}$ and the identity $j^! = j^*$ we obtain a distinguished triangle

$$(\pi_s \circ j)_!i_{y_s}^*\mathcal{S} \rightarrow v^*\pi_{s!}\mathcal{S} \rightarrow (\pi_s \circ i)_!i_y^*\mathcal{S} \xrightarrow{[1]}. \tag{5.1.8}$$

Now $(\pi_s \circ i)$ is an isomorphism of varieties and $(\pi_s \circ j)$ is a trivial fibration with fibre \mathbb{A}_\circ^1 . The direct image with compact supports along such a fibration sends the constant sheaf $\underline{\mathbb{O}}$ to the (shifted and twisted) constant sheaf $\underline{\mathbb{O}}[-2](-1)$. Our assumptions on \mathcal{S} now guarantee that the long exact sequence of cohomology of the triangle (5.1.8) has only zeros in odd degree, and extensions of the constant sheaf $\underline{\mathbb{O}}_{X_{y,\circ}^s}(-k)$ in degree $2k$. Pulling back to X_\circ via $\pi_s^! \cong \pi_s^*[2](1)$ now yields (i).

(ii) Assume that $f = (s, t, \dots, r)$ is a reduced expression for w^{-1} and let $f' = (t, \dots, r)$; then f' is a reduced expression for $sw^{-1} = (ws)^{-1}$. By induction the support of $\mathcal{BS}_{f',\circ}^\circ$ is contained in $\overline{X_{ws,\circ}}$ and $i_{ws}^*\mathcal{BS}_{f',\circ}^\circ = \underline{\mathbb{O}}_{X_{ws,\circ}}[2\ell(ws)](\ell(ws))$. Fix $y \in W^s$. The distinguished triangle (5.1.8) shows that:

- (a) if $i_{y_s}^*\mathcal{S} = i_y^*\mathcal{S} = 0$ then $i_{y_s}^*(\pi_s^!\pi_{s!}\mathcal{S}) = i_y^*(\pi_s^!\pi_{s!}\mathcal{S}) = 0$;
- (b) if $i_{y_s}^*\mathcal{S} = 0$ and $i_y^*\mathcal{S} \cong \underline{\mathbb{O}}[m](r)$ then $i_{y_s}^*(\pi_s^!\pi_{s!}\mathcal{S}) \cong \underline{\mathbb{O}}[m+2](r+1)$.

It follows that the support of $\mathcal{BS}_{f,\circ}^\circ$ is contained in $\overline{X_{w,\circ}}$ and that $i_w^*\mathcal{BS}_{f,\circ}^\circ \cong \underline{\mathbb{O}}_{X_{w,\circ}}[2\ell(w)](\ell(w))$ as claimed. □

5.2 Algebraic description of direct and inverse image functors

Let \mathcal{P} be a projective generator of the category $\text{Perv}_{(B)}(X, \mathbb{E})$.

LEMMA 5.2.1. *The object $\pi_{s[!]} \mathcal{P}$ is a projective generator of $\text{Perv}_{(B)}(X^s, \mathbb{E})$.*

Proof. In this proof we assume that $\mathbb{E} = \mathbb{O}$; the case $\mathbb{E} = \mathbb{F}$ is similar. By the remarks after the definition of $\pi_{s[!]}$ (see §4.1), $\pi_{s[!]} \mathcal{P}$ is projective. By the remarks in the proof of Corollary 2.3.3, it is sufficient to prove that for any $v \in W^s$, there exists $n > 0$ such that $\pi_{s[!]} \mathcal{P}^{\oplus n}$ surjects to IC_v . Fix such a v . As \mathcal{P} is a projective generator there exist $n > 0$ and a surjection

$$f : \mathcal{P}^{\oplus n} \rightarrow \text{IC}_{v^s} \cong \pi_s^{[!]} \text{IC}_v.$$

By adjunction we obtain a morphism $g : \pi_{s[!]} \mathcal{P}^{\oplus n} \rightarrow \text{IC}_v$ such that f factors as

$$\mathcal{P}^{\oplus n} \rightarrow \pi_s^{[!]} \pi_{s[!]} \mathcal{P}^{\oplus n} \xrightarrow{\pi_s^{[!]} g} \pi_s^{[!]} \text{IC}_v.$$

Hence, $\pi_s^{[!]} g$ is surjective. As the functor $\pi_s^{[!]}$ is exact and fully faithful (see [BBD82, Proposition 4.2.5]), and hence kills no object, we deduce that g is surjective. □

We set

$$A := \text{End}_{\text{Perv}_{(B)}(X, \mathbb{E})}(\mathcal{P}), \quad A^s := \text{End}_{\text{Perv}_{(B)}(X^s, \mathbb{E})}(\pi_{s[!]} \mathcal{P}),$$

so that we have equivalences

$$\begin{aligned} \Pi &:= \text{Hom}(\mathcal{P}, -) : \text{Perv}_{(B)}(X, \mathbb{E}) \xrightarrow{\sim} \text{Modf} - A, \\ \Pi^s &:= \text{Hom}(\pi_{s[!]} \mathcal{P}, -) : \text{Perv}_{(B)}(X^s, \mathbb{E}) \xrightarrow{\sim} \text{Modf} - A^s. \end{aligned}$$

We denote similarly the induced equivalences between derived categories. There is a natural algebra morphism $A \rightarrow A^s$ induced by the functor $\pi_{s[1]}$, and hence associated functors

$$\text{Modf} - A \xrightleftharpoons[\text{res}]{(-) \otimes_A A^s} \text{Modf} - A^s. \tag{5.2.2}$$

LEMMA 5.2.3. *The following diagram commutes up to natural transformations.*

$$\begin{array}{ccc} \text{Perv}_{(B)}(X, \mathbb{E}) & \xrightarrow[\sim]{\Pi} & \text{Modf} - A \\ \pi_s^{[1]} \updownarrow \pi_{s[1]} & & \text{res} \updownarrow (-) \otimes_A A^s \\ \text{Perv}_{(B)}(X^s, \mathbb{E}) & \xrightarrow[\sim]{\Pi^s} & \text{Modf} - A^s \end{array}$$

Proof. The fact that

$$\text{res} \circ \text{Hom}(\pi_{s[1]} \mathcal{P}, -) \cong \text{Hom}(\mathcal{P}, -) \circ \pi_s^{[1]}$$

follows from the property that $\pi_{s[1]}$ is left adjoint to $\pi_s^{[1]}$. Then the second isomorphism of functors follows from the first one by adjunction. \square

As recalled above, the functor $\pi_s^{[1]}$ is exact; we denote similarly the functor induced between derived categories. The functor $\pi_{s[1]}$ is right exact; we denote its left derived functor by $L\pi_{s[1]}$. The values of this functor can be computed using projective resolutions (which exist thanks to Corollary 2.3.3). In particular, it follows from Corollary 2.3.4(ii) that $L\pi_{s[1]}$ restricts to a functor between *bounded* derived categories. Note that the functor $L\pi_{s[1]}$ is left adjoint to $\pi_s^{[1]}$. The same remarks apply to the functors (5.2.2).

We denote by

$$\begin{aligned} \text{real} &: \mathcal{D}^b \text{Perv}_{(B)}(X, \mathbb{E}) \rightarrow \mathcal{D}^b_{(B)}(X, \mathbb{E}), \\ \text{real}^s &: \mathcal{D}^b \text{Perv}_{(B)}(X^s, \mathbb{E}) \rightarrow \mathcal{D}^b_{(B)}(X^s, \mathbb{E}) \end{aligned}$$

the realization functors. Recall that these functors are equivalences of categories (see Corollary 2.3.4(i) if $\mathbb{E} = \mathbb{O}$, and [BGS96, Corollary 3.3.2] if $\mathbb{E} = \mathbb{F}$).

PROPOSITION 5.2.4. *The following diagram commutes up to natural transformations.*

$$\begin{array}{ccccc} \mathcal{D}^b_{(B)}(X, \mathbb{E}) & \xleftarrow[\sim]{\text{real}} & \mathcal{D}^b \text{Perv}_{(B)}(X, \mathbb{E}) & \xrightarrow[\sim]{\Pi} & \mathcal{D}^b(\text{Modf} - A) \\ \pi_s^{[1]}[-1] \updownarrow \pi_{s[1]} & & \pi_s^{[1]} \updownarrow L\pi_{s[1]} & & \text{res} \updownarrow (-) \otimes_A^L A^s \\ \mathcal{D}^b_{(B)}(X^s, \mathbb{E}) & \xleftarrow[\sim]{\text{real}^s} & \mathcal{D}^b \text{Perv}_{(B)}(X^s, \mathbb{E}) & \xrightarrow[\sim]{\Pi^s} & \mathcal{D}^b(\text{Modf} - A^s) \end{array}$$

Proof. Commutativity of the right-hand square follows from Lemma 5.2.3. Let us consider the left-hand square. The isomorphism

$$\text{real} \circ \pi_s^{[1]} \cong \pi_s^{[1]}[-1] \circ \text{real}^s$$

follows from [Bei87, Lemma A.7.1]. The second isomorphism follows by adjunction, since real and real^s are equivalences. \square

Now we assume that there exists \mathcal{P}_\circ in $\text{Perv}_{(B_\circ)}(X_\circ, \mathbb{E})$ such that $\mathcal{P} = \text{For}(\mathcal{P}_\circ)$. (This is possible thanks to Lemma 3.3.4 or §4.4.) Then the algebra A , respectively A^s , is endowed with a natural automorphism ϕ , respectively ϕ^s . By Lemma 3.3.7, the functors $\text{Hom}_{X, \mathbb{E}}(\mathcal{P}, -)$

and $\text{Hom}_{X^s, \mathbb{E}}(\pi_{s[!]} \mathcal{P}, -)$ induce natural equivalences of categories

$$\begin{aligned} \Pi_o &: \text{Perv}_{(B_o)}(X_o, \mathbb{E}) \xrightarrow{\sim} \text{Modf}-(A, \phi), \\ \Pi_o^s &: \text{Perv}_{(B_o)}(X_o^s, \mathbb{E}) \xrightarrow{\sim} \text{Modf}-(A^s, \phi^s). \end{aligned}$$

The morphism $A \rightarrow A^s$ commutes with the automorphisms ϕ and ϕ^s , so that we obtain functors

$$\begin{aligned} \text{res} &: \text{Modf}-(A^s, \phi^s) \rightarrow \text{Modf}-(A, \phi), \\ (-) \otimes_A A^s &: \text{Modf}-(A, \phi) \rightarrow \text{Modf}-(A^s, \phi^s) \end{aligned}$$

which are compatible with the functors (5.2.2) in the obvious sense. The proof of the following result is similar to that of Lemma 5.2.3.

LEMMA 5.2.5. *The following diagram commutes up to natural transformations.*

$$\begin{array}{ccc} \text{Perv}_{(B_o)}(X_o, \mathbb{E}) & \xrightarrow[\sim]{\Pi_o} & \text{Modf}-(A, \phi) \\ \pi_s^{[!]} \updownarrow \pi_{s[!]} & & \text{res} \updownarrow (-) \otimes_A A^s \\ \text{Perv}_{(B_o)}(X_o^s, \mathbb{E}) & \xrightarrow[\sim]{\Pi_o^s} & \text{Modf}-(A^s, \phi^s) \end{array}$$

We denote by

$$\begin{aligned} \text{real}_o &: \mathcal{D}^b \text{Perv}_{(B_o)}(X_o, \mathbb{E}) \rightarrow \mathcal{D}^b_{(B_o)}(X_o, \mathbb{E}), \\ \text{real}_o^s &: \mathcal{D}^b \text{Perv}_{(B_o)}(X_o^s, \mathbb{E}) \rightarrow \mathcal{D}^b_{(B_o)}(X_o^s, \mathbb{E}) \end{aligned}$$

the realization functors. These functors are equivalences by Proposition 3.4.1. It is easy to prove that the functor $(-) \otimes_A A^s$ admits a left derived functor, which can be computed using resolutions in $\text{Modf}-(A, \phi)$ whose terms have a flat underlying right A -module. This derived functor will be denoted by

$$(-) \overset{L}{\otimes}_A A^s : \mathcal{D}^b(\text{Modf}-(A, \phi)) \rightarrow \mathcal{D}^b(\text{Modf}-(A^s, \phi^s)).$$

Using Lemma 5.2.5, one deduces (or one proves directly) that the functor $\pi_{s[!]}$ also admits a left derived functor

$$L\pi_{s[!]} : \mathcal{D}^b \text{Perv}_{(B_o)}(X_o, \mathbb{E}) \rightarrow \mathcal{D}^b \text{Perv}_{(B_o)}(X_o^s, \mathbb{E}).$$

The proof of the following result is similar to that of Proposition 5.2.4.

PROPOSITION 5.2.6. *The following diagram commutes up to natural transformations.*

$$\begin{array}{ccccc} \mathcal{D}^b_{(B_o)}(X_o, \mathbb{E}) & \xleftarrow[\sim]{\text{real}_o} & \mathcal{D}^b \text{Perv}_{(B_o)}(X_o, \mathbb{E}) & \xrightarrow[\sim]{\Pi_o} & \mathcal{D}^b(\text{Modf}-(A, \phi)) \\ \pi_s^{[-1]} \updownarrow \pi_{s[1]} & & \pi_s^{[!]} \updownarrow L\pi_{s[!]} & & \text{res} \updownarrow (-) \overset{L}{\otimes}_A A^s \\ \mathcal{D}^b_{(B_o)}(X_o^s, \mathbb{E}) & \xleftarrow[\sim]{\text{real}_o^s} & \mathcal{D}^b \text{Perv}_{(B_o)}(X_o^s, \mathbb{E}) & \xrightarrow[\sim]{\Pi_o^s} & \mathcal{D}^b(\text{Modf}-(A^s, \phi^s)) \end{array}$$

5.3 Existence of a decomposable projective resolution

From now on we will have to work simultaneously with coefficients \mathbb{O} and \mathbb{F} , but to distinguish the two cases. We add superscripts or subscripts to indicate which coefficients we are working with.

We fix an object $\mathcal{P}_o^\mathbb{O}$ in $\text{Perv}_{(B_o)}(X_o, \mathbb{O})$ such that $\mathcal{P}^\mathbb{O} := \text{For}(\mathcal{P}_o^\mathbb{O})$ is a projective generator of the category $\text{Perv}_{(B)}(X, \mathbb{O})$. For simplicity, we also assume that there exists a

surjection $\mathcal{P}_\circ^\mathbb{O} \twoheadrightarrow \mathrm{IC}_{e,\circ}^\mathbb{O}$. Set $\mathcal{P}_\circ^\mathbb{F} := \mathbb{F}(\mathcal{P}_\circ^\mathbb{O})$ and $\mathcal{P}^\mathbb{F} := \mathrm{For}(\mathcal{P}_\circ^\mathbb{F}) \cong \mathbb{F}(\mathcal{P}^\mathbb{O})$. One can easily check that $\mathcal{P}^\mathbb{F}$ is a projective generator of the category $\mathrm{Perv}_{(B)}(X, \mathbb{F})$.

Set

$$A_\mathbb{O} := \mathrm{End}_{X,\mathbb{O}}(\mathcal{P}^\mathbb{O}), \quad A_\mathbb{O}^s := \mathrm{End}_{X^s,\mathbb{O}}(\pi_{s[!]} \mathcal{P}^\mathbb{O}),$$

$$A_\mathbb{F} := \mathrm{End}_{X,\mathbb{F}}(\mathcal{P}^\mathbb{F}), \quad A_\mathbb{F}^s := \mathrm{End}_{X^s,\mathbb{F}}(\pi_{s[!]} \mathcal{P}^\mathbb{F})$$

(for any simple reflection s). Note that by Lemma 5.2.1 and Proposition 2.4.1(iv), $A_\mathbb{O}$ and $A_\mathbb{O}^s$ are \mathbb{O} -free (for any s) and the natural morphism

$$\mathbb{F} \otimes_\mathbb{O} A_\mathbb{O} \rightarrow A_\mathbb{F} \tag{5.3.1}$$

is an isomorphism. We denote by ϕ , respectively ϕ^s , the automorphism induced by the Frobenius on $A_\mathbb{O}$ and $A_\mathbb{F}$, respectively on $A_\mathbb{O}^s$ and $A_\mathbb{F}^s$. In addition, we use the same notation as in § 5.2.

To prove an analogue of isomorphism (5.3.1) for A^s , we need some preparation. We denote by $\mathrm{Proj}(X, \mathbb{O})$, $\mathrm{Proj}(X^s, \mathbb{O})$ the categories of projective objects in $\mathrm{Perv}_{(B)}(X, \mathbb{O})$, $\mathrm{Perv}_{(B)}(X^s, \mathbb{O})$, and similarly for coefficients \mathbb{F} . Note that the functors $\pi_{s[!]}$ and $\mathbb{F} = \mathbb{F} \otimes_\mathbb{O}^L (-)$ send projective perverse sheaves to projective perverse sheaves (see § 4.1 and Proposition 2.4.1).

LEMMA 5.3.2. *There exists an isomorphism of functors*

$$\pi_{s[!]} \circ \mathbb{F} \cong \mathbb{F} \circ \pi_{s[!]} : \mathrm{Proj}(X, \mathbb{O}) \rightarrow \mathrm{Proj}(X^s, \mathbb{F}).$$

Proof. First, let \mathcal{M} be in $\mathcal{D}_{(B)}^b(X, \mathbb{O})$ with ${}^p\mathcal{H}^i(\mathcal{M}) = 0$ for $i > 0$. Consider the canonical truncation triangle

$$\mathcal{N} \rightarrow \mathcal{M} \rightarrow {}^p\mathcal{H}^0(\mathcal{M}) \xrightarrow{[1]}$$

and its modular reduction

$$\mathbb{F}\mathcal{N} \rightarrow \mathbb{F}\mathcal{M} \rightarrow \mathbb{F}({}^p\mathcal{H}^0(\mathcal{M})) \xrightarrow{[1]}.$$

As the functor \mathbb{F} is right exact, the second and third terms in the latter triangle have canonically isomorphic 0th perverse cohomology. Hence there exists a functorial morphism

$$\mathbb{F}({}^p\mathcal{H}^0(\mathcal{M})) \rightarrow {}^p\mathcal{H}^0(\mathbb{F}({}^p\mathcal{H}^0(\mathcal{M}))) \cong {}^p\mathcal{H}^0(\mathbb{F}\mathcal{M}),$$

which is an isomorphism if the left-hand side is a perverse sheaf.

Now, if Q is in $\mathrm{Proj}(X, \mathbb{O})$ we obtain isomorphisms

$$\mathbb{F}(\pi_{s[!]} Q) := \mathbb{F}({}^p\mathcal{H}^0(\pi_{s!} Q[1])) \xrightarrow{\sim} {}^p\mathcal{H}^0(\mathbb{F}(\pi_{s!} Q)[1]) \cong {}^p\mathcal{H}^0(\pi_{s!}(\mathbb{F}Q)[1]) = \pi_{s[!]}(\mathbb{F}Q),$$

which finishes the proof. □

PROPOSITION 5.3.3. *There exists an isomorphism of \mathbb{F} -algebras $\mathbb{F} \otimes_\mathbb{O} A_\mathbb{O}^s \cong A_\mathbb{F}^s$ such that the diagram*

$$\begin{array}{ccc} \mathbb{F} \otimes_\mathbb{O} A_\mathbb{O} & \longrightarrow & \mathbb{F} \otimes_\mathbb{O} A_\mathbb{O}^s \\ \text{(5.3.1)} \downarrow \wr & & \downarrow \wr \\ A_\mathbb{F} & \longrightarrow & A_\mathbb{F}^s \end{array}$$

commutes, where horizontal morphisms are induced by the functor $\pi_{s[!]}$.

Proof. Our isomorphism can be defined as the composition

$$\mathbb{F} \otimes_\mathbb{O} \mathrm{End}_{\mathrm{Perv}_{(B)}(X,\mathbb{O})}(\pi_{s[!]} \mathcal{P}^\mathbb{O}) \xrightarrow{\sim} \mathrm{End}_{\mathrm{Perv}_{(B)}(X,\mathbb{F})}(\mathbb{F}(\pi_{s[!]} \mathcal{P}^\mathbb{O})) \xrightarrow{\sim} \mathrm{End}_{\mathrm{Perv}_{(B)}(X,\mathbb{F})}(\pi_{s[!]} \mathcal{P}^\mathbb{F})$$

where the first isomorphism follows from Proposition 2.4.1(iv), and the second one follows from Lemma 5.3.2. Commutativity of the diagram follows from functoriality. \square

We consider the following conditions:

$$A_{\mathbb{O}} \text{ is } q\text{-decomposable, and } A_{\mathbb{O}}^s \text{ is } q\text{-decomposable for any } s \in S. \tag{5.3.4}$$

Recall that, by Propositions 4.4.2 and 4.5.4, if the order of q in \mathbb{F} is strictly bigger than $2\ell(w_0)$ then one can choose $\mathcal{P}_{\circ}^{\mathbb{O}}$ such that condition (5.3.4) is satisfied. However, most of our results below hold under this condition, independently of the proofs in §4.

LEMMA 5.3.5. *Assume that condition (5.3.4) is satisfied. Then the $(A_{\mathbb{O}}, \phi)$ -module*

$$\text{Hom}_{X, \mathbb{O}}(\mathcal{P}^{\mathbb{O}}, \text{IC}_e^{\mathbb{O}})$$

is \mathbb{O} -free and q -decomposable. Moreover, the natural morphism

$$\mathbb{F} \otimes_{\mathbb{O}} \text{Hom}_{X, \mathbb{O}}(\mathcal{P}^{\mathbb{O}}, \text{IC}_e^{\mathbb{O}}) \rightarrow \text{Hom}_{X, \mathbb{F}}(\mathcal{P}^{\mathbb{F}}, \text{IC}_e^{\mathbb{F}})$$

is an isomorphism.

Proof. The fact that $\text{Hom}_{X, \mathbb{O}}(\mathcal{P}^{\mathbb{O}}, \text{IC}_e^{\mathbb{O}})$ is \mathbb{O} -free and the final claim follow from Proposition 2.4.1(ii) because $\text{IC}_e^{\mathbb{O}} = \Delta_e$. By our choice of $\mathcal{P}_{\circ}^{\mathbb{O}}$ we have a surjection of $(A_{\mathbb{O}}, \phi)$ -modules

$$A_{\mathbb{O}} \rightarrow \text{Hom}_{X, \mathbb{O}}(\mathcal{P}^{\mathbb{O}}, \text{IC}_e^{\mathbb{O}}).$$

Hence, the fact that $\text{Hom}_{X, \mathbb{O}}(\mathcal{P}^{\mathbb{O}}, \text{IC}_e^{\mathbb{O}})$ is q -decomposable follows from Lemma 3.1.3. \square

Recall the objects $\mathcal{BS}_{\circ}^{\mathbb{O}}$, $\mathcal{BS}_{\circ}^{\mathbb{F}}$ defined in §5.1. By definition we have an isomorphism $\mathbb{F}(\mathcal{BS}_{\circ}^{\mathbb{O}}) \cong \mathcal{BS}_{\circ}^{\mathbb{F}}$. Recall also the category $\text{Modf}_{\text{dec}}-(A_{\mathbb{O}}, \phi)$ defined in §3.1. We denote by

$$\text{Modf}_{\text{dec}}^{\text{proj}}-(A_{\mathbb{O}}, \phi) \subset \text{Modf}_{\text{dec}}-(A_{\mathbb{O}}, \phi)$$

the full additive subcategory with objects (P, ϕ_P) such that P is a projective $A_{\mathbb{O}}$ -module. Note that if (P, ϕ_P) is in $\text{Modf}_{\text{dec}}^{\text{proj}}-(A_{\mathbb{O}}, \phi)$, then P is \mathbb{O} -free and $\mathbb{F} \otimes_{\mathbb{O}} P$ is a projective $A_{\mathbb{F}}$ -module, endowed with a natural automorphism induced by ϕ_P . We denote by $\mathbb{F}(P, \phi_P)$ the corresponding object of $\text{Modf}-(A_{\mathbb{F}}, \phi)$.

The following result is the technical key to our proof of formality.

PROPOSITION 5.3.6. *Assume that condition (5.3.4) is satisfied. Then there exists a bounded complex M_{\circ}^{\bullet} of objects of $\text{Modf}_{\text{dec}}^{\text{proj}}-(A_{\mathbb{O}}, \phi)$ which satisfies the following conditions.*

(i) *The image of M_{\circ}^{\bullet} in $\mathcal{D}^b(\text{Modf}-(A_{\mathbb{O}}, \phi))$ coincides (up to isomorphism) with the image of $\mathcal{BS}_{\circ}^{\mathbb{O}}$ under the equivalence*

$$\Pi_{\circ}^{\mathbb{O}} \circ (\text{real}_{\circ}^{\mathbb{O}})^{-1} : \mathcal{D}_{(B_{\circ})}^b(X_{\circ}, \mathbb{O}) \xrightarrow{\sim} \mathcal{D}^b(\text{Modf}-(A_{\mathbb{O}}, \phi)).$$

(ii) *The image of $\mathbb{F}(M_{\circ}^{\bullet})$ in $\mathcal{D}^b(\text{Modf}-(A_{\mathbb{F}}, \phi))$ coincides (up to isomorphism) with the image of $\mathcal{BS}_{\circ}^{\mathbb{F}}$ under the equivalence*

$$\Pi_{\circ}^{\mathbb{F}} \circ (\text{real}_{\circ}^{\mathbb{F}})^{-1} : \mathcal{D}_{(B_{\circ})}^b(X_{\circ}, \mathbb{F}) \xrightarrow{\sim} \mathcal{D}^b(\text{Modf}-(A_{\mathbb{F}}, \phi)).$$

Proof. Let $f \in F$, and write $f = (s, t, \dots, r)$. By Proposition 5.2.6, the image of the object $\mathcal{BS}_{f, \circ}^{\mathbb{O}}$ of $\mathcal{D}_{(B_{\circ})}^b(X_{\circ}, \mathbb{O})$ under $\Pi_{\circ}^{\mathbb{O}} \circ (\text{real}_{\circ}^{\mathbb{O}})^{-1}$ is isomorphic to the object

$$\Pi_{\circ}^{\mathbb{O}}(\text{IC}_{e, \circ}^{\mathbb{O}}) \overset{L}{\otimes}_{A_{\mathbb{O}}} A_{\mathbb{O}}^r \overset{L}{\otimes}_{A_{\mathbb{O}}} \dots \overset{L}{\otimes}_{A_{\mathbb{O}}} A_{\mathbb{O}}^t \overset{L}{\otimes}_{A_{\mathbb{O}}} A_{\mathbb{O}}^s \tag{5.3.7}$$

of $\mathcal{D}^b(\text{Modf}-(A, \phi))$, where for any u , $A_\mathbb{O}^u$ is considered as an $A_\mathbb{O}$ -bimodule with compatible automorphism. By Lemma 5.3.5, $\Pi_\mathbb{O}^\mathbb{O}(\text{IC}_{e,\mathbb{O}}^\mathbb{O}) = \text{Hom}_{X,\mathbb{O}}(\mathcal{P}^\mathbb{O}, \text{IC}_e^\mathbb{O})$ is a q -decomposable, \mathbb{O} -free (right) $(A_\mathbb{O}, \phi)$ -module.

For any u that appears in f one can construct a resolution

$$Q_u^\bullet \xrightarrow{\text{qis}} A_\mathbb{O}^u$$

of $A_\mathbb{O}^u$ by objects of $\text{Modf}_{\text{dec}}^{\text{proj}}-(A_\mathbb{O} \otimes_\mathbb{O} A_\mathbb{O}^{\text{op}}, \phi \otimes \phi)$ which is concentrated in non-positive degrees (but possibly infinite). Indeed, by Lemma 3.1.8 and condition (5.3.4) there exist an object Q^0 in $\text{Modf}_{\text{dec}}^{\text{proj}}-(A_\mathbb{O} \otimes_\mathbb{O} A_\mathbb{O}^{\text{op}}, \phi \otimes \phi)$ and a surjection $Q^0 \twoheadrightarrow A_\mathbb{O}^u$. By Lemma 3.1.3 the kernel of this surjection is q -decomposable, and it is \mathbb{O} -free of finite rank, hence we can repeat the argument.

Then one can define the complex

$$'M_{f,\mathbb{O}}^\bullet := \Pi_\mathbb{O}^\mathbb{O}(\text{IC}_{e,\mathbb{O}}^\mathbb{O}) \otimes_{A_\mathbb{O}} Q_r^\bullet \otimes_{A_\mathbb{O}} \cdots \otimes_{A_\mathbb{O}} Q_t^\bullet \otimes_{A_\mathbb{O}} Q_s^\bullet.$$

Since a projective right $A_\mathbb{O} \otimes_\mathbb{O} A_\mathbb{O}^{\text{op}}$ -module is also projective as a left $A_\mathbb{O}$ -module, the image of $'M_{f,\mathbb{O}}^\bullet$ in $\mathcal{D}^b(\text{Modf}-(A_\mathbb{O}, \phi))$ is (5.3.7), and hence it is isomorphic to $\Pi_\mathbb{O}^\mathbb{O} \circ (\text{real}_\mathbb{O}^\mathbb{O})^{-1}(\mathcal{BS}_{f,\mathbb{O}}^\mathbb{O})$.

By Lemma 5.3.8 below, the underlying $A_\mathbb{O}$ -module of each $'M_{f,\mathbb{O}}^k$ is projective, and each $'M_{f,\mathbb{O}}^k$ is q -decomposable by Lemma 3.1.3. Indeed, it is a finite direct sum of torsion-free quotients of (\mathbb{O}, ϕ) -modules of the form

$$\Pi_\mathbb{O}^\mathbb{O}(\text{IC}_{e,\mathbb{O}}^\mathbb{O}) \otimes_\mathbb{O} Q_r^{k_r} \otimes_\mathbb{O} \cdots \otimes_\mathbb{O} Q_t^{k_t} \otimes_\mathbb{O} Q_s^{k_s},$$

the latter being \mathbb{O} -free and q -decomposable.

We claim that one can choose $k \ll 0$ such that $M_{f,\mathbb{O}}^\bullet := \tau^{\geq k}('M_{f,\mathbb{O}}^\bullet)$ is still a complex of objects of $\text{Modf}_{\text{dec}}^{\text{proj}}-(A_\mathbb{O}, \phi)$ whose image in the derived category is (5.3.7). Indeed, the complex $'M_{f,\mathbb{O}}^\bullet$ has bounded cohomology, hence if $k \ll 0$ the natural morphism $'M_{f,\mathbb{O}}^\bullet \rightarrow M_{f,\mathbb{O}}^\bullet$ is a quasi-isomorphism. As the category $\text{Modf}-A_\mathbb{O}$ has finite projective dimension, if $k \ll 0$ then the components of $M_{f,\mathbb{O}}^\bullet$ still have an image in $\text{Modf}-A_\mathbb{O}$ which is projective. Finally (again if $k \ll 0$), the only component of $M_{f,\mathbb{O}}^\bullet$ which is neither 0 nor a component of $'M_{f,\mathbb{O}}^\bullet$ is q -decomposable by Lemma 3.1.3.

Set

$$M_\mathbb{O}^\bullet := \bigoplus_{f \in F} M_{f,\mathbb{O}}^\bullet.$$

Then, by the remarks above, this complex satisfies condition (i). To prove that it satisfies condition (ii), one observes that the morphisms $Q_u^\bullet \rightarrow A_\mathbb{O}^u$ considered above are quasi-isomorphisms of complexes of free \mathbb{O} -modules; hence the induced morphisms $(\mathbb{F} \otimes_\mathbb{O} Q_u^\bullet) \rightarrow (\mathbb{F} \otimes_\mathbb{O} A_\mathbb{O}^u)$ are also quasi-isomorphisms. By the same argument, the morphisms $\mathbb{F}('M_{f,\mathbb{O}}^\bullet) \rightarrow \mathbb{F}(M_{f,\mathbb{O}}^\bullet)$ are quasi-isomorphisms. Finally, by Lemma 5.3.5 we have $\mathbb{F} \otimes_\mathbb{O} \Pi_\mathbb{O}^\mathbb{O}(\text{IC}_{e,\mathbb{O}}^\mathbb{O}) \cong \Pi_\mathbb{O}^\mathbb{F}(\text{IC}_{e,\mathbb{O}}^\mathbb{F})$ and by Proposition 5.3.3 we have $\mathbb{F} \otimes_\mathbb{O} A_\mathbb{O}^u \cong A_\mathbb{F}^u$. Hence, the same arguments as for \mathbb{O} show that $M_\mathbb{O}^\bullet$ also satisfies condition (ii). □

LEMMA 5.3.8. *Let R be any \mathbb{O} -algebra which is finitely generated and free over \mathbb{O} .*

Let Q be a projective (R, R) -bimodule and M be a right R -module which is \mathbb{O} -free. Then the right R -module $M \otimes_R Q$ is projective (hence also \mathbb{O} -free).

Proof. It suffices to prove that $M \otimes_R (R \otimes_\mathbb{O} R)$ is projective over R . However, we have

$$M \otimes_R (R \otimes_\mathbb{O} R) \cong M \otimes_\mathbb{O} R,$$

hence this fact is clear from our hypothesis on M . □

5.4 Dg-algebras of finite global dimension

Let R be a graded algebra, considered as a dg-algebra with differential $d = 0$. We assume that R is noetherian as an algebra, and that the category of finitely generated graded R -modules has finite projective dimension. There are *a priori* several ways to define the derived category of finitely generated right R -dg-modules. The goal of this subsection is to show that, under our assumptions, all of them coincide.

More precisely, we denote by D_1 the category obtained from the homotopy category of right R -dg-modules which are finitely generated as R -modules by inverting quasi-isomorphisms. We denote by D_2 the full subcategory of $\text{dgDer-}R$ whose objects are R -dg-modules which are isomorphic (in $\text{dgDer-}R$) to an R -dg-module which is finitely generated over R . (Note that it is not clear *a priori* that D_2 is a triangulated subcategory of $\text{dgDer-}R$.) We denote by D_3 the strictly full triangulated subcategory of $\text{dgDer-}R$ generated by finitely generated projective graded R -modules (considered as dg-modules with trivial differential). Finally, we denote by D_4 the full subcategory of $\text{dgDer-}R$ whose objects have their cohomology finitely generated over R . We have inclusions

$$D_3 \subset D_4, \quad D_2 \subset D_4$$

and a natural essentially surjective functor

$$D_1 \rightarrow D_2. \tag{5.4.1}$$

PROPOSITION 5.4.2. *We have $D_2 = D_3 = D_4$, and (5.4.1) is an equivalence of categories.*

Below we will denote by $\text{dgDerf-}R$ any of the equivalent triangulated categories D_i ($i = 1, 2, 3, 4$).

The main step in the proof of Proposition 5.4.2 is the following lemma.

LEMMA 5.4.3. *We have $D_4 \subset D_3$.*

Proof. We will prove that any object M in D_4 is in D_3 by induction on the projective dimension of $H^\bullet(M)$.

First, assume that $H^\bullet(M)$ has projective dimension 0, i.e. that it is projective. We claim that in this case there exists a quasi-isomorphism of R -dg-modules

$$H^\bullet(M) \xrightarrow{\text{qis}} M$$

where $H^\bullet(M)$ is considered as a dg-module with trivial differential. Indeed, as $H^\bullet(M)$ is projective, there exists a splitting $s : H^\bullet(M) \rightarrow \ker(d_M)$ of the natural morphism of graded R -modules $\ker(d_M) \rightarrow H^\bullet(M)$. Then the composition $H^\bullet(M) \rightarrow \ker(d_M) \hookrightarrow M$ is the desired quasi-isomorphism. As $H^\bullet(M)$ is in D_3 , we deduce that M is also in D_3 .

Now, assume that $H^\bullet(M)$ has projective dimension d . Choose a minimal length projective resolution

$$P^d \hookrightarrow \dots \rightarrow P^0 \xrightarrow{a} H^\bullet(M)$$

as a graded R -module. By the same argument as before, there exists a morphism of R -dg-modules $P^0 \rightarrow M$ which induces a in cohomology. We denote by N the cocone of this morphism, so that we have a distinguished triangle

$$N \rightarrow P^0 \rightarrow M \xrightarrow{[1]} .$$

Then $H^\bullet(N) \cong \ker(a)$ has projective dimension $d - 1$, hence N belongs to D_3 by induction. We deduce that M is in D_3 also. □

Proof of Proposition 5.4.2. By Lemma 5.4.3 we have $D_3 = D_4$. Now, as morphisms in $R\text{-dgDer}$ between projective graded R -modules (considered as dg-modules with $d = 0$) are simply morphisms of graded R -modules (because such objects are K -projective in the sense of [BL94, Definition 10.12.2.1]), one easily checks that $D_3 \subset D_2$. We deduce that $D_2 = D_3 = D_4$.

Now consider the functor $D_1 \rightarrow D_2 = D_3$. The same proof as in Lemma 5.4.3 shows that the category D_1 is also generated, as a triangulated category, by finitely generated projective graded R -modules (with trivial differential). Hence, it is enough to check that morphisms between such objects in D_1 and D_2 coincide; this is obvious. \square

5.5 Proof of formality

The following result is well known, and is usually attributed to Deligne, see [Del80, DGMS75].

LEMMA 5.5.1. *Let R^\bullet be a dg-algebra which is endowed with an additional \mathbb{Z} -grading*

$$R^i = \bigoplus_{j \in \mathbb{Z}} R^{i,j}$$

such that $d_R(R^{i,j}) \subset R^{i+1,j}$. Assume that for any $i \in \mathbb{Z}$ the cohomology $H^i(R^\bullet)$ is concentrated in degree $j = i$ (for this additional grading). Then R^\bullet is formal. More precisely, there exists a dg-subalgebra $R_{\triangleright}^\bullet \subset R^\bullet$ and quasi-isomorphisms

$$R^\bullet \xleftarrow{\text{qis}} R_{\triangleright}^\bullet \xrightarrow{\text{qis}} H^\bullet(R^\bullet),$$

where $H^\bullet(R^\bullet)$ is considered as a dg-algebra with trivial differential.

Proof. Denote by $d_j^i : R^{i,j} \rightarrow R^{i+1,j}$ the component of the differential in bidegree (i, j) . For any $i \in \mathbb{Z}$, set

$$R_{\triangleright}^i := \left(\bigoplus_{j > i} R^{i,j} \right) \oplus \ker(d_i^i).$$

Then $R_{\triangleright}^\bullet$ is a dg-subalgebra of R^\bullet . Moreover, the projection $R_{\triangleright}^i \rightarrow \ker(d_i^i)/\text{im}(d_i^{i-1})$ induces a quasi-isomorphism of dg-algebras $R_{\triangleright}^\bullet \xrightarrow{\text{qis}} H^\bullet(R^\bullet)$. \square

From now on we assume that condition (5.3.4) is satisfied. Let $M_\mathbb{O}^\bullet$ be as in Proposition 5.3.6, and let M^\bullet be the underlying complex of projective right $A_\mathbb{O}$ -modules. Consider the dg-algebras

$$E_\mathbb{O}^\bullet := \underline{\text{Hom}}_{A_\mathbb{O}}^\bullet(M^\bullet, M^\bullet), \quad E_\mathbb{F}^\bullet := \underline{\text{Hom}}_{A_\mathbb{F}}^\bullet(\mathbb{F} \otimes_\mathbb{O} M^\bullet, \mathbb{F} \otimes_\mathbb{O} M^\bullet)$$

of endomorphism complexes. Note that both of these dg-algebras are bounded, that $E_\mathbb{O}^\bullet$ is a finite-rank free \mathbb{O} -module, and that we have a natural isomorphism of dg-algebras

$$\mathbb{F} \otimes_\mathbb{O} E_\mathbb{O}^\bullet \cong E_\mathbb{F}^\bullet.$$

By the same considerations as in Remark 3.4.2, $E_\mathbb{O}^\bullet$ is endowed with a natural automorphism, so that it can be considered as an (\mathbb{O}, ϕ) -dg-algebra. Similarly, by the constructions of § 3.3 the graded \mathbb{O} -algebra $E_\mathbb{O}$ can be naturally considered as an (\mathbb{O}, ϕ) -algebra.

LEMMA 5.5.2. *There exist natural isomorphisms of graded algebras*

$$H^\bullet(E_\mathbb{O}^\bullet) \cong E_\mathbb{O}, \quad H^\bullet(E_\mathbb{F}^\bullet) \cong E_\mathbb{F}.$$

Moreover, the first isomorphism is an isomorphism of (\mathbb{O}, ϕ) -algebras.

Proof. By definition of $E_{\mathbb{O}}^{\bullet}$ we have an isomorphism of graded algebras

$$H^{\bullet}(E_{\mathbb{O}}^{\bullet}) \cong \text{Ext}_{\mathcal{D}^b(\text{Modf}-A_{\mathbb{O}})}^{\bullet}(M^{\bullet}, M^{\bullet}).$$

Using condition (i) in Proposition 5.3.6, we have an isomorphism

$$\text{real}^{\mathbb{O}} \circ (\Pi^{\mathbb{O}})^{-1}(M^{\bullet}) \cong \mathcal{BS}^{\mathbb{O}}.$$

We deduce the first isomorphism. The compatibility with automorphisms follows from Remark 3.4.2. The second isomorphism can be proved similarly. \square

LEMMA 5.5.3. *The algebras $E_{\mathbb{O}}$ and $E_{\mathbb{F}}$ have finite global dimension.*

Proof. We will prove in Corollary 6.4.2 below (using [Soe00]) that $E_{\mathbb{F}}$ has finite global dimension. Let us deduce the same property for $E_{\mathbb{O}}$. We claim that for M, N in $\mathcal{D}^b(\text{Modf}-E_{\mathbb{O}})$ there exists a natural isomorphism in the derived category of \mathbb{F} -vector spaces

$$\mathbb{F} \overset{L}{\otimes}_{\mathbb{O}} R \text{Hom}_{-E_{\mathbb{O}}}(M, N) \cong R \text{Hom}_{-E_{\mathbb{F}}}(\mathbb{F} \overset{L}{\otimes}_{\mathbb{O}} M, \mathbb{F} \overset{L}{\otimes}_{\mathbb{O}} N) \tag{5.5.4}$$

where $R \text{Hom}_{-E_{\mathbb{O}}}(M, N)$ is considered as an object in $\mathcal{D}(\text{Modf}-\mathbb{O})$. Indeed, one can assume that M is a bounded above complex of finitely generated projective $E_{\mathbb{O}}$ -modules, and then reduce the claim to the case $M = E_{\mathbb{O}}$, which is obvious. As \mathbb{O} has global dimension 1, any object of $\mathcal{D}(\text{Modf}-\mathbb{O})$ is isomorphic to its cohomology. Moreover, if M is in $\text{Modf}-\mathbb{O}$ then $M \neq 0$ implies $\mathbb{F} \overset{L}{\otimes}_{\mathbb{O}} M \neq 0$. Using these remarks, isomorphism (5.5.4) and the fact that $E_{\mathbb{F}}$ has finite global dimension, one easily checks that the same property holds for $E_{\mathbb{O}}$. \square

Using Lemma 5.5.3, one can define the categories $\text{dgDerf}-E_{\mathbb{O}}$ and $\text{dgDerf}-E_{\mathbb{F}}$ as in § 5.4.

PROPOSITION 5.5.5. *The dg-algebras $E_{\mathbb{O}}^{\bullet}$ and $E_{\mathbb{F}}^{\bullet}$ are formal. More precisely, there exists a dg-subalgebra $E_{\triangleright}^{\bullet} \subset E_{\mathbb{O}}^{\bullet}$ and quasi-isomorphisms*

$$E_{\mathbb{O}}^{\bullet} \xleftarrow{\text{qis}} E_{\triangleright}^{\bullet} \xrightarrow{\text{qis}} E_{\mathbb{O}} \tag{5.5.6}$$

such that the diagram

$$E_{\mathbb{F}}^{\bullet} \xleftarrow{\text{qis}} \mathbb{F} \otimes_{\mathbb{O}} E_{\triangleright}^{\bullet} \xrightarrow{\text{qis}} \mathbb{F} \otimes_{\mathbb{O}} E_{\mathbb{O}}$$

obtained by modular reduction also consists of quasi-isomorphisms.

Proof. We claim that the (\mathbb{O}, ϕ) -module $E_{\mathbb{O}}^{\bullet}$ is q -decomposable. Indeed, as $M_{\mathbb{O}}^{\bullet}$ is q -decomposable (see Proposition 5.3.6), the same holds for the \mathbb{O} -free (\mathbb{O}, ϕ) -module $F^{\bullet} := \underline{\text{Hom}}_{\mathbb{O}}^{\bullet}(M^{\bullet}, M^{\bullet})$. By definition $E_{\mathbb{O}}^{\bullet}$ is a sub- (\mathbb{O}, ϕ) -module of F^{\bullet} , and the quotient $F^{\bullet}/E_{\mathbb{O}}^{\bullet}$ has no π -torsion. We deduce from Lemma 3.1.3 that $E_{\mathbb{O}}^{\bullet}$ is indeed q -decomposable.

For any $i, j \in \mathbb{Z}$ we set

$$E_{j, \mathbb{O}}^i := \{e \in E^i \mid (\phi - q^{-j/2})^m \cdot e = 0 \text{ for } m \gg 0\}.$$

By q -decomposability this defines a grading on E^{\bullet} (which is concentrated in even degrees). Hence, using Lemma 5.5.2 and Theorem 5.1.5, we are in the situation of Lemma 5.5.1. We denote by $E_{\triangleright}^{\bullet}$ the subalgebra constructed in the proof of this lemma, so that we have a diagram (5.5.6). All the dg-algebras in this diagram are \mathbb{O} -free; indeed, $E_{\triangleright}^{\bullet}$ has no π -torsion since it is a submodule of the free \mathbb{O} -module $E_{\mathbb{O}}^{\bullet}$, and $E_{\mathbb{O}}$ is free by Theorem 5.1.5. Hence the diagram obtained by modular reduction also consists of quasi-isomorphisms. \square

Remark 5.5.7. In particular, it follows from Proposition 5.5.5 and Lemma 5.5.2 that the natural morphism $\mathbb{F} \otimes_{\mathbb{O}} E_{\mathbb{O}} \rightarrow E_{\mathbb{F}}$ is an isomorphism. This can also be proved directly using [JMW09, equation (2.13)] and the fact that $\mathcal{BS}_{\mathbb{O}}$ is parity.

Finally we can prove the main result of Part II.

THEOREM 5.5.8. *Assume that condition (5.3.4) is satisfied.*

There exist equivalences of triangulated categories

$$\mathcal{D}_{(B)}^b(X, \mathbb{O}) \cong \text{dgDerf} - E_{\mathbb{O}}, \quad \mathcal{D}_{(B)}^b(X, \mathbb{F}) \cong \text{dgDerf} - E_{\mathbb{F}}$$

where $E_{\mathbb{O}}$ and $E_{\mathbb{F}}$ are considered as dg-algebras with their natural grading and trivial differential.

Proof. We treat only the case of \mathbb{O} ; the case of \mathbb{F} is similar. As explained in § 5.2, there exists an equivalence of categories

$$\mathcal{D}_{(B)}^b(X, \mathbb{O}) \cong \mathcal{D}^b(\text{Mod} - A_{\mathbb{O}}).$$

Moreover, by Lemma 5.1.4 the category $\mathcal{D}_{(B)}^b(X, \mathbb{O})$ is generated (as a triangulated category) by the objects $\{\mathcal{BS}_f^{\mathbb{O}}, f \in F\}$. By construction (see Proposition 5.3.6), \mathcal{BS}_f corresponds, under this equivalence, to the complex M_f^{\bullet} of projective right $A_{\mathbb{O}}$ -modules. Hence, with the notation of § 1.3 we obtain

$$\mathcal{D}_{(B)}^b(X, \mathbb{O}) \cong \langle M_f^{\bullet}, f \in F \rangle_{\Delta} \subset \mathcal{D}(\text{Mod} - A_{\mathbb{O}}).$$

By the result recalled in § 1.3, the right-hand side is equivalent to

$$\langle 1_f E_{\mathbb{O}}^{\bullet}, f \in F \rangle_{\Delta} \subset \text{dgDer} - E_{\mathbb{O}}^{\bullet}$$

where 1_f is the idempotent of $E_{\mathbb{O}}^{\bullet}$ given by the projection to M_f^{\bullet} .

Now it follows from Proposition 5.5.5 that there is an equivalence of triangulated categories

$$\text{dgDer} - E_{\mathbb{O}}^{\bullet} \cong \text{dgDer} - E_{\mathbb{O}}.$$

This equivalence sends $1_f E_{\mathbb{O}}^{\bullet}$ to $e_f E_{\mathbb{O}}$, where $e_f \in E_{\mathbb{O}}$ is defined in § 5.1. We deduce an equivalence

$$\mathcal{D}_{(B)}^b(X, \mathbb{O}) \cong \langle e_f E_{\mathbb{O}}, f \in F \rangle_{\Delta} \subset \text{dgDer} - E_{\mathbb{O}}.$$

Finally, it is not difficult to check (using the category D_3 of § 5.4) that the right-hand side is equivalent to $\text{dgDerf} - E_{\mathbb{O}}$. □

5.6 Parabolic case

It is not difficult to generalize the results of § 5.5 to parabolic flag varieties. Here, for future reference, we explain the case of the varieties X^s for $s \in S$. First, we remark that by an obvious analogue of Lemma 5.5.3 the rings $E_{\mathbb{O}}^s$ and $E_{\mathbb{F}}^s$ have finite global dimension, so that we can consider the categories $\text{dgDerf} - E_{\mathbb{O}}^s$ and $\text{dgDerf} - E_{\mathbb{F}}^s$.

THEOREM 5.6.1. *Let $s \in S$, and assume that condition (5.3.4) is satisfied.*

There exist equivalences of triangulated categories

$$\mathcal{D}_{(B)}^b(X^s, \mathbb{O}) \cong \text{dgDerf} - E_{\mathbb{O}}^s, \quad \mathcal{D}_{(B)}^b(X^s, \mathbb{F}) \cong \text{dgDerf} - E_{\mathbb{F}}^s$$

where $E_{\mathbb{O}}^s$ and $E_{\mathbb{F}}^s$ are considered as dg-algebras with their natural grading and trivial differential.

Proof. First we treat the case of \mathbb{O} . Recall the resolution M_{\circ}^{\bullet} constructed in the proof of Proposition 5.3.6. Set

$$M_{s,\circ}^{\bullet} := M_{\circ}^{\bullet} \otimes_{A_{\mathbb{O}}} A_{\mathbb{O}}^s$$

and denote by M_s^\bullet the complex obtained by forgetting the automorphism ϕ . Then by construction $M_{s,\circ}^\bullet$ is a complex of objects of $\text{Modf}_{\text{dec}}^{\text{proj}}(A_\circ^s, \phi^s)$, and by Proposition 5.2.6 its image in $\mathcal{D}^b(\text{Modf}(A_\circ^s, \phi^s))$ is $\Pi_\circ^{s,\circ} \circ (\text{real}_\circ^{s,\circ})^{-1}(\pi_{s!}\mathcal{BS}_\circ^{\circ}[1])$. Then we consider the dg-algebra

$$E_\circ^{s,\bullet} := \underline{\text{Hom}}_{-A_\circ^s}^\bullet(M_s^\bullet, M_s^\bullet).$$

By construction it is endowed with an automorphism ϕ^s .

The same argument as for Lemma 5.5.2 shows that we have an isomorphism of (\circ, ϕ) -algebras $\mathbf{H}^\bullet(E_\circ^{s,\bullet}) \cong E_\circ^s$. Then the same argument as for Proposition 5.5.5 shows that there exists a sub-dg-algebra $E_{\triangleright}^{s,\bullet} \subset E_\circ^{s,\bullet}$ and quasi-isomorphisms.

$$E_\circ^{s,\bullet} \xleftarrow{\text{qis}} E_{\triangleright}^{s,\bullet} \xrightarrow{\text{qis}} E_\circ^s. \tag{5.6.2}$$

The rest of the proof is similar to that of Theorem 5.5.8, replacing E_\circ^\bullet by $E_\circ^{s,\bullet}$.

The case of \mathbb{F} is similar, using the dg-algebra

$$E_{\mathbb{F}}^{s,\bullet} := \underline{\text{Hom}}_{-A_{\mathbb{F}}^s}^\bullet(\mathbb{F} \otimes_\circ M_s^\bullet, \mathbb{F} \otimes_\circ M_s^\bullet)$$

and the quasi-isomorphisms

$$E_{\mathbb{F}}^{s,\bullet} \xleftarrow{\text{qis}} \mathbb{F} \otimes_\circ E_{\triangleright}^{s,\bullet} \xrightarrow{\text{qis}} \mathbb{F} \otimes_\circ E_\circ^s$$

obtained from (5.6.2) by modular reduction. □

Let us study the compatibility of the equivalences in Theorems 5.5.8 and 5.6.1. We set $\mathbb{E} = \circ$ or \mathbb{F} . For any $s \in S$, the functor $\pi_{s!}$ induces an algebra morphism $E_{\mathbb{E}} \rightarrow E_{\mathbb{E}}^s$; hence we have a restriction functor

$$\text{res} : \text{dgDerf} - E_{\mathbb{E}}^s \rightarrow \text{dgDerf} - E_{\mathbb{E}}.$$

PROPOSITION 5.6.3. *Assume that condition (5.3.4) is satisfied.*

The equivalences in Theorems 5.5.8 and 5.6.1 (for all $s \in S$) can be chosen so that for any $s \in S$ the following diagram commutes up to natural transformation.

$$\begin{array}{ccc} \mathcal{D}_{(B)}^b(X^s, \mathbb{E}) & \xrightarrow[\sim]{\text{Th. 5.6.1}} & \text{dgDerf} - E_{\mathbb{E}}^s \\ \pi_{s!}[-1] \downarrow & & \downarrow \text{res} \\ \mathcal{D}_{(B)}^b(X, \mathbb{E}) & \xrightarrow[\sim]{\text{Th. 5.5.8}} & \text{dgDerf} - E_{\mathbb{E}} \end{array}$$

Proof. We consider the equivalences constructed in the proofs of Theorems 5.5.8 and 5.6.1 (using the same notation), and we will show that with this choice, the diagram commutes up to natural transformation. For simplicity we only treat the case $\mathbb{E} = \circ$.

First, as the functor $\text{dgDerf} - E_\circ \rightarrow \text{dgDer} - E_\circ$ is fully faithful, it is enough to show that the following diagram commutes, where the horizontal arrows are the composition of our equivalences with the obvious embeddings.

$$\begin{array}{ccc} \mathcal{D}_{(B)}^b(X^s, \circ) & \hookrightarrow & \text{dgDer} - E_\circ^s \\ \pi_{s!}[-1] \downarrow & & \downarrow \text{res} \\ \mathcal{D}_{(B)}^b(X, \circ) & \hookrightarrow & \text{dgDer} - E_\circ \end{array}$$

Now by construction the horizontal arrows factor through fully faithful functors $\mathcal{D}_{(B)}^b(X^s, \mathbb{E}) \rightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet}$, respectively $\mathcal{D}_{(B)}^b(X, \mathbb{E}) \rightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{\bullet}$. Moreover, the functor $- \otimes_{A_{\mathbb{O}}} A_{\mathbb{O}}^s$ induces a morphism of (\mathbb{O}, ϕ) -dg-algebras $E_{\mathbb{O}}^{\bullet} \rightarrow E_{\mathbb{O}}^{s, \bullet}$ which, by Proposition 5.2.4, induces our morphism $E_{\mathbb{O}} \rightarrow E_{\mathbb{O}}^s$ in cohomology. Hence, we obtain morphisms which make the following diagram commutative.

$$\begin{array}{ccccc}
 E_{\mathbb{O}}^{\bullet} & \xleftarrow{\text{qis}} & E_{\triangleright}^{\bullet} & \xrightarrow{\text{qis}} & E_{\mathbb{O}} \\
 \downarrow & & \downarrow & & \downarrow \\
 E_{\mathbb{O}}^{s, \bullet} & \xleftarrow{\text{qis}} & E_{\triangleright}^{s, \bullet} & \xrightarrow{\text{qis}} & E_{\mathbb{O}}^s
 \end{array} \tag{5.6.4}$$

Then the following diagram commutes up to natural transformations, where all functors are restriction functors for the morphisms in (5.6.4).

$$\begin{array}{ccccc}
 \mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet} & \xrightarrow{\sim} & \mathbf{dgDer} - E_{\triangleright}^{s, \bullet} & \xleftarrow{\sim} & \mathbf{dgDer} - E_{\mathbb{O}}^s \\
 \text{res} \downarrow & & \downarrow \text{res} & & \downarrow \text{res} \\
 \mathbf{dgDer} - E_{\mathbb{O}}^{\bullet} & \xrightarrow{\sim} & \mathbf{dgDer} - E_{\triangleright}^{\bullet} & \xleftarrow{\sim} & \mathbf{dgDer} - E_{\mathbb{O}}
 \end{array}$$

Hence, it is enough to prove the commutativity of the following diagram.

$$\begin{array}{ccc}
 \mathcal{D}_{(B)}^b(X^s, \mathbb{O}) \hookrightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet} & & \\
 \pi_s^![-1] \downarrow & & \downarrow \text{res} \\
 \mathcal{D}_{(B)}^b(X, \mathbb{O}) \hookrightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{\bullet} & &
 \end{array}$$

Using Proposition 5.2.4, it is even enough to show the commutativity of the following diagram, where horizontal arrows are the functors considered in the proofs of Theorems 5.5.8 and 5.6.1.

$$\begin{array}{ccc}
 \mathcal{D}^b(\text{Modf} - A_{\mathbb{O}}^s) \hookrightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet} & & \\
 \text{res} \downarrow & & \downarrow \text{res} \\
 \mathcal{D}^b(\text{Modf} - A_{\mathbb{O}}) \hookrightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{\bullet} & &
 \end{array} \tag{5.6.5}$$

Now, consider the following diagram.

$$\begin{array}{ccccccc}
 \mathcal{D}^b \text{Modf} - A_{\mathbb{O}}^s & \xleftarrow{\sim} & \mathcal{K}^b \text{Projf} - A_{\mathbb{O}}^s & \xrightarrow{a} & \langle 1_f E_{\mathbb{O}}^{s, \bullet} \rangle_{\Delta}^{\text{dgHo}} & \xrightarrow{\sim} & \langle 1_f E_{\mathbb{O}}^{s, \bullet} \rangle_{\Delta}^{\text{dgDer}} \hookrightarrow \mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet} \\
 \text{res} \downarrow & & \downarrow \text{res} & & \downarrow \text{res} & & \downarrow \text{res} \\
 \mathcal{D}^b \text{Modf} - A_{\mathbb{O}} & \xleftarrow{\alpha} & \mathcal{K}^b \text{Modf} - A_{\mathbb{O}} & \xrightarrow{b} & \text{dgHo} - E_{\mathbb{O}}^{\bullet} & \xrightarrow{d} & \mathbf{dgDer} - E_{\mathbb{O}}^{\bullet} \\
 & \nwarrow \sim & \uparrow \beta & & \downarrow & & \downarrow \\
 & & \mathcal{K}^b \text{Projf} - A_{\mathbb{O}} & \xrightarrow{c} & \langle 1_f E_{\mathbb{O}}^{\bullet} \rangle_{\Delta}^{\text{dgHo}} & \xrightarrow{\sim} & \langle 1_f E_{\mathbb{O}}^{s, \bullet} \rangle_{\Delta}^{\text{dgDer}}
 \end{array}$$

Here, all unnamed functors are the obvious ones, as well as d , β and γ . The functor a is $\underline{\text{Hom}}_{A_{\mathbb{O}}^s}^{\bullet}(M_s^{\bullet}, -)$, and the functors b and c are given by $\underline{\text{Hom}}_{A_{\mathbb{O}}}^{\bullet}(M^{\bullet}, -)$, so that the upper path from $\mathcal{D}^b \text{Modf} - A_{\mathbb{O}}^s$ to $\mathbf{dgDer} - E_{\mathbb{O}}^{s, \bullet}$ is the upper horizontal arrow in (5.6.5), and moreover the lower path from $\mathcal{D}^b \text{Modf} - A_{\mathbb{O}}$ to $\mathbf{dgDer} - E_{\mathbb{O}}^{\bullet}$ is the lower horizontal arrow in (5.6.5).

The categories $\langle 1_f E_{\mathbb{O}}^{s,\bullet} \rangle_{\Delta}^{\text{dgHo}} \subset \text{dgHo} - E_{\mathbb{O}}^{s,\bullet}$ and $\langle 1_f E_{\mathbb{O}}^{s,\bullet} \rangle_{\Delta}^{\text{dgDer}} \subset \text{dgDer} - E_{\mathbb{O}}^{s,\bullet}$ are the triangulated subcategories generated by the objects $1_f E_{\mathbb{O}}^{s,\bullet}$ for $f \in F$, and similarly for $E_{\mathbb{O}}^{\bullet}$.

All the squares and triangles in our large diagram are obviously commutative, except for the square involving a and b . The commutativity of this diagram follows from the fact that the functor $- \otimes_{A_{\mathbb{O}}} A_{\mathbb{O}}^s$ is left adjoint to $\text{res} : \text{Mod} - A_{\mathbb{O}}^s \rightarrow \text{Mod} - A_{\mathbb{O}}$. We observe also that the functor $\beta\gamma^{-1}\alpha$ is isomorphic to the identity functor. Then the commutativity of (5.6.5) can be checked by diagram chasing. \square

5.7 Application to Koszulity

In this subsection we deduce from our results that, under certain assumptions, the ring $A_{\mathbb{F}}$ can be endowed with a Koszul grading. This statement is a modular analogue of [BGS96, Theorem 4.4.4] (in the case of the flag variety). It will not be used in the rest of the paper, except in § 7.3.

The proof will use the notion of a dgg-algebra; such an object is a \mathbb{Z}^2 -graded algebra endowed with a differential d of bidegree $(1, 0)$ which satisfies the Leibniz rule with respect to the first grading, and such that $d^2 = 0$. One has an obvious notion of a dgg-module over a dgg-algebra, and we denote the derived category of right dgg-modules over the dgg-algebra R by $\text{dgDer}^{\mathbb{Z}} - R$.

We also recall some well-known facts on graded algebras. Let R be a finite-dimensional graded \mathbb{F} -algebra. If M is in $\text{Modf} - R$, an object $M^{\mathbb{Z}}$ of $\text{Modf}^{\mathbb{Z}} - R$ is called a *graded lift* of M if there exists an isomorphism $v(M^{\mathbb{Z}}) \cong M$, where v is the obvious forgetful functor. Then every simple or indecomposable projective right R -module admits a graded lift, which is unique up to isomorphism and shift in the grading. Moreover, we have the following easy lemma, whose proof is left to the reader. (See e.g. [Soe00, Proposition 2.7.2] for a similar statement.)

LEMMA 5.7.1. *Let $M^{\mathbb{Z}}$ and $M_i^{\mathbb{Z}}$ ($i = 1, \dots, r$) be objects of $\text{Modf}^{\mathbb{Z}} - R$, such that $v(M_i^{\mathbb{Z}})$ is indecomposable for any i . Assume that there exists an isomorphism*

$$v(M^{\mathbb{Z}}) \cong \bigoplus_i v(M_i^{\mathbb{Z}})$$

in $\text{Modf} - R$. Then there exist integers k_i ($i = 1, \dots, r$) and an isomorphism

$$M^{\mathbb{Z}} \cong \bigoplus_i M_i^{\mathbb{Z}}\langle k_i \rangle$$

in $\text{Modf}^{\mathbb{Z}} - R$.

Now we come to our Koszulity result. Here our conventions are slightly different from those of [BGS96]: we define the Koszul dual of a Koszul ring $K = \bigoplus_n K_n$ as the graded ring $K^{\dagger} := \text{Ext}_{-K}^{\bullet}(K_0, K_0)$. It is also Koszul.

PROPOSITION 5.7.2. *Assume that condition (5.3.4) is satisfied. Assume moreover that for any $x \in W$ we have $\mathcal{E}_x^{\mathbb{F}} = \text{IC}_{x,\mathbb{F}}$.*

There exists a Koszul \mathbb{F} -algebra K and an equivalence of categories

$$\text{Modf} - K \cong \text{Perv}_{(B)}(X, \mathbb{F})$$

such that the Koszul dual ring satisfies

$$K^{\dagger} \cong \text{Ext}_{\mathcal{D}_{(B)}^{\bullet}(X,\mathbb{F})}(\text{IC}_X, \text{IC}_X)$$

as graded rings, where $\text{IC}_X := \bigoplus_{x \in W} \text{IC}_{x,\mathbb{F}}$.

Remark 5.7.3. (i) By [Soe00, Corollary 1.0.3], the assumption that $\mathcal{E}_x^{\mathbb{F}} = \text{IC}_x^{\mathbb{F}}$ for all x is equivalent to a particular case of Lusztig’s conjecture on characters of simple representations of reductive

groups over \mathbb{F} . See also [WB12, Proposition 3.11] for other conditions equivalent to this assumption.

(ii) By [AJS94, Lemma F.3], the proposition implies that the ring $A_{\mathbb{F}}$ can be endowed with a Koszul grading, as announced at the beginning of the subsection.

Proof. To make the proof clearer, we denote the graded ring $E_{\mathbb{F}}$ by $\tilde{E}_{\mathbb{F}}$ when it is considered as a dg-algebra, or as a dgg-algebra with bigrading concentrated on the diagonal in \mathbb{Z}^2 .

Recall the equivalence of categories $\Pi^{\mathbb{F}} : \text{Perv}_{(B)}(X, \mathbb{F}) \xrightarrow{\sim} \text{Modf} - A_{\mathbb{F}}$. The simple right $A_{\mathbb{F}}$ -modules are parametrized by W ; we set $S_x := \Pi^{\mathbb{F}}(\text{IC}_{x, \mathbb{F}})$. Let us denote by

$$\Psi : \mathcal{D}^b(\text{Modf} - A_{\mathbb{F}}) \xrightarrow{\sim} \text{dgDerf} - \tilde{E}_{\mathbb{F}}$$

the equivalence constructed in the proof of Theorem 5.5.8. By construction of this equivalence we have $\Psi(\Pi^{\mathbb{F}}(\mathcal{BS}_f^{\mathbb{F}})) = e_f \cdot \tilde{E}_{\mathbb{F}}$ for any $f \in F$. Now if f is a reduced expression for x^{-1} , by Proposition 5.1.2, $\mathcal{E}_x^{\mathbb{F}}[\ell(x)]$ is a direct summand in $\mathcal{BS}_f^{\mathbb{F}}$. We deduce that, under our assumption, $\Psi(S_x)$ is a direct summand in $\tilde{E}_{\mathbb{F}}[-\ell(x)]$.

Now we note that the action of the Frobenius defines (as in the proof of Proposition 5.5.5) a \mathbb{Z} -grading on the \mathbb{O} -algebra $A_{\mathbb{O}}$. Using isomorphism (5.3.1) we deduce a \mathbb{Z} -grading on $A_{\mathbb{F}}$. Each simple $A_{\mathbb{F}}$ -module S_x can be lifted to a graded right $A_{\mathbb{F}}$ -module \tilde{S}_x . It is clear, using eigenvalues of the Frobenius again, that from the resolution $M_{\mathbb{O}}^{\bullet}$ of Proposition 5.3.6 one can deduce a complex of projective graded right $A_{\mathbb{F}}$ -modules $M_{\mathbb{Z}}^{\bullet}$ and use it to construct an equivalence $\Psi^{\mathbb{Z}}$ which makes the following diagram commutative, where vertical arrows are the obvious forgetful functors and $\text{dgDerf}^{\mathbb{Z}} - \tilde{E}_{\mathbb{F}}$ is defined in the obvious way.

$$\begin{CD} \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}) @>\Psi^{\mathbb{Z}}>> \text{dgDerf}^{\mathbb{Z}} - \tilde{E}_{\mathbb{F}} \\ @VVV @VVV \\ \mathcal{D}^b(\text{Modf} - A_{\mathbb{F}}) @>\Psi>> \text{dgDerf} - \tilde{E}_{\mathbb{F}} \end{CD}$$

Using the observation that $\Psi(S_x)$ is a direct summand in $\tilde{E}_{\mathbb{F}}[-\ell(x)]$ and an argument similar to the one for Lemma 5.7.1, it is not difficult to check that there exists $n \in \mathbb{Z}$ such that $\Psi^{\mathbb{Z}}(\tilde{S}_x)\langle n \rangle$ is a direct summand in $\tilde{E}_{\mathbb{F}}[-\ell(x)]$ in the category $\text{dgDerf}^{\mathbb{Z}} - \tilde{E}_{\mathbb{F}}$.

One easily checks that the functor

$$\Omega : \text{dgDerf}^{\mathbb{Z}} - \tilde{E}_{\mathbb{F}} \rightarrow \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}})$$

which sends a dgg-module M to the complex $\Omega(M)$ such that $\Omega(M)^{i,j} = M^{i+j,j}$ is an equivalence of triangulated categories, which satisfies $\Omega(M\langle n \rangle) = \Omega(M)[n]\langle n \rangle$. The object $\Omega(\Psi^{\mathbb{Z}}(\tilde{S}_x))\langle n \rangle[n]$ is a direct summand in $E_{\mathbb{F}}[\ell(x)]$. Hence, replacing \tilde{S}_x by $\tilde{S}_x\langle n - \ell(x) \rangle$, we can assume that $\Omega(\Psi^{\mathbb{Z}}(\tilde{S}_x))$ is a projective graded right $E_{\mathbb{F}}$ -module. As the functor $\Omega \circ \Psi^{\mathbb{Z}}$ is an equivalence of categories, we deduce that

$$\text{Ext}_{\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}}^n(\tilde{S}_x, \tilde{S}_y\langle m \rangle) = 0 \quad \text{unless } n + m = 0.$$

For any $x \in W$, let \tilde{Q}_x be the projective cover of \tilde{S}_x in $\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}$, and set $\tilde{Q} := \bigoplus_x \tilde{Q}_x$. Define the graded ring

$$K := \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}}(\tilde{Q}, \tilde{Q}\langle n \rangle).$$

By construction, the underlying ungraded ring of K is Morita equivalent to $A_{\mathbb{F}}$, and the arguments in [Ric10, §9.2] show that K is a Koszul ring. The description of K^\dagger is clear by construction. \square

PART III. MODULAR CATEGORY \mathcal{O} AND KOSZUL DUALITY

6. Reminder on modular category \mathcal{O}

From now on we will no longer consider coefficients in \mathbb{O} . For simplicity, we sometimes drop the subscripts ‘ \mathbb{F} ’ in the notation. We assume from now on that G is semi-simple of adjoint type.

In order to arrive at the results announced in the introduction, we have to recall several results from [Soe00]. We denote by \mathbb{F} a field of characteristic ℓ , and by $G_{\mathbb{F}}^{\vee}$ the connected reductive algebraic group over \mathbb{F} which is Langlands dual to G (in particular simply connected). We assume that ℓ is bigger than the Coxeter numbers of the irreducible components of the root system of $G_{\mathbb{F}}^{\vee}$ (see [Jan03, §6.2]). Our choice of B and T determines a Borel subgroup $B_{\mathbb{F}}^{\vee} \subset G_{\mathbb{F}}^{\vee}$ and a maximal torus $T_{\mathbb{F}}^{\vee} \subset B_{\mathbb{F}}^{\vee}$.

6.1 Definitions

From now on we write \mathcal{O} for the regular subquotient category defined in [Soe00, §2.3] for the group $G_{\mathbb{F}}^{\vee}$. (It was denoted by $\mathcal{O}_0(\mathbb{F})$ in the introduction.) We write $(M_x)_{x \in W}$ for the standard objects in \mathcal{O} . The parametrization is chosen so that M_e is projective and M_{w_0} is simple. We consider the coinvariant algebra C as in [Soe00, §2.1] and the functor

$$\mathbb{V} : \mathcal{O} \rightarrow C\text{-Modf}$$

which is fully faithful on projective objects (see [Soe00, Theorem 2.6.1]). For each simple reflection $s \in S$ we consider the semi-regular subquotient category \mathcal{O}^s from [Soe00, §2.4] with standard objects $M_x^s = M_{xs}$. We denote by C^s the s -invariants in C . We have an exact functor

$$\mathbb{V}^s : \mathcal{O}^s \rightarrow C^s\text{-Modf}$$

as in [Soe00, proof of Theorem 2.6.2]. Finally, we consider the exact translation functors $T^s : \mathcal{O} \rightarrow \mathcal{O}^s$ and $T_s : \mathcal{O}^s \rightarrow \mathcal{O}$ from [Soe00, §2.5] and diagrams (see [Soe00, proof of Theorem 2.6.2]) which commute up to natural isomorphism.

$$\begin{array}{ccc}
 \mathcal{O} & \xrightarrow{\mathbb{V}} & C\text{-Modf} & & \mathcal{O}^s & \xrightarrow{\mathbb{V}^s} & C^s\text{-Modf} \\
 T^s \downarrow & & \downarrow \text{res} & & T_s \downarrow & & \downarrow C \otimes_{C^s} (-) \\
 \mathcal{O}^s & \xrightarrow{\mathbb{V}^s} & C^s\text{-Modf} & & \mathcal{O} & \xrightarrow{\mathbb{V}} & C\text{-Modf}
 \end{array} \tag{6.1.1}$$

Amongst other things this shows that \mathbb{V}^s is also fully faithful on projective objects in \mathcal{O}^s . Indeed, any projective object in \mathcal{O} is a direct summand of $T^s Q$ with Q projective in \mathcal{O} . Then using the adjunctions (T^s, T_s) and (T_s, T^s) from [Soe00, §2.5] and (6.1.1) we deduce isomorphisms, for $M \in \mathcal{O}^s$ projective,

$$\begin{aligned}
 \text{Hom}_{\mathcal{O}^s}(M, T^s Q) &\cong \text{Hom}_{\mathcal{O}}(T_s M, Q) \cong \text{Hom}_C(\mathbb{V} T_s M, \mathbb{V} Q) \cong \text{Hom}_C(C \otimes_{C^s} \mathbb{V}^s M, \mathbb{V} Q) \\
 &\cong \text{Hom}_{C^s}(\mathbb{V}^s M, \text{res} \mathbb{V} Q) \cong \text{Hom}_{C^s}(\mathbb{V}^s M, \mathbb{V}^s T^s Q).
 \end{aligned}$$

6.2 Projective objects and equivalences

We abbreviate $\Theta_s := T_s T^s$. Given any sequence $f = (s, t, \dots, r)$ of simple reflections, consider the projective object $P_f := \Theta_s \Theta_t \dots \Theta_r M_e$. If we denote by P_s the projective cover of M_s and if

our sequence f is a reduced expression for x^{-1} then

$$P_f = P_x \oplus \bigoplus_{y < x} P_y^{\oplus m(f,y)} \tag{6.2.1}$$

for some unknown multiplicities $m(f, y) \geq 0$. Furthermore, \mathcal{O} has finite homological dimension. As in § 5.1 let us fix a family $F = \{f_1, \dots, f_n\}$ of reduced expressions for the elements of W . It follows from (6.2.1) that every object of \mathcal{O} has a finite projective resolution in which every term is a finite direct sum of objects of the form P_f for some $f \in F$.

Let us write P_x^s for the projective cover of M_x^s in \mathcal{O}^s . Then, if f is a reduced expression for x^{-1} with $x \in W^s$ and if $P_f^s := T^s P_f$, we have an analogous decomposition

$$P_f^s = P_x^s \oplus \bigoplus_{\substack{y \in W^s \\ y < x}} P_y^{\oplus m^s(f,y)}$$

where $m^s(f, y) \geq 0$ are again unknown multiplicities. As before, \mathcal{O}^s has finite homological dimension, and every object has a finite projective resolution in which every term is a finite direct sum of objects of the form P_f^s for $f \in F$.

Set $P := \bigoplus_{f \in F} P_f$ and $E_{\mathbb{F}} := \text{End}_{\mathcal{O}}(P)$. We obtain an equivalence

$$\text{Hom}_{\mathcal{O}}(P, -) : \mathcal{O} \xrightarrow{\sim} \text{Modf} - E_{\mathbb{F}}.$$

If we denote by $e_f \in E_{\mathbb{F}}$ the projection to P_f then we have $P_f \mapsto e_f E_{\mathbb{F}}$ under this equivalence. Similarly, if we set $P^s := \bigoplus_{f \in F} P_f^s$ and $E_{\mathbb{F}}^s := \text{End}_{\mathcal{O}}(P^s)$, then we obtain an equivalence

$$\text{Hom}_{\mathcal{O}}(P^s, -) : \mathcal{O}^s \xrightarrow{\sim} \text{Modf} - E_{\mathbb{F}}^s$$

and if we denote by $e_f \in E_{\mathbb{F}}^s$ the projection to P_f^s then we have $P_f^s \mapsto e_f E_{\mathbb{F}}^s$.

Translation onto the wall yields a ring homomorphism $E_{\mathbb{F}} \rightarrow E_{\mathbb{F}}^s$ with $e_f \mapsto e_f$. It is straightforward to see that this is an injection, see e.g. (6.4.1) below (which explains why we do not decorate our idempotents with an upper index). Because $E_{\mathbb{F}}^s = \text{Hom}(T^s P, T^s P) \cong \text{Hom}(P, T_s T^s P)$, $E_{\mathbb{F}}^s$ is a projective right $E_{\mathbb{F}}$ -module. Similarly, the isomorphism $E_{\mathbb{F}}^s \cong \text{Hom}(T_s T^s P, P)$ shows that $E_{\mathbb{F}}^s$ is a projective left $E_{\mathbb{F}}$ -module. This bimodule gives us a commutative diagram up to natural isomorphism

$$\begin{array}{ccc} \mathcal{O}^s & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}}^s \\ T_s \downarrow & & \downarrow \text{res} \\ \mathcal{O} & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}} \end{array}$$

where the right-hand vertical map is the restriction under $E_{\mathbb{F}} \rightarrow E_{\mathbb{F}}^s$. The natural transformation is simply the adjunction isomorphism

$$\text{Hom}(T^s P, M) \xrightarrow{\sim} \text{Hom}(P, T_s M).$$

The right and left adjoints of T_s coincide, giving us two commutative diagrams up to natural isomorphism.

$$\begin{array}{ccc} \mathcal{O} & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}} \\ T^s \downarrow & & \downarrow (-) \otimes_{E_{\mathbb{F}}} E_{\mathbb{F}}^s \\ \mathcal{O}^s & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}}^s \end{array} \qquad \begin{array}{ccc} \mathcal{O} & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}} \\ T^s \downarrow & & \downarrow \text{Hom}_{-E_{\mathbb{F}}}(E_{\mathbb{F}}^s, -) \\ \mathcal{O}^s & \xrightarrow{\sim} & \text{Modf} - E_{\mathbb{F}}^s \end{array}$$

6.3 Standard objects

For later use, let us briefly explain how one can describe morphisms between standard objects in \mathcal{O} .

LEMMA 6.3.1. *For $x, y \in W$ we have*

$$\text{Hom}_{\mathcal{O}}(M_x, M_y) \cong \begin{cases} \mathbb{F} & \text{if } y \leq x, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, all non-zero morphisms between standard objects are injective.

Proof. By [Soe00, § 2.5.8], all objects M_y have the same socle M_{w_0} , where $w_0 \in W$ is the longest element, and moreover this simple object appears with multiplicity 1 in these modules. The last claim of the lemma follows directly from this.

The fact that the existence of an embedding $M_x \subset M_y$ implies that $y \leq x$ follows from [Soe00, § 2.3.4]. Finally, one can show the existence of such an embedding when $y \leq x$ using similar arguments as for existence of non-zero morphisms between Verma modules in usual category \mathcal{O} in characteristic zero. \square

6.4 Description of $E_{\mathbb{F}}$ in terms of the coinvariant algebra

With the help of (6.1.1) and the fully-faithfulness of \mathbb{V} and \mathbb{V}^s on morphisms between projective objects we can describe the rings $E_{\mathbb{F}} \subset E_{\mathbb{F}}^s$ more explicitly. For any expression $f = (s, t, \dots, r)$ let us consider the C -module

$$D_f := C \otimes_{C^s} C \otimes_{C^t} \cdots \otimes_{C^r} \mathbb{F}$$

(where $\mathbb{F} = \mathbb{V}(M_e)$ is the trivial C -module). If we set $D := \bigoplus_{f \in F} D_f$ and denote by $e_f \in \text{End}_C(D)$ the projection to D_f , then (6.1.1) gives an isomorphism

$$\mathbb{V}P \xrightarrow{\sim} D$$

and hence an isomorphism $E_{\mathbb{F}} \xrightarrow{\sim} \text{End}_C D$ which matches the idempotents e_f on both sides. Similarly, we obtain an isomorphism $\mathbb{V}^s P^s \xrightarrow{\sim} D$ (where D is now viewed as a module over C^s by restriction) and the following commutative diagram.

$$\begin{array}{ccc} E_{\mathbb{F}} & \xrightarrow{\sim} & \text{End}_C(D) \\ \downarrow & & \downarrow \\ E_{\mathbb{F}}^s & \xrightarrow{\sim} & \text{End}_{C^s}(D) \end{array} \tag{6.4.1}$$

In order to obtain a better compatibility with our geometric constructions we equip C with the doubled \mathbb{Z} -grading (so that $C^{\text{odd}} = 0$ and $C^2 \neq 0$). We immediately obtain \mathbb{Z} -gradings on D , $E_{\mathbb{F}}$ and $E_{\mathbb{F}}^s$ (using the isomorphisms in (6.4.1)). It is then easy to see that $\mathcal{D}^b(\text{Mod}^{\mathbb{Z}} - E_{\mathbb{F}})$ (respectively $\mathcal{D}^b(\text{Mod}^{\mathbb{Z}} - E_{\mathbb{F}}^s)$) is generated as a triangulated category by the shifts of $e_f E_{\mathbb{F}}$ (respectively $e_f E_{\mathbb{F}}^s$) for all $f \in F$.

We have now introduced two rings which are called $E_{\mathbb{F}}$, namely the ring $E_{\mathbb{F}} := \text{Ext}^{\bullet}(\mathcal{BS}^{\mathbb{F}}, \mathcal{BS}^{\mathbb{F}})$ in § 5.1 and the ring $E_{\mathbb{F}} = \text{End}_{\mathcal{O}}(P)$ above. Similarly, we have introduced two rings $E_{\mathbb{F}}^s$, namely the ring $E_{\mathbb{F}}^s := \text{Ext}^{\bullet}(\pi_{s!} \mathcal{BS}^{\mathbb{F}}, \pi_{s!} \mathcal{BS}^{\mathbb{F}})$ of § 5.1 and the ring $E_{\mathbb{F}}^s$ above. This was, however, intentional,

as we obtain the following commutative diagram of rings with horizontal isomorphisms.

$$\begin{array}{ccccc}
 \text{End}_{\mathcal{O}}(P) & \xrightarrow{\sim} & \text{End}_{\mathcal{O}}(D) & \xleftarrow{\sim} & \text{Ext}^{\bullet}(\mathcal{BS}^{\mathbb{F}}, \mathcal{BS}^{\mathbb{F}}) \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{End}_{\mathcal{O}^s}(T^s P) & \xrightarrow{\sim} & \text{End}_{\mathcal{O}^s}(D) & \xleftarrow{\sim} & \text{Ext}^{\bullet}(\pi_{s!}\mathcal{BS}^{\mathbb{F}}, \pi_{s!}\mathcal{BS}^{\mathbb{F}})
 \end{array}$$

The left-hand square is simply a copy of diagram (6.4.1), the right-hand vertical morphism is induced by $\pi_{s!}$ and the horizontal ring isomorphisms on the right-hand side are induced by hypercohomology (see [Soe00, Theorem 4.2.1]). In particular, our functors preserve the idempotents e_f for $f \in F$ in all six rings. Hence from now on we can use the notation $E_{\mathbb{F}}$ for all three rings in the upper row, and $E_{\mathbb{F}}^s$ for all three rings in the lower row. In particular, because \mathcal{O} and \mathcal{O}^s have finite global dimension (see § 6.2) we obtain the following result as promised.

PROPOSITION 6.4.2. *The rings $E_{\mathbb{F}}$ and $E_{\mathbb{F}}^s$ have finite global dimension.*

Remark 6.4.3. Note that in [Soe00] the flag varieties are defined over the complex numbers, while here we work over an algebraically closed field of positive characteristic, and with the étale topology. One can check that the arguments in [Soe00] also apply to our setting. Alternatively, one can deduce the statement in one setting from the other, see Remark 7.1.4(ii) below.

7. Modular Koszul duality

7.1 Statement

We briefly recall our notation in order to give a complete formulation of our main theorem. We let $R \supset R^+$ denote a root system together with a choice of positive roots, W the Weyl group and $S \subset W$ the simple reflections. Now, let us fix (as in § 6) a finite field \mathbb{F} of characteristic ℓ strictly bigger than the Coxeter numbers of the irreducible components of R (or equivalently R^{\vee}). We consider the following categories.

(i) To the dual root system R^{\vee} we associate the regular and semi-regular subquotient categories \mathcal{O} and \mathcal{O}^s for $s \in S$ with coefficients in \mathbb{F} . Recall that these categories are obtained as subquotients of the category of rational representations of $G_{\mathbb{F}}^{\vee}$, a semi-simple, simply-connected and split algebraic group over \mathbb{F} associated with R^{\vee} .

(ii) To the root system R we associate the flag variety $X = G/B$ and partial flag varieties $X^s = G/P_s$ for $s \in S$ and consider $\mathcal{D}_{(B)}^b(X, \mathbb{F})$ and $\mathcal{D}_{(B)}^b(X^s, \mathbb{F})$ the derived categories of Bruhat constructible étale sheaves on X and X^s with coefficients in \mathbb{F} . In order to define these varieties, we first choose a split connected reductive group G_{\circ} and Borel subgroup $B_{\circ} \subset G_{\circ}$ over a finite field \mathbb{F}_q of characteristic $\neq \ell$. The varieties in question are then obtained by base change from the corresponding versions over \mathbb{F}_q .

The following result is a more precise and expanded version of the first part of Theorem 1.2.1 from the introduction.

THEOREM 7.1.1 (‘Modular Koszul duality’). *Assume that the order of q in \mathbb{F}^{\times} is strictly bigger than $|R|$. Then there exist a finite-dimensional graded \mathbb{F} -algebra $E_{\mathbb{F}}$ with a complete set of mutually orthogonal idempotents $\{e_f \mid f \in F\}$, a finite-dimensional graded \mathbb{F} -algebra $E_{\mathbb{F}}^s$ containing $E_{\mathbb{F}}$ as a graded subalgebra for all simple reflections $s \in S$, and equivalences of*

categories such that all squares in the following diagram commute up to natural transformation.

$$\begin{array}{ccccccc}
 \mathcal{D}^b(\mathcal{O}^s) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E_{\mathbb{F}}^s) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}}^s) & \xrightarrow{\bar{v}} & \text{dgDerf} - E_{\mathbb{F}}^s \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X^s, \mathbb{F}) \\
 \downarrow T_s & & \downarrow \text{res} & & \downarrow \text{res} & & \downarrow \text{res} \\
 \mathcal{D}^b(\mathcal{O}) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E_{\mathbb{F}}) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}}) & \xrightarrow{\bar{v}} & \text{dgDerf} - E_{\mathbb{F}} \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X, \mathbb{F}) \\
 & & & & & & \downarrow \pi_s^![-1]
 \end{array}$$

(Here, the functors v and \bar{v} are defined as in § 1.1.) Moreover, the left equivalences are derived from equivalences of abelian \mathbb{F} -categories $\mathcal{O} \xrightarrow{\sim} \text{Modf} - E_{\mathbb{F}}$ and $\mathcal{O}^s \xrightarrow{\sim} \text{Modf} - E_{\mathbb{F}}^s$, and for all $f \in F$ we have

$$P_f \leftarrow e_f \cdot E_{\mathbb{F}} \mapsto \mathcal{BS}_f^{\mathbb{F}} \tag{7.1.2}$$

on the lower line.

In particular, the indecomposable projective objects in \mathcal{O} are mapped to the indecomposable parity sheaves on the flag variety. That is, for all $x \in W$ there exists a projective object $\tilde{P}_x \in \text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}}$ with

$$P_x \leftarrow \tilde{P}_x \mapsto \mathcal{E}_x^{\mathbb{F}}, \tag{7.1.3}$$

where $P_x \rightarrow M_x$ is the projective cover in \mathcal{O} and $\mathcal{E}_x^{\mathbb{F}}$ is the parity sheaf from § 5.1.

Remark 7.1.4. (i) The graded modules \tilde{P}_x are uniquely determined by properties (7.1.3), as follows, for example, from the reminder on graded rings in § 5.7.

(ii) Theorem 7.1.1 differs from Theorem 1.2.1 by the fact that our variety X is defined over $\overline{\mathbb{F}}_q$ instead of \mathbb{C} , and that we work with the étale topology instead of the classical topology. Let us briefly explain how one can deduce Theorem 1.2.1 from Theorem 7.1.1. First, one can replace the classical topology in Theorem 1.2.1 by the étale topology (still over \mathbb{C}) by the arguments in [BBD82, § 6.1.2]. If $\ell > |R| + 1$ then by Dirichlet’s theorem we can choose some prime number p whose order in \mathbb{F}^{\times} is strictly bigger than $|R|$, so that Theorem 7.1.1 applies to $\overline{\mathbb{F}}_p$. Choose a strictly henselian local ring $\mathfrak{R} \subset \mathbb{C}$ whose residue field is $\overline{\mathbb{F}}_p$. Then the flag varieties $X_{\overline{\mathbb{F}}_p}$ and $X_{\mathbb{C}}$ over $\overline{\mathbb{F}}_p$ and \mathbb{C} are obtained by base change from the flag variety $X_{\mathfrak{R}}$ over \mathfrak{R} . Moreover, we have inverse image functors

$$\mathcal{D}_{(B)}^b(X_{\mathbb{C}, \text{et}}, \mathbb{F}) \leftarrow \mathcal{D}_{(B)}^b(X_{\mathfrak{R}}, \mathbb{F}) \rightarrow \mathcal{D}_{(B)}^b(X_{\overline{\mathbb{F}}_p}, \mathbb{F})$$

(where all categories are defined as in § 2.1). Both of these functors are equivalences, which finishes the proof. Indeed these categories are generated by standard objects, as well as by costandard objects. Hence it is sufficient to prove that morphisms from a standard object to a shift of a costandard object coincide in all these categories. However, this easily follows from [Mil80, Corollary VI.4.20].

Proof of Theorem 7.1.1. Define the algebras $E_{\mathbb{F}}$ and $E_{\mathbb{F}}^s$ as in § 6. The commutativity of the left-hand square is established in § 6.2. The commutativity of the middle two squares is unproblematic. By Propositions 4.4.2 and 4.5.4 our assumptions on the order of q in \mathbb{F}^{\times} guarantee that condition (5.3.4) is satisfied. The equivalences on the right and the commutativity of the right-hand square are proved in Theorems 5.5.8, 5.6.1 and Proposition 5.6.3.

Property (7.1.2) is clear by construction. Let us deduce (7.1.3), by induction on the Bruhat order. If $x = e$ then the corresponding sequence of simple reflections is $f = \emptyset$, and $P_{\emptyset} = P_e$. Hence we can take $\tilde{P}_e = e_{\emptyset} \cdot E_{\mathbb{F}}$. Now let $x \in W$, and assume the result is known for all $y < x$. Let \bar{P}_x be a projective object in $\text{Modf}^{\mathbb{Z}} - E$ such that $v(\bar{P}_x)$ is sent to P_x . Let $f \in F$ be a reduced decomposition of x^{-1} . From (6.2.1) and Lemma 5.7.1 we deduce that there exist $m \in \mathbb{Z}$, graded

vector spaces V_y for $y < x$, and an isomorphism in $\text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}}$

$$e_f \cdot E_{\mathbb{F}} \cong \overline{P}_x \langle m \rangle \oplus \left(\bigoplus_{y < x} V_y \otimes_{\mathbb{F}} \tilde{P}_y \right).$$

Then, using induction and (7.1.2), the image of $\bar{v}(\overline{P}_x)[-m]$ in $\mathcal{D}_{(B)}^b(X, \mathbb{F})$ is the only direct summand \mathcal{F} in $\mathcal{BS}_f^{\mathbb{F}}$ such that $i_x^* \mathcal{F} \neq 0$; it follows that this image is $\mathcal{E}_x[\ell(x)]$. Hence the object $\tilde{P}_x := \overline{P}_x \langle m + \ell(x) \rangle$ satisfies (7.1.3). \square

The left adjoint of the three middle vertical arrows in the diagram in Theorem 7.1.1 is simply $-\otimes_E E^s$. (Recall that E^s is a projective left E -module, see §6.2.) If we add the left adjoints on the edges we obtain the following diagram, in which all squares commute up to natural isomorphism.

$$\begin{array}{ccccccc} \mathcal{D}^b(\mathcal{O}) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E) & \xrightarrow{\bar{v}} & \text{dgDerf} - E \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X, \mathbb{F}) \\ \downarrow T^s & & \downarrow -\otimes_E E^s & & \downarrow -\otimes_E E^s & & \downarrow -\otimes_E E^s \\ \mathcal{D}^b(\mathcal{O}^s) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E^s) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E^s) & \xrightarrow{\bar{v}} & \text{dgDerf} - E^s \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X^s, \mathbb{F}) \end{array}$$

Similarly, the right adjoint of res can be described as $\text{Hom}_{-E}(E^s, -)$, and adding all right adjoints we obtain the following commutative diagram.

$$\begin{array}{ccccccc} \mathcal{D}^b(\mathcal{O}) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E) & \xrightarrow{\bar{v}} & \text{dgDerf} - E \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X, \mathbb{F}) \\ \downarrow T^s & & \downarrow \text{Hom}_{-E}(E^s, -) & & \downarrow \text{Hom}_{-E}(E^s, -) & & \downarrow \text{Hom}_{-E}(E^s, -) \\ \mathcal{D}^b(\mathcal{O}^s) & \xrightarrow{\sim} & \mathcal{D}^b(\text{Modf} - E^s) & \xleftarrow{v} & \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - E^s) & \xrightarrow{\bar{v}} & \text{dgDerf} - E^s \xrightarrow{\sim} \mathcal{D}_{(B)}^b(X^s, \mathbb{F}) \end{array}$$

Comparison of both right-hand squares in the above diagrams (using the fact that $\pi_{s*} = \pi_{s!}$) allows us to deduce an isomorphism of \mathbb{Z} -graded (E, E^s) -bimodules

$$\text{Hom}_{-E}(E^s, E) \cong E^s \langle 2 \rangle.$$

Sometimes it is easier to think of the middle category above as a ‘graded version’. In order to emphasize this way of thinking we define

$$\tilde{\mathcal{O}} := \text{Modf}^{\mathbb{Z}} - E, \quad \tilde{\mathcal{O}}^s := \text{Modf}^{\mathbb{Z}} - E^s.$$

We can then regard our restriction functor res as a ‘graded version of translation onto the wall’ and denote it by \tilde{T}_s . This functor has a right adjoint \tilde{T}_*^s and a left adjoint $\tilde{T}_!^s$ which are related by $\tilde{T}_*^s = \tilde{T}_!^s \langle 2 \rangle$.

7.2 Standard objects

In this section we establish the last part of Theorem 1.2.1. More precisely, we prove the following theorem.

THEOREM 7.2.1 (‘Koszul dual of standard objects’). *For all $x \in W$ there exist graded right $E_{\mathbb{F}}$ -modules \tilde{M}_x in $\tilde{\mathcal{O}}$ with the following properties:*

- (i) $\tilde{M}_x \mapsto M_x$ in \mathcal{O} ;
- (ii) $\tilde{M}_x \mapsto \nabla_{x, \mathbb{F}}$ in $\mathcal{D}_{(B)}^b(X, \mathbb{F})$;
- (iii) if $xs > x$ there exists an embedding $\tilde{M}_{xs} \subset \tilde{M}_x \langle -1 \rangle$ of graded $E_{\mathbb{F}}$ -modules.

Remark 7.2.2. (i) The modules \widetilde{M}_x are uniquely determined by properties (i)–(ii), as follows, for example, from the reminder on graded rings in § 5.7.

(ii) More generally, one can show, as in Lemma 6.3.1, that there exists an embedding

$$\widetilde{M}_y\langle\ell(y)\rangle \subset \widetilde{M}_x\langle\ell(x)\rangle$$

between graded standard modules if and only if $y \geq x$ in W , and that these are the only homomorphisms of any degree (up to shift).

(iii) Using the fact that $\text{Hom}_{\mathcal{O}}(P_x, M_x) \cong \mathbb{F}$, Theorems 7.1.1 and 7.2.1, one can easily check that the surjection $P_x \rightarrow M_x$ lifts to a morphism of graded right $E_{\mathbb{F}}$ -modules $\widetilde{P}_x \rightarrow \widetilde{M}_x$.

Proof. If we examine $P_f \leftarrow e_f \cdot E_{\mathbb{F}} \mapsto \mathcal{BS}_f^{\mathbb{F}}$ in the case of the empty sequence $f = \emptyset$ we see that $M_e \leftarrow e_{\emptyset} \cdot E_{\mathbb{F}} \mapsto \text{IC}_{e, \mathbb{F}} = \nabla_{e, \mathbb{F}} = i_{e^*} \mathbb{F}_{\text{id}}$. Hence we can take $\widetilde{M}_e = e_{\emptyset} \cdot E_{\mathbb{F}}$. If $x < xs$ then by [Soe00, § 2.5.4] there is a short exact sequence

$$M_x \hookrightarrow T_s T^s M_x \twoheadrightarrow M_{xs}$$

and by [Soe00, § 2.5.2] we have $\text{Hom}_{\mathcal{O}}(M_x, T_s T^s M_x) \cong \mathbb{F}$. Hence, if we have already constructed a graded lift \widetilde{M}_x of M_x satisfying (i) and (ii) then we obtain a graded lift \widetilde{M}_{xs} of M_{xs} as the cokernel of the adjunction morphism after an appropriate shift in the grading. That is, we can define \widetilde{M}_{xs} via the short exact sequence

$$\widetilde{M}_x \hookrightarrow \widetilde{T}_s \widetilde{T}_1^s \widetilde{M}_x \twoheadrightarrow \widetilde{M}_{xs}\langle-1\rangle.$$

Property (i) for \widetilde{M}_{xs} now follows. Lemma 7.2.4 below implies that this object also satisfies (ii).

We now turn to (iii). We have $\text{Hom}_{\mathcal{O}}(M_x, M_x) = \mathbb{F}$, hence

$$\text{Hom}_{\mathcal{O}}(\widetilde{M}_x, \widetilde{M}_x\langle i \rangle) = 0 \quad \text{unless } i = 0. \tag{7.2.3}$$

Consider the composition of adjunction morphisms

$$\widetilde{M}_x \hookrightarrow \widetilde{T}_s \widetilde{T}_1^s \widetilde{M}_x \rightarrow \widetilde{M}_x\langle-2\rangle$$

(where we use $\widetilde{T}_1^s = \widetilde{T}_*^s\langle-2\rangle$). This composition is zero by (7.2.3). On the other hand, the second morphism is non-zero so that we obtain a non-zero morphism $\widetilde{M}_{xs}\langle-1\rangle \rightarrow \widetilde{M}_x\langle-2\rangle$. This morphism must be injective by Lemma 6.3.1. \square

LEMMA 7.2.4. *Given $s \in S$ and $x \in W^s$ we have a distinguished triangle*

$$\nabla_{x, \mathbb{F}} \rightarrow \pi_s^! \pi_{s!} \nabla_{x, \mathbb{F}} \rightarrow \nabla_{xs, \mathbb{F}}[1] \xrightarrow{[1]}$$

where the left-hand arrow is the adjunction morphism.

Proof. Using the isomorphisms of functors $\pi_s^! \cong \pi_s^*[2]$, $\pi_{s*} \cong \pi_{s!}$ and the fact that π_s restricted to X_x is an isomorphism we have

$$\pi_s^! \pi_{s!} i_{x*} \mathbb{F}_{X_x} \cong a_* \mathbb{F}_{X_{x, xs}}[2]$$

where $X_{x, xs} := X_x \sqcup X_{xs}$ and $a : X_{x, xs} \hookrightarrow X$ denotes the inclusion. If we write i (respectively j) for the closed (respectively open) inclusion of X_x (respectively X_{xs}) in $X_{x, xs}$ then we have a Gysin triangle

$$i_! i^! \mathbb{F}_{X_{x, xs}} \rightarrow \mathbb{F}_{X_{x, xs}} \rightarrow j_* j^* \mathbb{F}_{X_{x, xs}} \xrightarrow{[1]}$$

which we can rewrite as

$$i_* \mathbb{F}_{X_x}[-2] \rightarrow \mathbb{F}_{X_{x, xs}} \rightarrow j_* \mathbb{F}_{X_{xs}} \xrightarrow{[1]}.$$

If we now apply $a_*[\ell(x) + 2]$ we obtain the desired triangle except for the fact that it is not clear if the left-hand morphism so constructed coincides with the adjunction morphism. However, both morphisms generate the one-dimensional space $\text{Hom}(\nabla_{x,\mathbb{F}}, \pi_s^! \pi_{s!} \nabla_{x,\mathbb{F}})$ and the lemma follows. \square

7.3 Standard objects are Koszul modules

Let us consider again the setting and notation of § 5.7. In particular, we assume that $\mathcal{E}_x^{\mathbb{F}} = \text{IC}_{x,\mathbb{F}}$ for all $x \in W$, and we consider the Koszul rings K and K^\dagger .

In the proof of Proposition 5.7.2 we have considered a grading on the algebra $A_{\mathbb{F}}$. The functor

$$\Xi : \text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}} \rightarrow \text{Modf}^{\mathbb{Z}} - K$$

which sends M to the graded module whose n th component is

$$\text{Hom}_{\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}}(\tilde{Q}, M\langle n \rangle)$$

is an equivalence of categories, which satisfies $\Xi(M\langle n \rangle) = \Xi(M)\langle -n \rangle$.

Recall the objects \tilde{P}_x in $\tilde{\mathcal{O}}$ defined in Theorem 7.1.1, and set $\tilde{P} := \bigoplus_x \tilde{P}_x$.

LEMMA 7.3.1. *The graded ring K^\dagger is isomorphic to the graded ring whose n th component is*

$$\text{Hom}_{\tilde{\mathcal{O}}}(\tilde{P}, \tilde{P}\langle -n \rangle).$$

Proof. By Proposition 5.7.2 and its proof, the n th component of K^\dagger is

$$\text{Ext}^n(\text{IC}_X, \text{IC}_X) \cong \text{Ext}_{-A_{\mathbb{F}}}^n\left(\bigoplus_x S_x, \bigoplus_x S_x\right) \cong \text{Ext}_{\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}}^n\left(\bigoplus_x \tilde{S}_x, \bigoplus_x \tilde{S}_x\langle -n \rangle\right).$$

Now, using the equivalence $\Omega \circ \Psi^{\mathbb{Z}}$ considered in the proof of Proposition 5.7.2, we obtain an isomorphism

$$\text{Ext}_{\text{Modf}^{\mathbb{Z}} - A_{\mathbb{F}}}^n\left(\bigoplus_x \tilde{S}_x, \bigoplus_x \tilde{S}_x\langle -n \rangle\right) \cong \text{Hom}_{\tilde{\mathcal{O}}}\left(\bigoplus_x \Omega \circ \Psi^{\mathbb{Z}}(\tilde{S}_x), \bigoplus_x \Omega \circ \Psi^{\mathbb{Z}}(\tilde{S}_x)\langle -n \rangle\right).$$

Hence it is sufficient to prove that for every $x \in W$ we have an isomorphism

$$\Omega \circ \Psi^{\mathbb{Z}}(\tilde{S}_x) \cong \tilde{P}_x.$$

However, the object $\Omega \circ \Psi^{\mathbb{Z}}(\tilde{S}_x)$ is an indecomposable graded projective right $E_{\mathbb{F}}$ -module, hence is isomorphic to $\tilde{P}_y\langle j \rangle$ for some $y \in W$ and $j \in \mathbb{Z}$. As its image under $\Psi^{-1} \circ \bar{v} : \mathcal{D}^b \text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}} \rightarrow \mathcal{D}^b \text{Modf} - A_{\mathbb{F}}$ is S_x (by Theorem 7.1.1) we must have $y = x$ and $j = 0$. \square

Using Lemma 7.3.1 we can construct an equivalence of categories

$$\Upsilon : \text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}} \xrightarrow{\sim} \text{Modf}^{\mathbb{Z}} - K^\dagger$$

sending M to the graded module with n th component

$$\text{Hom}_{\text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}}}(\tilde{P}, M\langle -n \rangle).$$

This equivalence satisfies $\Upsilon(M\langle n \rangle) = \Upsilon(M)\langle n \rangle$.

Finally, we let

$$\kappa : \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - K) \xrightarrow{\sim} \mathcal{D}^b(\text{Modf}^{\mathbb{Z}} - K^\dagger)$$

be the composition

$$\mathcal{D}^b \text{Modf}^{\mathbb{Z}} - K \xrightarrow{\Xi^{-1}} \mathcal{D}^b \text{Modf}^{\mathbb{Z}} - A_{\mathbb{Z}} \xrightarrow{\Omega \circ \Psi^{\mathbb{Z}}} \mathcal{D}^b \text{Modf}^{\mathbb{Z}} - E_{\mathbb{F}} \xrightarrow{\Upsilon} \mathcal{D}^b \text{Modf}^{\mathbb{Z}} - K^\dagger.$$

This functor satisfies $\kappa(M\langle n \rangle) = \kappa(M)\langle -n \rangle[-n]$. We set

$$\mathfrak{P}_x := \Upsilon(\tilde{P}_x), \quad \mathfrak{M}_x := \Upsilon(\tilde{M}_x), \quad \mathfrak{S}_x := \Xi(\tilde{S}_x).$$

We let \mathfrak{L}_x be the unique simple quotient of \mathfrak{P}_x , and define

$$\mathfrak{N}_x := \kappa^{-1}(\mathfrak{M}_x), \quad \mathfrak{I}_x := \kappa^{-1}(\mathfrak{L}_x).$$

The following theorem summarizes all the results related to our ‘Koszul duality’ κ . Here we denote by \mathcal{I}_x the injective hull of $\text{IC}_{x,\mathbb{F}}$ in $\text{Perv}_{(B)}(X, \mathbb{F})$.

THEOREM 7.3.2. *Assume that condition (5.3.4) is satisfied. Assume moreover that for any $x \in W$ we have $\mathcal{E}_x^{\mathbb{F}} = \text{IC}_{x,\mathbb{F}}$.*

There exist dual Koszul rings K and K^\dagger and an equivalence κ fitting in the diagram

$$\begin{array}{ccc} \mathcal{D}^b(\text{Modf}-K) & \xleftarrow{v} \mathcal{D}^b(\text{Modf}^{\mathbb{Z}}-K) & \xrightarrow{\kappa} \mathcal{D}^b(\text{Modf}^{\mathbb{Z}}-K^\dagger) & \xrightarrow{v} \mathcal{D}^b(\text{Modf}-K^\dagger) \\ \downarrow \wr & & & \downarrow \wr \\ \mathcal{D}_{(B)}^b(X, \mathbb{F}) & & & \mathcal{D}^b(\mathcal{O}) \end{array}$$

where vertical arrows are induced by equivalences of abelian categories

$$\text{Modf}-K \cong \text{Perv}_{(B)}(X, \mathbb{F}), \quad \text{Modf}-K^\dagger \cong \mathcal{O}$$

and v are forgetful functors. The objects $\mathfrak{P}_x, \mathfrak{M}_x, \mathfrak{L}_x, \mathfrak{I}_x, \mathfrak{N}_x, \mathfrak{S}_x$ are all graded modules, which satisfy

$$\begin{array}{ccccccc} \text{IC}_x & \longleftarrow & \mathfrak{S}_x & \longleftarrow & \mathfrak{P}_x & \longrightarrow & P_x, \\ \nabla_x & \longleftarrow & \mathfrak{N}_x & \longleftarrow & \mathfrak{M}_x & \longrightarrow & M_x, \\ \mathcal{I}_x & \longleftarrow & \mathfrak{I}_x & \longleftarrow & \mathfrak{L}_x & \longrightarrow & L_x. \end{array}$$

Moreover, \mathfrak{M}_x is a Koszul K^\dagger -module in the sense of [BGS96, Definition 2.14.1].

Proof. We have already constructed all the functors in our diagram. By definition, $\mathfrak{P}_x, \mathfrak{M}_x, \mathfrak{L}_x$ and \mathfrak{S}_x are graded modules. We have also observed in the proof of Lemma 7.3.1 that $\kappa(\mathfrak{S}_x) \cong \mathfrak{P}_x$. By Theorem 7.2.1, \mathfrak{N}_x is sent to ∇_x in $\text{Perv}_{(B)}(X, \mathbb{F})$, hence it is a graded right K -module. Now we observe that

$$\text{Ext}_{\text{Modf}^{\mathbb{Z}}-K}^i(\mathfrak{S}_x, \mathfrak{I}_y\langle j \rangle) \cong \text{Ext}_{\text{Modf}^{\mathbb{Z}}-K^\dagger}^{i-j}(\mathfrak{P}_x, \mathfrak{L}_y\langle -j \rangle)$$

vanishes unless $i = j = 0$. This implies that \mathfrak{I}_x is the injective hull of \mathfrak{S}_x in $\text{Modf}^{\mathbb{Z}}-K$, which in turn implies that it is sent to \mathcal{I}_x in $\text{Perv}_{(B)}(X, \mathbb{F})$.

It remains to show that \mathfrak{M}_x is a Koszul module. However, for any i, j the \mathbb{F} -vector space

$$\text{Ext}_{\text{Modf}-K^\dagger}^i(\mathfrak{M}_x, \mathfrak{L}_y\langle j \rangle) \cong \text{Ext}_{\text{Modf}^{\mathbb{Z}}-K}^{i-j}(\mathfrak{N}_x, \mathfrak{I}_y\langle -j \rangle)$$

vanishes unless $i - j = 0$, since \mathfrak{I}_y is injective. We conclude by [BGS96, Proposition 2.14.2]. \square

ACKNOWLEDGEMENTS

We thank Annette Huber-Klawitter and Patrick Polo for helpful discussions. Some of this work was completed whilst G.W. visited the Université Blaise Pascal - Clermont-Ferrand II and S.R. visited the MPIM in Bonn. We would like to thank both institutions for their support.

REFERENCES

- AJS94 H. H. Andersen, J. C. Jantzen and W. Soergel, *Representations of quantum groups at a p th root of unity and of semisimple groups in characteristic p : independence of p* , *Astérisque* **220** (1994), 321.
- Bas68 H. Bass, *Algebraic K-theory* (W. A. Benjamin, Inc, New York–Amsterdam, 1968).
- Bei87 A. A. Beilinson, *On the derived category of perverse sheaves*, in *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, *Lecture Notes in Mathematics*, vol. 1289 (Springer, Berlin, 1987), 27–41.
- BBD82 A. Beilinson, J. Bernstein and P. Deligne, *Faisceaux pervers*, in *Analyse et topologie sur les espaces singuliers, I (Luminy, 1981)*, *Astérisque*, vol. 100 (Société Mathématique de France, Paris, 1982), 5–171.
- BGS96 A. Beilinson, V. Ginzburg and W. Soergel, *Koszul duality patterns in representation theory*, *J. Amer. Math. Soc.* **9** (1996), 473–527.
- BL94 J. Bernstein and V. Lunts, *Equivariant sheaves and functors*, *Lecture Notes in Mathematics*, vol. 1578 (Springer, Berlin, 1994).
- Cur88 C. W. Curtis, *A further refinement of the Bruhat decomposition*, *Proc. Amer. Math. Soc.* **102** (1988), 37–42.
- Del74 P. Deligne, *La conjecture de Weil. I*, *Publ. Math. Inst. Hautes Études Sci.* **43** (1974), 273–307.
- Del80 P. Deligne, *La conjecture de Weil. II*, *Publ. Math. Inst. Hautes Études Sci.* **52** (1980), 137–252.
- Deo85 V. V. Deodhar, *On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells*, *Invent. Math.* **79** (1985), 499–511.
- DGMS75 P. Deligne, P. Griffiths, J. Morgan and D. Sullivan, *Real homotopy theory of Kähler manifolds*, *Invent. Math.* **29** (1975), 245–274.
- Fei82 W. Feit, *The representation theory of finite groups*, *North-Holland Mathematical Library*, vol. 25 (North-Holland Publishing Co, Amsterdam, 1982).
- Fie11 P. Fiebig, *Sheaves on affine Schubert varieties, modular representations, and Lusztig’s conjecture*, *J. Amer. Math. Soc.* **24** (2011), 133–181.
- Fie12 P. Fiebig, *An upper bound on the exceptional characteristics for Lusztig’s character formula*, *J. Reine Angew. Math.* **673** (2012), 1–31.
- FW10 P. Fiebig and G. Williamson, *Parity sheaves, moment graphs and the p -smooth locus of Schubert varieties*, *Ann. Inst. Fourier (Grenoble)*, to appear, arXiv:1008.0719.
- Jan03 J. C. Jantzen, *Representations of algebraic groups*, *Mathematical Surveys and Monographs*, vol. 107, second edition (American Mathematical Society, Providence, RI, 2003).
- Jut09 D. Juteau, *Decomposition numbers for perverse sheaves*, *Ann. Inst. Fourier (Grenoble)* **59** (2009), 1177–1229.
- JMW09 D. Juteau, C. Mautner and G. Williamson, *Parity sheaves*, Preprint (2009), arXiv:0906.2994.
- KL80 D. Kazhdan and G. Lusztig, *Schubert varieties and Poincaré duality*, in *Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, *Proc. Sympos. Pure Math.*, XXXVI (American Mathematical Society, Providence, RI, 1980), 185–203.
- Kel94 B. Keller, *Deriving DG categories*, *Ann. Sci. Éc. Norm. Super. (4)* **27** (1994), 63–102.
- KW01 R. Kiehl and R. Weissauer, *Weil conjectures, perverse sheaves and l -adic Fourier transform*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 42 (Springer, Berlin, 2001).
- Lam01 T. Y. Lam, *A first course in noncommutative rings*, *Graduate Texts in Mathematics*, vol. 131, second edition (Springer, New York, 2001).

- Mil80 J. S. Milne, *Étale cohomology*, Princeton Mathematical Series, vol. 33 (Princeton University Press, Princeton, NJ, 1980).
- Ric10 S. Riche, *Koszul duality and modular representations of semisimple Lie algebras*, Duke Math. J. **154** (2010), 31–134.
- RW11 S. Riche and G. Williamson, *Formality of modular constructible sheaves on the flag variety*, unpublished preprint (2011).
- Ric89 J. Rickard, *Morita theory for derived categories*, J. Lond. Math. Soc. (2) **39** (1989), 436–456.
- Sch11 O. M. Schnürer, *Equivariant sheaves on flag varieties*, Math. Z. **267** (2011), 27–80.
- SGA4 $\frac{1}{2}$ P. Deligne, *Séminaire de Géométrie Algébrique du Bois-Marie – Cohomologie étale (SGA 4 $\frac{1}{2}$)*, Lecture Notes in Mathematics, vol. 569 (Springer, Berlin, 1977), avec la collaboration de J. F. Boutot, A. Grothendieck, L. Illusie et J. L. Verdier.
- SGA5 A. Grothendieck, *Séminaire de Géométrie Algébrique du Bois-Marie 1965–1966 – Cohomologie l -adique et fonctions L (SGA 5)*, Lecture Notes in Mathematics, vol. 589 (Springer, Berlin, 1977).
- Soe89 W. Soergel, *n-cohomology of simple highest weight modules on walls and purity*, Invent. Math. **98** (1989), 565–580.
- Soe00 W. Soergel, *On the relation between intersection cohomology and representation theory in positive characteristic*, J. Pure Appl. Algebra **152** (2000), 311–335; Commutative algebra, homological algebra and representation theory (Catania/Genoa/Rome, 1998).
- Soe11 W. Soergel, *Modulare Koszul-Dualität*, Preprint (2011), arXiv:1109.0563.
- WB12 G. Williamson and T. Braden, *Modular intersection cohomology complexes on flag varieties*, Math. Z. **272** (2012), 697–727.
- Yun09 Z. Yun, *Weights of mixed tilting sheaves and geometric Ringel duality*, Selecta Math. (N.S.) **14** (2009), 299–320.

Simon Riche simon.riche@math.univ-bpclermont.fr

Clermont Université, Université Blaise Pascal, Laboratoire de Mathématiques, BP 10448,
F-63000 Clermont-Ferrand, France

and

CNRS, UMR 6620, Laboratoire de Mathématiques, F-63177 Aubière, France

Wolfgang Soergel Wolfgang.Soergel@math.uni-freiburg.de

Mathematisches Institut, Universität Freiburg, Eckerstraße 1, D-79104 Freiburg, Germany

Geordie Williamson geordie@mpim-bonn.mpg.de

Max-Planck-Institut für Mathematik, Vivatsgasse 7, 53111, Bonn, Germany