

GROUP EXTENSIONS ARE QUASI-SIMPLY-FILTRATED

S.G. BRICK AND M.L. MIHALIK

A finitely presented group G is quasi-simply-filtrated (abbreviated qsf) if, given a finite complex with fundamental group G , the universal cover of the complex can be “approximated” by simply connected finite complexes. This notion is a generalisation of a concept of Casson’s used in the study of three-manifolds.

In this paper we show that any extension of a finitely presented infinite group by a finitely presented infinite group is qsf.

1. INTRODUCTION

Suppose X is a finite complex. Write \tilde{X} for its universal cover. We say that X is *quasi-simply-filtrated* (abbreviated qsf) if, given a connected finite subcomplex $C \subset \tilde{X}$, there is a simply connected finite complex D , and a cellular map $f : D \rightarrow \tilde{X}$ with $f|_{f^{-1}(C)} : f^{-1}(C) \rightarrow C$ a homeomorphism.

In [2] it is shown that if X_1 and X_2 are finite complexes with isomorphic fundamental groups then X_1 is qsf if and only if X_2 is qsf. Thus one can say that a finitely presented group G is qsf if some (and hence any) finite complex X , with $\pi_1(X) = G$, is qsf.

The qsf condition is developed in [2] (also see [5]) as a geometric generalisation of Andrew Casson’s C_2 condition on presentations of groups (see [3]). Both of these conditions can be used to show that certain closed irreducible 3-manifolds are covered by \mathbb{R}^3 . The advantage of the qsf condition, besides it being a geometric condition, is that it is an invariant of finitely presented groups whereas the validity of the C_2 condition may vary between different presentations of a group.

In [2] it is shown that all one-relator groups and all simply connected at infinity groups are qsf. And (see [2]) the qsf property is preserved by taking finite extensions, amalgamations, and HNN-extensions. In [4] all automatic and semihyperbolic groups are shown to be qsf.

The purpose of this paper is to expand the class of groups known to be qsf by proving the following result:

THEOREM. *If $1 \rightarrow H \rightarrow G \rightarrow L \rightarrow 1$ is a short exact sequence of finitely presented infinite groups, then G is qsf.*

Received 13th September, 1993.

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/94 \$A2.00+0.00.

2. PRELIMINARIES

Choose a finite presentation

$$\mathcal{P} = \langle A \cup B \mid R \cup S \cup T \rangle$$

of G where $\langle A \mid R \rangle$ is a presentation of H and each element of T is of the form $b^{-1}abw^{-1}$ where $a \in A^{\pm 1}$, $b \in B^{\pm 1}$ and $w = w(a, b)$ is some word in the generators A .

Take X to be the two-complex associated to \mathcal{P} , and, as above, write \tilde{X} for the universal cover of X . Let Y be the two-complex associated to the presentation $\langle A \mid R \rangle$ of H . Then $Y \subset X$ and lying above Y in \tilde{X} is a collection of components, each component being a copy of the universal cover \tilde{Y} . There is one component for each coset of H in G or, equivalently, each element of L . Write $l\tilde{Y}$ for a typical component.

We shall refer to those edges of \tilde{X} lying above edges that correspond to elements of $A^{\pm 1}$ as A -edges. An A -path or an A -loop is an edge path or loop consisting entirely of A -edges. We define B -edges in a similar fashion. And those two-cells of \tilde{X} lying above two-cells corresponding to relators in T will be called *conjugation cells*. Note that the attaching map of a typical conjugation cell is of the form $b_1^{-1}ab_2w$ where b_1 and b_2 are B -edges lying above the same edge in X , a is an A -edge, and w is an A -path lying above the path corresponding to the word $w = w(a, b)$ (thus we abuse notation slightly).

We shall work with subcomplexes of \tilde{X} that are connected, finite and intersect each copy, $l\tilde{Y}$, of \tilde{Y} in a connected (possibly empty) subset. Call such subcomplexes *admissible*.

Start with an arbitrary connected finite subcomplex $C \subset \tilde{X}$. Choose a basepoint $*$ in C . By adding finitely many edges, we may assume that C is admissible. We shall enlarge C to a subcomplex K by adjoining finitely many edges, and conjugation cells such that $\pi_1(K)$ can be killed by killing finitely many loops which are null-homotopic in $\tilde{X} \setminus C$. Then we can attach disks to each such loop, yielding a simply connected finite complex D , and map each of the disks into $\tilde{X} \setminus C$. Clearly, this will suffice to show that X is qsf.

So we need to study the generators of the fundamental group of finite subcomplexes. They are of two different types: conjugates of A -loops and other elements. Refer to those generators that are conjugates of A -loops as being of type 1 and the others as being of type 2.

We need two lemmas on the effect of adjoining A -edges and conjugation cells.

LEMMA 2.1. *Suppose Z is an admissible subcomplex of \tilde{X} and e is an A -edge that meets Z . Then $Z \cup e$ is admissible. And if $\pi_1(Z)$ is generated by $\{u_1, \dots, u_n\}$*

then $\pi_1(Z \cup e)$ is generated by $\{u_1, \dots, u_n, \tau\lambda\tau^{-1}\}$, where λ is an A -loop in the copy of \tilde{Y} containing e .

PROOF: We need only consider the case where both endpoints of e lie in Z . Since Z is admissible there is an A -path, p , in Z from the terminal point of e to the initial point of e . Let $\lambda = ep$. Take τ to be a path from the basepoint of Z to the initial point of e . Now the result is immediate. □

LEMMA 2.2. Suppose Z is an admissible subcomplex of \tilde{X} and Δ is a conjugation cell with $\partial\Delta = b_1^{-1}ab_2w^{-1}$, and $b_1 \subset Z$. Then $Z \cup \Delta$ is admissible. Further if $\pi_1(Z)$ is generated by $\{u_1, \dots, u_n\}$ then $\pi_1(Z \cup \Delta)$ is generated by $\{u_1, \dots, u_n, v_1, \dots, v_m\}$ where each v_i is a type 1 generator. Moreover, if $a \subset Z$ then each v_i is of the form $\tau\lambda\tau^{-1}$ where λ is an A -loop in the copy of \tilde{Y} containing the endpoint of b_1 .

PROOF: Applying lemma 2.1 repeatedly shows that $\pi_1(Z \cup a \cup w)$ is gotten from $\pi_1(Z)$ by adding some new type 1 generators.

If a is in Z , then these new generators arise from adjoining w , and are of the desired form, again by lemma 2.1.

If b_2 is not in Z then $Z \cup a \cup w$ is a strong deformation retract of $Z \cup \Delta$. Otherwise $\pi_1(Z \cup \Delta)$ is obtained from $\pi_1(Z \cup a \cup w)$ by adding a new relator. In either case, the lemma follows. □

For sake of brevity, we make the following convention: When we speak of adjoining conjugation cells to a subcomplex Z , we mean that Z is admissible and that the cells may be indexed $\Delta_1, \Delta_2, \dots, \Delta_t$ with attaching maps $\partial\Delta_i = b_{i1}^{-1}a_i b_{i2} w_i^{-1}$ such that $b_{i1} \subset Z$ and $b_{i1} \subset Z \cup \Delta_1 \cup \dots \cup \Delta_{i-1}$ for $i > 1$.

The above lemmas show that if a generating set for $\pi_1(Z)$ has been chosen then adding finitely many A -edges or conjugation cells does not create any new type 2 generators. Hence if by adding such edges and cells we are able to homotope a single arbitrary type 2 generator to "nicer form" then by adding finitely many A -edges and conjugation cells, we can homotope all of the type 2 generators to such nicer forms.

3. TYPE 2 GENERATORS

In this section we shall adjoin finitely many conjugation cells and A -edges to C , yielding an admissible subcomplex C_1 such that $\pi_1(C_1)$ has a set of generators with each type 2 generator being of the form $\delta_1\rho\delta_2$, where ρ is a loop null-homotopic in $\tilde{X} \setminus C$ and $\delta_1\delta_2$ is an A -loop.

Fix a generating set for $\pi_1(C)$. First observe that adjoining conjugation cells lets us move B -edges to the right of A -edges (where doing so replaces the A -edges with A -paths). See Figure 3.1. Note that the conjugation cell pictured lies above the cell in X corresponding to the relator $(b^{-1})^{-1}ab^{-1}w^{-1}$ where $w = w(a, b^{-1})$. Hence we may




Figure 3.2

replacing each non-empty $C_1 \cap I\tilde{Y}$ with a simply connected finite complex that maps into $I\tilde{Y}$, as well as attaching disks to each of the loops ρ and mapping them into $\tilde{X} \setminus C$. We leave any further details to the reader.

4. FINISHING UP

At this point we have an admissible subcomplex C_1 containing C with $\pi_1(C_1)$ having a generating set of the form

$$\{\tau_1 \lambda_1 \tau_1^{-1}, \dots, \tau_s \lambda_s \tau_s^{-1}, \delta_{11} \rho_1 \delta_{21}, \dots, \delta_{1t} \rho_t \delta_{2t}\}$$

where the λ_i 's and the $\delta_{1j} \delta_{2j}$'s are A -loops and the ρ_j 's are null-homotopic in $\tilde{X} \setminus C$. In this section, we shall adjoin conjugation cells and attach a B -edge so as to obtain a subcomplex K containing C such that $\pi_1(K)$ has a generating set of the form

$$\{\tau_1 \lambda_1 \tau_1^{-1}, \dots, \tau_s \lambda_s \tau_s^{-1}, \dots, \tau_r \lambda_r \tau_r^{-1}, \delta_{11} \rho_1 \delta_{21}, \dots, \delta_{1t} \rho_t \delta_{2t}\}$$

where all of the A -loops λ_i and $\delta_{1j} \delta_{2j}$ are freely homotopic in K to A -loops $\{u_k\}$ that themselves are null-homotopic in $\tilde{X} \setminus C$. As remarked in §2 above, we can attach disks to each loop u_k and to each ρ_j , yielding a simply connected finite complex D , and map each of the disks into $\tilde{X} \setminus C$. This will finish the proof that X is qsf.

Since C_1 is a finite subcomplex, there are only finitely many components $I\tilde{Y}$ that meet C_1 . As the group L is infinite, there is a B -edge b with initial vertex $v_1 \in C_1$ and terminal vertex v_2 in some component $I\tilde{Y}$ that does not meet C_1 . Consider the subcomplex $C_1 \cup b$. Clearly $\pi_1(C_1) = \pi_1(C_1 \cup b)$ and $C_1 \cup b$ is admissible.

Define a finite graph Γ as follows: the vertices of Γ are those components $I\tilde{Y}$ that meet $C_1 \cup b$, the edges of Γ are the B -edges of $C_1 \cup b$. Let \bar{v}_2 be the vertex that contains v_2 .

We shall be considering subcomplexes K obtained from $C_1 \cup b$ by adjoining conjugation cells. Given such a K and given a subtree \mathcal{T} of Γ containing the vertex \bar{v}_2 , we shall say that *its A -loops are carried by \mathcal{T}* if $\pi_1(K)$ has a set of generators of the form

$$\{\tau_1 \lambda_1 \tau_1^{-1}, \dots, \tau_s \lambda_s \tau_s^{-1}, \dots, \tau_r \lambda_r \tau_r^{-1}, \delta_{11} \rho_1 \delta_{21}, \dots, \delta_{1t} \rho_t \delta_{2t}\}$$

where all of the A -loops λ_i and $\delta_{1j} \delta_{2j}$ are freely homotopic in K to A -loops that are in $\bigcup \mathcal{T}^{(0)}$ (recall that the vertices are components $l\tilde{Y}$). Formally, we shall be working with 4-tuples $(K, \sigma, \mathcal{T}, \mathcal{S})$ where σ is the set of generators for $\pi_1(K)$ of the above form and \mathcal{S} is a set of A -loops in $\bigcup \mathcal{T}^{(0)}$ such that each of the A -loops λ_i and $\delta_{1j} \delta_{2j}$ are freely homotopic in K to some loop in \mathcal{S} . Call such 4-tuples *acceptable*.

Letting $K = C_1 \cup b$, we see that its A -loops are carried by a maximal tree of Γ . Thus there is an acceptable 4-tuple. We shall use induction on the size of the subtrees \mathcal{T} to prove that there is some such K with its A -loops carried by the singleton $\{\bar{v}_2\}$, that is, an acceptable 4-tuple $(K, \sigma, \{\bar{v}_2\}, \mathcal{S})$.

Suppose $(K, \sigma, \mathcal{T}, \mathcal{S})$ is an acceptable 4-tuple. Since \mathcal{T} is finite, there is at least one vertex different from \bar{v}_2 that is terminal in Γ , that is, has valency one. Let \bar{v} be such a vertex, and let \bar{b} be the B -edge in \mathcal{T} emanating from \bar{v} . And let \bar{v}' be the terminal vertex of \bar{b} in \mathcal{T} . We shall see how adjoining conjugation cells will yield a new acceptable 4-tuple with A -loops carried by $\mathcal{T} \setminus \bar{b}$.

Assume u is one of the A -loops in \mathcal{S} which is contained in the vertex \bar{v} . Let v be the initial vertex of \bar{b} when viewed as an edge in \tilde{X} . By the admissibility of K , there is an A -path p in K from v to the basepoint of u . Consider the A -loop $pup^{-1} = a_1 \cdots a_n$. By adjoining conjugation cells as pictured in Figure 4.1, and applying lemma 2.2, we can easily homotope the loop pup^{-1} to an A -loop in \bar{v}' . Note that this may create more type 1 generators of the form $\tau \lambda \tau^{-1}$, but the point is that, by lemma 2.2, each of the new A -loops, λ , is in \bar{v}' . Doing so for each such u , yields the desired 4-tuple, $(K', \sigma', \mathcal{T} \setminus \bar{b}, \mathcal{S}')$.




Figure 4.1

Hence, by induction, there is an acceptable 4-tuple $(K, \sigma, \{\bar{v}_2\}, S)$. Now each A -loop $u \in S$ is in a copy $l\tilde{Y}$ of \tilde{Y} disjoint from C (the copy $l\tilde{Y}$ corresponds to the vertex \bar{v}). Since $l\tilde{Y}$ is simply connected, u is null-homotopic in $l\tilde{Y} \subset \tilde{X} \setminus C$. As noted above, we can now conclude that X is qsf.

REFERENCES

- [1] S.G. Brick, 'Quasi-isometries and ends of groups', *J. Pure Appl. Algebra* (to appear).
- [2] S.G. Brick and M.L. Mihalik, 'Qsf property for groups and spaces', (preprint).
- [3] S.M. Gersten and J.R. Stallings, 'Casson's idea about 3-manifolds whose universal cover is \mathbb{R}^3 ', *Internat. J. of Algebra and Comput.* 1 (1991), 395–406.
- [4] M.L. Mihalik and S.T. Tschantz, 'Tame combings and the quasi-simply-filtered condition for groups', (preprint).
- [5] J. R. Stallings, 'Brick's quasi simple filtrations for groups and 3-manifolds', in *Geometric methods in group theory, University of Sussez, July, 1991* (to appear).

Department of Mathematics
 CSU-Fresno
 Fresno CA 93740
 United States of America
 e-mail: stephen@zimmer.csufresno.edu

Department of Mathematics
 Vanderbilt University
 Nashville TN 37240
 United States of America
 e-mail: mihalikm@ctrvax.vanderbilt.edu