

Evaluating the Potential Benefits of Advanced Automatic Crash Notification

Rebecca E. Plevin, MD;¹ Robert Kaufman, BS;² Laura Fraade-Blanar, PhD;² Eileen M. Bulger, MD¹

1. Department of Trauma Surgery, Harborview Medical Center, Seattle, Washington USA
2. Harborview Injury Prevention and Research Center, University of Washington, Seattle, Washington USA

Correspondence:

Laura Fraade-Blanar, PhD
Harborview Injury Prevention & Research Center
325 Ninth Ave, Box 359960
Seattle, Washington 98104-2499 USA
E-mail: lblanar@uw.edu

Conflicts of interest/funding: Work was performed for the Crash Injury Research and Engineering Network (CIREN) project at the University of Washington/Harborview Injury Prevention and Research Center (Seattle, Washington USA) in cooperation with the US Department of Transportation/National Highway Traffic Safety Administration (USDOT/NHTSA; Washington DC, USA). Funding has been provided by NHTSA under Cooperative Agreement Number DTNH22-10-H-00295. Views expressed are those of the authors and do not represent the views of NHTSA. The first author is additionally funded by NIH Post-Doctoral Training Grant GM007037.

Keywords: Emergency Medical Services; EMS communication systems; motor vehicles; time factors; traffic accidents

Abbreviations:

AACN: Advanced Automatic Collision Notification
CDC: Centers for Disease Control and Prevention
CIREN: Crash Injury Research and Engineering Network
EMS: Emergency Medical Services
FARS: Fatality Analysis Reporting System

Abstract

Objective: Advanced Automatic Collision Notification (AACN) services in passenger vehicles capture crash data during collisions that could be transferred to Emergency Medical Services (EMS) providers. This study explored how EMS response times and other crash factors impacted the odds of fatality. The goal was to determine if information transmitted by AACN could help decrease mortality by allowing EMS providers to be better prepared upon arrival at the scene of a collision.

Methods: The Crash Injury Research and Engineering Network (CIREN) database of the US Department of Transportation/National Highway Traffic Safety Administration (USDOT/NHTSA; Washington DC, USA) was searched for all fatal crashes between 1996 and 2012. The CIREN database also was searched for illustrative cases. The NHTSA's Fatal Analysis Reporting System (FARS) and National Automotive Sampling System Crashworthiness Data System (NASS CDS) databases were queried for all fatal crashes between 2000 and 2011 that involved a passenger vehicle. Detailed EMS time data were divided into prehospital time segments and analyzed descriptively as well as via multiple logistic regression models.

Results: The CIREN data showed that longer times from the collision to notification of EMS providers were associated with more frequent invasive interventions within the first three hours of hospital admission and more transfers from a regional hospital to a trauma center. The NASS CDS and FARS data showed that rural collisions with crash-notification times >30 minutes were more likely to be fatal than collisions with similar crash-notification times occurring in urban environments. The majority of a patient's prehospital time occurred between the arrival of EMS providers on-scene and arrival at a hospital. The need for extrication increased the on-scene time segment as well as total prehospital time.

Conclusion: An AACN may help decrease mortality following a motor vehicle collision (MVC) by alerting EMS providers earlier and helping them discern when specialized equipment will be necessary in order to quickly extricate patients from the collision site and facilitate expeditious transfer to an appropriate hospital or trauma center.

Plevin RE, Kaufman R, Fraade-Blanar L, Bulger EM. Evaluating the potential benefits of advanced automatic crash notification. *Prehosp Disaster Med.* 2017;32(2):156-164.

Introduction

Motor vehicle collisions (MVCs) are a considerable source of morbidity and mortality in the United States. In 2010, there were nearly 5.5 million MVCs in the US, with 32,885 resulting fatalities and 2.2 million associated traumatic injuries. While these numbers represent significant improvements over the last decade, there is still, on average, one death every 16 minutes related to MVCs.¹

ISS: injury severity score
MVC: motor vehicle collision
NASS CDS: National Automotive Sampling System Crashworthiness Data System
NHTSA: National Highway Traffic Safety Administration
TBI: traumatic brain injury

Received: December 22, 2015

Revised: May 25, 2016

Accepted: June 9, 2016

Online publication: January 31, 2017

doi:10.1017/S1049023X16001473

A fundamental principle of modern trauma systems is the delivery of the right patient to the right hospital at the right time. Previous work demonstrates that after trauma, decreased time-to-treatment correlates with improved outcomes.²⁻⁴ Furthermore, appropriate triage of severely injured patients to a Level I trauma center is associated with decreased morbidity and a 25% reduction in mortality.^{5,6} Emergency Medical Services (EMS) providers use the Guidelines for Field Triage of Trauma Patients, published by the Centers for Disease Control and Prevention (CDC; Atlanta, Georgia USA) National Expert Panel, to appropriately triage trauma patients.⁷ After assessing vital signs and the anatomic location of injuries, the mechanism of injury, vehicle intrusion, passenger ejection, and same-compartment occupant deaths are further evaluated as rules to identify patients who would require transfer to the highest level trauma center.⁷ An additional guideline was added that also examines the likelihood of severe injury based on telematics data.⁷

Studies have demonstrated that demographic and vehicle collision data can be utilized to create models that predict the likelihood of severe injury.⁸ Advanced Automatic Collision Notification (AACN) services in passenger vehicles capture data during a collision triggered by an airbag deployment that includes: change in velocity, principal direction of force, and seat belt use. Based on the potential utility of this data in guiding patient management, in 2011 “vehicle telemetry data consistent with a high risk of injury” was added as one criterion in step three of the triage guidelines.⁷ Demographic and collision data are often unavailable at the time of EMS dispatch, but preliminary models created with AACN collision data are equally sensitive at predicting severe injury as compared to the models that incorporate demographic with crash data into the algorithm.⁹

Other research has, in isolation, examined the impact of environmental factors on EMS response time. Studies have found that rural crashes in particular could benefit by improved crash notification, in terms of greater survivorship.^{10,11} Overall, mortality would be reduced and analyses using Fatality Analysis Reporting System (FARS) data from the US Department of Transportation/National Highway Traffic Safety Administration (USDOT/NHTSA; Washington DC, USA) showed decreases in fatalities when all crash-to-notification times were within one minute compared to later response times.^{12,13}

It is hypothesized that quicker notification time following a collision by utilizing automatic crash notification systems may improve occupant outcomes and help appropriately allocate resources to the scene for crashes involving severely injured patients. Using the NHTSA's Crash Injury Research and Engineering Network (CIREN) database,¹⁴ the National Automotive Sampling System Crashworthiness Data System (NASS CDS) database,¹⁵ and the FARS database,¹⁶ this study assessed the association between EMS response times and the odds of fatality, taking into account fixed crash times, and examined crash, EMS system, and environmental factors impacting EMS response times.

Methods

Crash Injury Research and Engineering Network (CIREN)

The NHTSA's CIREN database contains MVC data collected by six Level I trauma centers in the US. The database, compiled from 1996 to the present, includes detailed case reviews of serious MVCs. Approximately 650 crash elements and 250 medical elements are recorded for each occupant. The crash investigators document details of the crash scene (eg, tire skid marks and road conditions) in order to establish crash patterns and reconstruct the

mechanics of the collision. The involved vehicles are inspected and a crush profile of the damaged plane is obtained to determine the force of the crash. The restraint systems are examined and occupant contact marks (eg, skin and fabric transfers) are documented to correlate with injury mechanisms. Prehospital data are obtained from EMS records. Details of the patient's hospital course, including radiological findings, operations, and outcomes, are recorded. Each case is reviewed by a crash investigator, a research nurse, bioengineers, and several physicians (including a radiologist and a trauma surgeon) in order to establish a probable cause and mechanism of injury. The CIREN database was queried for all fatal crashes between 1996 and 2012 which had complete EMS times documented (n = 171). These cases were divided into categories based on crash-notification time, defined as the amount of time elapsed between the crash and notification of emergency medical personnel (ie, 911 being alerted). In order to assess whether notification time correlates with poor outcomes, prehospital time was divided into segments and descriptive analysis was performed to evaluate patient demographics, cause of serious injury and death, transport mode to the hospital, disposition to a trauma center versus a non-trauma hospital, and number of invasive procedures performed within 24 hours of arrival at a trauma center. This study includes two case studies as examples of patients who may have benefited from AACN. These case studies aim to illustrate how health outcomes are impacted by response times.

Fatal Analysis Reporting System (FARS)

The FARS includes data from all police-reported crashes on public roadways in the US that result in at least one fatality within 30 days of the crash. The FARS database was searched for all fatal crashes between 2000 and 2011 that involved at least one passenger vehicle. Because only one set of EMS response and transport times were available for each crash, detailed analysis was limited to the driver in single-occupant, single-vehicle crashes. Available time segments included crash-time-to-notification, notification-to-EMS-arrival, and EMS-arrival-on-scene to EMS-arrival-at-hospital.

The FARS data were examined via descriptive statistics with an emphasis on examining the role of crash factors that could contribute to prolonged prehospital times. Extrication and crash location (urban or rural) were of particular interest. Rather than mean times, medians were reported due to the distribution of the data. These data are intended to examine factors which impact EMS times rather than mortality, as all crashes were fatal.

National Automotive Sampling System Crashworthiness Data System (NASS CDS)

The NHTSA's NASS CDS is a nationwide crash data collection program. In order to be included in the NASS CDS, a crash must be police reported, involve a harmful event (property damage and/or personal injury) resulting from a crash, and involve at least one towed passenger car or light truck or van that was in transport on a traffic-way. Data are collected in 27 Primary Sampling Units throughout the US on vehicles involved in approximately 5,000 automobile crashes annually. Ten sampling strata are used that over-sample crashes resulting in fatalities and serious injuries. A crash investigation is performed and all injuries are documented and scored using the Abbreviated Injury Severity coding.^{17,18}

All fatal crashes in the US involving a passenger vehicle that occurred between 2000 and 2011 were identified in NASS CDS. Passenger-vehicle driver and occupant(s) at all ages were included

in the analysis. The NHTSA provided the CIREN team with time segment information beyond what was available in the public datasets. Time segments included: crash-to-EMS-notification, notification-to-EMS-arrival, EMS-arrival-to-scene-departure, and scene-departure-to-hospital-arrival. If multiple forms of transportation were used, the earliest time available was used for the EMS-arrival-to-scene time, and the latest time available for EMS-departure-from-scene.

The time segments in NASS CDS data were examined descriptively as well as via a logistic regression model, with death as the outcome. Time segments were modeled as a linear predictor, as an ordered categorical predictor with time grouped in intervals, and lastly as an ordered categorical model with dummy measures to give greater flexibility. Time segments for each prehospital period were divided into the following time intervals: zero to nine minutes, 10-19 minutes, 20-29 minutes, 30-59 minutes, and 60+ minutes. Results are reported as odd ratios of fatality in comparison to the zero to nine minute reference interval. To explore potential non-linearity of time in relation to odds of death and to identify potential change points in the direction and significance of odds of death, a spline model was implemented. A variety of cut points were explored.

All models were adjusted to account for differences in age, injury severity score (ISS), and collision severity using change in velocity during the collision (ΔV) as a marker of crash severity. Due to concerns over influential outlier values, a delta-beta analysis was conducted to assess change in odds ratio when outliers were excluded from the analysis. A sensitivity analysis including only those crashes with crash-to-notification times less than 12 hours also was conducted. Results of the delta-beta analysis and sensitivity analysis did not vary considerably from previous findings and thus did not suggest the presence of influential outliers. Because the sample no longer represents national-level data due to the large number of cases with missing time data and the variability of time segments due to crash locations, weights were not used for NASS CDS data. Consequently, results should not be interpreted as nationally representative.

For all three data sets, only cases with complete time data were used. Within the analyses for NASS/CDS and FARS, all included individuals arrived at the hospital alive. The data analysis for this paper was completed using Stata software, Version 13 (StataCorp; College Station, Texas USA).

This study was approved by the University of Washington Institutional Review Board, Human Subjects Division (Seattle, Washington USA).

Results

Crash Injury Research and Engineering Network (CIREN)

Data Analysis—The mean age and ISS were similar between the notification interval groups, with the exception of a slightly higher ISS in patients with a notification time of 20-29 minutes (Table 1). The most common cause of death was thoracic trauma (38% of fatalities overall), followed by traumatic brain injury (TBI; 31% of fatalities overall). The distribution of cause of death was similar between notification time intervals.

The mean change in velocity (mean ΔV) of fatal collisions was highest in the zero to nine minute interval group (48 kmph) and lowest in the 20-29 minute interval group (38 kmph). Although patients with a crash-notification time of 10-19 minutes were most likely to undergo an invasive procedure overall (60%),

notification times of greater than 20 minutes were associated with an increased likelihood of requiring an invasive or operative procedure within the first three hours of hospital admission, although numbers were small. As notification time increased, so did the proportion of patients who were first taken to a regional hospital and subsequently transferred to a trauma center (eight percent in the zero to nine minute group versus 40% in the 30+ minute group), presumably after being stabilized at the local facility.

Case Studies—Multiple case studies were examined in CIREN. Two were selected where, because of the location/environment of the collision, the vehicles were not discovered until the following day. Two of these cases are detailed below as illustrative of situations in which AACN might have benefited the occupants of the vehicles.

- **Case Study 1:** The first case study involved the 66-year-old, male front passenger of a 2003 four-door utility vehicle that was involved in a severe frontal impact with a large tree (Figures 1 and 2). The crash occurred on a two-lane rural highway in the late evening in a dark area with no streetlights. The vehicle went down a slight slope before impacting the tree, and was thus not visible from the roadway in the dark conditions. The passenger and driver remained in the vehicle until the following morning when someone noticed the vehicle off the roadway and called 911. The front passenger was found dead when medics arrived the following morning. His injuries included a right subarachnoid hemorrhage, C4 fracture, hyoid fracture, multiple fractures of the sternum with a large right hemothorax and right lower lobe lacerations, a comminuted left patella fracture, and a fracture of the symphysis pubis. Review of these injuries suggests that early crash notification and treatment, particularly of the large right hemothorax, may have improved this patient's outcome.
- **Case Study 2:** The second case study involved the 17-year-old, male rear-seat passenger in a four-door sedan that was involved in a severe frontal impact. He was wearing a manual lap/shoulder seatbelt. This crash occurred in the morning hours in a rural area on a two-lane blacktop county road. The vehicle drifted off the roadway, impacted a ditch/culvert with significant resulting damage (Figure 3), rotated out of the ditch, and came to rest in a resident's driveway. The vehicle appeared parked, and it wasn't until dawn when the resident noticed the driver and two passengers still in the car. This backseat passenger sustained bilateral intra-cerebral hemorrhage, facial fractures, bilateral pneumothoraces, a left radial/ulnar fracture, and two small bowel perforations and a serosal tear of the descending colon. The TBI resulted in a prolonged coma and 26 days in the intensive care unit. The delay in treating his small bowel injuries contributed to the development of bacteremia. Earlier notification and initiation of treatment for his brain and abdominal injuries could have improved his overall hospital course and outcome.

Fatal Analysis Reporting System (FARS)

Information was extracted for a total of 91,200 drivers involved in single-vehicle, single-occupant fatal collisions between 2000 and 2011. The patients included in these analyses had a mean age of 40 years and were largely male (76%). The majority had a crash-notification time of less than 10 minutes (Table 2). However, 10.1%

	Crash-Notification Time (minutes) ^a			
	0-9 (n = 95)	10-19 (n = 45)	20-29 (n = 10)	30+ (n = 20)
Mean Age, years	51	49	49	55
Mean ISS	43	43	53	41
% Transported to Medical Facility	72%	76%	80%	75%
Cause of Death:				
Thoracic	37 (39%)	17 (38%)	3 (30%)	8 (40%)
Traumatic Brain Injury (TBI)	29 (31%)	15 (33%)	3 (30%)	5 (25%)
Spinal	8 (8%)	3 (7%)	3 (30%)	2 (10%)
Abdominal	5 (5%)	1 (2%)	0%	1 (5%)
Complications	16 (17%)	9 (20%)	1 (10%)	4 (20%)
Location of Death:				
On-scene	28%	24%	20%	25%
In EMS Vehicle	1%	0%	0%	0%
Emergency Department	22%	18%	30%	10%
Radiology	2%	0%	0%	0%
Operating Room (OR)	7%	4%	0%	0%
Intensive Care Unit (ICU)	32%	47%	40%	50%
On Floor/After Hospitalization	6%	2%	10%	10%
At Outside Hospital	1%	4%	0%	5%
Transport Mode				
Dead On-scene	28%	24%	20%	25%
Airlift to Trauma Center	24%	47%	50%	25%
EMS	47%	29%	30%	50%
Mean ΔV , kmph (% of cases coded)	48 (51%)	46 (67%)	38 (40%)	45 (40%)
Invasive Procedure within 24hrs of Trauma Center Admission	41%	60%	40%	45%
Within First 3hrs of Admission	27%	24%	40%	40%
Taken to Outside Hospital, Then Transferred to Trauma Center	8 (8%)	1 (2%)	2 (20%)	8 (40%)

Plevin © 2017 Prehospital and Disaster Medicine

Table 1. Demographic, Crash, and Hospital Data by Crash-Notification Time

Abbreviations: EMS, Emergency Medical Services; ICU, intensive care unit; ISS, Injury Severity Score; OR, operating room; TBI, traumatic brain injury.

^a Crash-notification times are presented as median times, in minutes.

of collisions resulted in crash-notification times of 10-19 minutes, and 9.1% had notification times of over 19 minutes (Table 2).

In order to evaluate the relative contributions of each prehospital time segment to the total prehospital time, the total prehospital time was divided into quartiles (Figure 4). The median crash-notification time, notification-to-EMS-arrival time, and EMS-arrival-to-hospital-arrival time increased with each successive quartile. The time from EMS-arrival to arrival-at-receiving-hospital composed approximately

74% of the prehospital time, regardless of quartile (Figure 4). When the fatalities were divided according to location (urban or rural), the median times from crash-to-notification were similar (urban = two minutes versus rural = three minutes; Figure 5). However, the crash-notification time was greater than 10 minutes in a higher proportion of rural fatalities. Ninety percent of urban fatalities had crash-notification times of zero to 19 minutes; of the remaining 10%, only three percent had crash-notification times of greater than



Plevin © 2017 Prehospital and Disaster Medicine

Figure 1. Case Study 1: Exterior Damage to the Front of the Vehicle.



Plevin © 2017 Prehospital and Disaster Medicine

Figure 2. Case Study 1: Interior of Vehicle, Front-Right Passenger Position.

30 minutes. Of the rural fatalities, however, 76.7% had crash-notification times of less than 10 minutes, 11.9% were in the 10-19 minute group, 3.4% were in the 20-29 minute group, and eight percent were in the >30 minute group (Figure 6). There also were considerable differences in the other prehospital time segments (Figure 5). Rural location nearly doubled the time from crash notification to EMS arrival (urban=six minutes versus rural = 10 minutes) and the time from EMS arrival to hospital arrival



Plevin © 2017 Prehospital and Disaster Medicine

Figure 3. Case Study 2: Exterior Damage to the Vehicle.

increased significantly as well (urban=25 minutes versus rural=32 minutes).

Collisions which required extrication of the driver resulted, perhaps unsurprisingly, in increased times from EMS arrival on-scene to arrival at a hospital. In Figure 7, box plots show that collisions requiring patient extrication resulted in higher median, 25th percentile, and 75th percentile EMS-arrival-to-hospital-arrival times than collisions that did not require extrication. The interquartile ranges also are higher, indicating that the group requiring extrication contains more outliers (ie, long EMS-arrival-to-hospital-arrival times).

National Automotive Sampling System Crashworthiness Data System (NASS CDS)

Information from a total of 12,298 fatal crashes between 2000 and 2011 was extracted from the database. In collisions where the patients were transported by a single ambulance, entrapment increased the median total prehospital time. The time from crash-to-notification was three minutes regardless of whether the victims were trapped in the vehicle. The time from notification-to-arrival of EMS on-scene also was similar (no entrapment=six minutes versus entrapment=seven minutes), as was the time from scene-departure-to-arrival-at-hospital (no entrapment=11 minutes versus entrapment=12 minutes). When victims were trapped, EMS personnel spent a median of 22 minutes on-scene, in contrast to the 16 minutes needed when entrapment was not a factor (Figure 8). Earlier time periods (eg, crash-to-notification and notification-to-EMS-arrival) were very close for both groups.


A univariate analysis of the odds of fatality based on the total prehospital time showed that a one-minute increase in prehospital time was associated with a significant increase in the odds of death (OR=1.001; P value=.024). When prehospital time was broken into individual segments, a one-minute increase in crash-notification time, notification-to-EMS-arrival time, and EMS-arrival-to-scene-departure time were associated with increased odds of death, and a one-minute increase in scene-departure-to-hospital-arrival time was associated with decreased odds of death. However, only the odds ratios for the last two time periods, EMS-arrival-on-scene to departure and EMS-departure-from-scene to hospital arrival, were significant. When adjusted for age, ISS, and change in velocity, all time periods became non-significant with the exception of EMS-arrival-to-scene-departure time (OR 1.02 per minute; P<.001). An EMS-arrival-to-scene departure time of one

Crash-Notification Time ^a	Number of Patients	Percent of Total
0-9 minutes	36,367	80.8
10-19 minutes	4,528	10.1
19-29 minutes	1,236	2.7
> 30 minutes	2,888	6.4

Plevin © 2017 Prehospital and Disaster Medicine

Table 2. Distribution of Crash-Notification Times, from FARS Data
Abbreviation: FARS, Fatality Analysis Reporting System.

^aCrash-notification times are presented as median times, in minutes.



Plevin © 2017 Prehospital and Disaster Medicine


Figure 4. Total Prehospital Time by Quartile, from FARS Data. Each quartile is divided into time segments: crash notification, notification to EMS arrival, and EMS arrival to hospital arrival. Data are presented as median times, in minutes. Abbreviations: EMS, Emergency Medical Services; FARS, Fatality Analysis Reporting System.

hour or longer, compared to zero to nine minutes, was significantly associated with increased odds of death (OR 5.46; P = .029; Table 3).

Using a spline model and adjusting for patient age, ISS, change in velocity, and EMS mode of transportation (ie, Basic Life Support, Advanced Life Support, ambulance/airlift, or airlift only), the odds of death generally increased with longer crash-notification times. However, the odds ratio only became significant at and beyond 30 minutes. When crash-notification times were 30 minutes or greater, each subsequent one-minute increase in crash-notification time was associated with a significant increase in the odds of death (OR = 1.13 per minute; P value < .001). When crash-notification times were less than 30 minutes, there was no statistically significant association between crash-notification time and odds of death, although the results trended in the same direction.

Discussion

Three national databases were used to evaluate whether fatalities related to automobile collisions might decrease with the use of AACN systems. The results indicate that TBI and thoracic trauma



Plevin © 2017 Prehospital and Disaster Medicine

Figure 5. Median Total Prehospital Time, Urban versus Rural, from FARS Data. Each group is divided into time segments: crash notification, notification to EMS arrival, and EMS arrival to hospital arrival. Data are presented as median times, in minutes. Abbreviations: EMS, Emergency Medical Services; FARS, Fatality Analysis Reporting System.

are the most common causes of death following an automobile collision (as seen in Case Study 1). Furthermore, longer crash-notification times were associated with an increased likelihood of a patient being taken to a local hospital and later transferred to a trauma center for a higher level of care, and involved more invasive procedures performed in the first three hours of arrival at a trauma center. This suggests that longer times in the field after a collision are associated with some level of clinical decompensation, ultimately resulting in increased needs for higher-level care.

In the two case studies, examples of patients who sustained severe injuries and were not located until hours after the collision as a result of environmental factors were presented. Their most severe injuries (pneumothorax, bowel perforation, and intra-cerebral hemorrhage) are life threatening if untreated, but with timely care can be managed and result in at least partial, and often full recovery. An AACN might have benefited these patients by alerting EMS providers of their location and facilitating timely transfer to a trauma center.




Figure 6. Percentage of Fatal Crashes by Crash-Notification Time, from FARS Data.

Results are presented separately for crashes occurring in urban and rural locations. Crash-notification times are presented as median times, in minutes.

Abbreviation: FARS, Fatality Analysis Reporting System.




Figure 7. Box Plots of the Median, 25th, and 75th Percentile EMS Arrival-to-Hospital Arrival Times.

Separated into patients requiring extrication and those not requiring extrication. Data were obtained from the FARS database and is presented as median times, in minutes.

Abbreviations: EMS, Emergency Medical Services; FARS, Fatality Analysis Reporting System.

Analysis of FARS data demonstrated that the majority of a patient’s prehospital time occurs between the arrival of EMS providers on-scene and arrival at a receiving hospital. Similarly, the need for extrication led to an increase in time from EMS arrival to hospital arrival. This finding was mirrored in the NASS CDS data, which demonstrated that the need for extrication increased the amount of time between EMS arrival and departure to a hospital, as well as the overall prehospital time. The NASS CDS data also demonstrated that above a threshold crash-notification time of 30 minutes, there is a significantly increased likelihood of death with each additional minute of crash-notification time.

This study identified several areas where AACN would be beneficial. Over 30 years ago, Cowley described the “golden hour” after trauma, with patients who arrive at a trauma center during that time having more favorable outcomes.^{19,20} Although the




Figure 8. Median Prehospital Time by Entrapment, from NASS CDS Data.

Each group is divided into time segments: crash notification, notification to EMS arrival, EMS arrival to EMS departure, and EMS departure to hospital arrival. Data are presented as median times, in minutes.

Abbreviations: EMS, Emergency Medical Services; NASS CDS, National Automotive Sampling System Crashworthiness Data System.

importance of exact quantity of time within “golden hour” has been the subject of debate, the need for prompt and efficient care is undisputed.^{21,22} One of the benefits of AACN is that it notifies EMS of the occurrence and location of a crash immediately after it takes place (likely would have benefited the patients in Case Studies 1 and 2). In light of the significant increase in mortality with crash-notification times over 30 minutes, this basic functionality of AACN could have profound effects on patient outcomes. Since 88%-92% of crashes have a crash-notification time of less than 30 minutes (Tables 1 and 2), AACN would likely be most beneficial in specific circumstances where there is an increased risk of long crash-notification times, such as crashes occurring in rural locations.

Previous work has demonstrated that increased overall prehospital time is associated with higher mortality following rural MVCs (as seen in Case Study 1).²³ It was found that the time from EMS arrival to hospital arrival following rural crashes was longer than in crashes that occurred in urban settings. Furthermore, higher crash-notification time was associated with a larger percentage of fatalities in the rural setting when compared to urban crashes and, in NASS CDS data, dummy category analysis demonstrated that time on-scene was associated with increased odds of fatality after one hour. Taken in aggregate, these data suggest that interventions aimed at decreasing the amount of time EMS personnel spend on-scene may be particularly beneficial for patients in rural collisions by helping them arrive at a hospital within the first hour after injury. This agrees with previous studies which found that EMS interventions were beneficial until they reached over 30 minutes.³

Certain prehospital time points are fixed. For example, a rural crash will almost always have a longer EMS-to-scene-arrival time than an urban crash, and crashes requiring extrication will almost always have longer on-scene times compared to crashes where no extrication is needed. An AACN can minimize the non-fixed time point: the time between crash and notification. In addition to

Time Segment	OR	P Value	95% CI	
Crash-to-Notification:				
10 to 19 min	0.79	.371	0.47	1.32
20 to 29 min	0.77	.612	0.27	2.14
30 to 59 min	0.54	.391	0.13	2.22
1 hour+	4.53	.028	1.18	17.42
Notification-to-EMS-Arrival:				
10 to 19 min	0.91	.665	0.59	1.40
20 to 29 min	0.46	.32	0.10	2.14
30 to 59 min	1.13	.876	0.25	4.98
1 hour+	100% fatality			
EMS-Arrival to EMS-Departure:				
10 to 19 min	0.47	.001	0.30	0.75
20 to 29 min	0.44	.001	0.26	0.73
30 to 59 min	0.46	.031	0.23	0.93
1 hour+	5.64	.029	1.19	26.65
Departure-to-Hospital-Arrival:				
10 to 19 min	0.89	.544	0.61	1.30
20 to 29 min	1.00	.989	0.56	1.77
30 to 59 min	0.56	.296	0.19	1.66
1 hour+	100% fatality			

Plevin © 2017 Prehospital and Disaster Medicine

Table 3. Odds of Fatality by Time Segment,^a from NASS CDS Data

Abbreviations: EMS, Emergency Medical Services; ISS, Injury Severity Score; NASS CDS, National Automotive Sampling System Crashworthiness Data System; ΔV , change in velocity during the collision.

^aData are presented as odds ratios compared to the 0–9 minute interval for each time segment and are adjusted to account for differences in age, ISS, and ΔV .

notifying EMS providers of the occurrence and location of a crash, AACN transmits data specific to the nature of the collision, including change in velocity, seatbelt use, and direction of force. These variables were among those proposed by the CDC in 2008 for inclusion in AACN data transmission for field triage.²⁴ Since that time, they have been demonstrated to correlate with the severity of injury sustained in automobile collisions.^{8,25} These variables could alert providers to the need for multiple emergency crews, air transport, and specialized equipment for patient extrication. Advanced notification could decrease the time spent dispatching appropriate extrication resources and, therefore, decrease the time patients spend in the field.

Limitations

This analysis has several limitations. None of the three datasets are nationally representative. The CIREN cases are not selected in a statistically random way, but rather are selected based on the enrollment criteria for cases presenting to participating trauma centers. Thus, the cases represent a case series rather than a population-based sample, which limits the ability to make

inferences to nationwide rates. However, the CIREN database provides more in-depth injury and medical data than any other nationwide crash database. Only single-passenger vehicle crashes were analyzed in the FARS database because of the complexity of collecting detailed EMS time data on crashes involving multiple vehicles and occupants, and because such data could not be correlated to individual occupants. Furthermore, data on pedestrians involved in an automobile collision as well as occupants of large trucks, motorcycles, and other vehicles were excluded.

The NASS CDS data represent a probability sample survey of police-reported MVCs. Complete case analysis requires time data being available for each case, with individuals surviving at least until they reached the hospital. Because ΔV was used to adjust for crash severity and rollover events do not include a ΔV estimate that utilizes the horizontal force, even if it was less severe than the rollover, provided the crash severity estimate. In addition, crashes with non-horizontal element (rollover only) were not included as ΔV was missing. Also, due to variability in crash location and missing time periods within the database, it is impossible to weight the data. Consequently, these data are not representative at the

national level. In addition, fatality was the only outcome evaluated. Among those who survived, time elapsed could still impact the severity of injuries. Incomplete data reporting occurred in all three the databases, and in some cases, it was difficult to accurately pinpoint the beginning and end of each prehospital time segment.

Conclusion

After 30 minutes had elapsed following a MVC, every one-minute increase in crash-notification time significantly increased mortality.

Furthermore, the majority of a patient's prehospital time occurred between the arrival of EMS providers on-scene and arrival at a hospital. The need for extrication and rural location increased these time periods, as well as the overall prehospital time. An AACN may help decrease mortality following MVC by alerting EMS providers to a collision earlier and by helping providers discern when specialized or heavy equipment will be necessary in order to quickly extricate patients from the collision site and facilitate expeditious transfer to a trauma center.

References

- National Highway Traffic Safety Administration. Traffic Safety Facts; 2010 Data. <http://www-nrd.nhtsa.dot.gov/Pubs/811659.pdf>. Accessed August 10, 2015.
- Zafullah Arifin M, Gunawan W. Analysis of pre-surgery time as a prognostic factor in traumatic acute subdural hematoma. *J Neurosurg Sci*. 2013;57(3):277-280.
- Clark DE, Winchell RJ, Betensky R. Estimating the effect of emergency care on early survival after traffic crashes. *Accid Anal Prev*. 2013;60:141-147.
- Dinh MM, Bein K, Roncal S, Byrne CM, Petchell J, Brennan J. Redefining the golden hour for severe head injury in an urban setting: the effect of prehospital arrival times on patient outcomes. *Injury*. 2013;44(5):606-610.
- MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med*. 2006;354(4):366-378.
- Mackenzie EJ, Rivara FP, Jurkovich GJ, et al. The impact of trauma-center care on functional outcomes following major lower-limb trauma. *J Bone Joint Surg Am*. 2008;90(1):101-109.
- Sasser SM. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. *MMWR Recomm Rep*. 2012;61(RR-1):1-20.
- Kononen DW, Flannagan CAC, Wang SC. Identification and validation of a logistic regression model for predicting serious injuries associated with motor vehicle crashes. *Accid Anal Prev*. 2011;43(1):112-122.
- Ayong-Chee P, Mack CD, Kaufman R, Bulger E. Predicting severe injury using vehicle telemetry data. *J Trauma Acute Care Surg*. 2013;74(1):190-194; discussion 194-195.
- The potential impact of rural mayday systems on vehicular crash fatalities. [http://www-ncbi.nlm.nih.gov/ncbi.nlm.nih.gov/pmc/articles/PMC3256762/pdf/aam52_p085.pdf](http://www-ncbi.nlm.nih.gov/ncbi.nlm.nih.gov/ncbi.nlm.nih.gov/pmc/articles/PMC3256762/pdf/aam52_p085.pdf). Accessed August 7, 2015.
- Lahausse JA, Fildes BN, Page Y, Fitzharris MP. The potential for automatic crash notification systems to reduce road fatalities. *Ann Adv Automot Med*. 2008;52:85-92.
- Wu J, Subramanian R, Craig M, Starnes M, Longthorne A. The effect of earlier or automation collision notification on traffic mortality by survival analysis. *Traffic Inj Prev*. 2013;14(Suppl):S50-S57.
- Clark DE, Cushing BM. Predicted effect of automatic crash notification on traffic mortality. *Accid Anal Prev*. 2002;34(4):507-513.
- Crash Injury Research (CIREN); National Highway Traffic Safety Administration (NHTSA). <http://www.nhtsa.gov/CIREN>. Accessed August 10, 2015.
- National Automotive Sampling System (NASS); National Highway Traffic Safety Administration (NHTSA). <http://www.nhtsa.gov/NASS>. Accessed August 10, 2015.
- Fatality Analysis Reporting System (FARS); National Highway Traffic Safety Administration (NHTSA). <http://www.nhtsa.gov/FARS>. Accessed August 10, 2015.
- Revell M, Pynsent P, Abudu A, Fairbank J. Trauma scores and trauma outcome measures. *Trauma*. 2003;5(1):61-70.
- Association for the Advancement of Automotive Medicine - Abbreviated Injury Scale; 2005 revision. <http://www.aaam.org/about-ais.html>. Accessed August 10, 2015.
- Cowley RA. The resuscitation and stabilization of major multiple trauma patients in a trauma center environment. *Clin Med (Northfield Il)*. 1976;83:16-22.
- Cowley RA. A total emergency medical system for the State of Maryland. *Md State Med J*. 1975;24(7):37-45.
- Rogers FB, Rittenhouse KJ, Gross BW. The golden hour in trauma: dogma or medical folklore? *Injury*. 2015;46(4):525-527.
- Lerner EB, Moscatti RM. The golden hour: scientific fact or medical "urban legend?" *Acad Emerg Med*. 2001;8(7):758-760.
- Gonzalez RP, Cummings GR, Phelan HA, Mulekar MS, Rodning CB. Does increased Emergency Medical Services prehospital time affect patient mortality in rural motor vehicle crashes? A statewide analysis. *Am J Surg*. 2009;197(1):30-34.
- Centers for Disease Control and Prevention. Advanced Automatic Collision Notification and Triage of the Injured Patient. http://www.cdc.gov/injuryresponse/pdf/AACN_Report_FINAL-a.pdf. Accessed August 10, 2015.
- Augenstein J, Perdeck E, Stratton J, Digges K, Bahouth G. Characteristics of crashes that increase the risk of serious injuries. *Annu Proc Assoc Adv Automot Med*. 2003;47:561-576.