

**EXISTENCE OF NORMAL MEROMORPHIC FUNCTIONS
WITH A PERFECT SET AS THE SET
OF ESSENTIAL SINGULARITIES**

TOSHIKO KUROKAWA

§1. Introduction

1. We are interested in whether there is a Cantor set E admitting no exceptionally ramified or normal meromorphic functions with E as the set of essential singularities. As for an exceptionally ramified meromorphic function, we [2] have recently given the following result.

THEOREM A. *Let E be a Cantor set with successive ratios ξ_n satisfying the condition*

$$\xi_{n+1} = o(\xi_n^5),$$

then the domain complementary to E admits no exceptionally ramified meromorphic functions with E as the set of essential singularities.

However, for a normal meromorphic function, S. Toppila [4] proved that if the set F is an infinite closed set, there exists a normal meromorphic function in the domain F^c complementary to F with at least one essential singularity in F . In [4], he gave a normal meromorphic function in F^c with one essential singularity in F .

In this paper, using the analogous method in S. Toppila [4], we shall give a normal meromorphic function with a Cantor set as the set of essential singularities.

Our result is stated as follows:

THEOREM. *Let E be a Cantor set with successive ratios ξ_n such that*

$$(1) \quad \lim_{n \rightarrow \infty} \xi_n = 0$$

and

$$(2) \quad \xi_{n+1} = O(\xi_n).$$

Received June 20, 1984.

Then there exists a normal meromorphic function in the domain complementary to E with E as the set of essential singularities.

Thus it follows from Theorem that the conclusion of Theorem A is false if we assume that a function is normal, instead of exceptionally ramified.

§2. Proof of Theorem

2. We form a Cantor set with successive ratios ξ_n , $0 < \xi_n < 2/3$, in the usual manner. We remove first an open interval of length $(1 - \xi_1)$ from the interval $I_{0,1}$: $[-1/2, 1/2]$, so that on both sides there remains a closed interval of length $\xi_1/2 \equiv \ell_1$. The remained intervals are denoted by $I_{1,1}$ and $I_{1,2}$. Inductively we remove an open interval of length $(1 - 2\eta_n) \prod_{p=1}^{n-1} \eta_p$, with $\xi_p/2 \equiv \eta_p$, $p = 1, 2, 3, \dots$, from each $I_{n-1,k}$, $k = 1, 2, \dots, 2^{n-1}$, so that on both sides there remains a closed interval of length $\prod_{p=1}^n \eta_p \equiv \ell_n$. The remaining intervals are denoted by $I_{n,2k-1}$ and $I_{n,2k}$. By repeating this procedure endlessly, we obtain an infinite sequence of closed intervals $\{I_{n,k}\}_{n=0,1,2,\dots, k=1,2,\dots,2^n}$. The set given by

$$E = \bigcap_{n=0}^{\infty} \bigcup_{k=1}^{2^n} I_{n,k}$$

is said to be the Cantor set in the interval $I_{0,1}$ with successive ratios ξ_n .

Denoting by $z_{n,k}$ the midpoint of $I_{n,k}$ and setting $\alpha_{n,k} = z_{n,k} + i\ell_n/2$, we shall give an infinite product

$$f(z) = \prod_{n=1}^{\infty} \prod_{k=1}^{2^n} \frac{z - \alpha_{n,k}}{z - \bar{\alpha}_{n,k}}$$

Obviously this function f has the set E as the set of essential singularities. In order to prove Theorem it is enough to show that f is normal in the domain Ω complementary to E .

The proof of this is based on the following result due to O. Lehto and K. I. Virtanen [3].

THEOREM B. *A function f meromorphic in a domain G of hyperbolic type, is normal in G if and only if there exists a finite constant C so that for every $z \in G$*

$$\frac{|f'(z)|}{1 + |f(z)|^2} |dz| \leq C d\sigma_G(z),$$

where $d\sigma_G(z)$ denotes the hyperbolic element of length of G .

In order to estimate $|dz|/d\sigma_\rho(z)$, we need the following

LEMMA. *Let D be the domain complementary to the set $\{0, 1, \infty\}$. Then*

$$\lim_{w \rightarrow 0} \left(|w| \log \frac{1}{|w|} \right) \frac{d\sigma_D(w)}{|dw|} = \frac{1}{2}$$

(see C. Constantinescu [1]).

3. We first discuss $|dz|/d\sigma_\rho(z)$. By Lemma, there exists a positive number $\delta_0, 1/8 > \delta_0 > 0$, such that

$$(3) \quad \frac{|dw|}{d\sigma_D(w)} < 4|w| \log \frac{1}{|w|} \quad \text{in } w \in \{w \mid 0 < |w| < 4\delta_0\} \equiv R_0.$$

Applying the linear transformations $w = 1 - \zeta$ and $w = 1/\zeta$ to (3), we have

$$(4) \quad \frac{|dw|}{d\sigma_D(w)} < 4|w - 1| \log \frac{1}{|w - 1|} \\ \text{in } w \in \{w \mid 0 < |w - 1| < 4\delta_0\} \equiv R_1$$

and

$$(5) \quad \frac{|dw|}{d\sigma_D(w)} < 4|w| \log |w| \quad \text{in } w \in \{w \mid 1/4\delta_0 < |w| < \infty\} \equiv R_\infty,$$

respectively. Since the set $R \equiv \{w \mid |w| \geq \delta_0/4, |w - 1| \geq \delta_0/4, |w| \leq 4/\delta_0\}$ is compact, there exists a positive number C_1 such that

$$(6) \quad \frac{|dw|}{d\sigma_D(w)} < C_1 \quad \text{in } w \in R.$$

We now set

$$\begin{aligned} \hat{\gamma}_{n,k} &= \{z \mid |z - z_{n,k}| = \delta_0 \ell_{n-1}\}, \\ \check{\gamma}_{n,k} &= \{z \mid |z - z_{n,k}| = \ell_n / \delta_0\}, \\ \Gamma_{n,k} &= \{z \mid |z - z_{n,k}| = \sqrt{\ell_n \ell_{n-1}}\} \end{aligned}$$

and

$$(\Gamma_{n,k}) = \{z \mid |z - z_{n,k}| < \sqrt{\ell_n \ell_{n-1}}\},$$

for $n = 1, 2, 3, \dots, k = 1, 2, \dots, 2^n$. We denote by $S_{n,k}$ (resp. $T_{n,k}$) the closed ring domain bounded by $\hat{\gamma}_{n,k}$ (resp. $\check{\gamma}_{n,k}$) and $\Gamma_{n,k}$. The triply connected closed domain bounded by $\Gamma_{n,k}$, $\Gamma_{n+1,2k-1}$ and $\Gamma_{n+1,2k}$ (resp. $\check{\gamma}_{n,k}$, $\hat{\gamma}_{n+1,2k-1}$ and $\hat{\gamma}_{n+1,2k}$) is denoted by $\Delta_{n,k}$ (resp. $\Delta'_{n,k}$), where $\Delta_{0,1}$ denotes the

closed ring domain bounded by $\Gamma_{1,1}$ and $\Gamma_{1,2}$ in the extended complex plane $\hat{\mathbb{C}}$. Immediately we have

$$\Omega = \bigcup_{\substack{k=1,2,\dots,2^n \\ n=0,1,2,\dots}} \mathcal{A}_{n,k}.$$

Denoting by $a_{n,k}$ (resp. $b_{n,k}$) the left (resp. right) endpoint of $I_{n,k}$, we write

$$D_{n,k} = \begin{cases} \text{the domain complementary to the set } \{a_{n,k}, b_{n,k}, 1\}, \\ \text{if } k = 1, 2, \dots, 2^{n-1}, \\ \text{the domain complementary to the set } \{0, a_{n,k}, b_{n,k}\}, \\ \text{if } k = 2^{n-1} + 1, 2^{n-1} + 2, \dots, 2^n. \end{cases}$$

We take the conformal mapping $w = \phi_{n,k}(z)$ from $D_{n,k}$ onto D such that

$$\phi_{n,k}(a_{n,k}) = 0, \quad \phi_{n,k}(b_{n,k}) = 1$$

and

$$\begin{cases} \phi_{n,k}(1) = \infty, & \text{if } k = 1, 2, \dots, 2^{n-1}, \\ \phi_{n,k}(0) = \infty, & \text{if } k = 2^{n-1} + 1, 2^{n-1} + 2, \dots, 2^n. \end{cases}$$

From (1), there is a positive integer N , $N \geq 2$,

$$(7) \quad \xi_n < \delta_0^2/2, \quad \text{for } n \geq N.$$

We denote by Ω_0 the closed domain bounded by the circles $\{\Gamma_{N,k}\}_{k=1,2,\dots,2^N}$ in $\hat{\mathbb{C}}$. For every $z \in \Omega - \Omega_0$, we choose the integers n and k such that $z \in \mathcal{A}_{n,k}$. Since $d\sigma_\Omega(z) \geq d\sigma_{D_{n,k}}(z)$ and since the hyperbolic element of length is conformally invariant, we have for $z \in \mathcal{A}_{n,k}$

$$(8) \quad \frac{|dz|}{d\sigma_\Omega(z)} \leq \frac{|dz|}{d\sigma_{D_{n,k}}(z)} = \frac{|dz|}{|dw|} \cdot \frac{|dw|}{d\sigma_D(w)} < 9 \ell_n \frac{|dw|}{d\sigma_D(w)},$$

where $w = \phi_{n,k}(z)$.

By elementary computations, we have

$$\begin{aligned} \phi_{n,k}(\hat{r}_{n+1,2k-1}) &\subset \{w \mid \delta_0/4 < |w| < 4\delta_0\}, \\ \phi_{n,k}(\hat{r}_{n+1,2k}) &\subset \{w \mid \delta_0/4 < |w - 1| < 4\delta_0\} \end{aligned}$$

and

$$\phi_{n,k}(\check{r}_{n,k}) \subset \{w \mid 1/4\delta_0 < |w| < 4/\delta_0\},$$

in view of (7). Thus

$$\begin{aligned} \phi_{n,k}(S_{n+1,2k-1}) &\subset R_0, \\ \phi_{n,k}(S_{n+1,2k}) &\subset R_1, \\ \phi_{n,k}(T_{n,k}) &\subset R_\infty \end{aligned}$$

and

$$\phi_{n,k}(A'_{n,k}) \subset R.$$

Hence applying (3), (4), (5) and (6) to the image of $A_{n,k}$ under $w = \phi_{n,k}(z)$, we deduce from (8) that

$$(9) \quad \begin{cases} \frac{|dz|}{d\sigma_\rho(z)} < C_2 |z - a_{n,k}| \log \frac{3\ell_n}{|z - a_{n,k}|}, & \text{for } z \in S_{n+1,2k-1}, \\ \frac{|dz|}{d\sigma_\rho(z)} < C_3 |z - b_{n,k}| \log \frac{3\ell_n}{|z - b_{n,k}|}, & \text{for } z \in S_{n+1,2k}, \\ \frac{|dz|}{d\sigma_\rho(z)} < C_4 |z - a_{n,k}| \log \frac{2|z - a_{n,k}|}{\ell_n}, & \text{for } z \in T_{n,k}, \\ \frac{|dz|}{d\sigma_\rho(z)} < C_5 \ell_n, & \text{for } z \in A'_{n,k}, \end{cases}$$

where C_j are constant.

4. We next discuss the spherical derivative $\rho(f(z)) \equiv |f'(z)|/(1 + |f(z)|^2)$ of f . We have for $z \in A_{n,k}$, $n \geq N$,

$$\begin{aligned} \rho(f(z)) &\leq \frac{|f(z)|}{1 + |f(z)|^2} \sum_{\substack{h=1,2,\dots,2^m \\ m=1,2,3,\dots}} \frac{\ell_m}{|z - \alpha_{m,h}| |z - \bar{\alpha}_{m,h}|} \\ &\leq \frac{1}{2} \sum_{\substack{h=1,2,\dots,2^m \\ m=1,2,\dots,n \\ (m,h) \neq (n,k)}} \frac{\ell_m}{|z - \alpha_{m,h}| |z - \bar{\alpha}_{m,h}|} \\ &\quad + \frac{|f(z)|\ell_n}{(1 + |f(z)|^2) |z - \alpha_{n,k}| |z - \bar{\alpha}_{n,k}|} \\ &\quad + \frac{1}{2} \sum_{\substack{h=1,2,\dots,2^m \\ m=n+1,n+2,\dots}} \frac{\ell_m}{|z - \alpha_{m,h}| |z - \bar{\alpha}_{m,h}|} \\ &\equiv \text{I} + \text{II} + \text{III}. \end{aligned}$$

The second term II is simply estimated as follows:

We have

$$\begin{aligned} \text{II} &< \frac{\ell_n}{|z - \bar{\alpha}_{n,k}|^2} \prod_{(m,h) \neq (n,k)} \left| \frac{z - \alpha_{m,h}}{z - \bar{\alpha}_{m,h}} \right| < \frac{\ell_n}{|z - \bar{\alpha}_{n,k}|^2} < \frac{36}{\ell_n}, \\ &\text{for } z \in U_{n,k} \equiv \{z \mid |z - \alpha_{n,k}| \leq \ell_n/6\}, \end{aligned}$$

$$\begin{aligned} \text{II} < \ell_n / \left\{ |z - \alpha_{n,k}|^2 \prod_{(m,h) \neq (n,k)} \left| \frac{z - \alpha_{m,h}}{z - \bar{\alpha}_{m,h}} \right| \right\} < \frac{\ell_n}{|z - \alpha_{n,k}|^2} < \frac{36}{\ell_n}, \\ \text{for } z \in U'_{n,k} \equiv \{z \mid |z - \bar{\alpha}_{n,k}| \leq \ell_n/6\} \end{aligned}$$

and

$$\begin{aligned} \text{II} < \frac{\ell_n}{2|z - \alpha_{n,k}||z - \bar{\alpha}_{n,k}|} < 18/\ell_n, \\ \text{for } z \in A'_{n,k} - (U_{n,k} \cup U'_{n,k}), \end{aligned}$$

so that

$$(10) \quad \text{II} < 36/\ell_n, \quad \text{for } z \in A'_{n,k}.$$

For $z \in S_{n+1,2k-1} \cup S_{n+1,2k} \cup T_{n,k}$ we have immediately

$$(11) \quad \text{II} < \frac{\ell_n}{2|z - \alpha_{n,k}||z - \bar{\alpha}_{n,k}|}.$$

In order to estimate I and III, we take roughly a lower bound of $|z - \alpha_{m,h}|$ or $|z - \bar{\alpha}_{m,h}|$, $(m, h) \neq (n, k)$. We may without loss of generality suppose that $k = 1$, i.e. $z \in A_{n,1}$.

(i) If $\alpha_{m,h} \in (\Gamma_{p,2})$, $p = 1, 2, 3, \dots, n$, we have

$$(12) \quad |z - \alpha_{m,h}| \geq d(\Gamma_{n,1}, \Gamma_{p,2}) \geq d(\Gamma_{p,1}, \Gamma_{p,2}) \geq \ell_{p-1}/3,$$

where $d(\Gamma_{p,q}, \Gamma_{r,s})$ denotes the distance between $\Gamma_{p,q}$ and $\Gamma_{r,s}$.

(ii) If $\alpha_{m,h} \in (\Gamma_{n+1,j})$, $j = 1, 2$, we have

$$(13) \quad \begin{cases} |z - \alpha_{m,h}| \geq (\ell_n/\delta_0) - (\ell_n/2) > \ell_n, & \text{for } z \in T_{n,1}, \\ |z - \alpha_{m,h}| \geq d(\Gamma_{n+1,j}, \alpha_{m,h}) \geq \sqrt{\ell_n \ell_{n+1}}/3, & \text{for } z \in A_{n,1} - T_{n,1}. \end{cases}$$

(iii) For the others, i.e. $\alpha_{1,1}, \alpha_{2,1}, \dots, \alpha_{n-1,1}$, we have

$$(14) \quad |z - \alpha_{m,1}| \geq d(\Gamma_{n,1}, \alpha_{m,1}) \geq d(\Gamma_{m+1,1}, \alpha_{m,1}) \geq \ell_m/3,$$

for $m = 1, 2, \dots, n - 1$. Here we may substitute $\bar{\alpha}_{m,h}$ for $\alpha_{m,h}$ in (12), (13) and (14). We need also

$$(15) \quad \ell_p/\ell_q = \eta_p \eta_{p-1} \cdots \eta_{q+1} < (1/3)^{p-q} \quad (p > q).$$

Using (12) and (14) we deduce

$$\text{I} = \sum_{\substack{m=1 \\ (m,h) \neq (n,1)}}^n \frac{\ell_m}{2} \left(\sum_{h=1,2,\dots,2^m} \frac{1}{|z - \alpha_{m,h}||z - \bar{\alpha}_{m,h}|} \right)$$

$$\begin{aligned}
 &= \sum_{m=1}^{n-1} \frac{\ell_m}{2} \left\{ \sum_{p=1}^m \left(\sum_{\alpha_{m,h} \in (I_{p,2})} \frac{1}{|z - \alpha_{m,h}| |z - \bar{\alpha}_{m,h}|} \right) + \frac{1}{|z - \alpha_{m,1}| |z - \bar{\alpha}_{m,1}|} \right\} \\
 &\quad + \frac{\ell_n}{2} \sum_{p=1}^n \sum_{\alpha_{n,h} \in (I_{p,2})} \frac{1}{|z - \alpha_{n,h}| |z - \bar{\alpha}_{n,h}|} \\
 &\leq \sum_{m=1}^{n-1} \frac{\ell_m}{2} \left\{ \left(\frac{3}{\ell_m} \right)^2 + \left(\frac{3}{\ell_{m-1}} \right)^2 + 2 \left(\frac{3}{\ell_{m-2}} \right)^2 + 2^2 \left(\frac{3}{\ell_{m-3}} \right)^2 + \dots + 2^{m-1} \left(\frac{3}{\ell_0} \right)^2 \right\} \\
 &\quad + \frac{\ell_n}{2} \left\{ \left(\frac{3}{\ell_{n-1}} \right)^2 + 2 \left(\frac{3}{\ell_{n-2}} \right)^2 + \dots + 2^{n-1} \left(\frac{3}{\ell_0} \right)^2 \right\}.
 \end{aligned}$$

Also in view of (15) we have

$$(16) \quad I < C_6/\ell_{n-1} = C_6\eta_n/\ell_n.$$

Similarly we deduce from (12) and (13)

$$\begin{aligned}
 \text{III} &< \sum_{m=n+1}^{\infty} 9 \cdot 2^{m-n-1} \frac{\ell_m}{\ell_n^2} \\
 &\quad \times \left\{ \frac{1}{9} + \left(\frac{\ell_n}{\ell_{n-1}} \right)^2 + 2 \left(\frac{\ell_n}{\ell_{n-2}} \right)^2 + \dots + 2^{n-1} \left(\frac{\ell_n}{\ell_0} \right)^2 \right\}, \\
 &\hspace{15em} \text{for } z \in T_{n,1}, \\
 \text{III} &< \sum_{m=n+1}^{\infty} 9 \cdot 2^{m-n-1} \frac{\ell_m}{\ell_n \ell_{n+1}} \\
 &\quad \times \left\{ 1 + \frac{\ell_n \ell_{n+1}}{\ell_{n-1}^2} + 2 \frac{\ell_n \ell_{n+1}}{\ell_{n-2}^2} + \dots + 2^{n-1} \frac{\ell_n \ell_{n+1}}{\ell_0^2} \right\}, \\
 &\hspace{15em} \text{for } z \in A_{n,1} - T_{n,1},
 \end{aligned}$$

and so we have

$$(17) \quad \begin{cases} \text{III} < C_7\eta_n/\ell_n, & \text{for } z \in T_{n,1}, \\ \text{III} < C_8/\ell_n, & \text{for } z \in A_{n,1} - T_{n,1}, \end{cases}$$

in view of (2) and (15).

Thus summing (10), (11), (16) and (17), we have

$$(18) \quad \begin{cases} \rho(f(z)) < \frac{C_9}{\ell_n} + \frac{\ell_n}{2|z - \alpha_{n,k}| |z - \bar{\alpha}_{n,k}|}, & \text{for } z \in S_{n+1,2k-1} \cup S_{n+1,2k}, \\ \rho(f(z)) < \frac{C_{10}}{\ell_n} \eta_n + \frac{\ell_n}{2|z - \alpha_{n,k}| |z - \bar{\alpha}_{n,k}|}, & \text{for } z \in T_{n,k}, \\ \rho(f(z)) < C_{11}/\ell_n, & \text{for } z \in A'_{n,k}. \end{cases}$$

Hence combining (9) and (18), we deduce that

$$\begin{aligned} \rho(f(z)) \frac{|dz|}{d\sigma_\varrho(z)} &< C_2 \left(3C_9 + \frac{3\ell_n^2}{2|z - \alpha_{n,k}||z - \bar{\alpha}_{n,k}|} \right) \\ &\quad \times \frac{|z - a_{n,k}|}{3\ell_n} \log \frac{3\ell_n}{|z - a_{n,k}|}, \quad \text{for } z \in S_{n+1,2k-1}, \\ \rho(f(z)) \frac{|dz|}{d\sigma_\varrho(z)} &< C_3 \left(3C_9 + \frac{3\ell_n^2}{2|z - \alpha_{n,k}||z - \bar{\alpha}_{n,k}|} \right) \\ &\quad \times \frac{|z - b_{n,k}|}{3\ell_n} \log \frac{3\ell_n}{|z - b_{n,k}|}, \quad \text{for } z \in S_{n+1,2k}, \\ \rho(f(z)) \frac{|dz|}{d\sigma_\varrho(z)} &< C_4 \left(\frac{C_{10}|z - a_{n,k}|^2}{\ell_n^2} \eta_n + \frac{|z - a_{n,k}|^2}{2|z - \alpha_{n,k}||z - \bar{\alpha}_{n,k}|} \right) \\ &\quad \times \frac{\ell_n}{2|z - a_{n,k}|} \log \frac{2|z - a_{n,k}|}{\ell_n}, \quad \text{for } z \in T_{n,k} \end{aligned}$$

and

$$\rho(f(z)) \frac{|dz|}{d\sigma_\varrho(z)} < C_5 C_{11}, \quad \text{for } z \in 4'_{n,k}.$$

Using the simple inequalities:

$$\begin{aligned} 0 < x \log \frac{1}{x} &< 1/e, && \text{for } 0 < x < 1, \\ |z - a_{n,k}|/\ell_n &< 1, && \text{for } z \in S_{n+1,2k-1}, \\ |z - b_{n,k}|/\ell_n &< 1, && \text{for } z \in S_{n+1,2k}, \\ |z - \alpha_{n,k}| &> \ell_n/3, && \text{for } z \in S_{n+1,2k-1} \cup S_{n+1,2k}, \\ |z - \bar{\alpha}_{n,k}| &> \ell_n/3, && \text{for } z \in S_{n+1,2k-1} \cup S_{n+1,2k}, \\ \frac{1}{4} &< \left| \frac{z - a_{n,k}}{z - \alpha_{n,k}} \right| &< 4, && \text{for } z \in T_{n,k}, \\ \frac{1}{4} &< \left| \frac{z - a_{n,k}}{z - \bar{\alpha}_{n,k}} \right| &< 4, && \text{for } z \in T_{n,k} \end{aligned}$$

and

$$\frac{2}{3} \sqrt{\eta_n} < \frac{\ell_n}{|z - a_{n,k}|} < \frac{1}{4}, \quad \text{for } z \in T_{n,k},$$

we are able to prove that $\rho(f(z))(|dz|/d\sigma_\varrho(z))$ is bounded in $\Omega - \Omega_0$. Further, since $\rho(f(z))(|dz|/d\sigma_\varrho(z))$ is also bounded in a compact set Ω_0 , it is bounded in Ω . Thus by Theorem B, we deduce that f is normal in Ω . This completes the proof of Theorem.

REFERENCES

- [1] C. Constantinescu, Einige Anwendungen des hyperbolischen Masses, *Math. Nachr.*, **15** (1956), 155–172.
- [2] T. Kurokawa, Exceptionally ramified meromorphic functions with a non-enumerable set of essential singularities, *Nagoya Math. J.*, **88** (1982), 133–154.
- [3] O. Lehto and K. I. Virtanen, Boundary behaviour and normal meromorphic functions, *Acta Math.*, **97** (1957), 47–65.
- [4] S. Toppila, Linear Picard sets for entire functions, *Ann. Acad. Sci. Fennicae, Ser. I*, **1** (1975), 111–123.

Department of Mathematics
Faculty of Education
Mie University
Tsu 514, Japan