

## Letter to the Editor

**Cite this article:** Cleary A, Chivers S, Daubeny PE, and Simpson JM (2021) Impact of COVID-19 on patients with congenital heart disease. *Cardiology in the Young* **31**: 163–165. doi: [10.1017/S1047951120004345](https://doi.org/10.1017/S1047951120004345)

Received: 1 September 2020  
Revised: 5 November 2020  
Accepted: 9 November 2020  
First published online: 14 December 2020

**Author for correspondence:**

Professor John Simpson, Department of Paediatric Cardiology, Evelina London Children's Hospital, SE1 7EH, England, UK.  
Tel: +44 20 7188 4550; Fax: + 44 20 7188 2307.  
E-mail: [john.simpson@gstt.nhs.uk](mailto:john.simpson@gstt.nhs.uk)

<sup>1</sup>Department of Paediatric Cardiology, Evelina London Children's Hospital, UK; <sup>2</sup>Department of Paediatric Cardiology, Royal Brompton Hospital, UK and <sup>3</sup>National Heart and Lung Institute, Imperial College, London, UK

The Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-Cov-2) virus pandemic has infected 25.1 million people worldwide with 844,000 deaths (31 August, 2020) with numbers rising rapidly. In the adult population, coronary artery disease, heart failure, and arrhythmias have been found to be associated with a more severe disease course with COVID-19 as well as higher mortality.<sup>1</sup> With respect to CHD, both paediatric and adult, there is a paucity of data. Many professional groups have issued guidelines for patients and parents/carers based on judgement and physiological principles rather than hard data.

We conducted a PubMed search using the terms “congenital heart disease” and “COVID 19” and “congenital heart disease” and “SARS-CoV-2” up until 4 August, 2020. We excluded articles, which involved Paediatric Inflammatory Multisystem Syndrome Temporally Associated with SARS-CoV-2 (PIMS-TS) or Multisystem Inflammatory Syndrome in Children (MIS-C), as these patients are a different disease entity. We identified 19 papers of which 12 were guidelines, commentary, or opinion and 7 included patient data (Table 1). We included the 7 papers,<sup>2–8</sup> which contained actual patient data and this included a variety of case reports, case series, and retrospective reviews. We excluded the 12 papers<sup>9–20</sup> that did not include any patient data and were deemed guidelines, opinion pieces, or a commentary.

A total of 94 suspected or confirmed cases have been reported in 7 papers.<sup>2–8</sup> By far, the biggest published series is from Italy<sup>2</sup> with 76 patients included. Amongst the 94 reported cases of CHD and COVID-19 infection, 79 patients (84%) were adults and 15 (16%) were children. Only one death was reported from all papers. Simpson *et al*<sup>3</sup> described a 19-year-old male patient with hypertrophic cardiomyopathy, obesity, diabetes, and metabolic syndrome, who was supported with ECMO for COVID-19 infection but died. With respect to critical care, of all 94 patients, 3 patients required ECMO (3.2%), 6 received inotropes (6.4%), 6 received positive pressure ventilation (6.4%), and 3 were supported with non-invasive ventilation (3.2%). With regard to specific CHD groups, reported cases are heterogeneous with no obvious predilection for a particular form of CHD (Tables 2 and 3).

In April, 2020, the Center for Disease Control (CDC) in the United States of America released its data on the number of patients affected by COVID-19.<sup>21</sup> Of their cohort, 2572 (1.7%) were children. Of these 2572 children, information on underlying medical conditions was only available for 345 patients (13%), and of these 25 had pre-existing cardiovascular disease (1% of overall numbers, 7.3% of those in whom there was the information of underlying conditions). Overall three deaths were reported, but no data was given on the background of these patients and whether they had cardiovascular disease or not. Data on the outcome of adult patients with CHD could not be extracted because acquired and CHD were not differentiated.

From the relatively small amount of data available up until the time of our literature search and analysis (31 August, 2020), some tentative conclusions can be drawn. The mortality of patients with CHD affected by COVID-19 appears low, and the disease does not appear to impact a specific form of CHD. Data on adult outcomes will remain difficult to extract until CHD is coded separately from acquired cardiac disease. In children, the mortality from COVID-19 appears low from North American data.<sup>4,21</sup> Data sources may, however, be impacted by ascertainment bias to report more severe cases. Conversely, a lack of rigorous testing may underestimate the number of patients affected. The extent to which shielding measures have impacted CHD patients also remains unknown.

Since we conducted our literature review and analysis, an additional paper has been published by Lewis *et al*,<sup>22</sup> looking at the effect of COVID-19 on patients with CHD. (The data from this additional paper by Lewis and colleagues is not included in our analysis and our Tables 1–3). In the manuscript by Lewis and colleagues, from a population of >7000 adults and children under follow-up, 53 CHD patients presented with COVID-19-related symptoms over a 4-month period. Nineteen percent of the patients were <18 years and 43 (81%) had mild symptoms. Nine patients (seven adults and two children) had moderate/severe symptoms. The cohort included a broad range of CHD including tetralogy of Fallot/pulmonary valvar stenosis (n = 16, 30%), functionally univentricular physiology status post-Fontan palliation (n = 10, 19%), shunting defects (n = 6, 11%), congenital valve abnormality (n = 7, 13%),

© The Author(s), 2020. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1.** Demographic data of patients with CHD and COVID-19 infection in seven published papers.

	Number of patients	% of patients
Total number of patients	94	
Number of deaths	1	1.1%
<b>Patient category</b>		
Adults	79	84%
Paediatrics	15	16%
<b>Type of CHD</b>		
Known	91	97%
Unknown	3	3%
<b>Treatment</b>		
NIV	3	3.2%
Intubation	6	6.4%
Inotrope	6	6.4%
ECMO	3	3.2%

ECMO=extra corporeal membrane oxygenation; NIV=non-invasive ventilation

**Table 2.** Specific CHD lesions reported in an Italian paper.<sup>2</sup>

Italian paper (n = 76) <sup>(2)</sup>	
CHD diagnosis	% of people with diagnosis
ASD	9%
VSD	13%
PS	5%
AS/BAV	9%
CoA	12%
Ebstein	5%
DORV	4%
TOF	12%
Pulmonary atresia	10%
TGA	8%
ccTGA	3%
AVSD	7%
TCPC	10%

AS=Aortic stenosis; ASD=Atrial septal defect; AVSD=Atrioventricular septal defect; BAV=Bicuspid aortic valve; ccTGA=Congenitally corrected transposition of the great arteries; DORV=Double outlet right ventricle; PS=Pulmonary stenosis; TCPC=Total cavopulmonary connection; TGA=Transposition of the great arteries; TOF=Tetralogy of Fallot; VSD=Ventricular septal defect

atrioventricular canal defects (n = 7, 13%), and other diagnoses (n = 7, 13%). Fifty patients survived but there were three deaths (6%), at the age of 34, 65, and 69 years. Moderate/severe disease was significantly associated with the genetic syndrome and adult CHD physiological stages C and D.<sup>23</sup>

In the United Kingdom, the British Congenital Cardiac Association is conducting a nationwide, multicentre survey to gauge the impact of COVID-19 on CHD more comprehensively. This, we hope, will permit a better understanding of groups at risk which will impact preventative strategies.

**Table 3.** Specific CHD lesions reported in all other papers.<sup>3-8</sup>

All other papers combined (n = 18) <sup>3-8</sup>		
CHD diagnosis	Number of patients with diagnosis	% of patients with diagnosis
Unknown	3	17%
TGA	3	17%
TCPC	1	6%
Heart transplant	1	6%
AVSD	5	28%
TOF/AVSD	1	6%
Cardiomyopathy	3	17%
TOF	1	6%

AVSD=Atrioventricular septal defect; TCPC=Total cavopulmonary connection; TGA=Transposition of the great arteries; TOF=Tetralogy of Fallot

**Acknowledgements.** None.

**Financial support.** This research received no specific grant from any funding agency, commercial, or not-for-profit sectors.

**Conflicts of interest.** None.

**Ethical standards.** This paper did not involve human or animal experimentation.

## References

- Mehra MR, Desai SS, Kuy SR, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in Covid-19. *NEJM* 2020; 382: e102.
- Sabatino J, Ferrero P, Chessa M et al. COVID-19 and congenital heart disease: results from a nationwide survey. *J Clin Med*. 2020; 9: 1774.
- Simpson M, Collins C, Nash DB, Panesar LE, Oster ME. COVID-19 infection in children with pre-existing heart disease. *J Pediatr* 2020; S0022-3476(20)30968-9.
- Shekerdeman LS, Mahmood NR, Wolfe KK, et al. Characteristics and outcomes of children with coronavirus disease 2019 (COVID-19) infection admitted to US and Canadian Pediatric Intensive Care Units. *JAMA* 2020. doi: 10.1001/jamapediatrics.2020.1948.
- Ferrero P, Piazza I, Ciuffreda M. COVID-19 in adult patients with CHD: a matter of anatomy or comorbidities? *Cardiol Young*. 2020; 30: 1196-1198.
- Bezerra RF, Franchi SM, Khader H, et al. COVID-19 as a confounding factor in a child submitted to staged surgical palliation of hypoplastic left heart syndrome: one of the first reports of SARS-CoV-2 infection in patients with congenital heart disease. *J Thoracic Cardiovasc Surg* 2020; S0022-5223(20)31317-9.
- Salik I, Mehta B. Tetralogy of Fallot palliation in a COVID-19 positive neonate. *J Clin Anesth* 2020; 66: 109914.
- Krishnan US, Krishnan SS, Jain S et al. Severe acute respiratory syndrome associated with Coronavirus 2 infection in patients with down syndrome, congenital heart disease, and pulmonary hypertension: is down syndrome a risk factor? *J Pediatr*. 2020; S0022-3476(20)30830-1.
- Lastinger LT, Daniels CJ, Lee M, Sabanayagam A, Bradley EA. Triage and management of the ACHD patient with COVID-19: a single center approach. *Int J Cardiol*. 2020; 320: 178-182.
- Radke RM, Frenzel T, Baumgartner H, Diller G-P. Adult congenital heart disease and the COVID-19 pandemic. *Heart*. 2020; heartjnl-2020-317258.
- Alsaied T, Aboulhosn JA, Cotts TB, et al. Coronavirus Disease 2019 (COVID-19) Pandemic Implications in Pediatric and Adult Congenital Heart Disease. *JAHA* 2020; 120.017224.
- Anthony JM, Kasargod Prabhakar CR, Clift P, Hudsmith L. UK ACHD response to COVID-19: Which adult patients are being advised to follow shielding by UK centres? *Cardiol Young* 2020; 30: 1547-1548.

13. Brida M, Chessa M, Gu H, Gatzoulis MA, The globe on the spotlight: Coronavirus disease 2019 (Covid-19). *Int J Cardiol.* 2020; 310: 170–172.
14. Giordano R, Cantinotti M. Congenital heart disease in the era of COVID-19 pandemic. *Gen Thorac Cardiovasc Surg* 2020; 13.
15. Sanna G, Serrau G, Bassareo PP, Neroni P, Fanos V, Marcialis MA. Children's heart and COVID-19: up-to-date evidence in the form of a systematic review. *Eur J Pediatr.* 2020; 179: 1079–1087.
16. Gallego P, Ruperti-Repilado FJ, Schwerzmann M. Adults with congenital heart disease during the coronavirus disease 2019 (COVID-19) pandemic: are they at risk? *Rev Esp Cardiol.* 2020; 73: 795–798.
17. Gatzoulis MA. COVID-19 and congenital heart disease in perspective. *Eur Heart J.* 2020; 41: 1871–1872.
18. Hemphill NM, Kuan MTY, Harris KC, Reduced physical activity during COVID-19 pandemic in children with congenital heart disease. *Can J Cardiol.* 2020; 36: 1130–1134.
19. Tan W, Aboulhosn J. The cardiovascular burden of coronavirus disease 2019 (COVID-19) with a focus on congenital heart disease. *Int J Cardiol.* 2020; 309: 70–77.
20. Frogoudaki AA, Farmakis D, Tsounis D et al., Telephone based survey in adults with congenital heart disease during COVID-19 pandemic. *Cardiol J.* 2020.
21. Coronavirus Disease 2019 in Children — United States, February 12–April 2, 2020. *CDC* April 2020.
22. Lewis MJ, Anderson BR, Fremed M, et al. The impact of coronavirus disease 2019 (COVID-19) on patients with congenital heart disease across the lifespan: the experience of an Academic Congenital Heart Disease Center in New York City. *JAHA* 2020.
23. Stout KK, Daniels CJ, Aboulhosn JA, et al. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019; 139: e698–e800.