

## Public attitudes towards grief in animals

N McGrath<sup>†‡</sup>, J Walker<sup>‡</sup>, D Nilsson<sup>‡</sup> and C Phillips<sup>\*‡</sup>

<sup>†</sup> The Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush, Roslin, Midlothian EH25 9RG, UK

<sup>‡</sup> Centre for Animal Welfare and Ethics, School of Veterinary Science, University of Queensland, Gatton 4343, QLD, Australia

\* Contact for correspondence and requests for reprints: c.phillips@uq.edu.au

### Abstract

Animals under human management are often separated from conspecifics, which may lead to behaviour indicative of separation distress or grief. For the purposes of this paper, grief is considered as a biological response to separation, indicated by a bi-phasic 'protest-despair' behavioural response. It is reasonable to assume that only animals which are able to form complex social bonds can experience grief. Scientific experiments have suggested that some farm and laboratory animals experience distress or grief as a result of maternal separation and social isolation. However, little is known about whether the public believe that animals are capable of grief. Therefore, we surveyed 1,000 members of the public to establish what knowledge they have about grief in animals and to compare this to what we know in science. The survey revealed that 90% of the general public believed that some or all animals can experience grief, with 23% believing that all animals can grieve. They attributed grief more to companion animals and animals with higher level cognitive abilities than to farm animals and animals that may be feared. It is concluded that public belief about grief in animals extends beyond scientific evidence, and that educating people about scientific findings and management practices connected with grief and separation distress may improve the welfare of farm and laboratory animals.

**Keywords:** animals, animal welfare, emotion, grief, public attitudes, separation

### Introduction

In human management systems, animals are often abruptly separated from their mothers (Newberry & Swanson 2008; Enriquez *et al* 2011) and other conspecifics (Siebert *et al* 2011) either temporarily or permanently. Both separation and social isolation may have a negative physiological and emotional effect on animals, which could directly impact upon their welfare.

Bowlby describes grief as:

A peculiar amalgam of anxiety, anger, and despair following the experience of what is feared to be an irretrievable loss...

and he differentiates it from separation anxiety saying:

Anxiety is experienced when the loss is believed to be retrievable and hope remains (Bowlby 1961).

Bowlby's description relates to the human experience of grief, and predominantly the psychological aspects of this emotion, the nature of which cannot be definitively known or inferred in animals. However, as emotions act to govern behaviour in response to specific events, behavioural and physiological changes are often induced by these events. Grief is considered by Averill (1968) to be:

A biological reaction, the evolutionary function of which is to ensure group cohesiveness in species where a social form of existence is necessary for survival...

Which also

Comprises a stereotyped set of psychological and physiological reactions.

Humans exhibit emotional responses to loss including depression, anxiety and anger, but may also suffer from physiological symptoms, such as sleeplessness, loss of appetite, and decreased immunological resistance (Boccia *et al* 1997; Stroebe *et al* 2007). It is difficult to know for certain whether a non-human animal has a conscious emotional experience of grief similar to that of a human. Our understanding of the subjective experience of animals is limited by their communication capabilities, and the potential differences between our psychological responses to stresses and theirs. However, some of these physiological symptoms could compromise an animal's ability to cope with survival situations, impair growth and reproduction, and increase susceptibility to disease (Weary & Chua 2000).

It is reasonable to assume that only animals which are able to form complex social bonds with conspecifics can experience grief as a result of severance of these bonds. A social bond is defined by Newberry and Swanson (2001) as:

A mutual, affectionate, emotional attachment between two individuals that is relatively long lasting and survives temporary separations.

Attachment helps govern behaviour in parent-offspring relationships (Bowlby 1958). Bowlby's early work on grief in humans outlined a bi-phasic 'protest-despair' reaction (Bowlby 1960). Experiments have demonstrated that animals go through distinct affective phases after separation, initially vocalising and exhibiting agitation (Seay *et al* 1962). These early reactions indicate distress and a desire for reunion (Hofer 1984, 2006; Topal *et al* 2005). A severe and sustained response may lead to the second phase of inactivity and withdrawal similar to the despair and depression exhibited by humans (Panksepp & Watt 2011). Prolonged separation of some animals has been shown to lead to physiological changes, such as increased heart rate, and behaviour indicative of despair, such as withdrawal (Seay *et al* 1962; Kraemer 1992) and increased self-directed behaviour (Bard & Nadler 1983). However, there can be no certainty that the psychological impact is the same as in humans.

Most scientific accounts of grief in animals, including the classic protest-despair phase, relate to primates (Seay *et al* 1962; Bard & Nadler 1983; Codner & Nadler 1984; Reite *et al* 1989; Laudenslager *et al* 1990; Cronin *et al* 2011), although anecdotal evidence suggests that elephants (*Elephas, Loxodonta* spp) and dolphins (Delphinidae) also experience grief as a result of chronic separation from conspecifics (Fertl & Schiro 1994; Herzing 2000; Douglas-Hamilton *et al* 2006), and that dogs (*Canis* spp) and cats (*Felis catus*) also experience a sustained grief response (Schwartz 2003).

There is ample literature providing evidence that cows (*Bos* spp), pigs (*Sus scrofa domestica*), sheep (*Ovis aries*), goats (*Capra hircus*), horses (*Equus caballus*) and chickens (*Gallus domesticus*) experience separation distress as a result of maternal separation and temporary or chronic isolation (eg Weary *et al* 1999; Watts *et al* 2001; Siebert *et al* 2011; Ungerfeld *et al* 2011). Separation distress, in these cases, typically includes distress vocalisations, increased activity or locomotion, escape attempts, and increased heart-rate, or increased hypothalamic-pituitary activity indicating stress. However, none of these accounts detail the bi-phasic protest-despair reaction typical of grief. This does not mean that these animals are incapable of grieving, and some other reactions, such as chronic stereotypies in horses, may be indicative of a prolonged reaction to separation (Waran 2001). Neuroscientific studies also suggest that mammals have the brain mechanisms that mediate grief reactions (Panksepp & Watt 2011), and animals are often used in scientific models of anxiety and depression (D'Aquila *et al* 1994; Cryan *et al* 2005; David *et al* 2009).

Farm animal production systems often result in permanent separation of farm animals: for example, maternal separation during weaning, separation from peers during herd changes, or as a result of death or sale. Temporary separations may involve social isolation during routine procedures or when an animal is sick. Weaning is a natural process which involves a reduction in maternal care and provision of milk initiated by the mother (Jensen 2001; Jasper *et al* 2008). However, this process can be stressful for young animals, and typically there are behavioural responses, such as increased locomotion and vocalisation which indicate

distress, and function to solicit attention from the caregiver (Weary *et al* 2008). In addition, early weaning can initially reduce growth rates (Jasper *et al* 2008).

Although scientific studies provide evidence that animals suffer under modern husbandry practices, such as during artificial weaning (Weary *et al* 2008), or social isolation, they are unlikely to bring about improvements in welfare unless the public has the appropriate knowledge about current practices and scientific findings (Serpell 2004). Public opinion has been the driving force behind significant legislation changes (Lawrence 2008). However, to date, no study has examined public belief about whether animals can experience grief. Therefore, our aim was to explore public beliefs about grief in animals and specifically to reveal what types of animals people believe can grieve, and compare this with scientific findings. Establishing the gap between these two knowledge bases would then provide the basis for further education of the public.

Studies which examine public opinion on the types of emotions animals may feel have been conducted, but these have been limited to dogs and other pets (Rasmussen 1995; Morris *et al* 2008). This questionnaire analysed the demographic effects on responses in order to most effectively target future education of the public regarding scientific evidence for grief in animals and the management practices that may produce such an emotional challenge.

## Materials and methods

### Participants and procedure

A face-to-face survey was conducted with 1,000 participants during April and May 2011 in four city-centre locations in Brisbane, Queensland, Australia. Three researchers collected survey answers. A script was carefully written for the survey and all collectors used this to ensure accurate standardised delivery of the questionnaire to all participants. Simple random sampling (De Vaus 2002) was used to select participants. All aspects of this research were approved by The University of Queensland Behavioural & Social Sciences Ethical Review Committee (reference number 2011000356).

### Questionnaire design

A pilot study was conducted using 25 randomly chosen participants in Brisbane city centre in April. Changes were made to the structure of some of the questions afterwards, as well as the order in which questions were asked, in order to reduce bias from questions about pet ownership that may have influenced answers to subsequent questions about animals in general. The new order was then tested on five students at the University of Queensland to determine if it reduced the potential bias. The resulting format was kept due to a consistently less-biased response from these test subjects.

The questionnaire was divided into three sections and comprised a combination of: i) questions to determine people's beliefs as to whether animals could experience emotions, with a specific focus on grief; ii) questions about pet ownership; and iii) demographic questions. The effects of pet ownership will be addressed in a separate paper.

The questionnaire gave a definition of grief which was derived from the literature on human grief (Archer 1999; Granek 2010):

An emotional reaction to loss, including sorrow, distress, sadness, anxiety and depression, which causes behavioural, emotional, mental, physical and social symptoms.

Subjects were asked to consider this definition when answering further questions. Participants were asked whether they believed any animal species could grieve. Respondents who answered 'some' or 'none' were then asked to state the reasons they thought some or all animal species might not grieve. Participants who agreed that some or all animal species could grieve were asked to specify which, from a pre-defined list of species.

### Statistical analysis

All questionnaire data were entered into Microsoft Excel (Microsoft 2007) and then exported to Minitab Version 15 for statistical analysis.

The marital status category was collapsed and re-coded post-survey completion to reflect the Australian Census 2006 categories of 'Never married', 'Married', 'Separated and Divorced', 'Widowed' (ABS 2006a) and to reduce statistical error from the small numbers of participants in some categories.

Multivariate analysis was used to examine associations between demographics and beliefs about grief in animals. Logistic regression models have an advantage over cross-tabulation analysis as they control for other variables within the model. This ensures that significant differences are not 'masked' and any significance found is a true reflection of how that factor is affecting response when all other variables are controlled for:

$$Z = b_0 + b_1X_1 + b_2X_2 + b_3X_3 + b_4X_4 + b_5X_5 + b_6X_6 + b_7X_7$$

Where  $Z$  is the log odds of the dependent variable,  $b_0$  is a constant,  $b_1$  = logistic regression coefficient for age ( $X_1$ ),  $b_2$  = logistic regression coefficient for gender ( $X_2$ ),  $b_3$  = logistic regression coefficient for nationality ( $X_3$ ),  $b_4$  = logistic regression coefficient for marital status ( $X_4$ ),  $b_5$  = logistic regression coefficient for education level ( $X_5$ ),  $b_6$  = logistic regression coefficient for income ( $X_6$ ) and  $b_7$  = logistic regression coefficient for where people lived ( $X_7$ ). The results are presented as odds ratios (OR) and  $P$ -values.

### Comment on the methodology

Random sampling was used when conducting this face-to-face survey. We approached members of the public at random, and those approached did not know the nature of the survey before we began, hence a biased response was minimised. Face-to-face surveys have a much higher response rate (in this survey a 70% response rate was achieved) than postal or web surveys which typically have response rates of 45 or 34%, respectively (Shih & Fan 2008).

## Results

A total of 1,000 completed responses were collected. One response was discarded from the data after it was collated due to inconsistency of responses.

Out of the total 999 people surveyed, 901 (90%) respondents believed animals could grieve according to our definition, with 229 (23%) saying all animals could grieve and 672 (67%) saying some animals could grieve (Table 1). Very high numbers of these respondents believed dogs ( $n = 883$ ; 98%), chimpanzees (*Pan troglodytes*) ( $n = 872$ ; 97%), dolphins ( $n = 850$ ; 94%) and elephants ( $n = 848$ ; 94%) could grieve (Figure 1). The majority of respondents believed that cats ( $n = 797$ ; 88%), pigs ( $n = 662$ ; 73%), cows ( $n = 637$ ; 71%) and magpies (*Pica* spp) ( $n = 530$ ; 59%) could experience grief. Less than half the respondents believed that turtles (Testudines) ( $n = 443$ ; 49%), bats (Chiroptera) ( $n = 359$ ; 40%), chickens ( $n = 358$ ; 40%), and crocodiles (Crocodylidae) ( $n = 353$ ; 39%) could grieve. At the lower end of the scale, only 10% ( $n = 91$ ) of respondents believed that prawns (Dendrobrachiata) could grieve, and low numbers of respondents believed that stick insects (*Carausius morosus*) ( $n = 107$ ; 12%), ants (Formicidae) ( $n = 160$ ; 18%) and fish (Pisces) ( $n = 168$ ; 19%) could grieve.

### Demographic effects

When asked whether all, some or no animals could grieve, females were more likely than males to respond that all animals could grieve (OR = 1.49,  $P = 0.007$ ; Table 2). They were also more likely than males to say that cats (OR = 2.99,  $P < 0.001$ ), dogs (OR = 3.82,  $P = 0.04$ ), pigs (OR = 1.47,  $P = 0.02$ ), chickens (OR = 1.69,  $P < 0.001$ ), crocodiles (OR = 1.48,  $P = 0.004$ ), bats (OR = 1.31,  $P = 0.049$ ), stick insects (OR = 1.61,  $P = 0.001$ ), magpies (OR = 1.50,  $P = 0.006$ ), turtles (OR = 1.50,  $P = 0.004$ ), ants (OR = 1.60,  $P = 0.001$ ) and fish (OR = 1.58,  $P = 0.002$ ) could grieve. However, there was no statistically significant difference between males and females in response to whether dolphins, elephants, cows or prawns could grieve ( $P > 0.05$ ).

As age increased, respondents were more likely to be uncertain as to whether elephants could grieve (OR = 0.74,  $P = 0.02$ ) and less likely to believe that cats (OR = 0.78,  $P = 0.01$ ), cows (OR = 0.87,  $P = 0.03$ ), pigs (OR = 0.85,  $P = 0.02$ ), chickens (OR = 0.85,  $P = 0.004$ ), crocodiles (OR = 0.85,  $P = 0.005$ ), bats (OR = 0.86,  $P = 0.007$ ) or turtles (OR = 0.82,  $P = 0.001$ ) could grieve (Table 3).

## Discussion

Our survey showed there was a strong public belief that animals can experience grief. Other studies have found that people believed animals could experience guilt, shame (Rasmussen 1995), and jealousy (Morris *et al* 2008). Emotions are therefore not seen as an exclusively human domain.

**Table 1 Participant Demographics (n = 999) showing a comparison of survey sample sizes with Brisbane and National Statistics.**

Statistics		n	Survey sample	Brisbane Statistical Division at June 2010	Census 2006 Brisbane	Census 2006 Australia
Age	18–25	271	27.1%	15.3%*		
	26–35	215	21.5%	15.2%		
	36–45	165	16.5%	14.7%		
	46–55	144	14.4%	13.2%		
	56–65	126	12.6%	10.6%		
	66+	78	7.8%	11.1%		
Gender	Male	521	52.2%		49.1%	49.4%
	Female	478	47.8%		50.9%	50.6%
Nationality	Australian	679	68.0%		72.0%	70.9%
	New Zealand	50	5.0%		4.1%	2.0%
	British	73	7.3%		5.2%	4.3%**
	Other	197	19.7%		18.7*	22.8%
Marital status (Two missing values)	Never married	523	52.5%		48.1%	49.6%
	Married	372	37.3%		35.0%	33.2%
	Separated or divorced	72	7.2%		11.8%	11.3%
	Widowed	30	3.0%		5.1%	5.9%
Education level	Primary	6	0.6%			
	Secondary	372	37.2%		48.4%	
	Certificate	89	8.9%		9.9%	
	Diploma	106	10.6%		7.6%	
	Undergraduate	236	23.6%			
	Postgraduate	190	19.0%		30.9%	
Income level (90 missing values)	Less than \$10k	177	17.7%			
	\$10,000–\$30,000	195	19.5%			
	\$30,001–\$50,000	151	15.1%			
	\$50,001–\$70,000	142	14.2%			
	\$70,001–\$90,000	96	9.6%			
	\$90,001–\$120,000	86	8.6%			
	\$120,001–\$150,000	27	2.7%			
> \$150,001	35	3.5%				

\* ABS source includes persons aged 15–24.

\*\* ABS source refers to England rather than Great Britain as country of birth.


Source: (ABS 2006a,b, 2011).

### Which animals can grieve?

Our aim was to establish which specific types of animals people believed could grieve. Dogs were the most frequently cited animal, which may reflect the high incidence of pet-ownership amongst our respondents. Only 5% of respondents had never owned a pet. Dogs form attachments to their owners which have been suggested to be analogous to those of human infants to their mothers (Topal *et al* 2005). Selection traits for breeding have tended towards neoteny

(McGreevy & Nicholas 1999), increasing the likelihood that dogs are more reliant on us, and develop strong attachment bonds. Strong responses to separation have been documented in scientific studies with dogs (Lund & Jorgensen 1999; Mendl *et al* 2010; Soares *et al* 2010), in addition to prolonged greeting behaviour after temporary separation (Flannigan & Dodman 2001; Konok *et al* 2011). Schwartz (2003) also suggested dogs exhibit signs of depression and social withdrawal which may be indicative of grief.

Figure 1



Percentage of 'Yes', 'No' and 'Don't know' responses to 'Can these animals grieve?'

Humans are prone to anthropomorphise about dogs (Serpell 2003) which is a reflection of the close relationship that we have developed with this domesticated animal. However, pet-owners have been demonstrated to provide meaningful and consistent reports about the behaviour of their animals, and the implication is that they can interpret these behavioural cues to infer what their animals may feel, using their expert knowledge of individual animals (Morris *et al* 2008). Current public knowledge about whether dogs can experience grief is consistent with the previously described scientific evidence.

The majority of the public believed that chimpanzees could grieve. There is an enormous amount of literature supporting the occurrence of grief in primates, as defined by a bi-phasic protest-despair behavioural reaction (Seay *et al* 1962; Mineka & Suomi 1978; Suomi 1983). Primates form highly complex social groups, including strong mother-infant bonds, and also attachment between peers. Examples of behaviour symptomatic of grief have been recorded in many primates. Some studies have revealed examples of withdrawal and self-directed behaviour in gorilla (*Gorilla gorilla* spp), orangutan (*Pongo* spp) and chimpanzee infants separated from their mother (Codner & Nadler 1984). Other experiments have provided evidence for a two-stage protest-despair reaction to maternal separation in rhesus macaques (*Macaca mulatta*) (Seay *et al* 1962), pig-tailed macaques (*Macaca nemestrina*) (Laudenslager *et al* 1990) and bonnet macaques (*Macaca radiata*) (Reite *et al* 1989), and increased vocalisations and stress hormones in squirrel monkeys (*Saimiri sciureus*) (Wiener *et al* 1990) as well as distress vocalisations in marmosets (Callitrichidae) (Norcross & Newman 1999). Maternally deprived rhesus macaques have also been used as a non-human primate model for depression (Paul *et al* 2000).

Maternal separation has been shown to evoke strong behavioural reactions in chimpanzees (Cronin *et al* 2011). Cronin *et al* documented activity demonstrating the existence of a sustained mother-infant bond even after the death of the infant. In this account, a mother chimpanzee was observed carrying its dead infant's body around and then exhibiting various behaviours, such as approaching the body, touching the face of the dead infant, and maintaining visual contact. In addition, members of the peer group also attended to the dead infant, approaching and touching the body (Cronin *et al* 2011). Another study by Bard and Nadler (1983) demonstrated a bi-phasic separation distress reaction to peer social separation in chimpanzees. Mineka and Suomi (1978) have commented that the separation response depends on many variables, such as behaviour prior to separation and the nature of the separation and reunion environment. Individual differences are also important factors in determining the level of attachment and therefore the strength of separation response (Hennessy 1997).

That the public believe that chimpanzees can grieve is most likely a reflection of their knowledge of scientific findings which are widely disseminated in the media, including evidence of chimpanzees' advanced cognitive abilities (Call & Tomasello 2008). The implications of these findings may affect the protection chimpanzees are afforded in the wild, and also improve welfare where chimpanzees are held captive under human management. Primates, such as macaques and marmosets, are still used in pre-clinical pharmaceutical toxicology trials (Smith *et al* 2001). Laboratory housing and testing practices may involve isolation from conspecifics, either permanently or temporarily. Therefore, there is the potential that some laboratory primates may suffer from grief.

**Table 2 Gender effects.**

		All animals		Some animals		No animals		Odds ratio (OR)	P-value
		n	%	n	%	n	%		
Can any animal grieve?	M	99	19%	362	69%	60	12%	1.49	0.007
	F	130	27%	310	65%	38	8%		
		Yes		No		Don't know		Odds ratio (OR)	P-value
		n	%	n	%	n	%		
Can cats grieve?	M	389	84%	44	10%	28	6%	2.99	0.000
	F	408	93%	14	3%	18	4%		
Can dogs grieve?	M	449	97%	5	1%	7	2%	3.82	0.040
	F	434	99%	4	1%	2	0%		
Can crocodiles grieve?	M	155	34%	202	44%	104	22%	1.48	0.004
	F	198	45%	139	32%	103	23%		
Can bats grieve?	M	169	37%	177	38%	115	25%	1.31	0.049
	F	190	43%	134	31%	116	26%		
Can ants grieve?	M	67	15%	318	69%	76	16%	1.60	0.001
	F	93	21%	246	56%	101	23%		
Can fish grieve?	M	63	14%	310	67%	88	19%	1.58	0.002
	F	105	24%	242	55%	93	21%		
Can turtles grieve?	M	204	44%	148	32%	109	24%	1.50	0.004
	F	239	54%	96	22%	105	24%		
Can stick insects grieve?	M	41	9%	319	69%	101	22%	1.61	0.001
	F	66	15%	253	57%	121	28%		
Can magpies grieve?	M	248	54%	129	28%	84	18%	1.50	0.006
	F	282	64%	95	22%	63	14%		
Can chickens grieve?	M	152	33%	223	48%	86	19%	1.69	0.000
	F	206	47%	162	37%	72	16%		
Can pigs grieve?	M	328	71%	73	16%	60	13%	1.47	0.020
	F	334	76%	55	13%	51	11%		

Responses to questions showing significant ( $P < 0.05$ ) gender effects on the proportion of 'Yes', 'No' and 'Don't know' responses. F: female; M: male.

Elephants and dolphins were widely believed to be capable of grief. Elephants mainly live in family units, which are held together by a matriarch which is responsible for the transmission of information, such as location of food, to other members of the social group (Douglas-Hamilton *et al* 2006). Field studies have documented strong reactions of other family members to the death of a matriarch (Douglas-Hamilton *et al* 2006) and the death of a calf (Payne 2003). Elephants have strong mother-calf relationships and weaning occurs naturally when the calf is four-years old. In addition, there are strong co-operative relationships between females in the group, which includes allomothering (Bates *et al* 2008). Bates *et al* (2008) report that elephants give excited greetings on reunion with other group members, and that the attention elephants give to bones and carcasses is consistent with them comprehending and responding to death of a conspecific. Distress vocalisa-

tions from calves which elicit responses from mothers and other members of the family group (allomothers and siblings) have been documented (Lee 1987). In addition, studies have shown that elephants have advanced cognitive abilities (Byrne *et al* 2009).

Dolphins and whales (Cetacea) also form complex social groups which are often co-operative, especially during predation (Mann & Smuts 1999; Simmonds 2006). Strong associations develop between mothers and infants, female kin (Wells *et al* 1987) and other adult females (Smolker *et al* 1992), and research has shown that some dolphins give allo-maternal care (Mann & Smuts 1998). Separation between mother and infant dolphins is common when the young forage (Mann & Smuts 1998, 1999). Some anecdotal accounts reveal behaviour which may indicate that these animals experience grief as a result of social loss (Fertl & Schiro 1994; Herzog 2000; Rose 2000).

Table 3 Age effects.

	Age	All animals		Some animals		No animals		Odds ratio (OR)	P-value
		n	%	n	%	n	%		
Can any animal grieve?	18–25	78	29%	173	64%	20	7%	0.91	0.12
	26–35	48	22%	152	71%	15	7%		
	36–45	31	19%	112	68%	22	13%		
	46–55	29	20%	103	72%	12	8%		
	56–65	27	21%	82	65%	17	13%		
	66+	16	21%	50	64%	12	15%		
	Age	Yes		No		Don't know		Odds ratio (OR)	P-value
		n	%	n	%	n	%		
Can elephants grieve?	18–25	243	97%	5	2%	3	1%	0.74	0.018
	26–35	185	92%	5	3%	10	5%		
	36–45	136	95%	0	0%	7	5%		
	46–55	124	94%	2	2%	6	4%		
	56–65	101	93%	0	0%	8	7%		
	66+	59	89%	1	2%	6	9%		
Can cats grieve?	18–25	237	95%	13	5%	1	0%	0.78	0.011
	26–35	181	90%	9	5%	10	5%		
	36–45	127	89%	11	8%	5	3%		
	46–55	110	83%	13	10%	9	7%		
	56–65	91	83%	6	6%	12	11%		
	66+	51	77%	6	9%	9	14%		
Can cows grieve?	18–25	188	75%	48	19%	15	6%	0.87	0.027
	26–35	151	75%	29	15%	20	10%		
	36–45	102	72%	19	13%	22	15%		
	46–55	79	60%	26	20%	27	20%		
	56–65	79	72%	17	16%	13	12%		
	66+	38	58%	10	15%	18	27%		
Can pigs grieve?	18–25	198	79%	42	17%	11	4%	0.85	0.017
	26–35	147	74%	27	13%	26	13%		
	36–45	118	83%	13	9%	12	8%		
	46–55	88	67%	17	13%	27	20%		
	56–65	78	71%	16	15%	15	14%		
	66+	33	50%	13	20%	20	30%		
Can chickens grieve?	18–25	136	54%	99	40%	16	6%	0.85	0.004
	26–35	76	38%	85	43%	39	19%		
	36–45	60	42%	57	40%	26	18%		
	46–55	41	31%	62	47%	29	22%		
	56–65	33	30%	51	47%	25	23%		
	66+	12	18%	31	47%	23	35%		

Responses to questions showing significant ( $P < 0.05$ ) age effects on the proportion of 'Yes', 'No and 'Don't know' responses.

Table 3 (cont)

	Age	Yes		No		Don't know		Odds ratio (OR)	P-value
		n	%	n	%	n	%		
Can turtles grieve?	18–25	162	65%	64	25%	25	10%	0.82	0.001
	26–35	110	55%	52	26%	38	19%		
	36–45	63	44%	36	25%	44	31%		
	46–55	44	33%	49	37%	39	30%		
	56–65	47	43%	24	22%	38	35%		
	66+	17	26%	19	29%	30	45%		
Can crocodiles grieve?	18–25	146	58%	77	31%	28	11%	0.85	0.005
	26–35	73	37%	81	40%	46	23%		
	36–45	49	34%	55	39%	39	27%		
	46–55	41	31%	60	45%	31	24%		
	56–65	31	28%	38	35%	40	37%		
	66+	13	20%	30	45%	23	35%		
Can bats grieve?	18–25	135	54%	92	36%	24	10%	0.86	0.007
	26–35	75	38%	76	38%	49	24%		
	36–45	60	42%	38	27%	45	31%		
	46–55	39	30%	48	36%	45	34%		
	56–65	36	33%	34	31%	39	36%		
	66+	14	21%	23	35%	29	44%		

Dolphins are also thought to be highly intelligent (Kuczaj *et al* 2009), and are reported to be capable of expressing empathy towards humans (Mercado III & De Long 2010). As much scientific evidence about elephant and dolphin intelligence is made available to the public via the media, people may make associations between this perceived intelligence/empathy, and the ability to grieve.

Cats are not typically portrayed as social animals. However, they do form affiliations and even strong relationships with individuals which form part of a complex social group (Crowell-Davis *et al* 2004). Kittens will emit distress vocalisations when the mother leaves the nest as newborns (Schwartz 2003). Schwartz (2003) also suggests that symptoms of grief, such as persistent depression or anxiety, are apparent in cats. That the public believed strongly that cats could grieve could again be a reflection of the high incidence of pet-ownership in our studies. There was a gender effect in our results, which is supported by the literature. Specific studies about the relationship between women and cats show that females are more likely to see cats as sentient (Phillips *et al* 2010), and to grieve more when their cats die (Wrobel & Dye 2003).

Over 70% of participants believed that pigs and cows could experience grief. Cows are highly social animals (Rushen *et al* 1999). The natural weaning age of cattle is

7–14 months (Reinhardt & Reinhardt 1981), but dairy calves are weaned early, often within 24 h after birth (Marchant-Forde *et al* 2002). This is for a variety of reasons, including reducing the welfare complications associated with weaning after a strong bond has formed (Weary & Chua 2000), to enable the mother to return to oestrus quickly, and to increase milk production (Flower & Weary 2003). Studies have shown that dairy cows and calves have an increased separation response as they get older (Weary & Chua 2000), which includes locomotion and vocalisation, and increased cortisol in calves (Loberg *et al* 2008) indicating stress. However, work done by Hopster *et al* (1995) did not detect a cortisol response in mothers when separated from their calves, although the authors caution that observation time and sample size were both restricted.

Cows also form attachments to their peers (Flower & Weary 2003). Social isolation of dairy cows for brief periods has also been shown to produce symptoms indicative of stress, inducing increased heart rate, hypothalamic-pituitary axis activity and vocalisation (Rushen *et al* 1999). High frequency call rates increase with the amount of social contact prior to separation (Boissy & Le Neindre 1997).

Beef calves are not often raised in isolation. However, artificial weaning is usually abrupt and necessitated by the need to encourage the cow to return to reproductive cyclicity



(Enriquez *et al* 2011), even though the age they are weaned approaches those of natural conditions (Weary *et al* 2008). As a result of the longer period of suckling, the mother-young bond is stronger than in dairy cows (Enriquez *et al* 2011). Beef cattle and calves have been shown to increase activity and vocalise in response to weaning (Ungerfeld *et al* 2011) and isolation (Watts *et al* 2001) and to have increased heart rate and vocalisations when separated from peers (Boissy & Le Neindre 1997). Social isolation also occurs when veal calves are isolated in crates (Bouissou *et al* 2001) and during routine procedures.

Pigs are a very gregarious species, which live naturally in social groupings of two to four sows and their litters (Gonyou 2001). The natural weaning age for piglets is around 4–5 months. Under commercial conditions, however, pigs are weaned abruptly at around 3–4 weeks of age to facilitate higher production, despite potential costs to welfare (Weary & Chua 2000). Piglets are known to stop eating for 1–3 days after weaning (Metz & Gonyou 1990) and studies have shown high rates of vocalisation, with younger piglets having a stronger reaction (Weary & Fraser 1997; Weary *et al* 1999). Pigs are also isolated socially in common practice (Gonyou 2001), and research indicates that vocalisations by isolated piglets are a good predictor of their level of need (Weary *et al* 1997).

Other farm animals also react to maternal separation and isolation. Sheep and goats are social animals, living in social groupings (Lyons *et al* 1993) and developing strong mother-offspring bonds (Fisher & Matthews 2001) and attachments to peers (Lyons *et al* 1993). Adult sheep maintain close relationships with the rest of the herd. They are also weaned earlier than they would be under natural conditions, which causes stress (Fisher & Matthews 2001). Both sheep and goats produce vocalisations and increased activity during social isolation (Carbonaro *et al* 1992; Lyons *et al* 1993; Orgeur *et al* 1999; Rault *et al* 2011; Siebert *et al* 2011), and sheep also experience increases in plasma cortisol during isolation, indicating stress (Cockram *et al* 1994; Guesdon *et al* 2012).

However, there have been no scientific studies on any of the farm animals discussed above (cows, pigs, sheep or goats) which have resulted in evidence for a two-stage protest-despair response characteristic of grief. This may be due to the fact that these are prey animals and therefore a period of withdrawal or inactivity could compromise their survival. In addition, there has been no systematic investigation into the long-term reactions to extensive separation periods. Despite this, there is ample evidence that these animals do experience distress, and the results of our survey suggest that the public are aware of this. However, Hills' (1993) study on Australian attitudes towards farm animals revealed ambivalent dispositions in many types of respondents. She found that the urban public did not assign more value to farm animals than would be expected from a utilitarian viewpoint. Their unwillingness to reconcile the values that they assign to animals with the desire to eat them may stem from widespread cultural

acceptance of livestock farming in Australia, and also from the fact that they do not see animal issues as being very important in their lives (Hills 1993). This observation is supported by social research, in which respondents rated animal welfare lowest out of a list of ten major issues they felt the government should address (Mazur 2006). Serpell (2004) points out that referring to farm animals as 'food' or 'production' animals reduces them to instrumental objects. The one remaining farm animal species from our survey is the chicken, which is also a social species, and chicks will emit distress calls when separated from their mothers (Kent 1987) or from conspecifics (Panksepp *et al* 1978; Marx *et al* 2001). In commercial systems, the young are rarely kept with the mother after birth (Mench & Keeling 2001). Only 40% of the public attributed grief to chickens. Conversely, nearly 60% believed Australian magpies (*Gymnorhina tibicen*) could grieve. Magpies are iconic Australian birds which live in co-operative groups of between 2–20 birds (Brown & Farabaugh 1997) and which share territory with humans (Kaplan 2004). Humans, therefore, often have the opportunity to observe their behaviour. Anecdotally, during our survey, people attributed intelligent behaviour to magpies, however there is no scientific evidence that they suffer from grief. The difference in attitudes to these two bird species supports the previous contention that people assign values to farm animals according to a utilitarian viewpoint (Hills 1993).

Bats are highly social animals which live in colonies and some are known to be nursed by their mothers for a month after birth (Matsumura 1981). In addition, it has been noted that vampire bats (Desmodontinae) perform reciprocal altruism, regurgitating meals for individuals which have been unsuccessful in finding food, and also practice mutual self-grooming (Denault & McFarlane 1995). Research has found evidence of vocalisations emitted in mother-infant situations. The researchers have suggested that these are isolation calls, given by the infant when the mother is flying in and out of the nursery colony, possibly as cues for location, facilitating recognition by the mother (Barclay *et al* 1979; Wilkinson 2003). However, only 40% of respondents attributed grief to bats, which is lower than any of our other mammals, and which placed them on a similar level to crocodiles. Crocodiles, however, are known to exhibit parental care (Burghardt 1977) and also to emit distress calls (Vergne *et al* 2009). The mother-infant bond is strong in some crocodiles, with the infants staying with the mother after hatching. Research has suggested that specific calls attract the mother's protection and facilitate group cohesion (Vergne *et al* 2011).

Crocodiles and bats are sometimes seen as animals to be feared and our results may reflect this belief. Previous research shows differences in attitudes associated with the phylogenetic relatedness of animals to humans (Eddy *et al* 1993; Harrison 2010; Phillips *et al* 2010), and the appearance of animals (Serpell 2003; Schlegel & Rupf 2010). Kellert (1985) found that people appeared to have more negative attitudes towards predators which could be related

to fear. Davey *et al* (1998) also suggest that phylogenetically based predispositions predict fear attitudes. Kellert (1985) found that older people had more negative attitudes towards predators, and our results support this.

Of the other animals presented in our survey, only ants are a social species. Ants form colonies of related individuals that participate in co-operative behaviour (Bos *et al* 2012). However, no studies have investigated separation distress in ants. Ants also voluntarily separate from the group if they are sick and die in isolation (Bos *et al* 2012), rather than soliciting care from nest-mates. Public knowledge about ants therefore seems to be consistent with the lack of scientific evidence for grief or distress in this species.

### Demographic effects

Our results imply some gender differences in empathy for animals. Women were more likely than men to say that all animals could grieve, and although evidence from pet-ownership shows that men and women like companion animals equally (Herzog 2007), other studies suggest women have greater empathy for animals (Hills 1993; Taylor & Signal 2005; Signal & Taylor 2006). This includes greater concern for animal use (Pifer *et al* 1994; Knight *et al* 2004), farm animal welfare (Hills 1993), and animal rights (Kruse 1999; Phillips *et al* 2010). It would therefore be logical to target women's support for animal welfare and protection issues. However, it is worth noting that the gender effects were not large across all the species.

### Animal welfare implications

That the overwhelming majority of the respondents in this survey believe that some animals can experience grief is significant for the welfare of many animals. This belief may result in increased demand for better welfare standards and could produce benefits in terms of animal welfare improvement. However, in the long term, the most benefits for animals would be seen where scientific evidence demonstrates what the animals need, and public demand for improved welfare standards is in line with this scientific evidence. Thus, further education of the public on modern husbandry practices may be needed in order to advance the welfare of these animals, in combination with consumer preferences for practices that avoid complex and negative emotions such as grief. In addition, more advanced scientific techniques to elucidate the emotional state of animals are required, without resorting to highly invasive physiological studies.

### Conclusion

There is a growing body of evidence which suggests that some animals have the requisite neural mechanisms that mediate reward processing in humans and animals (Berridge & Kringelbach 2008; Haber & Knutson 2010) and emotions in humans (Panksepp 1998, 2011; Damasio *et al* 2000; Burgdorf & Panksepp 2006). In addition, electrical brain stimulation (ESB) studies have managed to evoke behaviour suggestive of specific emotional states in some animals, by stimulating the brain networks associated with emotions, which are homologous in humans and other mammals. These stimulations appear to be either rewarding or punishing to the

animal, and Panksepp (2011) has suggested that this is evidence that the behaviour is linked to an affective experience. The position that some animals may have a conscious experience of emotions has many proponents (Cabanac 1999; Bekoff 2007; Burghardt 2007). However, in giving animals the benefit of the doubt as to whether they have emotions, we are making an assumption that they may be experiencing something similar to our own experience. This projection of our own experience is also necessary with other humans, as we cannot always be confident that what another human is experiencing, or even reporting, is the same as our experiences in similar situations.

It is clear from our study that public beliefs about grief in some animals extend beyond current scientific findings. Although a grief response, defined by a bi-phasic protest-despair reaction, has not been reported in farm animals, such as cows and pigs, over 70% of respondents believed these animals could grieve in ways described for humans. These results have implications for the welfare of animals. For example, the public may desire welfare standards that acknowledge emotions for which there is currently little evidence.

People evidently believe that animals experience psychological grief to the loss of a conspecific, and this belief may stem from empathy, which is an ability to relate behaviour to feelings and to share the perceived emotional experience of an animal. However, it may also be due to people anthropomorphising, which involves projecting human-like experiences and behaviours on to animals (Wynne 2007). Both these possibilities have implications for the welfare of animals under our care. The existence of empathy may mean people increasingly give the benefit of the doubt to animals, and make or would like to make adjustments to management practices to ensure no animal suffers unnecessarily. However, in contrast to this, anthropomorphism may lead to the inappropriate treatment of animals which may, in turn, compromise their welfare. For example, owners' overprotective behaviour towards their companion animals may result in the animals developing neuroses, such as separation anxiety.

Our study suggests that there is scope for educating the public on specific human management practices connected with separation distress in order to improve welfare for animals under our care. Although potentially not as strong an emotional reaction as grief, separation distress impacts severely on the behaviour and physiology of animals, such that their welfare can be compromised. Public knowledge about these practices and the effects they have on some animals, such as farm animals, or laboratory animals, could help drive improvements in animal welfare, supported by the relevant scientific evidence.

### Acknowledgements

Firstly, our thanks go to the Morris Animal Foundation for funding this research. We are grateful to Sian Ringrose for proof-reading and commenting on our original questionnaire, and to Sue Mitchell for providing technical guidance on statistical analyses. We also thank Alejandra Montoya for her assistance during data collection.

## References

- ABS** 2006a *Australian Bureau of Statistics*. 2006 Census QuickStats: Brisbane (Statistical Division). Viewed 27th March, 2012. <http://www.censusdata.abs.gov.au/ABSNavigation/prenav/LocationSearch?locationLastSearchTerm=Brisbane&locationSearchTerm=Brisbane&newarea=305&submitbutton=View+QuickStats+%3E&collection=census&period=2006&areacode=305&geography=&method=Place+of+Usual+Residence&productlabel=&producttype=QuickStats&topic=&navmapdisplayed=false&javascript=true&breadcrumb=PL&topholder=0&leftholder=0&currentaction=104&action=401&textversion=false&subaction=1>
- ABS** 2006b *Australian Bureau of Statistics*. 2006 Census Quickstats: Australia. Viewed 11 June 2011 <http://www.censusdata.abs.gov.au/ABSNavigation/prenav/ProductSelect:newproducttype=QuickStats&btnSelectProduct=View+QuickStats+%3E&collection=Census&period=2006&areacode=0&geography=&method=&productlabel=&producttype=&topic=&navmapdisplayed=true&javascript=true&breadcrumb=LP&topholder=0&leftholder=0&currentaction=201&action=401&textversion=false>
- ABS** 2011 *Australian Bureau of Statistics*. National Regional Profile: Brisbane (Statistical Division). Population. Viewed 11 June 2011. <http://www.abs.gov.au/AUSSTATS/abs@nrp.nsf/Latestproducts/305Population/People12006-2010?opendocument&tabname=Summary&prodno=305&issue=2006-2010>
- Archer J** 1999 *The Nature of Grief: The Evolution and Psychology of Reactions to Loss*. Routledge: London, UK
- Averill JR** 1968 Grief: its nature and significance. *Psychological Bulletin* 70: 721-748. <http://dx.doi.org/10.1037/h0026824>
- Barclay RMR, Fenton MB and Thomas DW** 1979 Social-behaviour of the little brown bat, *Myotis lucifugus*. *Behavioral Ecology and Sociobiology* 6: 137-146. <http://dx.doi.org/10.1007/BF00292560>
- Bard KA and Nadler RD** 1983 The effect of peer separation in young chimpanzees (*Pan troglodytes*). *American Journal of Primatology* 5: 25-37. <http://dx.doi.org/10.1002/ajp.1350050104>
- Bates LA, Poole JH and Byrne RW** 2008 Elephant cognition. *Current Biology* 18: R544-R546. <http://dx.doi.org/10.1016/j.cub.2008.04.019>
- Bekoff M** 2007 *The Emotional Lives of Animals*. New World Library: Novato, USA
- Berridge KC and Kringelbach ML** 2008 Affective neuroscience of pleasure: reward in humans and animals. *Psychopharmacology* 199: 457-480. <http://dx.doi.org/10.1007/s00213-008-1099-6>
- Boccia ML, Scanlan JM, Laudenslager ML, Berger CL, Hijazi AS and Reite ML** 1997 Juvenile friends, behavior, and immune responses to separation in bonnet macaque infants. *Physiology & Behavior* 61: 191-198. [http://dx.doi.org/10.1016/S0031-9384\(96\)00370-8](http://dx.doi.org/10.1016/S0031-9384(96)00370-8)
- Boissy A and Le Neindre P** 1997 Behavioral, cardiac and cortisol responses to brief peer separation and reunion in cattle. *Physiology & Behavior* 61: 693-699. [http://dx.doi.org/10.1016/S0031-9384\(96\)00521-5](http://dx.doi.org/10.1016/S0031-9384(96)00521-5)
- Bos N, Lefevre T, Jensen AB and d'Ettorre P** 2012 Sick ants become unsociable. *Journal of Evolutionary Biology* 25: 342-351. <http://dx.doi.org/10.1111/j.1420-9101.2011.02425.x>
- Bouissou M-F, Boissy A, Le Neindre P and Veissier I** 2001 The social behaviour of cattle. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 113-145. CAB International: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0113>
- Bowlby J** 1958 The nature of the child's tie to his mother. *International Journal of Psycho-Analysis* 39: 350-373
- Bowlby J** 1960 Grief and mourning in infancy and early-childhood. *Psychoanalytic Study of the Child* 15: 9-52
- Bowlby J** 1961 Processes of mourning. *International Journal of Psychoanalysis* 42: 317-340
- Brown ED and Farabaugh SM** 1997 What birds with complex social relationships can tell us about vocal learning: vocal sharing in avian groups. In: Snowdon C and Hausberger M (eds) *Social Influences on Vocal Development* pp 98-127. Cambridge University Press: Cambridge, UK. <http://dx.doi.org/10.1017/CBO9780511758843.007>
- Burgdorf J and Panksepp J** 2006 The neurobiology of positive emotions. *Neuroscience and Biobehavioral Reviews* 30: 173-187. <http://dx.doi.org/10.1016/j.neubiorev.2005.06.001>
- Burghardt G** 1977 Of iguanas and dinosaurs: social behaviour and communication in neonate reptiles. *American Zoologist* 17: 177-190
- Burghardt G** 2007 Critical anthropomorphism, uncritical anthropomorphism and naive nominalism. *Comparative Cognitive Behavior Reviews* 2: 136-138
- Byrne RW, Bates LA and Moss CJ** 2009 Elephant cognition in primate perspective. *Comparative Cognition & Behavior Reviews* 4: 65-79
- Cabanac M** 1999 Emotion and Phylogeny. *Japanese Journal of Physiology* 49: 1-10. <http://dx.doi.org/10.2170/jjphysiol.49.1>
- Call J and Tomasello M** 2008 Does the chimpanzee have a theory of mind? 30 years later. *Trends in Cognitive Sciences* 12: 187-192. <http://dx.doi.org/10.1016/j.tics.2008.02.010>
- Carbonaro DA, Friend TH, Dellmeier GR and Nuti LC** 1992 Behavioral and physiological responses of dairy goats to isolation. *Physiology & Behavior* 51: 297-301. [http://dx.doi.org/10.1016/0031-9384\(92\)90144-Q](http://dx.doi.org/10.1016/0031-9384(92)90144-Q)
- Cockram MS, Ranson M, Imlah P, Goddard PJ, Burrells C and Harkiss GD** 1994 The behavioural, endocrine and immune responses of sheep to isolation. *Animal Production* 58: 389-399. <http://dx.doi.org/10.1017/S0003356100007339>
- Codner MA and Nadler RD** 1984 Mother-infant separation and reunion in the great apes. *Primates* 25: 204-217. <http://dx.doi.org/10.1007/BF02382392>
- Cronin KA, van Leeuwen EJC, Mulenga IC and Bodamer MD** 2011 Behavioral response of a chimpanzee mother toward her dead infant. *American Journal of Primatology* 73: 415-421. <http://dx.doi.org/10.1002/ajp.20927>
- Crowell-Davis SL, Curtis TM and Knowles RJ** 2004 Social organization in the cat: a modern understanding. *Journal of Feline Medicine and Surgery* 6: 19-28. <http://dx.doi.org/10.1016/j.jfms.2003.09.013>
- Cryan JF, Mombereau C and Vassout A** 2005 The tail suspension test as a model for assessing antidepressant activity: review of pharmacological and genetic studies in mice. *Neuroscience & Biobehavioral Reviews* 29: 571-625. <http://dx.doi.org/10.1016/j.neubiorev.2005.03.009>
- Damasio AR, Grabowski TJ, Bechara A, Damasio H, Ponto LLB, Parvizi J and Hichwa RD** 2000 Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nature Neuroscience* 3: 1049-1056. <http://dx.doi.org/10.1038/79871>
- D'Aquila PS, Brain P and Willner P** 1994 Effects of chronic mild stress on performance in behavioural tests relevant to anxiety and depression. *Physiology & Behavior* 56: 861-867. [http://dx.doi.org/10.1016/0031-9384\(94\)90316-6](http://dx.doi.org/10.1016/0031-9384(94)90316-6)

- Davey GCL, McDonald AS, Hirisave U, Prabhu GG, Iwawaki S, Jim CI, Merckelbach H, de Jong PJ, Leung PWL and Reimann BC** 1998 A cross-cultural study of animal fears. *Behaviour Research and Therapy* 36: 735-750. [http://dx.doi.org/10.1016/S0005-7967\(98\)00059-X](http://dx.doi.org/10.1016/S0005-7967(98)00059-X)
- David DJ, Samuels BA, Rainer Q, Wang J-W, Marsteller D, Mendez I, Drew M, Craig DA, Guiard BP, Guilloux J-P, Artymyshyn RP, Gardier AM, Gerald C, Antonijevic IA, Leonardo ED and Hen R** 2009 Neurogenesis-dependent and independent effects of fluoxetine in an animal model of anxiety/depression. *Neuron* 62: 479-493. <http://dx.doi.org/10.1016/j.neuron.2009.04.017>
- Denault LK and McFarlane DA** 1995 Reciprocal altruism between male vampire bats, *Desmodus rotundus*. *Animal Behaviour* 49: 855-856
- De Vaus DA** 2002 *Surveys in Social Research*. Allen & Unwin: Sydney, Australia
- Douglas-Hamilton I, Bhalla S, Wittermyer G and Vollrath F** 2006 Behavioural reactions of elephants towards a dying and deceased matriarch. *Applied Animal Behaviour Science* 100: 87-102. <http://dx.doi.org/10.1016/j.applanim.2006.04.014>
- Eddy TJ, Gallup GG and Povinelli DJ** 1993 Attribution of cognitive states to animals. Anthropomorphism in comparative perspective. *Journal of Social Issues* 49: 87-101. <http://dx.doi.org/10.1111/j.1540-4560.1993.tb00910.x>
- Enriquez D, Hotzel MJ and Ungerfeld R** 2011 Minimising the stress of weaning of beef calves: a review. *Acta Veterinaria Scandinavica* 53: 28. <http://dx.doi.org/10.1186/1751-0147-53-28>
- Fertl D and Schiro A** 1994 Carrying of dead calves by free-ranging Texas bottlenose dolphins (*Tursiops truncatus*). *Aquatic Mammals* 20: 53-56
- Fisher A and Matthews L** 2001 The social behaviour of sheep. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 211-245. CABI Publishing: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0211>
- Flannigan G and Dodman NH** 2001 Risk factors and behaviors associated with separation anxiety in dogs. *Journal of the American Veterinary Medical Association* 219: 460-466. <http://dx.doi.org/10.2460/javma.2001.219.460>
- Flower FC and Weary DM** 2003 The effects of early separation on the dairy cow and calf. *Animal Welfare* 12: 339-348
- Gonyou HW** 2001 The social behaviour of pigs. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 147-176. CABI Publishing: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0147>
- Granek L** 2010 Grief as pathology: the evolution of grief theory in psychology from Freud to the present. *History of Psychology* 13: 46-73. <http://dx.doi.org/10.1037/a0016991>
- Guesdon V, Ligout S, Delagrance P, Spedding M, Levy F, Laine A-L, Malpoux B and Chaillou E** 2012 Multiple exposures to familiar conspecific withdrawal is a novel robust stress paradigm in ewes. *Physiology & Behavior* 105: 203-208. <http://dx.doi.org/10.1016/j.physbeh.2011.08.011>
- Haber SN and Knutson B** 2010 The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology* 35: 4-26. <http://dx.doi.org/10.1038/npp.2009.129>
- Harrison MA** 2010 Anthropomorphism, empathy, and perceived communicative ability vary with phylogenetic relatedness to humans. *Journal of Social, Evolutionary, and Cultural Psychology* 4: 34-48
- Hennessy MB** 1997 Hypothalamic-pituitary-adrenal responses to brief social separation. *Neuroscience and Biobehavioral Reviews* 21: 11-29. [http://dx.doi.org/10.1016/S0149-7634\(96\)00013-9](http://dx.doi.org/10.1016/S0149-7634(96)00013-9)
- Herzing DL** 2000 A trail of grief. In: Bekoff M (ed) *The Smile of a Dolphin: Remarkable Accounts of Animal Emotions*. Random House: New York, USA
- Herzog HA** 2007 Gender differences in human-animal interactions: a review. *Anthrozoös* 20: 7-21. <http://dx.doi.org/10.2752/089279307780216687>
- Hills AM** 1993 The motivational bases of attitudes toward animals. *Society & Animals* 1: 111-128. <http://dx.doi.org/10.1163/156853093X00028>
- Hofer MA** 1984 Relationships as regulators: a psychobiologic perspective on bereavement. *Psychosomatic Medicine* 46: 183-197
- Hofer MA** 2006 Psychobiological roots of early attachment. *Current Directions in Psychological Science* 15: 84-88. <http://dx.doi.org/10.1111/j.0963-7214.2006.00412.x>
- Hopster H, O'Connell JM and Blokhuis HJ** 1995 Acute effects of cow-calf separation on heart rate, plasma cortisol and behaviour in multiparous dairy cows. *Applied Animal Behaviour Science* 44: 1-8. [http://dx.doi.org/10.1016/0168-1591\(95\)00581-C](http://dx.doi.org/10.1016/0168-1591(95)00581-C)
- Jasper J, Budzynska M and Weary DM** 2008 Weaning distress in dairy calves: acute behavioural responses by limit-fed calves. *Applied Animal Behaviour Science* 110: 136-143. <http://dx.doi.org/10.1016/j.applanim.2007.03.017>
- Jensen P** 2001 Parental behaviour. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 59-81. CABI Publishing: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0059>
- Kaplan G** 2004 *Australian magpie: biology and behaviour of an unusual songbird*. CSIRO Publishing: Australia
- Kellert SR** 1985 Public perceptions of predators, particularly the wolf and coyote. *Biological Conservation* 31: 167-189. [http://dx.doi.org/10.1016/0006-3207\(85\)90047-3](http://dx.doi.org/10.1016/0006-3207(85)90047-3)
- Kent JP** 1987 Experiments on the relationship between the hen and the chick (*Gallus gallus*): the role of the auditory mode in recognition and the effects of maternal separation. *Behaviour* 102: 1-14. <http://dx.doi.org/10.1163/156853986X00018>
- Knight S, Vrij A, Cherryman V and Nunukoosing K** 2004 Attitudes towards animal use and belief in animal mind. *Anthrozoös* 17: 43-62. <http://dx.doi.org/10.2752/089279304786991945>
- Konok V, Doka A and Miklosi A** 2011 The behavior of the domestic dog (*Canis familiaris*) during separation from and reunion with the owner: a questionnaire and an experimental study. *Applied Animal Behaviour Science* 135: 300-308. <http://dx.doi.org/10.1016/j.applanim.2011.10.011>
- Kraemer GW** 1992 A psychobiological theory of attachment. *Behavioral and Brain Sciences* 15: 493-541. <http://dx.doi.org/10.1017/S0140525X00069752>
- Kruse CR** 1999 Gender, views of nature, and support for animal rights. *Society & Animals* 7: 179-198. <http://dx.doi.org/10.1163/156853099X00077>
- Kuczaj SA II, Gory JD and Xitco MJ Jr** 2009 How intelligent are dolphins? A partial answer based on their ability to plan their behavior when confronted with novel problems. *Japanese Journal of Animal Psychology* 59: 99-115. <http://dx.doi.org/10.2502/j.anip.59.1.9>

- Laudenslager ML, Held PE, Boccia ML, Reite ML and Cohen JJ** 1990 Behavioral and immunological consequences of brief mother-infant separation: a species comparison. *Developmental Psychobiology* 23: 247-264. <http://dx.doi.org/10.1002/dev.420230305>
- Lawrence AB** 2008 Applied animal behaviour science: past, present and future prospects. *Applied Animal Behaviour Science* 115: 1-24. <http://dx.doi.org/10.1016/j.applanim.2008.06.003>
- Lee PC** 1987 Allomothering among African elephants. *Animal Behaviour* 35: 278-291. [http://dx.doi.org/10.1016/S0003-3472\(87\)80234-8](http://dx.doi.org/10.1016/S0003-3472(87)80234-8)
- Loberg JM, Hernandez CE, Thierfelder T, Jensen MB, Berg C and Lidfors L** 2008 Weaning and separation in two steps: a way to decrease stress in dairy calves suckled by foster cows. *Applied Animal Behaviour Science* 111: 222-234. <http://dx.doi.org/10.1016/j.applanim.2007.06.011>
- Lund JD and Jorgensen MC** 1999 Behaviour patterns and time course of activity in dogs with separation problems. *Applied Animal Behaviour Science* 63: 219-236. [http://dx.doi.org/10.1016/S0168-1591\(99\)00011-8](http://dx.doi.org/10.1016/S0168-1591(99)00011-8)
- Lyons DM, Price EO and Moberg GP** 1993 Social grouping tendencies and separation-induced distress in juvenile sheep and goats. *Developmental Psychobiology* 26: 251-259. <http://dx.doi.org/10.1002/dev.420260503>
- Mann J and Smuts BB** 1998 Natal attraction: allomaternal care and mother-infant separations in wild bottlenose dolphins. *Animal Behaviour* 55: 1097-1113. <http://dx.doi.org/10.1006/anbe.1997.0637>
- Mann J and Smuts BB** 1999 Behavioral development in wild bottlenose dolphin newborns (*Tursiops* spp). *Behaviour* 136: 529-566. <http://dx.doi.org/10.1163/156853999501469>
- Marchant-Forde JN, Marchant-Forde RM and Weary DM** 2002 Responses of dairy cows and calves to each other's vocalisations after early separation. *Applied Animal Behaviour Science* 78: 19-28. [http://dx.doi.org/10.1016/S0168-1591\(02\)00082-5](http://dx.doi.org/10.1016/S0168-1591(02)00082-5)
- Marx G, Leppelt J and Ellendorff F** 2001 Vocalisation in chicks (*Gallus gallus*) during stepwise social isolation. *Applied Animal Behaviour Science* 75: 61-74. [http://dx.doi.org/10.1016/S0168-1591\(01\)00180-0](http://dx.doi.org/10.1016/S0168-1591(01)00180-0)
- Matsumura S** 1981 Mother-infant communication in a horse-shoe bat (*Rhinolophus ferrumequinum nippon*): vocal communication in 3 week-old infants. *Journal of Mammalogy* 62: 20-28. <http://dx.doi.org/10.2307/1380474>
- Mazur N** 2006 *Social research to support the Australian animal welfare strategy*. ENVision Environmental Consulting: Australia
- McGreevy PD and Nicholas FW** 1999 Some practical solutions to welfare problems in dog breeding. *Animal Welfare* 8: 329-341
- Mench J and Keeling L** 2001 The social behaviour of domestic birds. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 177-209. CABI Publishing: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0177>
- Mendl M, Brooks J, Basse C, Burman O, Paul E, Blackwell E and Casey R** 2010 Dogs showing separation-related behaviour exhibit a 'pessimistic' cognitive bias. *Current Biology* 20: R839-R840. <http://dx.doi.org/10.1016/j.cub.2010.08.030>
- Mercado III E and De Long C** 2010 Dolphin cognition: representations and processes in memory and perception. *International Journal of Comparative Psychology* 23: 344-378
- Metz JHM and Gonyou HW** 1990 Effect of age and housing conditions on the behavioral and hemolytic reaction of piglets to weaning. *Applied Animal Behaviour Science* 27: 299-309. [http://dx.doi.org/10.1016/0168-1591\(90\)90126-X](http://dx.doi.org/10.1016/0168-1591(90)90126-X)
- Mineka S and Suomi SJ** 1978 Social separation in monkeys. *Psychological Bulletin* 85: 1376-1400. <http://dx.doi.org/10.1037/0033-2909.85.6.1376>
- Morris P, Doe C and Godsell E** 2008 Secondary emotions in non-primate species? Behavioural reports and subjective claims by animal owners. *Cognition & Emotion* 22: 3-20. <http://dx.doi.org/10.1080/02699930701273716>
- Newberry R and Swanson J** 2001 Breaking social bonds. In: Keeling LJ and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 307-331. CABI Publishing: Wallingford, UK. <http://dx.doi.org/10.1079/9780851993973.0307>
- Newberry R and Swanson J** 2008 Implications of breaking mother-young social bonds. *Applied Animal Behaviour Science* 110: 3-23. <http://dx.doi.org/10.1016/j.applanim.2007.03.021>
- Norcross JL and Newman JD** 1999 Effects of separation and novelty on distress vocalizations and cortisol in the common marmoset (*Callithrix jacchus*). *American Journal of Primatology* 47: 209-222. [http://dx.doi.org/10.1002/\(SICI\)1098-2345\(1999\)47:3<209::AID-AJP3>3.0.CO;2-0](http://dx.doi.org/10.1002/(SICI)1098-2345(1999)47:3<209::AID-AJP3>3.0.CO;2-0)
- Orgeur P, Bernard S, Naciri M, Nowak R, Schaal B and Levy F** 1999 Psychobiological consequences of two different weaning methods in sheep. *Reproduction Nutrition Development* 39: 231-244. <http://dx.doi.org/10.1051/rnd:19990208>
- Panksepp J** 1998 *Affective Neuroscience: The Foundations of Human and Animal Emotions*. Oxford University Press: Oxford, UK
- Panksepp J** 2011 Toward a cross-species neuroscientific understanding of the affective mind: do animals have emotional feelings? *American Journal of Primatology* 73: 545-561. <http://dx.doi.org/10.1002/ajp.20929>
- Panksepp J, Vilberg T, Bean NJ, Coy DH and Kastin AJ** 1978 Reduction of distress vocalizations in chicks by opiate-like peptides. *Brain Research Bulletin* 3: 663-667. [http://dx.doi.org/10.1016/0361-9230\(78\)90014-X](http://dx.doi.org/10.1016/0361-9230(78)90014-X)
- Panksepp J and Watt D** 2011 Why does depression hurt? Ancestral primary-process separation-distress (PANIC/GRIEF) and diminished brain reward (SEEKING) processes in the genesis of depressive affect. *Psychiatry* 74: 5-13. <http://dx.doi.org/10.1521/psyc.2011.74.1.5>
- Paul IA, English JA and Halaris A** 2000 Sucrose and quinine intake by maternally deprived and control rhesus monkeys. *Behavioural Brain Research* 112: 127-134. [http://dx.doi.org/10.1016/S0166-4328\(00\)00173-X](http://dx.doi.org/10.1016/S0166-4328(00)00173-X)
- Payne K** 2003 Sources of social complexity in the three elephant species. In: de Waal FBM and Tyack PL (eds) *Animal Social Complexity: Intelligence, Culture and Individualized Societies* pp 57-85. Harvard University Press: Cambridge, MA, USA
- Phillips C, Izmirlı S, Aldavood J, Alonso M, Choe B, Hanlon A, Handziska A, Illmann G, Keeling L, Kennedy M, Lee G, Lund V, Mejdell C, Pelagic V and Rehn T** 2010 An international comparison of female and male students' attitudes to the use of animals. *Animals* 1: 7-26. <http://dx.doi.org/10.3390/ani1010007>
- Pifer L, Shimizu K and Pifer R** 1994 Public attitudes toward animal research: some international comparisons. *Society & Animals* 2: 95-113. <http://dx.doi.org/10.1163/156853094X00126>

- Rasmussen JL** 1995 Differences and similarities in humans' perceptions of the thinking and feeling of a dog and a boy. *Society & Animals* 3: 117-137. <http://dx.doi.org/10.1163/156853095X00116>
- Rault JL, Boissy A and Boivin X** 2011 Separation distress in artificially reared lambs depends on human presence and the number of conspecifics. *Applied Animal Behaviour Science* 132: 42-50. <http://dx.doi.org/10.1016/j.applanim.2011.02.011>
- Reinhardt V and Reinhardt A** 1981 Natural sucking performance and age of weaning in Zebu cattle (*Bos indicus*). *Journal of Agricultural Science* 96: 309-312. <http://dx.doi.org/10.1017/S0021859600066089>
- Reite M, Kaemingk K and Boccia ML** 1989 Maternal separation in bonnet monkey infants: altered attachment and social support. *Child Development* 60: 473-480. <http://dx.doi.org/10.2307/1130991>
- Rose N** 2000 A death in the family. In: Bekoff M (ed) *The Smile of a Dolphin: Remarkable Accounts of Animal Emotions*. Random House/Discovery Books: New York, USA
- Rushen J, Boissy A, Terlouw EMC and de Passillé AMB** 1999 Opioid peptides and behavioral and physiological responses of dairy cows to social isolation in unfamiliar surroundings. *Journal of Animal Science* 77: 2918-2924
- Schlegel J and Rupf R** 2010 Attitudes towards potential animal flagship species in nature conservation: a survey among students of different educational institutions. *Journal for Nature Conservation* 18: 278-290. <http://dx.doi.org/10.1016/j.jnc.2009.12.002>
- Schwartz S** 2003 Separation anxiety syndrome in dogs and cats. *Journal of the American Veterinary Medical Association* 222: 1526-1532. <http://dx.doi.org/10.2460/javma.2003.222.1526>
- Seay B, Hansen E and Harlow HF** 1962 Mother-infant separation in monkeys. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 3: 123-132. <http://dx.doi.org/10.1111/j.1469-7610.1962.tb02047.x>
- Serpell JA** 2003 Anthropomorphism and anthropomorphic selection: beyond the 'cute response'. *Society & Animals* 11: 83-100. <http://dx.doi.org/10.1163/156853003321618864>
- Serpell JA** 2004 Factors influencing human attitudes to animals and their welfare. *Animal Welfare* 13: S145-S151
- Shih T-H and Fan X** 2008 Comparing response rates from web and mail surveys: a meta-analysis. *Field Methods* 20: 249-271. <http://dx.doi.org/10.1177/1525822X08317085>
- Siebert K, Langbein J, Schoen P-C, Tuchscherer A and Puppe B** 2011 Degree of social isolation affects behavioural and vocal response patterns in dwarf goats (*Capra hircus*). *Applied Animal Behaviour Science* 131: 53-62. <http://dx.doi.org/10.1016/j.applanim.2011.01.003>
- Signal TD and Taylor N** 2006 Attitudes to animals: demographics within a community sample. *Society & Animals* 14: 147-157. <http://dx.doi.org/10.1163/156853006776778743>
- Simmonds MP** 2006 Into the brains of whales. *Applied Animal Behaviour Science* 100: 103-116. <http://dx.doi.org/10.1016/j.applanim.2006.04.015>
- Smith D, Trennery P, Farningham D and Klapwijk J** 2001 The selection of marmoset monkeys (*Callithrix jacchus*) in pharmaceutical toxicology. *Laboratory Animals* 35: 117-130. <http://dx.doi.org/10.1258/0023677011911444>
- Smolker RA, Richards AF, Connor RC and Pepper JW** 1992 Sex differences in patterns of association among Indian-Ocean bottlenose dolphins. *Behaviour* 123: 38-69. <http://dx.doi.org/10.1163/156853992X00101>
- Soares GM, Pereira JT and Paixao RL** 2010 Exploratory study of separation anxiety syndrome in apartment dogs. *Ciencia Rural* 40: 548-553. <http://dx.doi.org/10.1590/S0103-84782010000300008>
- Stroebe M, Schut H and Stroebe W** 2007 Health outcomes of bereavement. *Lancet* 370: 1960-1973. [http://dx.doi.org/10.1016/S0140-6736\(07\)61816-9](http://dx.doi.org/10.1016/S0140-6736(07)61816-9)
- Suomi SJ** 1983 Models of depression in primates. *Psychological Medicine* 13: 465-468. <http://dx.doi.org/10.1017/S0033291700047887>
- Taylor N and Signal TD** 2005 Empathy and attitudes to animals. *Anthrozoös* 18: 18-27. <http://dx.doi.org/10.2752/089279305785594342>
- Topal J, Gacsi M, Miklosi A, Viranyi Z, Kubinyi E and Csanyi V** 2005 Attachment to humans: a comparative study on hand-reared wolves and differently socialized dog puppies. *Animal Behaviour* 70: 1367-1375. <http://dx.doi.org/10.1016/j.anbehav.2005.03.025>
- Ungerfeld R, Hotzel MJ, Scarsi A and Quintans G** 2011 Behavioral and physiological changes in early-weaned multiparous and primiparous beef cows. *Animal* 5: 1270-1275. <http://dx.doi.org/10.1017/S1751731111000334>
- Vergne AL, Aubin T, Taylor P and Mathevon N** 2011 Acoustic signals of baby black caimans. *Zoology* 114: 313-320. <http://dx.doi.org/10.1016/j.zool.2011.07.003>
- Vergne AL, Pritz MB and Mathevon N** 2009 Acoustic communication in crocodylians: from behaviour to brain. *Biological Reviews* 84: 391-411. <http://dx.doi.org/10.1111/j.1469-185X.2009.00079.x>
- Waran N** 2001 The social behaviour of horses. In: Keeling L and Gonyou HW (eds) *Social Behaviour in Farm Animals* pp 247-274. <http://dx.doi.org/10.1079/9780851993973.0247>
- Watts JM, Stookey JM, Schmutz SM and Waltz CS** 2001 Variability in vocal and behavioural responses to visual isolation between full-sibling families of beef calves. *Applied Animal Behaviour Science* 70: 255-273. [http://dx.doi.org/10.1016/S0168-1591\(00\)00163-5](http://dx.doi.org/10.1016/S0168-1591(00)00163-5)
- Weary DM, Appleby MC and Fraser D** 1999 Responses of piglets to early separation from the sow. *Applied Animal Behaviour Science* 63: 289-300. [http://dx.doi.org/10.1016/S0168-1591\(99\)00021-0](http://dx.doi.org/10.1016/S0168-1591(99)00021-0)
- Weary DM and Chua B** 2000 Effects of early separation on the dairy cow and calf 1. Separation at 6 h, 1 day and 4 days after birth. *Applied Animal Behaviour Science* 69: 177-188. [http://dx.doi.org/10.1016/S0168-1591\(00\)00128-3](http://dx.doi.org/10.1016/S0168-1591(00)00128-3)
- Weary DM and Fraser D** 1997 Vocal response of piglets to weaning: effect of piglet age. *Applied Animal Behaviour Science* 54: 153-160. [http://dx.doi.org/10.1016/S0168-1591\(97\)00066-X](http://dx.doi.org/10.1016/S0168-1591(97)00066-X)
- Weary DM, Jasper J and Hotzel MJ** 2008 Understanding weaning distress. *Applied Animal Behaviour Science* 110: 24-41. <http://dx.doi.org/10.1016/j.applanim.2007.03.025>

- Weary DM, Ross S and Fraser D** 1997 Vocalizations by isolated piglets: a reliable indicator of piglet need directed towards the sow. *Applied Animal Behaviour Science* 53: 249-257. [http://dx.doi.org/10.1016/S0168-1591\(96\)01173-2](http://dx.doi.org/10.1016/S0168-1591(96)01173-2)
- Wells RS, Irvine AB and Scott MD** 1987 The social structure of free-ranging bottlenose dolphins. In: Genoways H (ed) *Current Mammalogy* pp 247-305. Plenum: New York, USA
- Wiener SG, Bayart F, Faull KF and Levine S** 1990 Behavioral and physiological responses to maternal separation in squirrel-monkeys (*Saimiri sciureus*). *Behavioral Neuroscience* 104: 108-115. <http://dx.doi.org/10.1037/0735-7044.104.1.108>
- Wilkinson GS** 2003 Social and vocal complexity in bats. In: De Waal FBM and Tyack PL (eds) *Animal Social Complexity: Intelligence, Culture and Individualized Societies*. Harvard University Press: Cambridge, MA, USA
- Wrobel TA and Dye AL** 2003 Grieving pet death: normative, gender, and attachment issues. *Omega Journal of Death and Dying* 47: 385-393. <http://dx.doi.org/10.2190/QYV5-LLJ1-T043-U0F9>
- Wynne CDL** 2007 What are animals? Why anthropomorphism is still not a scientific approach to behavior. *Comparative Cognitive Behavior Reviews* 2: 125-135