



# COMPOSITIO MATHEMATICA

## The $K3$ category of a cubic fourfold

Daniel Huybrechts

Compositio Math. **153** (2017), 586–620.

[doi:10.1112/S0010437X16008137](https://doi.org/10.1112/S0010437X16008137)



FOUNDATION  
COMPOSITIO  
MATHEMATICA



LONDON  
MATHEMATICAL  
SOCIETY  
EST. 1865



# The K3 category of a cubic fourfold

Daniel Huybrechts

## ABSTRACT

Smooth cubic hypersurfaces  $X \subset \mathbb{P}^5$  (over  $\mathbb{C}$ ) are linked to K3 surfaces via their Hodge structures, due to the work of Hassett, and via a subcategory  $\mathcal{A}_X \subset D^b(X)$ , due to the work of Kuznetsov. The relation between these two viewpoints has recently been elucidated by Addington and Thomas. In this paper, both aspects are studied further and extended to twisted K3 surfaces, which in particular allows us to determine the group of autoequivalences of  $\mathcal{A}_X$  for the general cubic fourfold. Furthermore, we prove finiteness results for cubics with equivalent K3 categories and study periods of cubics in terms of generalized K3 surfaces.

## 1. Introduction

As shown by Kuznetsov [Kuz10, Kuz06], the bounded derived category  $D^b(X)$  of coherent sheaves on a smooth cubic hypersurface  $X \subset \mathbb{P}^5$  contains, as the semi-orthogonal complement of three line bundles, a full triangulated subcategory

$$\mathcal{A}_X := \langle \mathcal{O}, \mathcal{O}(1), \mathcal{O}(2) \rangle^\perp \subset D^b(X)$$

that behaves in many respects like the bounded derived category  $D^b(S)$  of coherent sheaves on a K3 surface  $S$ . In fact, for certain special cubics  $\mathcal{A}_X$  is equivalent to  $D^b(S)$  or, more generally, to the derived category  $D^b(S, \alpha)$  of  $\alpha$ -twisted sheaves on a K3 surface  $S$ . Kuznetsov also conjectured that  $\mathcal{A}_X$  is of the form  $D^b(S)$  if and only if  $X$  is rational. Neither of the two implications has been verified until now, although Addington and Thomas recently have shown in [AT14] that the conjecture is (generically) equivalent to a conjecture attributed to Hassett [Has00] describing rational cubics in terms of their periods.

**1.1** This paper is not concerned with the rationality of cubic fourfolds, but with basic results on  $\mathcal{A}_X$ . Ideally, one would like to have a theory for  $\mathcal{A}_X$  that parallels the theory for  $D^b(S)$  and  $D^b(S, \alpha)$ . In particular, one would like to have analogues of the following results and conjectures.

- For a given twisted K3 surface  $(S, \alpha)$  there exist only finitely many isomorphism classes of twisted K3 surfaces  $(S', \alpha')$  with  $D^b(S, \alpha) \simeq D^b(S', \alpha')$ .
- Two twisted K3 surfaces  $(S, \alpha), (S', \alpha')$  are derived equivalent, i.e. there exists a  $\mathbb{C}$ -linear exact equivalence  $D^b(S, \alpha) \simeq D^b(S', \alpha')$ , if and only if there exists an orientation-preserving Hodge isometry  $\tilde{H}(S, \alpha, \mathbb{Z}) \simeq \tilde{H}(S', \alpha', \mathbb{Z})$ .
- The group of linear exact autoequivalences of  $D^b(S, \alpha)$  admits a natural representation  $\rho: \text{Aut}(D^b(S, \alpha)) \rightarrow \text{Aut}(\tilde{H}(S, \alpha, \mathbb{Z}))$ , which is surjective up to index two. Moreover, up to finite

---

Received 13 September 2015, accepted in final form 26 August 2016, published online 2 March 2017.

2010 Mathematics Subject Classification 14J28, 14F05 (primary).

Keywords: cubic fourfold, K3 surfaces, Hodge theory, derived categories.

This work was supported by the SFB/TR 45 ‘Periods, Moduli Spaces and Arithmetic of Algebraic Varieties’ of the DFG (German Research Foundation).

This journal is © Foundation Compositio Mathematica 2017.

index  $\text{Aut}(\mathbb{D}^b(S, \alpha))$  is conjecturally described as a fundamental group of a certain Deligne–Mumford stack.

Most of the theory for untwisted K3 surfaces is due to Mukai [Muk87] and Orlov [Orl97], whereas the basic theory of twisted K3 surfaces was developed in [HS05, HS06]. See also [Huy06, Huy09] for surveys and further references. Originally, the generalization to twisted K3 surfaces was motivated by the existence of non-fine moduli spaces [Că100]. However, more recently it has become clear that allowing twists has quite unexpected applications, e.g. to the Tate conjecture [Cha16, LMS14]. Crucial for the purpose of this article is the observation proved together with Macrì and Stellari [HMS08] that  $\text{Ker}(\rho) = \mathbb{Z}[2]$  for many twisted K3 surfaces  $(S, \alpha)$ . Note that for untwisted projective K3 surfaces the kernel is always highly non-trivial and, in particular, not finitely generated. The conjectural description of  $\text{Aut}(\mathbb{D}^b(S))$  has in this case only been achieved for K3 surfaces of Picard rank one [BB13].

**1.2** As a direct attack on  $\mathcal{A}_X$  is difficult, we follow Addington and Thomas [AT14] and reduce the study of  $\mathcal{A}_X$  via deformation to the case of (twisted) K3 surfaces. Central to our discussion is the Hodge structure  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  associated with  $\mathcal{A}_X$  introduced in [AT14] as the analogue of the Mukai–Hodge structure  $\tilde{H}(S, \mathbb{Z})$  of weight two on the full cohomology  $H^*(S, \mathbb{Z})$  of a K3 surface  $S$  or of the twisted version  $\tilde{H}(S, \alpha, \mathbb{Z})$  introduced in [HS05]. For example, any FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  induces a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ , cf. Proposition 3.4. This suffices to prove the following theorem.

**THEOREM 1.1.** *For any given smooth cubic  $X \subset \mathbb{P}^5$  there are only finitely many cubics  $X' \subset \mathbb{P}^5$  up to isomorphism for which there exists a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$ . See Corollary 3.5.*

Recall that due to a result of Bondal and Orlov a smooth cubic  $X \subset \mathbb{P}^5$  itself does not admit any non-isomorphic Fourier–Mukai partners. This is no longer true if  $\mathbb{D}^b(X)$  is replaced by its K3 category  $\mathcal{A}_X$ . In particular, there exist FM-equivalences  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  that do not extend to equivalences  $\mathbb{D}^b(X) \simeq \mathbb{D}^b(X')$ . However, we will also see that general cubics  $X$  and  $X'$ , i.e. those for which  $\text{rk } H^{2,2}(X, \mathbb{Z}) = 1$ , admit a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  if and only if  $X \simeq X'$ , see Theorem 1.5 or Corollary 3.6.

The following can be seen as an easy analogue of the result of Bayer and Bridgeland [BB13] describing  $\text{Aut}(\mathbb{D}^b(S))$  for general K3 surfaces  $S$  (namely those with  $\rho(S) = 1$ ) or rather of [HMS09] describing this group for general non-projective K3 surfaces or twisted projective K3 surfaces  $(S, \alpha)$  without  $(-2)$  classes (see § 6.1).

**THEOREM 1.2.** (i) *For the very general<sup>1</sup> smooth cubic  $X \subset \mathbb{P}^5$  the group  $\text{Aut}_s(\mathcal{A}_X)$  of symplectic FM-autoequivalences is infinite cyclic with*

$$\text{Aut}_s(\mathcal{A}_X)/\mathbb{Z} \cdot [2] \simeq \mathbb{Z}/3\mathbb{Z}.$$

*Alternatively, the group of all FM-autoequivalences  $\text{Aut}(\mathcal{A}_X)$  is infinite cyclic containing  $\mathbb{Z} \cdot [1]$  as a subgroup of index three.*

(ii) *Moreover, the induced action on  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  of any FM-autoequivalence  $\Phi : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$  of a non-special cubic preserves the natural orientation.*

---

<sup>1</sup> A property holds for the *very general* cubic if it holds for cubics in the complement of countably many proper closed subsets of the space of cubics under consideration. It holds for the *generic* cubic if it holds for a Zariski open, dense subset.

In fact, for every smooth cubic  $\text{Aut}_s(\mathcal{A}_X)$  contains an infinite cyclic group with  $\mathbb{Z} \cdot [2]$  as a subgroup of index three, see Corollary 3.13. The theory of twisted K3 surfaces is crucial for the theorem, as eventually the problem is reduced to [HMS08] which deals with general twisted K3 surfaces.

The group  $\text{Aut}_s(\mathcal{A}_X)$  of an arbitrary cubic is described by an analogue of Bridgeland’s conjecture, see Conjecture 3.15.

**1.3** In [Has00] Hassett showed that in the moduli space  $\mathcal{C}$  of smooth cubics, the set of those cubics  $X$  for which there exists a primitive positive plane  $K_d \subset H^{2,2}(X, \mathbb{Z})$  of discriminant  $d$  containing the class  $c_1(\mathcal{O}(1))^2$  is an irreducible divisor  $\mathcal{C}_d \subset \mathcal{C}$ . Moreover,  $\mathcal{C}_d$  is not empty if and only if

$$(*) \quad d \equiv 0, 2 \pmod{6} \text{ and } d > 6.$$

Cubics parametrized by the divisors  $\mathcal{C}_d$  are called special. Hassett also introduced the numerical condition

$$(**) \quad d \text{ is even and } d/2 \text{ is not divisible by nine or any prime } p \equiv 2 \pmod{3}.$$

Hassett then proved that  $(**)$  is equivalent to the orthogonal complement of the corresponding lattice  $K_d$  in  $H^4(X, \mathbb{Z})$  being (up to sign) Hodge isometric to the primitive Hodge structure  $H^2(S, \mathbb{Z})_{\text{prim}}$  of a polarized K3 surface. In [AT14] the condition was shown to be equivalent to the existence of a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(S, \mathbb{Z})$  for some K3 surface  $S$ . We prove the following twisted version of it (cf. Proposition 2.10).

**THEOREM 1.3.** *For a smooth cubic  $X \subset \mathbb{P}^5$  the Hodge structure  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  is Hodge isometric to the Hodge structure  $\tilde{H}(S, \alpha, \mathbb{Z})$  of a twisted K3 surface  $(S, \alpha)$  if and only if  $X \in \mathcal{C}_d$  with*

$$(**') \quad d \text{ is even and in the prime factorization } d/2 = \prod p_i^{n_i} \text{ one has } n_i \equiv 0 \pmod{2} \text{ for all primes } p_i \equiv 2 \pmod{3}.$$

Obviously, if  $d$  satisfies  $(**)$ , then  $k^2d$  satisfies  $(**')$  for all integers  $k$ . Conversely, any  $d$  satisfying  $(**')$  can be written (in general, non-uniquely) as  $k^2d_0$  with  $d_0$  satisfying  $(**)$ .

Also note that for  $X \in \mathcal{C}_d$  with  $d$  satisfying  $(**')$  the transcendental part  $T(X) \subset H^{2,2}(X, \mathbb{Z})$  is Hodge isometric (up to sign) to  $T(S, \alpha)$  of a twisted K3 surface  $(S, \alpha)$  (cf. [HS05]):

$$T(X)(-1) \simeq T(S, \alpha) \simeq \text{Ker}(\alpha : T(S) \rightarrow \mathbb{Q}/\mathbb{Z}). \tag{1.1}$$

As the main result of [AT14], Addington and Thomas proved that at least generically  $(**)$  is equivalent to  $\mathcal{A}_X \simeq \text{D}^b(S)$  for some K3 surface  $S$ . The following twisted version of it will be proved in § 6.2.

**THEOREM 1.4.** (i) *If  $\mathcal{A}_X \simeq \text{D}^b(S, \alpha)$  for some twisted K3 surface  $(S, \alpha)$ , then  $X \in \mathcal{C}_d$  with  $d$  satisfying  $(**')$ .*

(ii) *Conversely, if  $d$  satisfies  $(**')$ , then there exists a Zariski open dense set  $\emptyset \neq U \subset \mathcal{C}_d$  such that for all  $X \in \mathcal{C}_d$  there exists a twisted K3 surface  $(S, \alpha)$  and an equivalence  $\mathcal{A}_X \simeq \text{D}^b(S, \alpha)$ .*

Non-special cubics are determined by their associated K3 category  $\mathcal{A}_X$  and for general special cubics  $\mathcal{A}_X$  is determined by its Hodge structure (see Corollary 3.6 and § 6.3).

---

<sup>2</sup> This condition was originally stated as:  $d \equiv 0, 2 \pmod{6}$  and  $d$  not divisible by four, nine or any prime  $2 \neq p \equiv 2 \pmod{3}$ . The reformulation has been suggested by the referee.

THEOREM 1.5. *Let  $X$  and  $X'$  be two smooth cubics.*

(i) *Assume  $X$  is not special, i.e. not contained in any  $\mathcal{C}_d \subset \mathcal{C}$ . Then there exists a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  if and only if  $X \simeq X'$ .*

(ii) *For  $d$  satisfying (\*\*') and a Zariski dense open set of cubics  $X \in \mathcal{C}_d$ , there exists a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  if and only if there exists a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ .*

(iii) *For arbitrary  $d$  and very general  $X \in \mathcal{C}_d$  there exists a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  if and only if there exists a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ .*

We will also see that arguments of Addington [Add16] can be adapted to show that (\*\*') is in fact equivalent to the Fano variety of lines on  $X$  being birational to a moduli space of twisted sheaves on some K3 surface, see Proposition 4.1.

**1.4** There are a few fundamental issues concerning  $\mathcal{A}_X$  that we do not know how to address and that prevent us from developing the theory in full. First, this paper only deals with FM-equivalences  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$ , i.e. those for which the composition  $D^b(X) \rightarrow \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'} \hookrightarrow D^b(X')$  is a Fourier–Mukai transform. One would expect this to be the case for all equivalences, but the classical result of Orlov [Orl97] and its generalization by Canonaco and Stellari [CS07] do not apply to this situation. Secondly, it is not known whether  $\mathcal{A}_X$  always admits bounded t-structures or stability conditions. This is problematic when one wants to study FM-partners of  $\mathcal{A}_X$  as moduli spaces of (stable) objects in  $\mathcal{A}_X$ . As in [AT14], the lack of stability is also the crucial stumbling block to use deformation theory to prove statements as in Theorem 1.4 for all cubics and not only for generic or very general ones.

**1.5** The plan of the paper is as follows. Section 2 deals with all issues related to the lattice theory and the abstract Hodge theory. In particular, natural (countable unions of) codimension-one subsets  $D_{K3} \subset D_{K3'}$  of the period domain  $D \subset \mathbb{P}(A_2^\perp \otimes \mathbb{C})$  are studied at great length. They parametrize periods that induce Hodge structures that are Hodge isometric to  $\tilde{H}(S, \mathbb{Z})$  and  $\tilde{H}(S, \alpha, \mathbb{Z})$ , respectively, and which are described in terms of the numerical conditions (\*\*) and (\*\*'). In particular, Theorem 1.3 is proved. We also provide a geometric description of all periods  $x \in D$  in terms of generalized K3 surfaces, see Proposition 2.17.

In §3 we extend results in [AT14] from equivalences  $\mathcal{A}_X \simeq D^b(S)$  to the twisted case and prove the finiteness of FM-partners for  $\mathcal{A}_X$ , see Theorem 1.1. Moreover, we produce an action of the universal cover of  $SO(A_2)$  on  $\mathcal{A}_X$  for all cubics (Remark 3.14) and formulate an analogue of Bridgeland’s conjecture (cf. Conjecture 3.15).

The short §4 shows that (\*\*') is equivalent to  $F(X)$  being birational to a moduli space of stable twisted sheaves on a K3 surface. In §5 we adapt the deformation theory of [AT14] to the twisted case. Finally, in §6 we conclude the proofs of Theorems 1.2, 1.4, and 1.5.

## 2. Lattice theory and period domains

We start by discussing the relevant lattice theory. To make the reading self-contained, we will also recall results due to Hassett and to Addington and Thomas on the way.

There are two kinds of lattices, those related to K3 surfaces,  $\Lambda, \tilde{\Lambda}$ , etc., and those attached to cubic fourfolds,  $I_{21,2}, K_d$ , etc. The two types are linked by a lattice  $A_2^\perp$  of signature  $(2, 20)$  and two embeddings

$$I_{2,21} \longleftarrow A_2^\perp \longrightarrow \tilde{\Lambda}.$$

The induced maps between the associated period domains allows one to relate periods of cubic fourfolds to periods of (generalized) K3 surfaces.

**2.1** By  $U$  we shall denote the hyperbolic plane, i.e.  $\mathbb{Z}^2$  with the intersection form  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . The K3 lattice  $\Lambda$  and the extended K3 lattice  $\tilde{\Lambda}$  are by definition the unique even, unimodular lattice of signature  $(3, 19)$  and  $(4, 20)$ , respectively. So,

$$\Lambda \simeq E_8(-1)^{\oplus 2} \oplus U^{\oplus 3} \quad \text{and} \quad \tilde{\Lambda} \simeq \Lambda \oplus U.$$

Next,  $A_2$  denotes the standard root lattice of rank two, i.e. there exists a basis  $\lambda_1, \lambda_2$  with respect to which the intersection matrix is given by  $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ . The lattice  $A_2$  is even and of signature  $(2, 0)$ . Moreover, its discriminant group is  $A_{A_2} := A_2^*/A_2 \simeq \mathbb{Z}/3\mathbb{Z}$  and, in particular,  $A_2$  is not unimodular.

Due to [Nik79, Theorem 1.14.4], there exist embeddings

$$A_2 \hookrightarrow \Lambda \quad \text{and} \quad A_2 \hookrightarrow \tilde{\Lambda},$$

which are both unique up to the action of  $O(\Lambda)$  and  $O(\tilde{\Lambda})$ , respectively. Note that all such embeddings are automatically primitive. In the following we will fix once and for all one such embedding  $A_2 \hookrightarrow \Lambda \hookrightarrow \tilde{\Lambda}$  and consider the orthogonal complement of  $A_2 \subset \tilde{\Lambda}$  as a fixed primitive sublattice

$$A_2^\perp \subset \tilde{\Lambda} \tag{2.1}$$

of signature  $(2, 20)$ . Its isomorphism type does not depend on the chosen embedding of  $A_2$ . It can be described explicitly as the orthogonal complement of the embedding  $A_2 \hookrightarrow \tilde{\Lambda}$  given by

$$A_2 \hookrightarrow U \oplus U \hookrightarrow \tilde{\Lambda}, \quad \lambda_1 \mapsto e' + f', \quad \lambda_2 \mapsto e + f - e', \tag{2.2}$$

where  $e, f$  and  $e', f'$  denote the standard bases of the two copies of the hyperbolic plane. Thus,<sup>3</sup>

$$A_2^\perp \simeq E_8(-1)^{\oplus 2} \oplus U^{\oplus 2} \oplus A_2(-1). \tag{2.3}$$

*Remark 2.1.* For later use we recall that the group of isometries  $O(A_2)$  of the lattice  $A_2$  is isomorphic to  $\mathfrak{S}_3 \times \mathbb{Z}/2\mathbb{Z}$ . Here, the Weyl group  $\mathfrak{S}_3$  permutes the unit vectors  $e_i \in \mathbb{R}^3$ , where  $A_2 \hookrightarrow \mathbb{R}^3$  via  $\lambda_1 = e_1 - e_2$  and  $\lambda_2 = e_2 - e_3$ , and the generator of  $\mathbb{Z}/2\mathbb{Z}$  acts by  $-\text{id}$ . In fact,  $\mathfrak{S}_3 \subset O(A_2)$  is the kernel of the natural map  $O(A_2) \rightarrow O(A_{A_2}) \simeq \mathbb{Z}/2\mathbb{Z}$  (use the aforementioned  $A_{A_2} \simeq \mathbb{Z}/3\mathbb{Z}$ ). The sign  $\mathfrak{S}_3 \rightarrow \mathbb{Z}/2\mathbb{Z}$  can be identified with the determinant  $O(A_2) \rightarrow \{\pm 1\}$ . Thus, the group of orientation-preserving isometries of  $A_2$  acting trivially on  $A_{A_2}$  is just  $\mathfrak{A}_3 \simeq \mathbb{Z}/3\mathbb{Z}$ , where the generator can be chosen to act by  $\lambda_1 \mapsto -\lambda_1 - \lambda_2, \lambda_2 \mapsto \lambda_1$ .

**2.2** Next, consider the unique odd, unimodular lattice

$$I_{2,21} := \mathbb{Z}^{\oplus 2} \oplus \mathbb{Z}(-1)^{\oplus 21} \simeq E_8(-1)^{\oplus 2} \oplus U^{\oplus 2} \oplus \mathbb{Z}(-1)^{\oplus 3}$$

of signature  $(2, 21)$  and an element  $h \in I_{2,21}$  with  $(h)^2 = -3$ , e.g.  $h = (1, 1, 1) \in \mathbb{Z}(-1)^{\oplus 3}$ . Then the primitive sublattice  $h^\perp \subset I_{2,21}$  is of signature  $(2, 20)$  and using (2.3) one finds

$$h^\perp \simeq A_2^\perp.$$

<sup>3</sup> In [Has00] the last summand is instead described as a lattice with intersection matrix  $\begin{pmatrix} -2 & -1 \\ -1 & -2 \end{pmatrix}$ , which is of course isomorphic to  $A_2(-1)$ .

In the following, we will always consider  $A_2^\perp$  with two fixed embeddings as above:

$$I_{2,21} \longleftarrow A_2^\perp \longrightarrow \tilde{\Lambda}.$$

Following Hassett [Has00], we now consider all primitive, negative-definite sublattices

$$h \in K_d \subset I_{2,21}$$

of rank two containing  $h$ . Here, the index  $d = \text{disc}(K_d)$  denotes the discriminant of  $K_d$ , which is necessarily positive. Using [Nik79, § 1.5] one finds that up to the action of the subgroup of  $O(I_{2,21})$  fixing  $h$  the lattice  $K_d \subset I_{2,21}$  is uniquely determined by  $d$ , see [Has00, Proposition 3.2.4] for the details.

Furthermore,  $d \equiv 0, 2 \pmod{6}$  and the generator  $v$  of  $K_d \cap A_2^\perp$  (unique up to sign) satisfies

$$-(v)^2 = \begin{cases} d/3 & \text{if } d \equiv 0 \pmod{6}, \\ 3d & \text{if } d \equiv 2 \pmod{6}. \end{cases} \tag{2.4}$$

More precisely, Hassett shows that up to isometry of  $A_2^\perp$  the vector  $v$  is given as

$$v = e_1 - (d/6)f_1 \quad \text{and} \quad v = 3(e_1 - ((d-2)/6)f_1) + \mu_1 - \mu_2, \tag{2.5}$$

respectively. Here,  $e_1, f_1$  is the standard basis of one of the copies of  $U$  in (2.3) and  $\mu_1, \mu_2$  denotes the standard basis of  $A_2(-1)$ . Viewing  $v \in A_2^\perp \subset \tilde{\Lambda}$  as an element of  $\tilde{\Lambda}$  leads to a lattice

$$A_2 \oplus \mathbb{Z} \cdot v \subset \tilde{\Lambda}$$

of rank three and signature  $(2, 1)$ . As it turns out, this is a primitive sublattice for  $d \equiv 0 \pmod{6}$  and it is of index three in its saturation for  $d \equiv 2 \pmod{6}$ . This follows from [Has00, Proposition 3.2.2] asserting that in the two cases  $(v \cdot A_2^\perp) = \mathbb{Z}$  and  $3\mathbb{Z}$ , respectively. Altogether this yields the following lemma.

LEMMA 2.2. *The saturation  $\Gamma_d \subset \tilde{\Lambda}$  of  $A_2 \oplus \mathbb{Z} \cdot v$  satisfies*

$$\text{disc}(\Gamma_d) = d.$$

*Proof.* This can either be proved by a direct computation or by observing that  $\text{disc}(\Gamma_d)$  equals the discriminant of  $\Gamma_d^\perp \subset \tilde{\Lambda}$ , which is isomorphic to  $v^\perp \subset A_2^\perp$ . Similarly,  $d = \text{disc}(K_d)$  equals the discriminant of the lattice  $\langle v, h \rangle^\perp$ , which again is just  $v^\perp \subset A_2^\perp$ .  $\square$

In our discussion, the lattices  $K_d$  will take a back seat, as it will be more natural to work with the generator  $v \in A_2^\perp \cap K_d$  directly.

**2.3** We shall be interested in the period domains

$$D \subset \mathbb{P}(A_2^\perp \otimes \mathbb{C}) \quad \text{and} \quad Q \subset \mathbb{P}(\tilde{\Lambda} \otimes \mathbb{C}),$$

defined by the two conditions  $(x \cdot x) = 0$  and  $(x \cdot \bar{x}) > 0$ . Observe that

$$\dim D = 20 \quad \text{and} \quad \dim Q = 22$$



and that  $Q$  is connected while  $D$  has two connected components. Using the embedding (2.1), we can write  $D = \mathbb{P}(A_2^\perp \otimes \mathbb{C}) \cap Q$  as part of the commutative diagram

$$\begin{array}{ccc} D & \xrightarrow{\quad} & Q \\ \downarrow & & \downarrow \\ \mathbb{P}(A_2^\perp \otimes \mathbb{C}) & \hookrightarrow & \mathbb{P}(\tilde{\Lambda} \otimes \mathbb{C}). \end{array}$$

Thus, points  $x \in D$  correspond to Hodge structures of weight two on the lattice  $A_2^\perp$ , but also to Hodge structures on  $\tilde{\Lambda}$  with  $A_2$  contained in the  $(1, 1)$  part. In fact, for very general points  $x \in D$  the integral  $(1, 1)$  part of the corresponding Hodge structure is the lattice  $A_2$ .

We shall refer to  $D$  as the period domain of cubic fourfolds, although only an open subset really corresponds to smooth cubics. More concretely, for a smooth cubic  $X \subset \mathbb{P}^5$  and any marking, i.e. an isometry,  $\varphi : h^\perp \xrightarrow{\sim} A_2^\perp$  (up to sign), one defines the associated period as the image  $x := [\varphi_{\mathbb{C}}(H^{3,1}(X))] \in D$ . A description of the image of the period map, allowing cubics with ADE singularities, has been given by Laza [Laz10] and Looijenga in [Loo09]. Points in  $Q$  are thought of as periods of generalized K3 surfaces, cf. § 2.8.

For later use we state the following technical observation.

**LEMMA 2.3.** *The Hodge structure on  $\tilde{\Lambda}$  defined by an arbitrary  $x \in D$  admits a Hodge isometry that reverses any given orientation of the four positive directions.*

*Proof.* Consider a transposition  $g := (12) \in \mathfrak{S}_3 \subset O(A_2)$ . Then  $g$  acts trivially on the discriminant  $A_{A_2}$  (see Remark 2.1) and can, therefore, be extended to  $\tilde{g} \in O(\tilde{\Lambda})$  acting trivially on  $A_2^\perp$ . Thus, the Hodge structure defined by  $x$  admits a Hodge isometry  $\tilde{g}$ , which preserves the orientation of the two positive directions given by the  $(2, 0)$  and  $(0, 2)$  parts. On the other hand, by construction, it reverses the orientation of the two positive directions in  $A_2$ .  $\square$

*Remark 2.4.* This result is the analogue of the observation that any Hodge structure on  $\tilde{\Lambda}$  containing a hyperbolic plane  $U$  in its  $(1, 1)$  part admits an orientation-reversing Hodge isometry. This assertion applies to the Hodge structure  $\tilde{H}(S, \mathbb{Z})$  of a K3 surface  $S$ , but it is not clear whether also  $\tilde{H}(S, \alpha, \mathbb{Z})$  of a twisted K3 surface  $(S, \alpha)$  (see below) admits an orientation-reversing Hodge isometry. The latter would be important for adapting the arguments in [HMS09] to the description of the image of  $\text{Aut}(D^b(S, \alpha)) \rightarrow \text{Aut}(\tilde{H}(S, \alpha, \mathbb{Z}))$ , see [Rei14] for partial results.

**2.4** Let us now turn to the geometric interpretation of certain periods in  $Q$ . Recall that for a K3 surface  $S$  the extended K3 (or Mukai) lattice  $\tilde{H}(S, \mathbb{Z})$  is abstractly isomorphic to  $\tilde{\Lambda}$ . Moreover,  $\tilde{H}(S, \mathbb{Z})$  comes with a natural Hodge structure of weight two defined by

$$\tilde{H}^{2,0}(S) := H^{2,0}(S) \quad \text{and} \quad \tilde{H}^{1,1}(S) := H^{1,1}(S) \oplus (H^0 \oplus H^4)(S, \mathbb{C}).$$

For a Brauer class  $\alpha \in \text{Br}(S) \simeq H^2(S, \mathbb{G}_m) \simeq H^2(S, \mathcal{O}_S^*)_{\text{tors}}$  we have introduced in [Huy05] the weight-two Hodge structure  $\tilde{H}(S, \alpha, \mathbb{Z})$ . As a lattice this is still isomorphic to  $\tilde{\Lambda}$  and its Hodge structure is determined by

$$\tilde{H}^{2,0}(S, \alpha) := \mathbb{C} \cdot (\sigma + B \wedge \sigma) \quad \text{and} \quad \tilde{H}^{1,1}(S, \alpha) := \exp(B) \cdot \tilde{H}^{1,1}(S).$$

Here,  $0 \neq \sigma \in H^{2,0}(S)$  and  $B \in H^2(S, \mathbb{Q})$  maps to  $\alpha$  under the exponential map

$$H^2(S, \mathbb{Q}) \longrightarrow H^2(S, \mathcal{O}_S) \xrightarrow{\text{exp}} H^2(S, \mathcal{O}_S^*).$$

The isomorphism type of the Hodge structure is independent of the choice of  $B$ .



DEFINITION 2.5. A period  $x \in Q$  is of *K3 type* (respectively *twisted K3 type*) if there exists a K3 surface  $S$  (respectively a twisted K3 surface  $(S, \alpha \in \text{Br}(S))$ ) such that the Hodge structure on  $\tilde{\Lambda}$  defined by  $x$  is Hodge isometric to  $\tilde{H}(S, \mathbb{Z})$  (respectively  $H(S, \alpha, \mathbb{Z})$ ).

The sets of periods of K3 type and twisted K3 type will be denoted

$$Q_{\text{K3}} \subset Q_{\text{K3}'} \subset Q.$$

There is also a geometric interpretation for points outside  $Q_{\text{K3}'}$  in terms of symplectic structures [Huy05], but those are *a priori* inaccessible by algebro-geometric techniques (see, however, § 2.8).

For the following recall that the twisted hyperbolic plane  $U(n)$  is the rank-two lattice with intersection matrix  $\begin{pmatrix} 0 & n \\ n & 0 \end{pmatrix}$ . The standard isotropic generators will be denoted  $e_n, f_n$  or simply  $e, f$ . Part (i) of the next lemma is well known.

LEMMA 2.6. Consider a period point  $x \in Q$ . Then:

- (i)  $x \in Q_{\text{K3}}$  if and only if there exists an embedding  $U \hookrightarrow \tilde{\Lambda}$  into the  $(1, 1)$  part of the Hodge structure defined by  $x$ ;
- (ii)  $x \in Q_{\text{K3}'}$  if and only if there exists a (not necessarily primitive) embedding  $U(n) \hookrightarrow \tilde{\Lambda}$  for some  $n \neq 0$  into the  $(1, 1)$  part of the Hodge structure defined by  $x$ .

*Proof.* We prove the second assertion; the first one is even easier. Start with a twisted K3 surface  $(S, \alpha)$  and pick a lift  $B \in H^2(S, \mathbb{Q})$  of  $\alpha$ . Then the algebraic part  $\tilde{H}^{1,1}(S, \alpha, \mathbb{Z}) = \exp(B) \cdot \tilde{H}^{1,1}(S, \mathbb{Q}) \cap \tilde{H}(S, \mathbb{Z})$  contains the lattice  $(\mathbb{Z} \cdot (1, B, B^2/2) \cap \tilde{H}(S, \mathbb{Z})) \oplus H^4(S, \mathbb{Z})$ , which is isomorphic to  $U(n)$  for  $n$  minimal with  $n(1, B, B^2/2) \in \tilde{H}(S, \mathbb{Z})$ .

Conversely, assume  $U(n) \subset \tilde{\Lambda}$  is of type  $(1, 1)$  with respect to  $x$ . Choosing  $n$  minimal, we can assume that the standard isotropic generator  $e_n = e$  is primitive in  $\tilde{\Lambda}$ . But then  $e \in U(n)$  can be completed to a sublattice of  $\tilde{\Lambda}$  isomorphic to the hyperbolic plane  $U = \langle e, f \rangle$ , which therefore induces an orthogonal decomposition (usually different from that defining  $\tilde{\Lambda}$ )

$$\tilde{\Lambda} \simeq \Lambda \oplus U. \tag{2.6}$$

With respect to (2.6) the second basis vector  $f_n \in U(n)$  can be written as  $f_n = \gamma + nf + ke$  with  $\gamma \in \Lambda$ . Similarly, a generator of the  $(2, 0)$  part of the Hodge structure determined by  $x$  is orthogonal to  $e$  and hence of the form  $\sigma + \lambda e$  for some  $\sigma \in \Lambda \otimes \mathbb{C}$  and  $\lambda \in \mathbb{C}$ . However, it is also orthogonal to  $f_n$  and so  $(\gamma, \sigma) + n\lambda = 0$ . Now set  $B := -(1/n)\gamma$ . Then  $\sigma + \lambda e = \sigma + B \wedge \sigma$ , where  $B \wedge \sigma$  stands for  $(B, \sigma)e$ .

Eventually, the surjectivity of the period map implies that  $\sigma \in \Lambda \otimes \mathbb{C}$  can be realized as the period of some K3 surface  $S$ , i.e. there exists an isometry  $H^2(S, \mathbb{Z}) \simeq \Lambda$  identifying  $H^{2,0}(S)$  with  $\mathbb{C} \cdot \sigma \subset \Lambda \otimes \mathbb{C}$ . Here one uses  $(\sigma, \sigma) = (\sigma + \lambda e, \sigma + \lambda e) = 0$  and  $(\sigma, \bar{\sigma}) = (\sigma + \lambda e, \bar{\sigma} + \bar{\lambda} e) > 0$ . Mapping  $H^4(S, \mathbb{Z})$  to  $\mathbb{Z} \cdot e \subset U \subset \Lambda \oplus U$  in (2.6) and defining  $\alpha \in \text{Br}(S)$  as the Brauer class induced by  $B$  under  $\Lambda \otimes \mathbb{Q} \simeq H^2(S, \mathbb{Q}) \rightarrow H^2(S, \mathbb{G}_m)$  yields a Hodge isometry between  $\tilde{H}(S, \alpha, \mathbb{Z})$  and the Hodge structure defined by  $x$  on  $\tilde{\Lambda}$ .  $\square$

COROLLARY 2.7. The sets  $Q_{\text{K3}} \subset Q_{\text{K3}'} \subset Q$  can be described as the intersections of  $Q$  with countably many linear subspaces of codimension two:

$$Q_{\text{K3}} = Q \cap \bigcup U^\perp \subset Q_{\text{K3}'} = Q \cap \bigcup U(n)^\perp \subset Q.$$

Here, the first union is over all embeddings  $U \hookrightarrow \tilde{\Lambda}$  and the second over all  $U(n) \hookrightarrow \tilde{\Lambda}$  with arbitrary  $n \neq 0$ .

**2.5** However, it will turn out that the further intersection with  $D$  yields countable unions of codimension-one subsets. These intersections are denoted by

$$D_{K3} := D \cap Q_{K3} \subset D_{K3'} := D \cap Q_{K3'} \subset D$$

and will be viewed as the sets of cubic periods that define generalized K3 periods of K3 type and of twisted K3 type, respectively. So:

- $x \in D_{K3}$  if and only if there exists a K3 surface  $S$  such that the Hodge structure on  $\tilde{\Lambda}$  defined by  $x$  is Hodge isometric to  $\tilde{H}(S, \mathbb{Z})$ ;
- $x \in D_{K3'}$  if and only if there exists a twisted K3 surface  $(S, \alpha \in \text{Br}(S))$  such that the Hodge structure on  $\tilde{\Lambda}$  defined by  $x$  is Hodge isometric to  $\tilde{H}(S, \alpha, \mathbb{Z})$ .

We remark for later use that for very general  $x \in D_{K3}$  (or  $x \in D_{K3'}$ ) the algebraic part  $\tilde{H}^{1,1}(S, \mathbb{Z})$  (respectively  $\tilde{H}^{1,1}(S, \alpha, \mathbb{Z})$ ) is of rank three.

We will first explain that  $D_{K3'}$  is a countable union of hyperplane sections. A second proof for the same assertion that also works for  $D_{K3}$  is provided in §2.6.

**PROPOSITION 2.8.** *The set of twisted K3 periods in  $D$  can also be described as the countable union of hyperplane sections:*

$$D_{K3'} = D \cap \bigcup e^\perp.$$

Here, the union runs over all  $0 \neq e \in \tilde{\Lambda}$  with  $(e)^2 = 0$ .

*Proof.* One inclusion follows from the fact that any  $U(n)$  contains an isotropic vector. For the converse, assume  $x \in e^\perp$  for some primitive isotropic  $0 \neq e \in \tilde{\Lambda}$ . Then  $e \notin A_2^\perp$ , as otherwise the positive plane corresponding to  $x$  would be contained in the orthogonal complement of  $e$  in  $A_2^\perp$ , which has only one positive direction left. Hence, there exists  $a \in A_2$  with  $(a.e) \neq 0$ . Let  $f := (a.e)a - ((a)^2/2)e$ , which satisfies  $(f)^2 = 0$  and  $(f.e) = (a.e)^2 =: n$ . Hence,  $e, f$  span a twisted hyperbolic plane  $U(n)$  in the  $(1, 1)$  part of the Hodge structure defined by  $x$ .  $\square$

There is yet another class of hyperplane sections of  $D$  that is of importance to us. We let

$$D_{\text{sph}} := D \cap \bigcup \delta^\perp,$$

where the union is over all  $\delta \in \tilde{\Lambda}$  with  $(\delta)^2 = -2$  and call it the set of *periods with spherical classes*. Indeed,  $x \in D$  is contained in  $D_{\text{sph}}$  if and only if the Hodge structure on  $\tilde{\Lambda}$  defined by  $x$  admits an integral  $(1, 1)$  class  $\delta$  with  $(\delta)^2 = -2$  and those classes typically appear as Mukai vectors of spherical objects, see Example 3.11.

Note that there are natural inclusions

$$D_{K3} \subset D_{\text{sph}} \subset D,$$

for every hyperbolic plane  $U$  contains a  $(-2)$  class. However,  $D_{K3'}$  is not contained in  $D_{\text{sph}}$  and, more precisely, the inclusions

$$D_{K3} \subsetneq D_{K3'} \cap D_{\text{sph}} \subsetneq D_{K3'}, D_{\text{sph}}$$

are all proper, see Example 2.14 and Proposition 2.15.

**2.6** It is instructive to study the sets  $D_{K3} \subset D_{K3'}$  and  $D_{K3} \subset D_{\text{sph}}$  from a more cubic perspective, i.e. in terms of the lattices  $K_d$ .

For any  $h \in K_d \subset I_{2,21}$  as in § 2.2 one introduces the hyperplane section

$$D \cap K_d^\perp \subset \mathbb{P}(A_2^\perp \otimes \mathbb{C})$$

of all cubic periods orthogonal to  $K_d \cap A_2^\perp$ . In other words,  $D \cap K_d^\perp$  is the set of cubic periods for which the generator  $v$  of  $K_d \cap A_2^\perp$  is of type  $(1, 1)$ , i.e.  $D \cap K_d^\perp = D \cap v^\perp$ . Then one defines

$$D_d := D \cap \bigcup K_d^\perp,$$

where the union runs over all  $h \in K_d \subset I_{2,21}$  as above. So, for each positive  $d \equiv 0, 2 \pmod{6}$  the set  $D_d$  is a countable union of hyperplane sections of  $D$ . Dividing  $D_d$  by the subgroup  $\tilde{O}(h^\perp) = O(I_{2,21}, h) \subset O(I_{2,21})$  of elements fixing  $h$  yields Hassett's irreducible divisor

$$\mathcal{C}_d := D_d / \tilde{O}(h^\perp).$$

Consider the following conditions for an even integer  $d > 6$ :

- (\*)  $d \equiv 0, 2 \pmod{6}$ ;
- (\*\*)  $d$  is even and  $d/2$  is not divisible by 9 or any prime  $p \equiv 2 \pmod{3}$ ;
- (\*\*')  $d$  is even and in the prime factorization  $d/2 = \prod p_i^{n_i}$  one has  $n_i \equiv 0 \pmod{2}$  for all primes  $p_i \equiv 2 \pmod{3}$ .

Obviously, (\*\*) implies (\*\*'). More precisely, if  $d$  satisfies (\*\*), then (\*\*') holds for all  $k^2d$ .

*Remark 2.9.* Conditions (\*) and (\*\*) have first been introduced and studied by Hassett [Has00]. He shows that  $D_d$  is not empty if and only if (\*) is satisfied. Moreover,  $d$  satisfies (\*\*) if and only if for all cubics  $X$  with period  $x$  contained in  $D_d$  there exists a polarized K3 surface  $(S, H)$  such that its primitive cohomology  $H^2(S, \mathbb{Z})_{\text{pr}}$  is Hodge isometric to the Hodge structure on  $K_d^\perp$  defined by  $x$ . To get polarized K3 surfaces and not only quasi-polarized ones, one has to use a result of Voisin [Voi86, § 4, Proposition 1] saying that  $H^{2,2}(X, \mathbb{Z})_{\text{pr}}$  does not contain any class of square two.

On the level of lattices this boils down to the observation that for  $v \in A_2^\perp$  as in (2.5), say for  $d \equiv 0 \pmod{6}$ , its orthogonal complement in  $A_2^\perp$  is isometric to  $E_8(-1)^{\oplus 2} \oplus U \oplus A_2(-1) \oplus \mathbb{Z}(d/3)$ . And indeed for  $d$  satisfying (\*\*')  $A_2(-1) \oplus \mathbb{Z}(d/3) \simeq U \oplus \mathbb{Z}(-d)$  (see [Nik79, Corollary 1.10.2, 1.13.3] or [Huy16, Theorem 14.1.5]) and, therefore,  $v^\perp \simeq E_8(-1)^{\oplus 2} \oplus U^{\oplus 2} \oplus \mathbb{Z}(-d)$ , which is the transcendental lattice of a very general polarized K3 surface of degree  $d$ . A similar argument holds for  $d \equiv 2 \pmod{6}$ .

**PROPOSITION 2.10.** *With the above notation one has*

$$D_{K3} = \bigcup_{(**)} D_d \quad \text{and} \quad D_{K3'} = \bigcup_{(**')} D_d,$$

where  $d$  runs through all  $d$  satisfying (\*\*) respectively (\*\*').

*Proof.* The first equality is due to Addington and Thomas [AT14, Theorem 3.1]. Indeed, they show that  $x \in D_d$  with  $d$  satisfying  $(**)$  if and only if there exists a hyperbolic plane  $U \subset \tilde{\Lambda}$  which is of type  $(1, 1)$  with respect to  $x$ . The latter is in turn equivalent to  $x \in D_{K3}$ , see Lemma 2.6.<sup>4</sup>

Maybe surprisingly, the second assertion is easier to prove. We include the elementary argument. Due to Corollary 2.8 we know  $D_{K3'} = D \cap \bigcup e^\perp$  with  $0 \neq e \in \tilde{\Lambda}$  isotropic. So for one inclusion one has to show that each  $D \cap e^\perp$  is of the form  $D_d$  with  $d$  satisfying  $(**')$ . Decompose  $e = e_1 + e_2 \in (A_2 \oplus A_2^\perp) \otimes \mathbb{Q}$ . Let then  $v \in A_2^\perp$  such that  $\mathbb{Q} \cdot e_2 \cap A_2^\perp = \mathbb{Z} \cdot v$  and define  $K_d \subset I_{2,21}$  as the saturation of the sublattice spanned by  $v \in A_2^\perp \subset I_{2,21}$  and  $h$ . We have to show that the discriminant  $d$  of  $K_d$  satisfies  $(**')$ .

Assume first that  $A_2 \oplus \mathbb{Z} \cdot v \subset \tilde{\Lambda}$  is primitive. Then  $d \equiv 0 \pmod{6}$  and  $d = -3(v)^2$ , see § 2.2. As  $e \in A_2 \oplus \mathbb{Z} \cdot v$  in this case, the quadratic equation  $2(x_1^2 + x_2^2 - x_1x_2) + (v)^2x^2 = 0$  admits an integral solution. However, it is a classical result that

$$2n = (w)^2 \tag{2.7}$$

for some  $w \in A_2$  if and only if  $n = \prod p_i^{n_i}$  with  $n_i \equiv 0 \pmod{2}$  for all  $p_i \equiv 2 \pmod{3}$ , see [Kne02].<sup>5</sup> But clearly this holds for  $n = -(v)^2/2$  if and only if  $d = 6n$  satisfies  $(**')$ .

Next assume that  $A_2 \oplus \mathbb{Z} \cdot v \subset \tilde{\Lambda}$  has index three in its saturation. Hence,  $d \equiv 2 \pmod{6}$  and  $3d = -(v)^2$ . Then argue as before, but now with the isotropic vector  $3e \in A_2 \oplus \mathbb{Z} \cdot v$  and with  $n = -(v)^2/2 = 3d/2$ .

Running the argument backwards proves the reverse inclusion. □

So, in particular, although  $Q_{K3} \subset Q_{K3'} \subset Q$  are countable unions of codimension-two subsets, their restrictions  $D_{K3} \subset D_{K3'} \subset D$  to  $D$  are countable unions of codimension-one subsets. For  $D_{K3'}$  we have observed this already in § 2.5.

*Remark 2.11.* As mentioned in [AT14, Add16] and explained to me by Addington, condition  $(**)$  is in fact equivalent to the existence of a primitive  $w \in A_2$  with  $d = (w)^2$ . And, as has become clear in the above proof, condition  $(**')$  is equivalent to the existence of a (not necessarily primitive)  $w \in A_2$  with  $d = (w)^2$ .

The first values of  $d > 6$  that satisfy the various conditions are

(*)	8	12	14	18	20	24	26	30	32	36	38	42	44	48
(**)			14				26				38	42		
(**')	8		14	18		24	26		32		38	42		

<sup>4</sup>The ‘only if’ direction is a consequence of Hassett’s original result saying that  $x \in D_d$  with  $d$  satisfying  $(**)$  if and only if there exists a polarized K3 surface  $(S, H)$  such that  $H^2(S, \mathbb{Z})_{\text{prim}}$  is Hodge isometric to the Hodge structure on  $K_d^\perp$  given by  $x$ . As the orthogonal complement of  $H^2(S, \mathbb{Z})_{\text{prim}} \subset \tilde{H}(S, \mathbb{Z}) \simeq \tilde{\Lambda}$  contains a hyperbolic plane, by [Nik79, Theorem 1.14.4] this Hodge isometry extends to a Hodge isometry of  $\tilde{H}(S, \mathbb{Z})$  with the Hodge structure on  $\tilde{\Lambda}$  given by  $x$ . For the other direction one has to show that any Hodge isometry between  $\tilde{H}(S, \mathbb{Z})$  and the one on  $\tilde{\Lambda}$  given by  $x$  can be used to get a Hodge isometry between the Hodge structure on  $K_d^\perp \cap A_2^\perp \subset \tilde{\Lambda}$  and  $H^2(S, \mathbb{Z})_{\text{prim}}$  for some polarization on  $S$ .

<sup>5</sup>For example, a prime  $p$  can be written as  $x^2 + 3y^2$  if and only if  $p = 3$  or  $p \equiv 1 \pmod{3}$ , see [Cox89]. Since  $4(x^2 + xy + y^2) = (2x + y)^2 + 3y^2$  and  $(x_1^2 + 3y_1^2) \cdot (x_2^2 + 3y_2^2) = (x_1x_2 - 3y_1y_2)^2 + 3(x_1y_2 + x_2y_1)^2$ , this proves one direction. The other one uses a computation with Hilbert symbols to determine when  $-nx_1^2 + x_2^2 + 3x_3^2 = 0$  has a rational solution.

*Example 2.12.* For certain  $d$  the condition that the period  $x \in D$  of a cubic  $X$  is contained in  $D_d$  has a geometric interpretation, see [Has00, § 4]. For example,  $x \in D_8$  if and only if  $X$  contains a plane  $\mathbb{P}^2 \subset X$ , or if  $X$  is a Pfaffian cubic, then  $x \in D_{14}$ .

Let  $x \in D_d$  with  $d$  satisfying (\*\*'). Then there exists a twisted K3 surface  $(S, \alpha)$  such that the Hodge structure defined by  $x$  is Hodge isometric to  $\tilde{H}(S, \alpha, \mathbb{Z})$ . If  $x \in D_d$  is a very general point of  $D_d$ , then  $\text{rk}(\tilde{H}^{1,1}(S, \alpha, \mathbb{Z})) = 3$  and  $A_2 \oplus \mathbb{Z} \cdot v \subset \tilde{H}^{1,1}(S, \alpha, \mathbb{Z})$  is of index one or three, respectively.

LEMMA 2.13. *For the order of the Brauer class  $\alpha$  one has*

$$\text{ord}(\alpha)^2 \mid d.$$

*Proof.* Let  $\ell := \text{ord}(\alpha)$ . As proved in [Huy05], the transcendental lattice  $T(S, \alpha)$  is isometric to the kernel of the natural map  $T(S) \twoheadrightarrow (1/\ell)\mathbb{Z}/\mathbb{Z}$  defined by  $\alpha$ . Hence,

$$|\text{disc}(T(S, \alpha))| = |\text{disc}(T(S))| \cdot \text{ord}(\alpha)^2.$$

On the other hand,  $\text{disc}(T(S, \alpha)) = \text{disc}(\tilde{H}^{1,1}(S, \alpha, \mathbb{Z})) = d$  by Lemma 2.2. □

Clearly,  $d/\text{ord}(\alpha)^2$  still satisfies (\*\*') (but not necessarily (\*\*)). As mentioned earlier, any  $d$  satisfying (\*\*') can be written (not always uniquely) as  $d = k^2 \cdot d_0$  with  $d_0$  satisfying (\*\*). For any such factorization one can indeed choose  $(S, \alpha)$  as above such that in addition  $\text{ord}(\alpha) = k$ . In particular, then the untwisted Hodge structure  $\tilde{H}(S, \mathbb{Z})$  defines a point in  $D_{d_0}$ . This is best seen by starting with  $D_{d_0}$  and then choosing globally a B-field which for the very general  $S$  in  $D_{d_0}$  defines a Brauer class of order  $k$ .

**2.7** We will need to say a few things about the spherical locus  $D_{\text{sph}}$ , as this will be crucial later.

*Example 2.14.* (i) Consider  $d = 24$  which obviously satisfies (\*\*') but not (\*\*), i.e.  $D_d \subset D_{K3'}$  but  $D_d \not\subset D_{K3}$ . Also,  $D_d \subset D_{\text{sph}}$ . Indeed, if  $v$  generates  $A_2^\perp \cap K_d$ , then  $(v)^2 = -8$  and hence there exists  $\delta \in A_2 \oplus \mathbb{Z} \cdot v$  with  $(\delta)^2 = -2$ , e.g.  $2\lambda_1 + \lambda_2 + v$ . So, as mentioned before, one has a proper inclusion

$$D_{K3} \subsetneq D_{K3'} \cap D_{\text{sph}}.$$

(ii) Consider  $d = 12$  which does not satisfy (\*\*'). So,  $D_{12} \not\subset D_{K3'}$ , but  $D_{12} \subset D_{\text{sph}}$ . Indeed, in this case  $v$  in (2.4) satisfies  $(v)^2 = -4$  and, therefore,  $(\lambda_i + v)^2 = -2$ . Hence,

$$D_{\text{sph}} \not\subset D_{K3'}.$$

It would be interesting to find a numerical condition (†) such that  $D_{\text{sph}} = \bigcup D_d$  with the union over all  $d$  satisfying (†). The best we have to offer at this time is the following proposition.

PROPOSITION 2.15. *Assume  $D_d \subset D_{K3'}$  and  $9 \mid d$ . Then  $D_d \not\subset D_{\text{sph}}$ .*

*Proof.* Consider a fixed  $K_d$  and the corresponding generator  $v$  of  $K_d \cap A_2^\perp$ . As  $9 \mid d$ , clearly  $d \equiv 0 \pmod{6}$  and so  $A_2 \oplus \mathbb{Z} \cdot v \subset \tilde{\Lambda}$  is primitive. If there were a  $(-2)$ -class  $\delta \in \tilde{\Lambda}$  with  $D \cap K_d^\perp = D \cap v^\perp \subset D \cap \delta^\perp$ , then  $\delta \in A_2 + \mathbb{Z} \cdot v$  and so  $\delta = w + kv$  for some  $w \in A_2$  and  $k \in \mathbb{Z}$ . But then  $-2 = (w)^2 - k^2 d/3$ . However, if  $9 \mid d$ , then  $k^2 d/3 \equiv 0 \pmod{3}$  and hence  $(w)^2 = 2m$  with  $m \equiv 2 \pmod{3}$ , which contradicts (2.7). □

The following immediate consequence is crucial for the proof of Theorem 1.2, see § 6.1.

COROLLARY 2.16. *The locus of twisted K3 periods  $D_{K3'}$  contains infinitely many hyperplane sections  $D_d$  with  $D_d \not\subset D_{\text{sph}}$ .*

**2.8** In [Huy05] we have shown that points in  $Q$  can be understood as periods of generalized K3 surfaces. It is useful to distinguish three types.<sup>6</sup>

(i) Periods of ordinary K3 surfaces are parametrized by  $Q_{K3}$ . Up to the action of  $O(\tilde{\Lambda})$ , the set of these periods is the intersection of  $Q$  with the linear codimension-two subspace  $\mathbb{P}(\Lambda \otimes \mathbb{C}) \subset \mathbb{P}(\tilde{\Lambda} \otimes \mathbb{C})$ .

(ii) More generally, one can consider periods of the form  $\sigma + B \wedge \sigma$ , where  $\sigma \in \Lambda \otimes \mathbb{C}$  is an ordinary period and  $B \in \Lambda \otimes \mathbb{C}$  (but not necessarily  $B \in \Lambda \otimes \mathbb{Q}$ ). Up to the action of  $O(\tilde{\Lambda})$ , these periods are parametrized by the intersection of  $Q$  with the linear subspace of codimension one  $\mathbb{P}((\Lambda \oplus \mathbb{Z} \cdot f) \otimes \mathbb{C}) \subset \mathbb{P}(\tilde{\Lambda} \otimes \mathbb{C})$ . Here,  $f$  is viewed as the generator of  $H^4$ . Note that, by definition,  $Q_{K3'}$  is the subset of periods for which  $B$  can be chosen in  $\Lambda \otimes \mathbb{Q}$ .

(iii) Periods of the form  $\exp(B + i\omega) = 1 + (B + i\omega) + ((B^2 - \omega^2)/2 + (B \cdot \omega)i)$  are geometrically interpreted as periods associated with complexified symplectic forms. Here, the first and third summands are considered in  $U \simeq H^0 \oplus H^4$ . Periods of this type are parametrized by an open dense subset of  $Q$ .

In particular, all cubic periods parametrized by  $D \subset Q$  should have an interpretation in terms of these three types. This has been discussed above for type (i) and has led to consider the intersection  $D_{K3} = D \cap Q_{K3}$ . For type (ii) with  $B$  rational the intersection with the cubic period domain gives  $D_{K3'}$ . It is now natural to ask whether the remaining periods, so the periods in  $D \setminus D_{K3'}$ , are of type (ii) with  $B$  not rational or rather of type (iii), i.e. related to complexified symplectic forms. It is the latter, as shown by the following proposition.

**PROPOSITION 2.17.** *The Hodge structure of a cubic period  $x \in D$  is Hodge isometric to the Hodge structure of a twisted projective K3 surface  $(S, \alpha)$ , i.e.  $x \in D_{K3'}$ , or to the Hodge structure associated with  $\exp(B + i\omega)$ . Furthermore, if the Hodge structure of  $x$  is Hodge isometric to a Hodge structure of the type  $\sigma + B \wedge \sigma$ , then  $B$  can be chosen rational.*

*Proof.* One first observes that, analogously to Lemma 2.6(ii), a period  $x \in Q$  is of the type (ii) if and only if the integral  $(1, 1)$  part of the Hodge structure associated with  $x$  contains an isotropic direction. Indeed, if  $x$  is of type (ii), i.e. of the form  $\sigma + B \wedge \sigma$ , then  $H^4$  provides an isotropic direction of type  $(1, 1)$ . For the converse use that any isotropic direction can be completed to a hyperbolic plane  $U$  as a direct summand of  $\tilde{\Lambda}$ . Now regard  $U$  as  $H^0 \oplus H^4$  with  $H^4$  as the given isotropic direction, which is of type  $(1, 1)$ . Hence,  $x$  is indeed of type (ii).

Now let  $x \in D \cap Q$  be of type (ii). It is enough to show that then  $x \in D_{K3'}$ . The integral  $(1, 1)$ -part of the Hodge structure associated with  $x$  contains  $A_2$  and an isotropic direction, say  $\mathbb{Z} \cdot f$ . Then conclude by Proposition 2.8. □

Note that both subsets,

$$D \subset Q \quad \text{and} \quad Q_{K3} \subset Q,$$

are of codimension two and that they both parametrize periods that can be interpreted in complex geometric terms (in contrast to the ‘symplectic periods’ of the form  $\exp(B + i\omega)$ ). In fact, periods in  $D$  are even algebro-geometric in the sense that essentially all of them are associated with cubic fourfolds  $X \subset \mathbb{P}^5$ , whereas most K3 surfaces are of course not projective.

In categorical language one would want to interpret the inclusion  $D \subset Q$  for points in the complement of  $D_{K3'}$  as saying that the cubic K3 category  $\mathcal{A}_X$  associated with the cubic fourfold

---

<sup>6</sup>The discussion has been prompted by a question of Ben Bakker.



$X \subset \mathbb{P}^5$  corresponding to  $x \in D \setminus D_{K3'}$  is equivalent to the derived Fukaya category  $\mathrm{DFuk}(B + i\omega)$  associated with a complexified symplectic form  $B + i\omega$ . Deciding which symplectic structures occur here is in principle possible, but establishing an equivalence

$$\mathcal{A}_X \simeq \mathrm{DFuk}(B + i\omega)$$

will be difficult even in special cases.

The categorical interpretation of  $D_{K3} \subset Q$  is the content of [AT14], where it is proved that at least for a Zariski open dense set of periods  $x \in D_{K3}$  the cubic K3 category  $\mathcal{A}_X$  really is equivalent to  $D^b(S)$  of the K3 surface  $S$  realizing the Hodge structure associated with  $x$ . This paper deals with the categorical interpretation of  $D_{K3'} \subset Q$ .

*Remark 2.18.* The period domain  $Q \subset \mathbb{P}(\tilde{\Lambda} \otimes \mathbb{C})$  contains  $D \subset Q$  as a codimension two subset, but it also contains natural codimension one subspaces. For example, for a K3 surface  $S$  and the Mukai vector  $v = (1, 0, 1 - n) \in \tilde{H}^{1,1}(S, \mathbb{Z})$  the hyperplane section  $Q \cap v^\perp$  can be seen as the period domain for deformations of the Hilbert scheme  $S^{[n]}$ . Note, however, that from a categorical point of view the situation is different, even when one restricts to the codimension-two part that corresponds to projective deformations of the Hilbert scheme. In [MM15] it is explained how the non-full subcategory  $D^b(S) \subset D^b(S^{[n]})$  deforms sideways.

### 3. The cubic K3 category

Let  $X \subset \mathbb{P}^5$  be a smooth cubic hypersurface. The *cubic K3 category* associated with  $X$  is the category

$$\mathcal{A}_X := \langle \mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2) \rangle^\perp := \{E \in D^b(X) \mid \mathrm{Hom}(\mathcal{O}_X(i), E[*]) = 0 \text{ for } i = 0, 1, 2\}.$$

The category has first been studied by Kuznetsov in [Kuz10], see also the more recent [Kuz15]. It behaves in many respects like the derived category  $D^b(S)$  of a K3 surface  $S$ . In particular, the double shift  $E \mapsto E[2]$  defines a Serre functor of  $\mathcal{A}_X$  (see [Kuz09], [Kuz04, Corollary 4.3] and [KM09, Remark 4.2]) and the dimension of Hochschild homology of  $\mathcal{A}_X$  and of  $D^b(S)$  coincide, see [Kuz10, Kuz09].

*Example 3.1.* Due to the work of Kuznetsov [Kuz10, Kuz06], certain cubic K3 categories  $\mathcal{A}_X$  are known to be equivalent to bounded derived categories  $D^b(S, \alpha)$  of twisted K3 surfaces  $(S, \alpha)$ . For example, if the period  $x \in D$  of a cubic  $X$  is contained in  $D_8$ , then  $X$  contains a plane and for generic choices there exists a twisted K3 surface  $(S, \alpha)$  with  $\mathcal{A}_X \simeq D^b(S, \alpha)$ . Similarly, if  $X$  is a Pfaffian cubic and hence  $x \in D_{14}$ , then  $\mathcal{A}_X \simeq D^b(S)$  for the K3 surface  $S$  naturally associated with the Pfaffian  $X$ .

*Remark 3.2.* Despite the almost perfect analogy between the cubic K3 category  $\mathcal{A}_X$  and the derived category  $D^b(S)$  of K3 surfaces, certain fundamental issues are more difficult for  $\mathcal{A}_X$ . For example, to the best of my knowledge no  $\mathcal{A}_X$ , which is not equivalent to the derived category  $D^b(S, \alpha)$  of some twisted K3 surface  $(S, \alpha)$ , has yet been endowed with a bounded t-structure, let alone a stability condition. See [Tod14, Tod13] for a discussion of special stability conditions on certain  $\mathcal{A}_X$  of the form  $D^b(S, \alpha)$ .

The semi-orthogonal decomposition  $D^b(X) = \langle \mathcal{A}_X, {}^\perp\mathcal{A}_X \rangle$  with  ${}^\perp\mathcal{A}_X = \langle \mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2) \rangle$  comes with the full embedding  $i_* : \mathcal{A}_X \hookrightarrow D^b(X)$  (which is often suppressed in the notation) and the left and right adjoint functors  $i^*, i^! : D^b(X) \rightarrow \mathcal{A}_X$ , see [Kuz09, § 3] for a survey.



**3.1** For a K3 surface  $S$  the Mukai lattice  $\tilde{H}(S, \mathbb{Z})$  is endowed with the Hodge structure determined by  $\tilde{H}^{2,0}(S) = H^{2,0}(S)$  and by requiring  $\tilde{H}^{2,0} \perp \tilde{H}^{1,1}$ . Using the natural isomorphism  $K_{\text{top}}(S) \simeq H^*(S, \mathbb{Z})$  this Hodge structure can also be regarded as a Hodge structure on  $K_{\text{top}}(S)$ .

In [AT14] Addington and Thomas introduce a similar Hodge structure associated with the category  $\mathcal{A}_X$ , defined on  $K_{\text{top}}(\mathcal{A}_X)$  and denoted by  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$ . Here,  $K_{\text{top}}(\mathcal{A}_X) \subset K_{\text{top}}(X)$  is the orthogonal complement of  $\{[\mathcal{O}], [\mathcal{O}(1)], [\mathcal{O}(2)]\}$  with respect to the pairing  $\chi(\alpha, \beta) = \langle v(\alpha), v(\beta) \rangle$  defined in terms of the Mukai vector  $v : K_{\text{top}}(X) \rightarrow H^*(X, \mathbb{Q})$  and the Mukai pairing on  $H^*(X, \mathbb{Q})$ . It is not difficult to see that one has in fact a semi-orthogonal direct sum decomposition

$$K_{\text{top}}(\tilde{X}) = K_{\text{top}}(\mathcal{A}_X) \oplus \langle [\mathcal{O}_X], [\mathcal{O}_X(1)], [\mathcal{O}_X(2)] \rangle.$$

As  $H^*(X, \mathbb{Z})$  is torsion free,  $K_{\text{top}}(X)$  and

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) := K_{\text{top}}(\mathcal{A}_X)$$

are as well. The Hodge structure is then defined by  $\tilde{H}^{2,0}(\mathcal{A}_X) := v^{-1}(H^{3,1}(X))$  and the condition  $\tilde{H}^{2,0} \perp \tilde{H}^{1,1}$ . Furthermore,  $N(\mathcal{A}_X)$  and the transcendental lattice  $T(\mathcal{A}_X)$  of  $\mathcal{A}_X$  are introduced in terms of this Hodge structure as  $\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$  and its orthogonal complement  $\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})^\perp$ , respectively. As a lattice  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  is independent of  $X$  and by [AT14] any equivalence  $\mathcal{A}_X \simeq \text{D}^b(S)$  (see Example 3.1) induces a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(S, \mathbb{Z})$  (cf. Proposition 3.3). In particular,  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  for all smooth cubics is abstractly isomorphic to  $\Lambda$ .

As explained in [AT14, Proposition 2.3], the classes  $\lambda_j := [i^* \mathcal{O}_\ell(j)] \in \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ , for a line  $\ell \subset X$  and  $j = 1, 2$ , can be viewed as the standard generators of a lattice  $A_2 \subset \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ . Moreover, the Mukai vector  $v : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) = K_{\text{top}}(\mathcal{A}_X) \rightarrow H^*(X, \mathbb{Q})$  induces an isometry (up to sign)

$$\langle \lambda_1, \lambda_2 \rangle^\perp \xrightarrow{\sim} h^\perp = H^4(X, \mathbb{Z})_{\text{prim}}.$$

In particular, any marking  $\varphi : h^\perp \xrightarrow{\sim} A_2^\perp$  induces a marking  $\langle \lambda_1, \lambda_2 \rangle \oplus \langle \lambda_1, \lambda_2 \rangle^\perp \xrightarrow{\sim} A_2 \oplus A_2^\perp$  and further a marking

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{\Lambda}. \tag{3.1}$$

Conversely, any marking (3.1) inducing the standard identification  $\langle \lambda_1, \lambda_2 \rangle \xrightarrow{\sim} A_2$  yields a marking  $H^4(X, \mathbb{Z})_{\text{prim}} \xrightarrow{\sim} A_2^\perp$ . In this sense, (an open set of) points  $x \in D$  will be considered as periods of cubic K3 categories  $\mathcal{A}_X$  via their Hodge structures  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$ .

Note that the positive directions of  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  come with a natural orientation, given by the real and imaginary parts of  $\tilde{H}^{2,0}(\mathcal{A}_X)$  and the oriented basis  $\lambda_1, \lambda_2$  of  $A_2 \subset \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ .

**3.2** As we are also interested in equivalences  $\text{D}^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X$ , we collect a few relevant facts dealing with the topological K-theory of twisted K3 surfaces  $(S, \alpha)$ . As it turns out, the topological setting does not require any substantially new arguments. In order to speak of twisted sheaves or bundles, let us fix a class  $B \in H^2(S, \mathbb{Q})$  which under the exponential map  $H^2(S, \mathbb{Q}) \rightarrow H^2(S, \mathcal{O}_S^*)$  is mapped to  $\alpha$ . Next choose a Čech representative  $\{B_{ijk}\}$  of  $B \in H^2(S, \mathbb{Q})$  and consider the associated Čech representative  $\{\alpha_{ijk} := \exp(B_{ijk})\}$  of  $\alpha$ . This allows one to speak of  $\{\alpha_{ijk}\}$ -twisted sheaves and bundles, in the holomorphic as well as in the topological setting.

As explained in [HS05, Proposition 1.2], any  $\{\alpha_{ijk}\}$ -twisted bundle  $E$  can be ‘untwisted’ to a bundle  $E_B$  by changing the transition functions  $\varphi_{ij}$  of  $E$  to  $\exp(a_{ij}) \cdot \varphi_{ij}$ , where the continuous functions  $a_{ij}$  satisfy  $-a_{ij} + a_{ik} - a_{jk} = B_{ijk}$ . The process can be reversed and so the categories of

$\{\alpha_{ijk}\}$ -twisted topological bundles is equivalent to the category of untwisted topological bundles. In particular,

$$K_{\text{top}}(S, \alpha) \simeq K_{\text{top}}(S)$$

which composed with the Mukai vector yields an isomorphism  $K_{\text{top}}(S, \alpha) \simeq \tilde{H}(S, \alpha, \mathbb{Z})$  that identifies the image of  $K(S, \alpha) \rightarrow K_{\text{top}}(S, \alpha)$  with  $\tilde{H}^{1,1}(S, \alpha, \mathbb{Z})$ .

The next result is the twisted version of the observation by Addington and Thomas mentioned earlier.

**PROPOSITION 3.3.** *Any linear, exact equivalence  $\mathcal{A}_X \simeq \text{D}^b(S, \alpha)$  induces a Hodge isometry*

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(S, \alpha, \mathbb{Z}).$$

*Proof.* By results due to Orlov in the untwisted case and to Canonaco and Stellari [CS07] in the twisted case, any fully faithful functor  $\Phi : \text{D}^b(S, \alpha) \rightarrow \text{D}^b(X)$  is of Fourier–Mukai type, i.e.  $\Phi \simeq \Phi_{\mathcal{E}}$  for some  $\mathcal{E} \in \text{D}^b(S \times X, \alpha^{-1} \boxtimes 1)$ . Therefore,  $\Phi$  induces a homomorphism  $\Phi_{\mathcal{E}}^K : K_{\text{top}}(S, \alpha) \rightarrow K_{\text{top}}(X)$ , see [HvdB07, Remark 3.4].

If  $\Phi$  is induced by an equivalence  $\text{D}^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X$ , then  $\Phi_{\mathcal{E}}^K : K_{\text{top}}(S, \alpha) \xrightarrow{\sim} K_{\text{top}}(\mathcal{A}_X)$  is an isomorphism and in fact a Hodge isometry  $\tilde{H}(S, \alpha, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z})$ . The compatibility with the Hodge structure follows from the twisted Chern character  $\text{ch}^{-\alpha \boxtimes 1}(\mathcal{E})$  of the Mukai kernel being of Hodge type. See [HS05, § 1] for the notion of twisted Chern characters. That the quadratic form is respected as well is proved by mimicking the argument for FM-equivalences, see e.g. [Huy06, § 5.2].

(We are suppressing a number of technical details here. As explained before, the actual realization of the Hodge structure  $\tilde{H}(S, \alpha, \mathbb{Z})$  depends on the choice of a  $B \in H^2(S, \mathbb{Q})$  lifting  $\alpha$ . Similarly, the Chern character  $\text{ch}^{-\alpha \boxtimes 1}(\mathcal{E})$  also actually depends on  $B$ .)  $\square$

**3.3** The above result generalizes to *FM-equivalences*  $\mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$ , i.e. to equivalences for which the composition  $\text{D}^b(X) \rightarrow \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'} \rightarrow \text{D}^b(X')$  admits a Fourier–Mukai kernel. It has been conjectured that in fact any linear exact equivalence is a FM-equivalence, but the existing results do not cover our case.

**PROPOSITION 3.4.** *Any FM-equivalence  $\mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$  induces a Hodge isometry*

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z}).$$

*Proof.* The argument is an easy modification of the above.  $\square$

The following improves upon a result in [BMMS12, Proposition 6.3] where it is shown that for a cubic  $X \in \mathcal{C}_8$ , so containing a plane, there exist at most finitely many (up to isomorphism) cubics  $X_1, \dots, X_n \in \mathcal{C}_8$  with  $\mathcal{A}_X \simeq \mathcal{A}_{X_1} \simeq \dots \simeq \mathcal{A}_{X_n}$ .

**COROLLARY 3.5.** *For any given smooth cubic  $X \subset \mathbb{P}^5$  there exist up to isomorphism only finitely many smooth cubics  $X' \subset \mathbb{P}^5$  admitting a FM-equivalence  $\mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$ .*

*Proof.* The proof follows the argument for the analogous statement for K3 surfaces [BM01] closely, but needs a modification at one point that shall be explained.

Due to the proposition, it suffices to prove that up to isomorphism there are only finitely many cubics  $X'$  such that there exists a Hodge isometry  $\varphi : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ . Any such Hodge isometry induces a Hodge isometry  $\varphi_T : T(\mathcal{A}_X) \xrightarrow{\sim} T(\mathcal{A}_{X'})$  and an isometry of lattices  $N(\mathcal{A}_X) \xrightarrow{\sim} N(\mathcal{A}_{X'})$ . We may assume

$$T(\mathcal{A}_X) \subset A_2^\perp \subset \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \quad \text{and} \quad A_2 \subset N(\mathcal{A}_X)$$

and similarly for  $X'$ . Note however that these inclusions need not be respected by  $\varphi$ . The orthogonal complement of  $T(\mathcal{A}_X)^\perp \subset A_2^\perp$  is just  $N(\mathcal{A}_X) \cap A_2^\perp$  and the two inclusions of  $A_2^\perp$  induce two Hodge structures on  $A_2^\perp$ . Note that if the Hodge isometry  $\varphi_T$  can be extended to a Hodge isometry  $A_2^\perp \xrightarrow{\sim} A_2^\perp$ , which can be interpreted as a Hodge isometry  $H^4(X, \mathbb{Z})_{\text{prim}} \simeq H^4(X', \mathbb{Z})_{\text{prim}}$ , then the global Torelli theorem [Voi86, Voi08] implies that  $X \simeq X'$ .

We first show that the set of isomorphism classes of lattices  $\Gamma$  occurring as  $N(\mathcal{A}_{X'}) \cap A_2^\perp$  is finite. The required lattice theory is slightly more involved than the original in [BM01]. Let us fix two even lattices  $\Lambda_1$  and  $\Lambda$  (in our situation,  $\Lambda_1 = T(\mathcal{A}_X)$  and  $\Lambda = A_2^\perp$ ). We show that up to isomorphisms there exist only finitely many lattices  $\Lambda_2$  occurring as the orthogonal complement of some primitive embedding  $\Lambda_1 \hookrightarrow \Lambda$ . For unimodular  $\Lambda$  this is standard, but the proof can be tweaked to cover the more general statement. Of course, it suffices to show that only finitely many discriminant forms  $(A_{\Lambda_2}, q_{\Lambda_2})$  can occur. Now  $G := \Lambda / (\Lambda_1 \oplus \Lambda_2)$  is naturally a finite subgroup of  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2)$  of index  $d = |\text{disc}(\Lambda)|$ . The first projection from  $G \subset \Lambda^* / (\Lambda_1 \oplus \Lambda_2) \subset A_{\Lambda_1} \oplus A_{\Lambda_2}$  defines an isomorphism of  $G$  with a finite subgroup of  $A_{\Lambda_1}$ . This leaves only finitely many possibilities for the finite groups  $G$  and  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2)$ . Note that  $\Lambda / (\Lambda_1 \oplus \Lambda_2) \subset A_{\Lambda_1} \oplus A_{\Lambda_2}$  is isotropic but not necessarily the bigger  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2) \subset A_{\Lambda_1} \oplus A_{\Lambda_2}$ . However, the restriction of the quadratic form to  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2)$  takes values only in  $(2/d^2)\mathbb{Z}/2\mathbb{Z}$ . For fixed  $G \subset A_{\Lambda_1}$  the restriction of  $q_{\Lambda_1}$  to  $G$  can be extended in at most finitely many ways to a quadratic form on  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2)$  with values in  $(2/d^2)\mathbb{Z}/2\mathbb{Z}$ . Now use the other projection  $\Lambda^* / (\Lambda_1 \oplus \Lambda_2) \twoheadrightarrow A_{\Lambda_2}$  to see that there are only finitely many possibilities for the group  $A_{\Lambda_2}$  and also for the quadratic form  $q_{\Lambda_2}$ .

To conclude the proof, we can assume that  $\Gamma$  is fixed. For two Fourier–Mukai partners realizing the fixed  $\Gamma$ , any Hodge isometry  $T(\mathcal{A}_{X_1}) \simeq T(\mathcal{A}_{X_2})$  can be extended to a Hodge isometry  $T(\mathcal{A}_{X_1}) \oplus \Gamma \simeq T(\mathcal{A}_{X_2}) \oplus \Gamma$ . As the finite index overlattices  $T(\mathcal{A}_{X_i}) \oplus \Gamma \subset H^4(X_i, \mathbb{Z})_{\text{prim}}$  are all contained in  $(T(\mathcal{A}_{X_i}) \oplus \Gamma)^*$ , there are only finitely many choices for them, which allows one to reduce to the case that the Hodge isometry extends to a Hodge isometry  $H^4(X_1, \mathbb{Z})_{\text{prim}} \simeq H^4(X_2, \mathbb{Z})_{\text{prim}}$ .  $\square$

Two very general cubics have FM-equivalent K3 categories only if they are isomorphic.

**COROLLARY 3.6.** *Let  $X$  be a smooth cubic with  $\text{rk } H^{2,2}(X, \mathbb{Z}) = 1$ . For a smooth cubic  $X'$  there exists a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  if and only if  $X \simeq X'$ .*

*Proof.* The assumption implies that  $N(\mathcal{A}_X) \simeq A_2$ . As any FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  induces a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ , also  $N(\mathcal{A}_{X'}) \simeq A_2$ . Moreover, the natural inclusions of the transcendental lattices  $T(\mathcal{A}_X) \subset A_2^\perp$  and  $T(\mathcal{A}_{X'}) \subset A_2^\perp$  are in fact equalities and the induced Hodge isometry  $T(\mathcal{A}_X) \simeq T(\mathcal{A}_{X'})$  can therefore be read as a Hodge isometry  $H^4(X, \mathbb{Z})_{\text{prim}} \simeq H^4(X', \mathbb{Z})_{\text{prim}}$ , which by the global Torelli theorem [Voi86] implies that  $X \simeq X'$ .  $\square$

Note that, in contrast, very general projective K3 surfaces  $S$ , i.e. such that  $\rho(S) = 1$ , usually have non-isomorphic FM-partners, see [Ogu02, Ste04]. The result may also be compared to the

main result of [BMMS12] showing that for all cubic threefolds  $Y \subset \mathbb{P}^4$  the full subcategory  $\langle \mathcal{O}, \mathcal{O}(1) \rangle^\perp \subset \mathrm{D}^b(Y)$  determines  $Y$ .

*Remark 3.7.* In principle, it should be possible to count FM-partners of  $\mathcal{A}_X$  for very general special cubics  $X \in \mathcal{C}_d$  (i.e.  $\mathrm{rk} H^{2,2}(X, \mathbb{Z}) = 2$ ). On the level of Hodge theory, this amounts to counting the number of Hodge structures on  $\tilde{\Lambda}$  parametrized by  $D$  which are Hodge isometric to  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  up to those that are Hodge isometric on  $A_2^\perp$ . The arguments should follow [HLOY04, Theorem 1.4], see also [Ste04], with the additional problem that  $A_2^\perp$  is not unimodular.

As an immediate consequence of Lemma 2.2 we also note the following.

**COROLLARY 3.8.** *Let  $X$  be a special cubic defining a very general point in  $\mathcal{C}_d$ . Then*

$$\mathrm{rk}(\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})) = 3 \quad \text{and} \quad \mathrm{disc}(\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})) = d.$$

*Remark 3.9.* Suppose  $d$  satisfies (\*\*') and is written as  $d = k^2 d_0$ . Then  $d_0$  also satisfies (\*\*'). The most interesting case is when in fact  $d_0$  satisfies (\*\*). Then for very general  $X \in \mathcal{C}_d$ , there exists a twisted K3 surface  $(S, \alpha)$  with  $\alpha$  of order  $k$  and such that  $\mathcal{A}_X \simeq \mathrm{D}^b(S, \alpha)$ , see Lemma 2.13. Moreover, there also exists a cubic  $X' \in \mathcal{C}_{d_0}$  such that  $\mathcal{A}_{X'} \simeq \mathrm{D}^b(S)$ . So, a K3 surface  $S$  of the proper degree, with its various Brauer classes, is often related to more than one smooth cubic  $X$ .

**3.4** We are interested in the group  $\mathrm{Aut}(\mathcal{A}_X)$  of isomorphism classes of FM-equivalences  $\Phi : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$ . As any FM-equivalence  $\Phi$  induces a Hodge isometry

$$\Phi^H : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z}),$$

there is a natural homomorphism

$$\rho : \mathrm{Aut}(\mathcal{A}_X) \longrightarrow \mathrm{Aut}(\tilde{H}(\mathcal{A}_X, \mathbb{Z})), \quad \Phi \longmapsto \Phi^H. \tag{3.2}$$

Here,  $\mathrm{Aut}(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$  denotes the group of Hodge isometries. We say that  $\Phi$  is *symplectic* if the induced action on  $\tilde{H}^{2,0}(\mathcal{A}_X)$ , or equivalently on  $T(\mathcal{A}_X)$ , is the identity. The subgroup of symplectic autoequivalences shall be denoted by  $\mathrm{Aut}_s(\mathcal{A}_X)$ . Thus, (3.2) induces

$$\rho : \mathrm{Aut}_s(\mathcal{A}_X) \longrightarrow \mathrm{Aut}(\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})).$$

*Remark 3.10.* By  $\mathrm{Aut}^+(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$  one denotes the subgroup of Hodge isometries preserving a given orientation of the four positive directions. We expect that  $\mathrm{Im}(\rho) = \mathrm{Aut}^+(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$ . This is known if  $\mathcal{A}_X \simeq \mathrm{D}^b(S)$ , see [HMS09], and one inclusion can be proved for non-special cubics, see Theorem 1.2.

*Example 3.11.* The most important autoequivalences of K3 categories, responsible for the complexity of the groups  $\mathrm{Aut}(\mathrm{D}^b(S))$  and  $\mathrm{Aut}(\mathcal{A}_X)$ , are spherical twists. Associated with any spherical object  $A \in \mathcal{A}_X$ , i.e.  $\mathrm{Ext}^*(A, A) \simeq H^*(S^2)$ , there exists a FM-equivalence

$$T_A : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$$

that sends  $E \in \mathcal{A}_X$  to the cone  $T_A(E)$  of the evaluation map  $R\mathrm{Hom}(A, E) \otimes A \longrightarrow E$ . This is indeed a FM-equivalence: its kernel can be described as the cone of the composition  $A' \boxtimes A \xrightarrow{\mathrm{tr}} \mathcal{O}_\Delta \longrightarrow (\mathrm{id}, i)^*(\mathcal{O}_\Delta)$ . Here,  $(\mathrm{id}, i)^*$  is the left adjoint  $\mathrm{D}^b(X \times X) \longrightarrow \mathrm{D}^b(X) \boxtimes \mathcal{A}_X$  and

$A^\vee \in \mathcal{A}_X(-2)$  is the image of the classical dual of  $A$  in  $D^b(X)$  under the left adjoint of  $\mathcal{A}_X(-2) \subset D^b(X)$ . (With these choices the cone is contained in  $\mathcal{A}_X(-2) \boxtimes \mathcal{A}_X$  and would indeed induce a functor  $D^b(X) \rightarrow \mathcal{A}_X$  that is trivial on  ${}^\perp \mathcal{A}_X$ .)

The action of the spherical twist  $T_A : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$  on  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  is given by the reflection  $s_\delta : v \mapsto v + \langle v, \delta \rangle \cdot \delta$ , where  $\delta \in \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$  is the Mukai vector of  $A$ .

In [Kuz04, § 4] Kuznetsov considers the functor

$$\Psi : \mathcal{A}_X \rightarrow \mathcal{A}_X, \quad E \mapsto i^*(i_*E \otimes \mathcal{O}_X(1))[-1],$$

which turns out to be an equivalence satisfying  $\Psi^3 \simeq [-1]$ . Clearly, by construction  $\Psi$  is a FM-equivalence. In fact, for the proof that  $\mathcal{A}_X$  is a K3 category this functor is crucial. Define

$$\Phi_0 := \Psi[1],$$

which satisfies  $\Phi_0^3 \simeq [2]$ .

**PROPOSITION 3.12.** *The autoequivalence  $\Phi_0 : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$  is symplectic and the induced action  $\Phi_0^H : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z})$  corresponds to the element in  $O(A_2)$  that is given by the cyclic permutation of the roots  $\lambda_1, \lambda_2, -\lambda_1 - \lambda_2$ .*

*Proof.* As the action on cohomology is independent of the specific cubic  $X \subset \mathbb{P}^5$ , we can assume that the transcendental lattice  $T(\mathcal{A}_X) \subset \tilde{H}(\mathcal{A}_X, \mathbb{Z})$  is of odd rank. However,  $\pm \text{id}$  are the only Hodge isometries of an irreducible Hodge structure of weight two of K3 type of odd rank, cf. [Huy16, Corollary 3.3.5], and, as  $\Phi_0^3 \simeq [2]$  acts trivially on  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$ , we must have  $\Phi_0^H = \text{id}$  on  $T(\mathcal{A}_X)$ , i.e.  $\Phi_0$  is symplectic.

If  $X$  is a cubic with  $A_2 \simeq \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ , then  $\Phi_0^H$  corresponds to an element in  $O(A_2)$ . As  $\Phi_0$  is symplectic,  $\Phi_0^H = \text{id}$  on  $A_2^\perp$  and hence  $\Phi_0^H = \text{id}$  on the discriminant group  $A_{A_2}$ . Therefore,  $\Phi_0^H \in \mathfrak{S}_3$ , see Remark 2.1. For a cubic  $X$  such that  $\mathcal{A}_X \simeq D^b(S)$ , we know that  $\Phi_0^H$  must be orientation preserving by [HMS09] and thus  $\Phi_0^H \in \mathfrak{A}_3 \simeq \mathbb{Z}/3\mathbb{Z}$  in general.

It remains to show that  $\Phi_0^H \neq \text{id}$ . One way to see this relies on a direct computation. Another possibility is to use the recent result of Bayer and Bridgeland [BB13] confirming Bridgeland’s conjecture in [Bri08] in the case of a K3 surface  $S$  of Picard rank one. More precisely, due to [BB13, Theorem 1.4] for a K3 surface  $S$  with  $\rho(S) = 1$  the subgroup of  $\text{Aut}(D^b(S))$  of autoequivalences acting trivially on  $\tilde{H}(S, \mathbb{Z})$  is the product of  $\mathbb{Z}[2]$  and the free group generated by squares of spherical twists  $T_E^2$  associated with spherical vector bundles  $E$  on  $S$ . (That this is a reformulation of Bridgeland’s original conjecture for  $\rho(S) = 1$  had also been observed by Kawatani [Kaw12].) Hence, if  $\Phi_0^H = \text{id}$ , then  $\Phi_0 = (*_i T_{E_i}^2) \circ [2k]$ , but then clearly  $\Phi_0^3$  could not be isomorphic to the double shift  $[2]$ . □

**COROLLARY 3.13.** *For every smooth cubic  $X \subset \mathbb{P}^5$  the group of symplectic FM-autoequivalences  $\text{Aut}_s(\mathcal{A}_X)$  contains an infinite cyclic group  $\mathbb{Z} \subset \text{Aut}_s(\mathcal{A}_X)$  generated by  $\Phi_0$  such that*

$$\mathbb{Z} \cdot [2] \subset \mathbb{Z}$$

*is a subgroup of index three and such that the natural map  $\rho : \text{Aut}_s(\mathcal{A}_X) \rightarrow \text{Aut}(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$  defines an isomorphism of the quotient  $\mathbb{Z}/\mathbb{Z} \cdot [2]$  with the subgroup  $\mathfrak{A}_3 \subset O(A_2) \subset O(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$  of alternating permutations of the roots  $\lambda_1, \lambda_2, -\lambda_1 - \lambda_2$  of  $A_2$ .*

*Remark 3.14.* The subgroup  $\mathrm{SO}(A_2) \subset \mathrm{O}(A_2)$  of orientation-preserving isometries of  $A_2$  is  $\mathfrak{A}_3 \times \mathbb{Z}/2\mathbb{Z} \simeq \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ , see Remark 2.1. Its action can be ‘lifted’ to an action on  $\mathcal{A}_X$  via the natural extension

$$0 \longrightarrow \mathbb{Z} \cdot [2] \longrightarrow (\mathbb{Z} \cdot \Phi_0 \times \mathbb{Z} \cdot [1]) / (\Phi_0^3 - [2]) \longrightarrow \mathrm{SO}(A_2) \longrightarrow 0,$$

which can be seen as induced by the universal cover of  $\mathrm{SO}(A_2 \otimes \mathbb{R})$ . Clearly, the group in the middle is still infinite cyclic.

Inspired by Bridgeland’s conjecture for K3 surfaces in [Bri08], we state the following conjecture (see [Huy14, § 5.4] explaining this reformulation).

CONJECTURE 3.15. There exists an isomorphism

$$\mathrm{Aut}_s(\mathcal{A}_X) \simeq \pi_1^{\mathrm{st}}[P_0/\mathrm{O}].$$

Here,  $P \subset \mathbb{P}(\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z}) \otimes \mathbb{C})$  is the period domain defined analogously to  $D$  and  $Q$  in § 2.3 and  $P_0 := P \setminus \bigcup \delta^\perp$ , with the union over all  $(-2)$  classes  $\delta \in \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ . Moreover,  $\mathrm{O} \subset \mathrm{O}(\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z}))$  is the subgroup of all isometries acting trivially on the discriminant. However, contrary to the case of untwisted K3 surfaces we do not even have a natural map between these two groups at the moment.

**3.5** The cubic K3 category  $\mathcal{A}_X$  can also be described as a category of graded matrix factorizations, see [Orl09]. More precisely, there exists an exact linear equivalence

$$\mathcal{A}_X \simeq \mathrm{MF}(W, \mathbb{Z}).$$

Here,  $W \in R := k[x_0, \dots, x_5]$  is a cubic polynomial defining  $X$ . The objects of  $\mathrm{MF}(W, \mathbb{Z})$  are pairs  $(K \xrightarrow{\alpha} L, L \xrightarrow{\beta} K(3))$ , where  $K$  and  $L$  are finitely generated, free, graded  $R$ -modules and  $\alpha, \beta$  are graded  $R$ -module homomorphisms with  $\beta \circ \alpha = W \cdot \mathrm{id} = \alpha \circ \beta$ . Recall that  $K(n)$  for a graded  $R$ -module  $K = \bigoplus K_i$  is the graded module with  $K(n)_i = K_{n+i}$ . Homomorphisms in  $\mathrm{MF}(W, \mathbb{Z})$  are the obvious ones modulo those that are homotopic to zero (everything  $\mathbb{Z}/2\mathbb{Z}$ -periodic).

The *shift functor* that makes  $\mathrm{MF}(W, \mathbb{Z})$  a triangulated category is given by

$$(K \xrightarrow{\alpha} L, L \xrightarrow{\beta} K(3))[1] = (L \xrightarrow{-\beta} K(3), K(3) \xrightarrow{-\alpha} L(3)).$$

Viewing  $\mathcal{A}_X$  as the category of graded matrix factorizations allows one to describe  $\Phi_0$  in Proposition 3.12 alternatively as follows. Consider the *grade shift functor*

$$\begin{aligned} \Phi_0 : \mathrm{MF}(W, \mathbb{Z}) &\xrightarrow{\sim} \mathrm{MF}(W, \mathbb{Z}) \\ (K \xrightarrow{\alpha} L, L \xrightarrow{\beta} K(3)) &\mapsto (K(1) \xrightarrow{\alpha} L(1), L(1) \xrightarrow{\beta} K(4)). \end{aligned}$$

Then, obviously,

$$\Phi_0^3 \simeq [2].$$

Note that  $\Phi_0$  constructed in this way coincides with the FM-equivalence of Proposition 3.12, see [BFK12, Proposition 5.8].



#### 4. The Fano variety

For the sake of completeness, let us also mention the recent results of Addington [Add16] building upon an observation of Hassett [Has00], see also [MS12]. For this consider the Fano variety of lines  $F(X)$ , which, due to work of Beauville and Donagi [BD85], is a four-dimensional irreducible holomorphic symplectic variety deformation equivalent to  $\text{Hilb}^2(\text{K3})$ .

- For a smooth cubic  $X$  and its period  $x \in D$  the following two conditions are equivalent:
  - (i)  $x \in D_d$  such that  $d$  satisfies (\*\*);
  - (ii)  $F(X)$  is birational to a moduli space of stable sheaves  $M(v)$  on some K3 surface  $S$ .
- For a smooth cubic  $X$  and its period  $x \in D$  the following two conditions are equivalent:
  - (iii)  $x \in D_d$  such that there exist integers  $n$  and  $a$  with  $da^2 = 2(n^2 + n + 1)$ ;
  - (iv)  $F(X)$  is birational to the Hilbert scheme  $\text{Hilb}^2(S)$  of some K3 surface  $S$ .

Obviously, condition (iv) implies condition (ii) or, equivalently and after a moment's thought, condition (iii) implies condition (i). See [GS14] for a discussion of the relation between rationality of the cubic  $X$  and condition (iii) (or, equivalently, condition (iv)).

PROPOSITION 4.1. *For the period  $x$  of a smooth cubic  $X$  the following two conditions are equivalent:*

- (i)  $x \in D_d$  with  $d$  satisfying (\*\*');
- (ii)  $F(X)$  is birational to a moduli space of stable twisted sheaves on some K3 surface.

*Proof.* The argument is an adaptation of Addington's proof [Add16]. Note however that in the twisted case the transcendental lattice cannot play the same role as in the untwisted case. This was observed in [HS05], where it was shown that twisted K3 surfaces  $(S, \alpha)$ ,  $(S', \alpha')$  with Hodge isometric transcendental lattices,  $T(S, \alpha) \simeq T(S', \alpha')$ , need not be derived equivalent.

Following Markman [Mar11] for every hyperkähler manifold  $Y$  deformation equivalent to  $\text{Hilb}^2(S)$  of a K3 surface  $S$  there exists a distinguished primitive embedding  $H^2(Y, \mathbb{Z}) \subset \tilde{\Lambda}$  orthogonal to a vector  $v \in \tilde{\Lambda}$  with  $(v, v) = 2$ . The Hodge structure of  $H^2(Y, \mathbb{Z})$  extends to a Hodge structure on  $\tilde{\Lambda}$  such that  $v$  is of type  $(1, 1)$ . Moreover,  $Y$  and  $Y'$  are birational if and only if there exists a Hodge isometry  $H^2(Y, \mathbb{Z}) \simeq H^2(Y', \mathbb{Z})$  that extends to a Hodge isometry  $\tilde{\Lambda} \simeq \tilde{\Lambda}$ . For a moduli space  $M(v)$  of  $\alpha$ -twisted stable sheaves on a K3 surface  $S$  with primitive  $v \in \tilde{H}^{1,1}(S, \alpha, \mathbb{Z})$  such that  $(v, v) = 2$  the universal family induces the distinguished embedding (see [Yos06, Theorem 3.19])

$$H^2(M(v), \mathbb{Z}) \simeq v^\perp \hookrightarrow \tilde{H}(S, \alpha, \mathbb{Z}).$$

Similarly, and this is the other crucial input, Addington shows in [Add16, Corollary 8] that for the Fano variety of lines the universal family of lines induces this distinguished embedding

$$H^2(F(X), \mathbb{Z}) \simeq \lambda_1^\perp \hookrightarrow \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{\Lambda}.$$

Hence,  $F(X)$  and  $M(v)$  are birational if and only if there exists a Hodge isometry

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(S, \alpha, \mathbb{Z}) \tag{4.1}$$

for some twisted K3 surface  $(S, \alpha)$  that restricts to  $H^2(F(X), \mathbb{Z}) \simeq H^2(M(v), \mathbb{Z})$ . Due to Proposition 2.10, the existence of a Hodge isometry (4.1) is equivalent to  $x \in D_d$  with  $d$  satisfying (\*\*'). This proves that condition (ii) implies condition (i).



Conversely, for a Hodge isometry (4.1) consider a primitive vector  $v \in \tilde{H}^{1,1}(S, \alpha, \mathbb{Z})$  (the image of  $\lambda_1$ ) in the orthogonal complement of  $H^2(F(X), \mathbb{Z}) \hookrightarrow \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}^2(S, \alpha, \mathbb{Z})$  and the induced moduli space  $M(v)$  of stable  $\alpha$ -twisted sheaves. Write  $v = (r, \ell, s)$ . If  $r \neq 0$ , then for  $v$  or  $-v$  the moduli space  $M(v)$  is indeed non-empty. For  $r = 0$  observe that  $(v)^2 > 0$  and, hence,  $(\ell)^2 > 0$ . Again by passing to  $-v$  if necessary, one can assume that  $(\ell.H) > 0$  for the polarization  $H$ . That the moduli space is non-empty in this case was shown in [Yos09, Corollary 3.5]. (Note that for  $r = 0$  twisted sheaves can also be considered as untwisted ones.) In [Add16] the case  $r = 0$  is dealt with by a reflection associated with  $\mathcal{O}$ , which does not work in the twisted situation.

To conclude, compose the Hodge isometry  $H^2(F(X), \mathbb{Z}) \simeq v^\perp$ , given by the choice of  $v$ , with  $H^2(M(v), \mathbb{Z}) \simeq v^\perp$ , induced by the universal family as above. By construction, it extends to a Hodge isometry  $\tilde{\Lambda} \simeq \tilde{\Lambda}$  and, therefore,  $F(X)$  and  $M(v)$  are birational.  $\square$

### 5. Deformation theory

This section contains two results on the deformation theory of equivalences  $D^b(S, \alpha) \simeq \mathcal{A}_X$  respectively  $\mathcal{A}_{X'} \simeq \mathcal{A}_X$  that are crucial for the main results of the paper. The techniques have been developed by Toda [Tod09], Huybrechts *et al.* [HMS09], Huybrechts and Thomas [HT10], and in the present setting by Addington and Thomas [AT14]. We follow [AT14] quite closely and often only indicate the additional difficulties and how to deal with them.

**5.1** We first consider FM-equivalences  $\mathcal{A}_{X'} \simeq \mathcal{A}_X$  between the K3 categories of two cubics  $X$  and  $X'$  and study under which condition they deform sideways with  $X$  and  $X'$ .

**THEOREM 5.1.** *Consider two families of smooth cubics  $\mathcal{X}, \mathcal{X}' \rightarrow T$  over a smooth base  $T$  and with distinguished fibres  $X := \mathcal{X}_0$  and  $X' := \mathcal{X}'_0$ , respectively. Assume  $\Phi : \mathcal{A}_{X'} \xrightarrow{\sim} \mathcal{A}_X$  is a FM-equivalence inducing a Hodge isometry  $\varphi : \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z})$  that remains a Hodge isometry  $\varphi_t : \tilde{H}(\mathcal{A}_{\mathcal{X}'_t}, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_{\mathcal{X}_t}, \mathbb{Z})$  under parallel transport for all  $t \in T$ .*

*Then  $\Phi$  deforms sideways to FM-equivalences  $\Phi_t : \mathcal{A}_{\mathcal{X}'_t} \xrightarrow{\sim} \mathcal{A}_{\mathcal{X}_t}$  for all  $t$  in a Zariski open neighbourhood  $0 \in U \subset T$ .*

*Proof.* The argument is a variant of the deformation theory in [AT14]. We only indicate the necessary modifications.

As, by assumption,  $\Phi$  is a FM-equivalence, the composition

$$\Phi_P : D^b(X') \longrightarrow \mathcal{A}_{X'} \xrightarrow[\Phi]{\sim} \mathcal{A}_X \hookrightarrow D^b(X)$$

is a FM-functor with some kernel  $P \in D^b(X' \times X)$  contained in  $\mathcal{A}_{X'}(-2) \boxtimes \mathcal{A}_X$ . It suffices to show that  $P$  deforms to  $P_t \in D^b(\mathcal{X}'_t \times \mathcal{X}_t)$  for  $t$  in some open neighbourhood  $0 \in U \subset T$ , because the conditions for  $\Phi_{P_t}$  to factorize via a functor  $\Phi_t : \mathcal{A}_{\mathcal{X}'_t} \rightarrow \mathcal{A}_{\mathcal{X}_t}$  and for this functor  $\Phi_t$  to define an equivalence are both Zariski open. Indeed,  $\Phi_t$  takes values in  $\mathcal{A}_{\mathcal{X}_t}$  if and only if its composition with the projection  $D^b(\mathcal{X}_t) \rightarrow {}^\perp \mathcal{A}_{\mathcal{X}_t} = \langle \mathcal{O}_{\mathcal{X}_t}, \mathcal{O}_{\mathcal{X}_t}(1), \mathcal{O}_{\mathcal{X}_t}(2) \rangle$  is trivial. The composition, however, is again of FM-type and the vanishing of a FM-kernel is a Zariski open condition. Similarly, whether  $\Phi_t$  is an equivalence can be detected by composing it with its adjoints and then checking whether the natural map to the kernel of the identity is an isomorphism, again a Zariski open condition.

The crucial part is to understand the first-order deformations, the higher-order obstructions are dealt with by the  $T^1$ -lifting property, see [AT14, § 7.2] and [HMS09, § 3.2]. First note that by results of Kuznetsov [Kuz09] one has

$$HH^*(\mathcal{A}_{X'}) \simeq \text{Ext}_{X' \times X}^*(P, P) \simeq HH^*(\mathcal{A}_X).$$

This allows one to compare the first-order deformations

$$\kappa_{X'} \in H^1(T_{X'}) \subset HH^2(X') \quad \text{and} \quad \kappa_X \in H^1(T_X) \subset HH^2(X)$$

corresponding to some tangent vector  $v \in T_0T$  of  $T$  at 0. Due to a result of Toda [Tod09] (cf. [AT14, Theorem 7.1]) it suffices to show that under  $HH^2(X') \rightarrow \text{Ext}_{X' \times X}^2(P, P)$  respectively  $HH^2(X) \rightarrow \text{Ext}_{X' \times X}^2(P, P)$  the classes  $\kappa_{X'}$  and  $\kappa_X$  are mapped to the same class. For this consider the following diagram (cf. [AT14, Proposition 6.2]).

$$\begin{array}{ccccccc}
 HH_2(X') & \xrightarrow{\sim} & HH_2(\mathcal{A}_{X'}) & \xrightarrow[\Phi_P^{HH^*}]{\sim} & HH_2(\mathcal{A}_X) & \xrightarrow{\sim} & HH_2(X) \\
 \kappa_{X'} \downarrow & & \downarrow \alpha & & \downarrow \bar{\kappa}_X & & \downarrow \kappa_X \\
 HH_0(X') & \xrightarrow{\sim} & HH_0(\mathcal{A}_{X'}) & \xrightarrow[\Phi_P^{HH^*}]{\sim} & HH_0(\mathcal{A}_X) & \xrightarrow{\sim} & HH_0(X) \\
 \downarrow & & & & & & \downarrow \\
 H^*(X') & \xrightarrow[\Phi_P^H]{\sim} & & & & & H^*(X) \\
 & & & & & & \uparrow \\
 & & \tilde{H}^*(\mathcal{A}_{X'}) & \xrightarrow[\varphi]{\sim} & \tilde{H}^*(\mathcal{A}_X) & & 
 \end{array}$$

By  $H^*(X) \simeq HH_*(X)$  we denote the HKR isomorphism (see [Cál05]) post-composed with  $\sqrt{\text{td}} \wedge (\ )$  and so, in particular,  $HH_2(X) \simeq H^1(\Omega_X^3)$  with chosen generator  $\sigma_X$ . Similarly for  $X'$ , where we choose the generator  $\sigma_{X'} \in H^1(\Omega_{X'}^3) \simeq HH_2(X')$  such that its image yields  $\sigma_X$ . Furthermore,  $\bar{\kappa}_X$  denotes the image of  $\kappa_X$  under the projection  $HH^2(X) \rightarrow HH^2(\mathcal{A}_X)$ , see [Kuz09], and  $\alpha := \Phi^{HH^*}(\bar{\kappa}_X)$ .

We aim at showing that (1) is commutative. For this note first that (4) is induced by the FM-transform  $\Phi_P : D^b(X') \rightarrow D^b(X)$  and hence commutative due to [MS09]. The commutativity of (2) is obvious, as Hochschild (co)homology is respected by equivalences, and commutativity of (3) is the analogue of [AT14, Proposition 6.1]. (Recall that  $\Phi_P$  does not necessarily induce a map  $\Phi_P^{HH^*}$ , as it is not fully faithful.)

The first-order version of the assumption on the Hodge isometry  $\varphi$  is the statement that the diagram

$$\begin{array}{ccc}
 H^1(T_{X'}) & \xleftarrow{T_0T} & H^1(T_X) \\
 \downarrow \sigma_{X'} & & \downarrow \sigma_X \\
 H^{2,2}(X')_{\text{prim}} & & H^{2,2}(X)_{\text{prim}} \\
 \downarrow & & \downarrow \\
 \tilde{H}^{1,1}(\mathcal{A}_{X'}) & \xrightarrow[\varphi]{\sim} & \tilde{H}^{1,1}(\mathcal{A}_X)
 \end{array}$$

is commutative. Using the ring-module isomorphism  $(HH^*, HH_*) \simeq (H^*(\wedge^* T), H^*(\Omega^*))$  for  $X'$ , this implies that the image in  $H^*(X')$  of  $\sigma_{X'} \in HH_2(X')$  under contraction with  $\kappa_{X'}$  is mapped

to the image of  $\sigma_X$  under contraction with  $\kappa_X$ . As  $HH_2(X')$  is one-dimensional, this shows that also (1) is commutative.

Therefore, in the diagram

$$\begin{CD} HH^2(X') @>>> HH^2(\mathcal{A}_{X'}) @>\simeq>> HH^2(\mathcal{A}_X) \\ @V\sigma_{X'}VV @VV\sigma_{\mathcal{A}_{X'}}V @VV\sigma_{\mathcal{A}_X}V \\ HH_0(X') @>>> HH_0(\mathcal{A}_{X'}) @>\simeq>> HH_0(\mathcal{A}_X) \end{CD}$$

the image of  $\kappa_{X'} \in HH^2(X')$  under the two compositions  $HH^2(X') \rightarrow HH_0(\mathcal{A}_{X'}) \simeq HH_0(\mathcal{A}_X)$  coincide. As the contraction  $HH^2(\mathcal{A}) \hookrightarrow HH_0(\mathcal{A})$  is injective (as for K3 surfaces), this implies that the image of  $\kappa_{X'}$  under  $HH^2(X') \rightarrow HH^2(\mathcal{A}_X)$  is indeed  $\bar{\kappa}_X$  as claimed.

As in [AT14], the deformation of  $P$  to first and then, by  $T^1$ -lifting property, to higher order is unique, for  $\text{Ext}^1_{X' \times X}(P, P) \simeq HH^1(\mathcal{A}_X) = 0$  by [Kuz09].  $\square$

**5.2** We now come to the more involved situation of equivalences  $D^b(S, \alpha) \simeq \mathcal{A}_X$  and their deformations.

**THEOREM 5.2.** *Consider two families  $\mathcal{X}, \mathcal{S} \rightarrow T$  of smooth cubics and K3 surfaces, respectively, over a smooth base  $T$ . Denote the distinguished fibres by  $X := \mathcal{X}_0, S := \mathcal{S}_0$  and let  $\alpha_t \in \text{Br}(\mathcal{S}_t)$  be a deformation of a Brauer class  $\alpha := \alpha_0$  on  $S$ . Assume  $\Phi : D^b(S, \alpha) \xrightarrow{\simeq} \mathcal{A}_X$  is an equivalence inducing a Hodge isometry  $\varphi : \tilde{H}(S, \alpha, \mathbb{Z}) \xrightarrow{\simeq} \tilde{H}(\mathcal{A}_X, \mathbb{Z})$  that remains a Hodge isometry  $\varphi_t : \tilde{H}(\mathcal{S}_t, \alpha_t, \mathbb{Z}) \xrightarrow{\simeq} \tilde{H}(\mathcal{A}_{X_t}, \mathbb{Z})$  under parallel transport for all  $t \in T$ .*

*Then  $\Phi$  deforms sideways to equivalences  $\Phi_t : D^b(\mathcal{S}_t, \alpha_t) \xrightarrow{\simeq} \mathcal{A}_{X_t}$  for all  $t$  in a Zariski open neighbourhood  $0 \in U \subset T$ .*

*Proof.* Let us fix representatives  $\alpha_t = \{\alpha_{t,ijk}\}$  for the Brauer classes on  $\mathcal{S}_t$  and a family  $E_t$  of locally free  $\{\alpha_{t,ijk}\}$ -twisted sheaves on the fibres  $\mathcal{S}_t$  in a Zariski open neighbourhood of  $0 \in U \subset T$ .

The proof now consists of copying [AT14, §§6, 7]. However, the techniques have to be adapted to the twisted case, which sometimes causes additional problems as certain fundamental issues related to Hochschild (co)homology have not been addressed in the twisted setting. For certain parts we choose *ad hoc* arguments to reduce to the untwisted case, for others we rely on Reinecke [Rei14].

Section 6 in [AT14] deals with Hochschild (co)homology. For a twisted variety  $(Z, \alpha)$  one defines  $HH^n(Z, \alpha) := \text{Ext}^n_{(Z, \alpha^{-1}) \times (Z, \alpha)}(\mathcal{O}_\Delta, \mathcal{O}_\Delta)$ . Here,  $(Z, \alpha^{-1}) \times (Z, \alpha)$  denotes the twisted variety  $(Z \times Z, \alpha^{-1} \boxtimes \alpha)$ . Note that  $\mathcal{O}_\Delta$  is indeed an  $(\alpha^{-1} \boxtimes \alpha)$ -twisted sheaf. Similarly, one defines  $HH_n(Z, \alpha) := \text{Ext}^{d-n}_{(Z, \alpha^{-1}) \times (Z, \alpha)}(\Delta_* \omega_Z^{-1}, \mathcal{O}_\Delta)$ , where  $d = \dim(Z)$ . Composition makes  $HH_*(Z, \alpha)$  a right  $HH^*(Z, \alpha)$ -module. Moreover, there are natural isomorphisms

$$\begin{aligned} HH^n(Z, \alpha) &= \text{Ext}^n_{(Z, \alpha^{-1}) \times (Z, \alpha)}(\mathcal{O}_\Delta, \mathcal{O}_\Delta) \simeq \text{Ext}^n_Z(\Delta^* \mathcal{O}_\Delta, \mathcal{O}_Z) \\ &\simeq \text{Ext}^n_{Z \times Z}(\mathcal{O}_\Delta, \mathcal{O}_\Delta) = HH^n(Z) \end{aligned}$$

and

$$\begin{aligned} HH_n(Z, \alpha) &= \text{Ext}^{d-n}_{(Z, \alpha^{-1}) \times (Z, \alpha)}(\Delta_* \omega_Z^{-1}, \mathcal{O}_\Delta) \simeq \text{Ext}^{d-n}_Z(\mathcal{O}_Z, \Delta^* \mathcal{O}_\Delta) \\ &\simeq \text{Ext}^{d-n}_{Z \times Z}(\Delta_* \omega_Z^{-1}, \mathcal{O}_\Delta) = HH_n(Z). \end{aligned}$$

In particular, the HKR isomorphisms post-composed with  $\mathrm{td}^{-1/2} \lrcorner(\ )$  respectively  $\mathrm{td}^{1/2} \wedge(\ )$  yield isomorphisms

$$I : HH^n(Z, \alpha) \xrightarrow{\sim} \bigoplus_{i+j=n} H^i(\Lambda^j T_Z) \quad \text{and} \quad I : HH_n(Z, \alpha) \xrightarrow{\sim} \bigoplus_{j-i=n} H^i(\Omega_Z^j).$$

Note that these isomorphisms are again compatible with the ring and module structures on both sides, which follows from the fact that the isomorphisms  $HH^*(Z, \alpha) \simeq HH^*(Z)$  and  $HH_*(Z, \alpha) \simeq HH_*(Z)$  are. The latter is a consequence of the functoriality properties of  $\Delta_!$ ,  $\Delta_*$  and  $\Delta^*$ .

For a twisted K3 surface  $(S, \alpha)$  one has  $HH_2(S, \alpha) \simeq H^0(\omega_S) = \mathbb{C} \cdot \sigma_S$  and the following diagram commutes.

$$\begin{array}{ccc} HH^2(S, \alpha) & \xrightarrow[\ I]{\sim} & H^0(\Lambda^2 T_S) \oplus H^1(T_S) \oplus H^2(\mathcal{O}_S) \\ \downarrow \cdot \sigma_S & & \downarrow \lrcorner \sigma_S \\ HH_0(S, \alpha) & \xrightarrow[\ I]{\sim} & H^{0,0} \oplus H^{1,1}(S) \oplus H^{2,2}(S) \end{array}$$

Let us now consider the fully faithful functor  $\Phi_P : D^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X \hookrightarrow D^b(X)$  between the twisted K3 surface  $(S, \alpha)$  and the smooth cubic  $X$ , where  $P \in D^b((S, \alpha^{-1}) \times X)$ . Then as in [AT14, § 6.1] one obtains natural maps

$$\Phi_P^{HH^*} : HH^*(X) \longrightarrow HH^*(S, \alpha) \quad \text{and} \quad \Phi_P^{HH_*} : HH_*(S, \alpha) \longrightarrow HH_*(X)$$

compatible with the module structures, i.e.  $\Phi_P^{HH^*}(a) \circ c = \Phi_P^{HH^*}(a \circ \Phi_P^{HH_*}(c))$  for all  $a \in HH_*(S, \alpha)$  and  $c \in HH^*(X)$ . This has been checked by Reinecke in [Rei14, § 4].

The remaining input in the proof of [AT14, Proposition 6.2] is the commutativity of the untwisted version of the following diagram.

$$\begin{array}{ccc} HH_0(S, \alpha) & \xrightarrow{\Phi_P^{HH_*}} & HH_0(X) \\ I^B \downarrow \wr & & \wr \downarrow I \\ \bigoplus H^{p,p}(S) & \xrightarrow{\Phi_P^{H,B}} & \bigoplus H^{p,p}(X) \end{array} \tag{5.1}$$

Note that defining the induced action on cohomology requires the lift of  $\alpha$  to a class  $B \in H^2(S, \mathbb{Q})$ , see [Huy05, HS05]. Moreover, the usual HKR isomorphism  $I$  post-composed with  $\mathrm{td}^{1/2} \wedge(\ )$  needs to be twisted further to  $I^B := \exp(B) \circ I$ .

In principle, one could try to prove the commutativity of (5.1) by rewriting the existing untwisted theory, in particular [Că105, MS09], for the twisted situation. Instead, we follow Yoshioka [Yos06] and reduce everything to the untwisted case by pulling back to a Brauer–Severi variety. We briefly review his approach and explain how to apply it to our situation.

Following [Yos06] we pick a locally free  $\alpha = \{\alpha_{ijk}\}$ -twisted sheaf  $E = \{E_i, \varphi_{ij}\}$  on a twisted variety  $(Z, \alpha)$  and associate to it the projective bundle  $\pi : Y := \mathbb{P}(E) \longrightarrow Z$ , which naturally comes with a  $\pi^* \alpha^{-1}$ -twisted line bundle  $L := \mathcal{O}_\pi(1)$ . The pull-back of any  $\alpha$ -twisted sheaf  $F = \{F_i, \psi_{ij}\}$  tensored with  $L$  then naturally leads to the untwisted sheaf  $\tilde{F} := \pi^* F \otimes L$ .

Analogously, any  $\alpha^{-1}$ -twisted sheaf  $F$  can be turned into the untwisted sheaf  $\pi^*F \otimes L^*$ . The construction yields a functor  $D^b(Z, \alpha) \rightarrow D^b(Y)$  which, in fact, defines an equivalence of  $D^b(Z, \alpha)$  with a full subcategory

$$\tilde{(\ )} : D^b(Z, \alpha) \xrightarrow{\sim} D^b(Y/Z) \subset D^b(Y).$$

The construction applied to  $E$  itself yields the sheaf  $G := \tilde{E}$  that corresponds to the unique non-trivial extension class in  $\text{Ext}_Y^1(\mathcal{T}_\pi, \mathcal{O}_Y)$  and  $D^b(Y/Z) \subset D^b(Y)$  can alternatively be described as the full subcategory of all objects  $H$  for which the adjunction map  $\pi^*\pi_*(G^* \otimes H) \rightarrow G^* \otimes H$  is an isomorphism. Analogously,  $D^b(Z, \alpha^{-1})$  is equivalent to the full subcategory of objects  $H$  for which  $\pi^*\pi_*(G \otimes H) \xrightarrow{\sim} G \otimes H$ .

We apply this construction to the twisted K3 surface  $(S, \alpha)$  and consider  $Y = \mathbb{P}(E) \rightarrow S$  as above. Assume  $\alpha$  is of order  $r$  and choose a lift  $B = (1/r)B_0$  with  $B_0 \in H^2(S, \mathbb{Z})$  of it. The FM-kernel of our given equivalence  $\Phi_P : D^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X \subset D^b(X)$ , which is an object in  $D^b((S, \alpha^{-1}) \times X)$ , is turned into the untwisted sheaf  $\tilde{P} := \pi^*P \otimes (L^* \boxtimes \mathcal{O})$  on  $Y \times X$ . This leads to the following commutative diagram.

$$\begin{CD} D^b(S, \alpha) @>\pi_1^*>> D^b((S, \alpha) \times X) @>\otimes P>> D^b(S \times X) @>\pi_{2*}>> D^b(X) \\ @VV\tilde{(\ )}V @VV\tilde{(\ )}V @VV\pi^*V @VV=V \\ D^b(Y/S) @>\pi_1^*>> D^b((Y \times X)/(S \times X)) @>\otimes \tilde{P}>> D^b((Y \times X)/(S \times X)) @>\pi_{2*}>> D^b(X) \end{CD}$$

Therefore, the FM-functor  $\Phi_P : D^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X \subset D^b(X)$  can be written as the composition  $\Phi_P = \Phi_{\tilde{P}} \circ \Phi_Q$  of a twisted FM-functor  $\Phi_Q := \tilde{(\ )}$ , with  $Q = (\mathcal{O}_S \boxtimes L)|_{\Gamma_\pi}$ , and an untwisted FM-functor  $\Phi_{\tilde{P}}$ :

$$\Phi_P : D^b(S, \alpha) \xrightarrow{\Phi_Q} D^b(Y) \xrightarrow{\Phi_{\tilde{P}}} D^b(X). \tag{5.2}$$

This allows one to decompose the diagram (5.1) as

$$\begin{CD} HH_0(S, \alpha) @>\Phi_Q^{HH^*}>> HH_0(Y) @>\Phi_{\tilde{P}}^{HH^*}>> HH_0(X) \\ @V I^B \wr VV @VV \wr V @VV \wr V \\ \bigoplus H^{p,p}(S) @>\Phi_Q^{H,B}>> \bigoplus H^{p,p}(Y) @>\Phi_{\tilde{P}}^H>> \bigoplus H^{p,p}(X). \end{CD} \tag{5.3}$$

The right-hand square is induced by the usual untwisted FM-functor  $\Phi_{\tilde{P}}$  and its commutativity therefore follows from the result of Macrì and Stellari [MS09, Theorem 1.2]. Hence, it suffices to prove the commutativity of the left-hand square (which does not involve the cubic  $X$  anymore). For greater clarity we split this further by decomposing  $\Phi_Q$  as

$$\Phi_Q : D^b(S, \alpha) \xrightarrow{\pi^*} D^b(Y, \pi^*\alpha) \xrightarrow{L \otimes} D^b(Y).$$

Let us first consider  $\pi^* : D^b(S, \alpha) \rightarrow D^b(Y, \pi^*\alpha)$  and the induced diagram

$$\begin{array}{ccc}
 HH_0(S, \alpha) & \longrightarrow & HH_0(Y, \pi^*\alpha) \\
 \downarrow \wr & & \downarrow \wr \\
 HH_0(S) & \longrightarrow & HH_0(Y) \\
 \downarrow I & & \downarrow I \\
 H^*(S) & \xrightarrow{\sqrt{\text{td}_\pi} \cdot \pi^*} & H^*(Y) \\
 \downarrow \exp(B) & & \downarrow \exp(\pi^*B) \\
 H^*(S) & \xrightarrow{\sqrt{\text{td}_\pi} \cdot \pi^*} & H^*(Y)
 \end{array}$$

$I^B$  on the left and  $I^{\pi^*B}$  on the right, with curved arrows connecting the top and bottom rows.

Note that the usual  $\sqrt{\text{td}_\pi} \cdot \pi^*$  on the bottom is indeed the map on cohomology induced by the functor  $\pi^* : D^b(S, \alpha) \rightarrow D^b(Y, \pi^*\alpha)$ , which *a priori* depends on the choice of the lifts of  $\alpha$  and  $\pi^*\alpha$  to classes in  $H^2(S, \mathbb{Q})$  and  $H^2(Y, \mathbb{Q})$ , respectively, for which we choose  $B$  and  $\pi^*B$ . The commutativity of the upper and the lower squares is trivial. The commutativity of the middle square is an easy case of [MS09, Theorem 1.2]. Next consider  $\Psi := L \otimes (\ ) : D^b(Y, \alpha) \rightarrow D^b(Y)$  and the following induced diagram (where  $\psi$  is defined by the requirement of commutativity).

$$\begin{array}{ccc}
 HH_0(Y, \pi^*\alpha) & \xrightarrow{\Psi^{HH_*}} & HH_0(Y) \\
 \downarrow \wr & \circlearrowleft & \downarrow = \\
 HH_0(Y) & \xrightarrow{\psi} & HH_0(Y) \\
 \downarrow I & & \downarrow I \\
 H^*(Y) & & H^*(Y) \\
 \downarrow \exp(\pi^*B) & & \downarrow \\
 H^*(Y) & \xrightarrow{\Psi^{H, \pi^*B}} & H^*(Y)
 \end{array}$$

By definition,  $\Psi^{H, \pi^*B}$  is given by multiplication with  $\text{ch}^{\pi^*(-B)}(L) = \exp(-\pi^*B) \cdot \exp(c_1(L))$ . Here, use that  $L^r$  is an untwisted line bundle and define  $c_1(L) := (1/r)c_1(L^r) \in H^{1,1}(Y, \mathbb{Q})$ . See [HS05, § 1] for the conventions concerning twisted Chern classes. In particular,  $\Psi^{H, \pi^*B} \circ \exp(\pi^*B) = \exp(c_1(L))$  and, therefore, it suffices to prove the commutativity of the diagram

$$\begin{array}{ccc}
 HH_0(Y) & \xrightarrow{\psi} & HH_0(Y) \\
 \downarrow I & & \downarrow I \\
 H^*(Y) & \xrightarrow{\exp(c_1(L))} & H^*(Y)
 \end{array} \tag{5.4}$$

which no longer depends on  $B$  and is a special case of Lemma 5.3 below.

This concludes the proof of the commutativity of the diagram (5.1) and hence of [AT14, Proposition 6.2] in our twisted setting. More precisely, if a first-order deformation of  $X$  in  $D_d$  given by a class  $\kappa_X \in H^1(T_X)$  corresponds via the interpretation of  $D_d$  as period domain for  $X$  and  $S$  to a first-order deformation  $\kappa_S \in H^1(T_S)$ , then  $\Phi^{HH^2} : HH^2(X) \rightarrow HH^2(S, \alpha)$  sends  $\kappa_X$  to  $\kappa_S$ .

To conclude the proof one has to prove that the kernel  $P \in D^b((S, \alpha^{-1}) \times X)$  deforms sideways, for which we again apply Yoshioka’s untwisting technique. Instead of attempting to deform the twisted  $P$  sideways with  $(S, \alpha) \times X$  we deform the untwisted  $\tilde{P}$ . As the condition describing the full subcategory  $D^b((S, \alpha^{-1}) \times X) \simeq D^b((Y \times X)/(S \times X)) \subset D^b(Y \times X)$  is open, any deformation of  $\tilde{P}$  will automatically induce a deformation of  $P$ .<sup>7</sup> The decomposition (5.2) leads to a diagram

$$\begin{array}{ccccc}
 HH^2(X) = \text{Ext}_{X \times X}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}) & \longrightarrow & \text{Ext}_{Y \times X}^2(\tilde{P}, \tilde{P}) & \xrightarrow{\sim} & \text{Ext}_{(S, \alpha^{-1}) \times X}^2(P, P) \\
 & & \uparrow & & \uparrow \wr \\
 & & \text{Ext}_{Y \times Y}^2(\mathcal{O}_{\Delta_Y}, \mathcal{O}_{\Delta_Y}) & \dashrightarrow & \text{Ext}_{(S, \alpha^{-1}) \times (S, \alpha)}^2(\mathcal{O}_{\Delta_S}, \mathcal{O}_{\Delta_S}) \\
 & & \parallel & & \parallel \\
 & & HH^2(Y) & \dashrightarrow & HH^2(S, \alpha).
 \end{array}$$

Recall that  $\Phi_R^{HH^*}$  is defined for any FM-functor  $\Phi_R$ , whereas in order to define  $\Phi_R^{HH^*}$  one needs  $\Phi_R$  to be fully faithful, which is the case for  $\Phi_P$  and  $\Phi_Q = (\tilde{\phantom{P}})$ . So, both maps in

$$\begin{array}{ccccc}
 HH^2(X) & \longrightarrow & HH^2(S, \alpha) & \longleftarrow & HH^2(Y) \\
 \kappa_X \longmapsto & & \kappa_S & \longleftarrow & \kappa_Y
 \end{array}$$

are well defined, where as above  $\kappa_X \in H^1(T_X) \subset HH^2(X)$  corresponds to  $\kappa_S \in H^1(T_S) \subset HH^2(S, \alpha)$  (via their periods or, equivalently, via  $\Phi^{HH^2}$ ) and  $\kappa_Y$  is determined by our pre-chosen deformation  $E_t$  of  $E$ .

Now by [HT10] the obstruction  $o(\tilde{P})$  can be expressed as

$$o(\tilde{P}) = (\kappa_Y, \kappa_X) \circ \text{At}(\tilde{P}).$$

(Unfortunately, an analogous formula in the twisted case is not available.) The crucial [AT14, Theorem 7.1], which goes back to Toda [Tod09], proves that in the untwisted case  $o(P) = 0$  if  $\kappa_X$  is mapped to  $\kappa_S$  under  $HH^2(X) \rightarrow HH^2(S)$ . However, in the twisted situation one has to face the additional problem that there is no natural map  $HH^2(X) \rightarrow HH^2(Y)$ . Nevertheless, the argument in [AT14] goes through essentially unchanged as follows. Using the same notation, one writes

$$o(\tilde{P}) = \pi_1^* \kappa_Y \circ \text{At}_Y(\tilde{P}) + \pi_2^* \kappa_X \circ \text{At}_X(\tilde{P}) \in \text{Ext}^2(\tilde{P}, \tilde{P}).$$

The first term is the image of  $\pi_1^* \kappa_Y \circ \text{At}_1(\mathcal{O}_{\Delta_Y}) \in \text{Ext}^2(\mathcal{O}_{\Delta_Y}, \mathcal{O}_{\Delta_Y}) = HH^2(Y)$  which is just  $\kappa_Y$ , whereas the second one is the image of  $-\pi_1^* \kappa_X \circ \text{At}_2(\mathcal{O}_{\Delta_X}) \in \text{Ext}^2(\mathcal{O}_{\Delta_X}, \mathcal{O}_{\Delta_X}) = HH^2(X)$  which is just  $-\kappa_X$ . Hence, to compare  $\kappa_X$  and  $\kappa_Y$  we do not need a map  $HH^2(X) \rightarrow HH^2(Y)$  (which simply does not exist naturally), as we only need to compare their images in  $\text{Ext}^2(\tilde{P}, \tilde{P}) \simeq HH^2(S, \alpha)$ . Therefore, it suffices to ensure that under  $HH^2(X) \rightarrow HH^2(S, \alpha)$  the class  $\kappa_X$  is mapped to  $\kappa_S$ , which was verified above.

<sup>7</sup> This is confirmed by the observation that under the natural isomorphisms

$$\text{Ext}_{(S, \alpha^{-1}) \times X}^2(P, P) \simeq \text{Ext}_{(Y, \pi^* \alpha^{-1}) \times X}^2(\pi^* P, \pi^* P) \simeq \text{Ext}_{Y \times X}^2(\tilde{P}, \tilde{P})$$

the obstruction  $o(P) \in \text{Ext}_{(S, \alpha^{-1}) \times X}^2(P, P)$  to deform  $P$  sideways to first order is first mapped to  $o(\pi^* P)$  and then to  $o(\tilde{P}) - \text{id}_{\pi^* P} \otimes o(\mathcal{O}_\pi(-1))$ . The latter, however, equals the obstruction  $o(\tilde{P}) \in \text{Ext}_{Y \times X}^2(\tilde{P}, \tilde{P})$  for  $\tilde{P}$ , because  $\mathcal{O}_\pi(-1)$  clearly deforms sideways.



This concludes the argument proving that the FM-kernel  $P$  deforms to first order with  $(S, \alpha) \times X$ . The arguments in [AT14, §7.2] proving the existence of deformations of  $P$  to all orders apply verbatim. Note that at the very end of the argument one needs to apply a result of Lieblich [Lie06] saying that the space of objects with no negative self-Exts in the derived category is an Artin stack of locally finite presentation. Again, the result as such does not seem to be in the literature for the twisted situation, but once again it can be deduced from the untwisted case by Yoshioka’s trick.  $\square$

It remains to check the commutativity of (5.4) which is a general fact. Consider a smooth variety  $Z$  and  $\alpha_{ijk} := \beta_{ij} \cdot \beta_{jk} \cdot \beta_{ki}$  with  $\beta_{ij} \in \mathcal{O}_{U_{ij}}^*$ . The associated Brauer class  $\alpha \in H^2(Z, \mathcal{O}_Z^*)$  is of course trivial and hence  $D^b(Z, \{\alpha_{ijk}\})$  and  $D^b(Z)$  are equivalent categories and an explicit equivalence can be given by ‘untwisting by  $\{\beta_{ij}\}$ ’, i.e. by  $E = \{E_i, \varphi_{ij}\} \mapsto \{E_i, \varphi_{ij} \cdot \beta_{ij}^{-1}\}$ . Note that changing  $\beta_{ij}$  by a cocycle  $\{\delta_{ij}\}$ , which would correspond to an untwisted line bundle say  $M$ , the equivalence would be modified by  $M^* \otimes (\ )$ .

Assume furthermore that  $\alpha_{ijk}^r = 1$ . Then  $\{\beta_{ij}^r\}$  is a cocycle defining a line bundle  $H$  and we define  $c_1(\beta) := (1/r)c_1(H) \in H^{1,1}(Z)$ . Explicitly,  $c_1(\beta) = \{d \log \beta_{ij}\}$ .

LEMMA 5.3. *The ‘untwisting by  $\{\beta_{ij}\}$ ’, i.e. the equivalence*

$$\Phi : D^b(Z, \{\alpha_{ijk}\}) \xrightarrow{\sim} D^b(Z), \quad E = \{E_i, \varphi_{ij}\} \mapsto \{E_i, \varphi_{ij} \cdot \beta_{ij}^{-1}\},$$

induces a commutative diagram

$$\begin{array}{ccc} HH_*(Z, \{\alpha_{ijk}\}) & \xrightarrow{\Phi^{HH_*}} & HH_*(Z) \\ \text{HKR} \downarrow & & \downarrow \text{HKR} \\ H^*(Z) & \xrightarrow{\exp(c_1(\beta))} & H^*(Z). \end{array}$$

The commutativity of (5.4) then follows from the observation that  $L \otimes (\ )$  can be written as the composition of the ‘untwisting by  $\{\beta_{ij}\}$ ’ as above with the equivalence  $\mathcal{L} \otimes (\ )$ . Here,  $\mathcal{L}$  is the untwisted line bundle given by  $\{\psi_{ij} \cdot \beta_{ij}\}$ , where  $L$  itself is the  $\{\alpha_{ijk}^{-1}\}$ -twisted line bundle given by  $\{\psi_{ij}\}$ .

Indeed, for  $\Psi := \mathcal{L} \otimes (\ ) : D^b(Z) \xrightarrow{\sim} D^b(Z)$  the commutativity of

$$\begin{array}{ccc} HH_*(Z) & \xrightarrow{\Psi^{HH_*}} & HH_*(Z) \\ \text{HKR} \downarrow & & \downarrow \text{HKR} \\ H^*(Z) & \xrightarrow{\exp(c_1(\mathcal{L}))} & H^*(Z) \end{array}$$

is an easy special case of<sup>8</sup> [MS09, Theorem 1.2], which can be proved by a direct calculation. The proof of the lemma is a variant of this computation.

*Proof.* Consider the universal Atiyah class  $\text{At} : \mathcal{O}_\Delta \rightarrow \Delta_*\Omega_Z[1]$ . Twisted with a line bundle of the form  $M \boxtimes M^*$  it yields a map  $\text{At}_M : \mathcal{O}_\Delta \rightarrow \Delta_*\Omega_Z[1]$ . The usual formula  $c_1(E \otimes M) = c_1(E) + \text{rk } E \cdot c_1(M)$  corresponds to the universal formula  $\text{At}_M = \alpha + \Delta_*c_1(M)$ , which can be checked

<sup>8</sup> Note that  $\text{td}^{1/2} \wedge$  can be dropped here and in the lemma, as it commutes with  $\exp(c_1(\mathcal{L}))$ .

by using arguments of [BF08, BF03] or a direct cocycle computation. Here,  $c_1(M)$  is viewed as a map  $\mathcal{O}_Z \rightarrow \Omega_Z[1]$  which can be pushed forward via  $\Delta$ . Similarly, the exponential  $\exp(\text{At}) : \mathcal{O}_\Delta \rightarrow \bigoplus \Delta_* \Omega_Z^i[i]$  (see [Că105]) twisted with  $M \boxtimes M^*$  is given by  $\exp(\text{At})_M = \Delta_* \exp(c_1(M)) \circ \exp(\text{At})$ .

Let now  $f \in HH_j(Z) = \text{Ext}_{Z \times Z}^j(\Delta_! \mathcal{O}_Z, \Delta_* \mathcal{O}_Z)$  and denote by  $F \in \text{Ext}_Z^j(\mathcal{O}_Z, \Delta^* \Delta_* \mathcal{O}_Z)$  its image under  $\text{Ext}_{Z \times Z}^j(\Delta_! \mathcal{O}_Z, \Delta_* \mathcal{O}_Z) \simeq \text{Ext}_Z^j(\mathcal{O}_Z, \Delta^* \Delta_* \mathcal{O}_Z)$ . So if  $\eta : (\Delta_! \Delta^*) \Delta_* \mathcal{O}_Z \rightarrow \Delta_* \mathcal{O}_Z$  denotes adjunction, then  $f = \eta \circ \Delta_! F$ . Due to [Că105, Proposition 4.4], the latter is under the HKR isomorphism given by  $\exp(\text{At})$ , so

$$\eta : (\Delta_! \Delta^*) \Delta_* \mathcal{O}_Z \simeq \bigoplus \Delta_* (\Omega_Z^i[i] \otimes \omega_Z^{-1}[-d]) \simeq \bigoplus \Delta_* (\Omega_Z^{d-i})^*[i-d] \xrightarrow{\exp(\text{At})} \Delta_* \mathcal{O}_Z.$$

The image of  $f$  under  $\mathcal{L} \otimes ( \ )$  is given by tensoring with  $\mathcal{L} \boxtimes \mathcal{L}^*$ . The push-forward  $\Delta_! F$  remains unchanged by tensoring with  $\mathcal{L} \boxtimes \mathcal{L}^*$  and by the above  $\eta$  changes by composing with  $\Delta_* \exp(c_1(\mathcal{L}))$ .

Literally the same argument applies to the untwisting by  $\{\beta_{ij}\}$  for which one has to observe that the universal Atiyah class  $\text{At} : \mathcal{O}_\Delta \rightarrow \Delta_* \Omega_Z[1]$  on  $(Z, \{\alpha_{ijk}^{-1}\}) \times (Z, \{\alpha_{ijk}\})$  under untwisting by  $\{\beta_{ij}\}$  becomes  $\text{At} + \Delta_* c_1(\beta) : \mathcal{O}_\Delta \rightarrow \Delta_* \Omega_Z[1]$  on  $Z \times Z$ .  $\square$

### 6. Proofs

#### 6.1 Proof of Theorem 1.2

(i) According to Corollary 3.13, for every smooth cubic  $X \subset \mathbb{P}^5$  there exists a distinguished FM-autoequivalence  $\Phi_0 : \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_X$  of infinite order which acts as the identity on  $T(\mathcal{A}_X)$ , so it is symplectic, and such that  $\Phi_0^3$  is the double shift  $E \mapsto E[2]$ . We have to show that for the very general cubic every symplectic FM-equivalence  $\Phi$  is a power of  $\Phi_0$ .

As  $\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z}) \simeq A_2$  for very general  $X$  and  $\Phi^H = \text{id}$  on  $T(\mathcal{A}_X) = A_2^\perp$ , the induced action  $\Phi^H$  is contained in  $O(A_2)$ . Clearly, any Hodge isometry of  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  that is the identity on  $A_2^\perp$  stays a Hodge isometry for all deformations of  $X$ . Therefore, applying Theorem 5.1,  $\Phi$  deforms to FM-autoequivalences  $\Phi_t : \mathcal{A}_{\mathcal{X}_t} \simeq \mathcal{A}_{\mathcal{X}_t}$  for cubics  $\mathcal{X}_t$  in a Zariski open neighbourhood  $U \subset \mathcal{C}$  of  $X$  inside the moduli space of smooth cubics.

Then for all but finitely many  $d$  satisfying (\*\*) the intersection  $U \cap \mathcal{C}_d$  is non-empty (and open) and, therefore, by [AT14, Theorem 1.1] there exists  $t \in U$  such that  $\mathcal{A}_{\mathcal{X}_t} \simeq D^b(S)$  for some K3 surface  $S$ . Due to [HMS09, Theorem 2], autoequivalences of  $D^b(S)$  are orientation preserving and hence  $\Phi^H \in \mathfrak{A}_3 \simeq \mathbb{Z}/3\mathbb{Z}$ , cf. Remark 2.1. Thus, by composing with some power of  $\Phi_0$ , we may assume that  $\Phi^H = \text{id}$ .

Now apply Corollary 2.16 and Theorem 1.4, to be proved below, to conclude that there exists  $t \in U$  such that  $\mathcal{A}_{\mathcal{X}_t} \simeq D^b(S, \alpha)$  for a twisted K3 surface  $(S, \alpha)$  not admitting any  $(-2)$  class. Indeed,

$$(\mathcal{C}_{K3'} \cap U) \setminus \mathcal{C}_{\text{sph}} \neq \emptyset,$$

where  $\mathcal{C}_{K3'} := \bigcup_{(**)} \mathcal{C}_d \subset \mathcal{C}$  and  $\mathcal{C}_{\text{sph}} \subset \mathcal{C}$  is the image of  $D_{\text{sph}}$ . By [HMS08, Theorem 2], we know that then  $\Phi_t$  is isomorphic to an even shift  $E \mapsto E[2k]$ . It is easy to see that  $k$  is independent of  $t$ .

The locus of points  $U_0 \subset U$  such that  $\Phi_t \simeq [2k]$  for  $t \in U_0$  is Zariski open and by the above non-empty. Therefore, for every  $X \in \mathcal{C}$  in the intersection of all  $U_0 \subset \mathcal{C}$  the assertion holds. But this intersection is certainly countable, as FM-kernels are parametrized by countably many products of Quot schemes.

(ii) Now consider a non-special cubic  $X$ , i.e.  $X \in \mathcal{C} \setminus \bigcup \mathcal{C}_d$ , and an arbitrary  $\Phi \in \text{Aut}(\mathcal{A}_X)$ . By composing with the shift functor [1] if necessary, we may assume that  $\Phi^H$  acts trivially on the discriminant group  $A_{A_2} \simeq A_{A_2^\perp}$ . But then the induced Hodge isometry of  $T(\mathcal{A}_X) \simeq A_2^\perp$  extends to a Hodge isometry of  $H^4(X, \mathbb{Z})$  that respects  $h$ . By the global Torelli theorem [Voi86, Loo09, Cha12] it is therefore induced by an automorphism  $f \in \text{Aut}(X)$ , which clearly acts trivially on the orthogonal complement of  $h^\perp \subset H^*(X, \mathbb{Z})$  and hence as the identity on  $A_2 \subset \tilde{H}(\mathcal{A}_X, \mathbb{Z})$ . Moreover, since  $f$  respects  $H^{3,1}(X)$ , the action of  $f$  in  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$  preserves the orientation.

So, composing, if necessary,  $\Phi$  with the shift functor and an automorphism, we reduce to the case  $\Phi \in \text{Aut}_s(\mathcal{A}_X)$ . As  $X$  is non-special, i.e.  $A_2 \simeq \tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z})$ , we can deform  $\Phi$  sideways as above until it can be interpreted as an autoequivalence of a category of the form  $D^b(S)$ , which implies that it is orientation preserving. This eventually proves that for every non-special cubic the image of  $\rho : \text{Aut}(\mathcal{A}_X) \rightarrow \text{Aut}(\tilde{H}(\mathcal{A}_X, \mathbb{Z}))$  is the subgroup of orientation-preserving Hodge isometries.

*Remark 6.1.* We expect the first assertion in Theorem 1.2 to hold for every non-special cubic, i.e. for all  $X \in \mathcal{C} \setminus \bigcup \mathcal{C}_d$ , but this would require to show that if  $\Phi \in \text{Aut}(\mathcal{A}_X)$  deforms to the identity functor and  $\tilde{H}^{1,1}(\mathcal{A}_X, \mathbb{Z}) \simeq A_2$ , then  $\Phi \simeq \text{id}$ . The techniques of [HMS08] should be useful here, but they require the existence of stability conditions.

Furthermore, one would also expect that any  $\Phi \in \text{Aut}(\mathcal{A}_X)$  of any cubic preserves the natural orientation.

**6.2 Proof of Theorem 1.4**

Assertion (i) follows from Theorem 1.3 and Proposition 3.3. For the converse, fix  $d$  satisfying (\*\*'). Then for any smooth cubic  $X \in \mathcal{C}_d$  there exists a Hodge isometry

$$\varphi : \tilde{H}(S, \alpha, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \tag{6.1}$$

for some twisted K3 surface  $(S, \alpha)$ . In fact, this Hodge isometry can be chosen globally over the period domain  $D_d$  (or some appropriately constructed covering  $\tilde{\mathcal{C}}_d$  of  $\mathcal{C}_d$ , see [AT14]). The aim is to show that generically this Hodge isometry is induced by an equivalence  $\mathcal{A}_X \simeq D^b(S, \alpha)$  (up to changing the orientation).

The starting point for the argument is [AT14, Theorem 4.1], which is based on Kuznetsov’s work [Kuz10] and on the description of the image of the period map for cubic fourfolds due to Laza [Laz10] and Looijenga [Loo09]. Combined, these results show that for every  $d$  satisfying (\*\*') (but in fact (\*) is enough) there exists a smooth cubic  $X \in \mathcal{C}_8 \cap \mathcal{C}_d$ , a K3 surface  $S_0$  and an equivalence

$$\Phi_0 : \mathcal{A}_X \xrightarrow{\sim} D^b(S_0).$$

By [AT14] or Proposition 3.3, any such  $\Phi_0$  induces a Hodge isometry  $\Phi_0^H : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(S_0, \mathbb{Z})$  (usually completely unrelated to (6.1)). Consider now the composition

$$\psi := \Phi_0^H \circ \varphi : \tilde{H}(S, \alpha, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(S_0, \mathbb{Z}).$$

By modifying  $\varphi$  (globally over  $D_d$ ) if necessary (use Lemma 2.3), we may assume that  $\psi$  preserves the orientation and then [HS06] applies and shows that there exists an equivalence  $\Psi : D^b(S, \alpha) \xrightarrow{\sim} D^b(S_0)$  with  $\Psi^H = \psi$ . Then the equivalence

$$\Phi := \Phi_0^{-1} \circ \Psi : D^b(S, \alpha) \xrightarrow{\sim} D^b(S_0) \xrightarrow{\sim} \mathcal{A}_X$$

satisfies  $\Phi^H = \varphi$ .

We can now forget about  $S_0$  and only keep  $X$  and  $S$  and the equivalence  $\Phi = \Phi_P$  with  $P \in D^b((S, \alpha^{-1}) \times X)$ . Then consider the two families  $\mathcal{X}_t$  and  $(\mathcal{S}_t, \alpha_t)$  over  $D_d$  (or rather  $\tilde{\mathcal{C}}_d$  in order to use the Zariski topology) of cubics respectively twisted K3 surfaces with  $X = \mathcal{X}_0, S = \mathcal{S}_0$ , for which  $\varphi$  defines Hodge isometries  $\tilde{H}(\mathcal{S}_t, \alpha_t, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_{\mathcal{X}_t}, \mathbb{Z})$  for all  $t$ . As  $\Phi : D^b(S, \alpha) \xrightarrow{\sim} \mathcal{A}_X$  induces  $\varphi$ , Theorem 5.2 applies and shows that  $\Phi$  can be deformed to equivalences  $\Phi_t : D^b(\mathcal{S}_t, \alpha_t) \xrightarrow{\sim} \mathcal{A}_{\mathcal{X}_t}$  for all  $t$  in a Zariski open neighbourhood of  $0 \in \tilde{\mathcal{C}}_d$ .

**6.3 Proof of Theorem 1.5**

The first assertion of the theorem has been proved already as Corollary 3.6. For assertions (ii) and (iii) recall that any FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$  induces a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$ , cf. Proposition 3.4. So it remains to prove the converse for generic  $X \in \mathcal{C}_d$  with  $d$  satisfying (\*\*') respectively very general  $X \in \mathcal{C}_d$  for arbitrary  $d$ . The first case is easy, as then, by Theorem 1.4,  $\mathcal{A}_X \simeq D^b(S, \alpha)$  and  $\mathcal{A}_{X'} \simeq D^b(S', \alpha')$  for twisted K3 surfaces  $(S, \alpha)$  respectively  $(S', \alpha')$ . The assertion then follows from [HS06] and Lemma 2.3.

For the second case consider the correspondence

$$Z := \{(X, X', \varphi) \mid X \in \mathcal{C}_d \text{ and } \varphi : \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow{\sim} \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})\}$$

of smooth cubics  $X, X'$  with  $X \in \mathcal{C}_d$  and a Hodge isometry  $\varphi$ . (Note that with  $X$  also  $X' \in \mathcal{C}_d$ .) This correspondence consists of countably many components  $Z_i \subset Z$  and for the image of a component  $Z_0 \subset Z$  under the first projection  $\pi : Z \rightarrow \mathcal{C}_d$  one either has  $\pi(Z_0) \subset \mathcal{C}_d \cap \bigcup_{d' \neq d} \mathcal{C}_{d'}$  or  $\pi(Z_0) \subset \mathcal{C}_d$  is dense.

As we are interested in very general  $X \in \mathcal{C}_d$  only, we may assume that we are in the latter situation. Then by [AT14, Theorem 1.1], cf. §6.1, one finds a  $(X, X', \varphi) \in Z_0$  for which there exist K3 surfaces  $S$  and  $S'$  and FM-equivalences

$$\Psi : \mathcal{A}_X \xrightarrow{\sim} D^b(S) \quad \text{and} \quad \Psi' : \mathcal{A}_{X'} \xrightarrow{\sim} D^b(S'). \tag{6.2}$$

By Proposition 3.3,  $\Psi$  and  $\Psi'$  induce Hodge isometries  $\Psi^H$  respectively  $\Psi'^H$ , which composed with  $\varphi$  yield a Hodge isometry

$$\varphi_0 : \tilde{H}(S, \mathbb{Z}) \xrightarrow[\Psi^{-1}]{\sim} \tilde{H}(\mathcal{A}_X, \mathbb{Z}) \xrightarrow[\varphi]{\sim} \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z}) \xrightarrow[\Psi'^H]{\sim} \tilde{H}(S', \mathbb{Z}).$$

We may assume that  $\varphi_0$  is orientation preserving and, thus, induced by a FM-equivalence  $\Phi_0 : D^b(S) \xrightarrow{\sim} D^b(S')$ . Composing the latter with the equivalences (6.2) yields a FM-equivalence  $\Phi : \mathcal{A}_X \simeq \mathcal{A}_{X'}$  inducing  $\varphi$ . Now use Theorem 5.1 to deform  $\Phi$  sideways to FM-equivalences  $\Phi_t : \mathcal{A}_{X_t} \xrightarrow{\sim} \mathcal{A}_{X'_t}$  for all points  $(X_t, X'_t, \varphi_t \equiv \varphi)$  in a Zariski dense open subset  $U_0 \subset Z_0$ .

Hence, for all  $X \in \bigcap \pi(U_i)$ , with the intersection over all components  $Z_i \subset Z$  (dominating  $\mathcal{C}_d$ ), the existence of a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \simeq \tilde{H}(\mathcal{A}_{X'}, \mathbb{Z})$  implies the existence of a FM-equivalence  $\mathcal{A}_X \simeq \mathcal{A}_{X'}$ .

ACKNOWLEDGEMENTS

I would like to thank Nick Addington and Sasha Kuznetsov for very helpful discussions during the preparation of the paper. I am also grateful to Ben Bakker, Daniel Halpern-Leistner, Jørgen Rennemo, Paolo Stellari, Andrey Soldatenkov, and Richard Thomas for comments and suggestions. I enjoyed several discussions with Alex Perry, in particular on the possibility of proving a result like Theorem 1.1, for which he has also found a proof. Thanks to Emanuel Reinecke and Pawel Sosna for a long list of comments on the first version and to the referee for a very careful reading and innumerable suggestions.

## REFERENCES

- Add16 N. Addington, *On two rationality conjectures for cubic fourfolds*, Math. Res. Lett. **23** (2016), 1–13.
- AT14 N. Addington and R. Thomas, *Hodge theory and derived categories of cubic fourfolds*, Duke Math. J. **163** (2014), 1885–1927.
- BB13 A. Bayer and T. Bridgeland, *Derived automorphism groups of K3 surfaces of Picard rank 1*, Duke Math. J., to appear. Preprint (2013), [arXiv:1310.8266](https://arxiv.org/abs/1310.8266).
- BFK12 M. Ballard, D. Favero and L. Katzarkov, *Orlov spectra: bounds and gaps*, Invent. Math. **189** (2012), 359–430.
- BD85 A. Beauville and R. Donagi, *La variété des droites d'une hypersurface cubique de dimension 4*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), 703–706.
- BMMS12 M. Bernardara, E. Macrì, S. Mehrotra and P. Stellari, *A categorical invariant for cubic threefolds*, Adv. Math. **229** (2012), 770–803.
- Bri08 T. Bridgeland, *Stability conditions on K3 surfaces*, Duke Math. J. **141** (2008), 241–291.
- BM01 T. Bridgeland and A. Maciocia, *Complex surfaces with equivalent derived categories*, Math. Z. **236** (2001), 677–697.
- BF03 R.-O. Buchweitz and H. Flenner, *A semiregularity map for modules and applications to deformations*, Compositio Math. **137** (2003), 135–210.
- BF08 R.-O. Buchweitz and H. Flenner, *The global decomposition theorem for Hochschild (co-)homology of singular spaces via the Atiyah–Chern character*, Adv. Math. **217** (2008), 243–281.
- Că100 A. Căldăraru, *Derived categories of twisted sheaves on Calabi–Yau manifolds*, PhD thesis, Cornell (2000).
- Că105 A. Căldăraru, *The Mukai pairing II. The Hochschild–Kostant–Rosenberg isomorphism*, Adv. Math. **194** (2005), 34–66.
- CS07 A. Canonaco and P. Stellari, *Twisted Fourier–Mukai functors*, Adv. Math. **212** (2007), 484–503.
- Cha12 F. Charles, *A remark on the Torelli theorem for cubic fourfolds*, Preprint (2012), [arXiv:1209.4509](https://arxiv.org/abs/1209.4509).
- Cha16 F. Charles, *Birational boundedness for holomorphic symplectic varieties, Zarhin's trick for K3 surfaces, and the Tate conjecture*, Ann. of Math. (2) **184** (2016), 487–526.
- Cox89 D. Cox, *Primes of the form  $x^2 + ny^2$ . Fermat, class field theory and complex multiplication* (John Wiley & Sons, New York, 1989).
- GS14 S. Galkin and E. Shinder, *The Fano variety of lines and rationality problem for a cubic hypersurface*, Preprint (2014), [arXiv:1405.5154](https://arxiv.org/abs/1405.5154).
- Has00 B. Hassett, *Special cubic fourfolds*, Compositio Math. **120** (2000), 1–23.
- HvdB07 L. Hille and M. van den Bergh, *Fourier–Mukai transforms*, in *Handbook of tilting theory*, London Mathematical Society Lecture Note Series, vol. 332 (Cambridge University Press, Cambridge, 2007), 147–177.
- HLOY04 S. Hosono, B. Lian, K. Oguiso and S.-T. Yau, *Fourier–Mukai number of a K3 surface*, CRM Proc. Lecture Notes **38** (2004), 177–192.
- Huy05 D. Huybrechts, *Generalized Calabi–Yau structures, K3 surfaces, and B-fields*, Int. J. Math. **19** (2005), 13–36.
- Huy06 D. Huybrechts, *Fourier–Mukai transforms in algebraic geometry*, Oxford Mathematical Monographs (Oxford University Press, Oxford, 2006).
- Huy09 D. Huybrechts, *The global Torelli theorem: classical, derived, twisted*, in *Algebraic geometry–Seattle 2005. Part 1*, Proceedings of Symposia in Pure Mathematics, vol. 80 (American Mathematical Society, Providence, RI, 2009), 235–258.

- Huy14 D. Huybrechts, *Introduction to stability conditions*, in *Moduli spaces*, London Mathematical Society Lecture Notes Series, vol. 411 (Cambridge University Press, Cambridge, 2014), 179–229.
- Huy16 D. Huybrechts, *Lectures on K3 surfaces*, Cambridge Studies in Advanced Mathematics (Cambridge University Press, Cambridge, 2016), <http://www.math.uni-bonn.de/people/huybrech/K3.html>.
- HMS08 D. Huybrechts, E. Macrì and P. Stellari, *Stability conditions for generic K3 categories*, *Compositio Math.* **144** (2008), 134–162.
- HMS09 D. Huybrechts, E. Macrì and P. Stellari, *Derived equivalences of K3 surfaces and orientation*, *Duke Math. J.* **149** (2009), 461–507.
- HS05 D. Huybrechts and P. Stellari, *Equivalences of twisted K3 surfaces*, *Math. Ann.* **332** (2005), 901–936.
- HS06 D. Huybrechts and P. Stellari, *Proof of Căldăraru’s conjecture*, in *Moduli spaces and arithmetic geometry*, Advanced Studies in Pure Mathematics, vol. 45, (2006), 31–42.
- HT10 D. Huybrechts and R. Thomas, *Deformation-obstruction theory for complexes via Atiyah and Kodaira–Spencer classes*, *Math. Ann.* **346** (2010), 545–569.
- Kaw12 K. Kawatani, *A hyperbolic metric and stability conditions on K3 surfaces with  $\rho = 1$* , Preprint (2012), [arXiv:1204.1128](https://arxiv.org/abs/1204.1128).
- Kne02 M. Kneser, *Quadratische formen* (Springer, 2002).
- Kuz04 A. Kuznetsov, *Derived categories of cubic and  $V_{14}$  threefolds*, *Proc. Steklov Inst. Math.* **3** (2004), 171–194; [arXiv:math/0303037](https://arxiv.org/abs/math/0303037).
- Kuz06 A. Kuznetsov, *Homological projective duality for Grassmannians of lines*, Preprint (2006), [arXiv:math.AG/0610957](https://arxiv.org/abs/math.AG/0610957).
- Kuz09 A. Kuznetsov, *Hochschild homology and semiorthogonal decompositions*, Preprint (2009), [arXiv:0904.4330](https://arxiv.org/abs/0904.4330).
- Kuz10 A. Kuznetsov, *Derived categories of cubic fourfolds*, in *Cohomological and geometric approaches to rationality problems*, Progress in Mathematics, vol. 282 (Springer, Berlin, 2010), 219–243.
- Kuz15 A. Kuznetsov, *Calabi–Yau and fractional Calabi–Yau categories*, Preprint (2015), [arXiv:1509.07657](https://arxiv.org/abs/1509.07657).
- KM09 A. Kuznetsov and D. Markushevich, *Symplectic structures on moduli spaces of sheaves via the Atiyah class*, *J. Geom. Phys.* **59** (2009), 843–860.
- Laz10 R. Laza, *The moduli space of cubic fourfolds via the period map*, *Ann. of Math. (2)* **172** (2010), 673–711.
- Lie06 M. Lieblich, *Moduli of complexes on a proper morphism*, *J. Algebraic Geom.* **15** (2006), 175–206.
- LMS14 M. Lieblich, D. Maulik and A. Snowden, *Finiteness of K3 surfaces and the Tate conjecture*, *Ann. Sci. Éc. Norm. Supér.* **47** (2014), 285–308.
- Loo09 E. Looijenga, *The period map for cubic fourfolds*, *Invent. Math.* **177** (2009), 213–233.
- Mar11 E. Markman, *A survey of Torelli and monodromy results for holomorphic-symplectic varieties*, in *Complex and differential geometry*, Proceedings in Mathematics, vol. 8 (Springer, Berlin, 2011), 257–322.
- MS09 E. Macrì and P. Stellari, *Infinitesimal derived Torelli theorem for K3 surfaces, (Appendix by S. Mehrotra)*, *Int. Math. Res. Not. IMRN* **2009** (2009), 3190–3220.
- MS12 E. Macrì and P. Stellari, *Fano varieties of cubic fourfolds containing a plane*, *Math. Ann.* **354** (2012), 1147–1176.
- MM15 E. Markman and S. Mehrotra, *Integral transforms and deformations of K3 surfaces*, Preprint (2015), [arXiv:1507.03108](https://arxiv.org/abs/1507.03108).



- Muk87 S. Mukai, *On the moduli space of bundles on K3 surfaces. I. Vector bundles on algebraic varieties (Bombay, 1984)*, Tata Inst. Fund. Res. Stud. Math. **11** (1987), 341–413.
- Nik79 V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), 111–177.
- Ogu02 K. Oguiso, *K3 surfaces via almost-primes*, Math. Res. Lett. **9** (2002), 47–63.
- Orl97 D. Orlov, *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. **84** (1997), 1361–1381.
- Orl09 D. Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, in *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin*, Vol. II, Progress in Mathematics, vol. 270 (Springer, Berlin, 2009), 503–531.
- Rei14 E. Reinecke, *Autoequivalences of twisted K3 surfaces*, Master thesis, Bonn (2014), <http://www.math.uni-bonn.de/people/huybrech/ReineckeMA.pdf>.
- Ste04 P. Stellari, *Some remarks about the FM-partners of K3 surfaces with Picard numbers 1 and 2*, Geom. Dedicata **108** (2004), 1–13.
- Tod09 Y. Toda, *Deformations and Fourier–Mukai transforms*, J. Differential Geom. **81** (2009), 197–224.
- Tod13 Y. Toda, *Gepner type stability condition via Orlov/Kuznetsov equivalence*, Preprint (2013), [arXiv:1308.3791](https://arxiv.org/abs/1308.3791).
- Tod14 Y. Toda, *Gepner type stability conditions on graded matrix factorizations*, Algebr. Geom. **1** (2014), 613–665.
- Voi86 C. Voisin, *Théorème de Torelli pour les cubiques de  $\mathbb{P}^5$* , Invent. Math. **6** (1986), 577–601.
- Voi08 C. Voisin, *Correction à : ‘Théorème de Torelli pour les cubiques de  $\mathbb{P}^5$ ’*, Invent. Math. **172** (2008), 455–458.
- Yos06 K. Yoshioka, *Moduli spaces of twisted sheaves on a projective variety*, in *Moduli Spaces and Arithmetic Geometry*, Advanced Studies in Pure Mathematics, vol. 45 (World Scientific, Singapore, 2006), 1–30.
- Yos09 K. Yoshioka, *Stability and the Fourier–Mukai transform. II*, Compositio Math. **145** (2009), 112–142.

Daniel Huybrechts [huybrech@math.uni-bonn.de](mailto:huybrech@math.uni-bonn.de)

Mathematisches Institut, Universität Bonn, Endenicher Allee 60, 53115 Bonn, Germany