

NOTES FOR AUTHORS

The *Bulletin of Entomological Research* publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the *Bulletin* is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

Page Format. The *Bulletin* is printed in a two-column format (column width of three inches) with a text area of 170×240 mm.

Text. Papers should be typed, one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

On acceptance, word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with less type-setting errors. Further instructions regarding coding of word-processor documents are available on request.

Abstract. Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words.

Tables. Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format.

Illustrations. Text figures, line drawings and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and Table captions should be typed on a separate sheet in the following format:

Figs 23–26. Figs 23–24, *Urophora* eggs: 23, *U. hispanica*; 24, *U. stigma*. Figs 25–26, spermathecae: 25, *U. maura*; 26, *U. stigma*; scale lines = 0.05 mm.

References. References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 in Waage, J. & Greathead, D. (Eds) *Insect parasitoids*. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) *Ecological methods with particular reference to the study of insect populations*. 2nd edn. 524 pp. London, Chapman & Hall.

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, *Metopolophium dirhodum* (Walker) (Hemiptera: Aphididae), in the UK. *Bulletin of Entomological Research* 79, 373–380.

Citation of authors in the text should appear in the form “Wilson (1986)” or “(Wilson, 1986)”. More than one author should be cited as “(Holloway *et al.*, 1987; Walker & Huddleston, 1988)”.

Offprints. 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. The original manuscript and artwork plus two copies should be submitted to:

The Editors
Bulletin of Entomological Research
CAB International Institute of Entomology
56, Queen's Gate
London
SW7 5JR, United Kingdom.

Bulletin of Entomological Research

| | |
|---|-----|
| Guest Editorial..... | 359 |
| Bosque-Pérez, N. A. & Mareck, J. H. Distribution and species composition of lepidopterous maize borers in southern Nigeria | 363 |
| Esser, J. R. Factors influencing oviposition, larval growth and mortality in <i>Chrysomya megacephala</i> (Diptera: Calliphoridae), a pest of salted dried fish in south-east Asia..... | 369 |
| Farrell, J. A. & Stufkens, M. W. The impact of <i>Aphidius rhopalosiphi</i> (Hymenoptera: Aphidiidae) on populations of the rose grain aphid (<i>Metopolophium dirhodum</i>) (Hemiptera: Aphididae) on cereals in Canterbury, New Zealand | 377 |
| Fay, H. A. C., MacQueen, A. & Doube, B. M. Impact of fauna on mortality and size of <i>Haematobia</i> spp. (Diptera: Muscidae) in natural dung pads in Australia and South Africa..... | 385 |
| Gomulski, L. Polyandry in nulliparous <i>Anopheles gambiae</i> mosquitoes (Diptera: Culicidae) | 393 |
| Hargrove, J. W. & Langley, P. A. Sterilizing tsetse (Diptera: Glossinidae) in the field: a successful trial..... | 397 |
| Kyorku, C., Brightwell, R. & Dransfield, R. D. Traps and odour baits for the tsetse fly, <i>Glossina longipennis</i> (Diptera: Glossinidae)..... | 405 |
| Löhr, B., Varela, A. M. & Santos, B. Exploration for natural enemies of the cassava mealybug, <i>Phenacoccus manihoti</i> (Homoptera: Pseudococcidae), in South America for the biological control of this introduced pest in Africa..... | 417 |
| McClay, A. S., McFadyen, R. E. & Bradley, J. D. Biology of <i>Bucculatrix parthenica</i> Bradley sp.n. (Lepidoptera: Bucculatricidae) and its establishment in Australia as a biological control agent for <i>Parthenium hysterophorus</i> (Asteraceae) | 427 |
| Obeng-Ofori, D. & Coaker, T. H. Some factors affecting responses of four stored product beetles (Coleoptera: Tenebrionidae & Bostrichidae) to pheromones | 433 |
| Obeng-Ofori, D. & Coaker, T. H. <i>Tribolium</i> aggregation pheromone: monitoring, range of attraction and orientation behaviour of <i>T. castaneum</i> (Coleoptera: Tenebrionidae)..... | 443 |
| Peiris, H. T. R. & Hemingway, J. Mechanisms of insecticide resistance in a temephos selected <i>Culex quinquefasciatus</i> (Diptera: Culicidae) strain from Sri Lanka | 453 |
| Phillips, A., Sabatini, A., Milligan, P. J. M., Boccolini, D., Broomfield, G. & Molyneux, D. H. The <i>Anopheles maculipennis</i> complex (Diptera: Culicidae): comparison of the cuticular hydrocarbon profiles determined in adults of five Palaearctic species | 459 |
| Price, R. E. & Brown, H. D. Reproductive performance of the African migratory locust, <i>Locusta migratoria migratorioides</i> (Orthoptera: Acrididae), in a cereal crop environment in South Africa | 465 |
| Ridsdill-Smith, T. J. & Hayles, L. Stages of bush fly, <i>Musca vetustissima</i> (Diptera: Muscidae), killed by scarabaeine dung beetles (Coleoptera: Scarabaeidae) in unfavourable cattle dung..... | 473 |
| Williams, B. G., Dransfield, R. D. & Brightwell, R. Tsetse fly (Diptera: Glossinidae) population dynamics and the estimation of mortality rates from life-table data | 479 |
| Book Reviews..... | 487 |
| Index of Authors (Volume 80)..... | 491 |