





Research Article

The SISAL webApp: exploring the speleothem climate and environmental archives of the world

István Gábor Hatvani^{1,2,†} , Zoltán Kern^{1,2,†} , Péter Tanos³, Micah Wilhelm⁴, Franziska A. Lechleitner⁵ 
and Nikita Kaushal^{6*} 

¹Institute for Geological and Geochemical Research, Research Centre for Astronomy and Earth Sciences, ELKH, Budapest, Budaörsi út 45, H-1112, Hungary; ²CSFK, MTA Centre of Excellence, Budapest, Konkoly Thege Miklós út 15-17., H-1121, Hungary; ³Department of Geology, Institute of Geography and Earth Sciences, ELTE, Budapest, Hungary; ⁴Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland; ⁵Department of Chemistry, Biochemistry and Pharmaceutical Sciences and Oeschger Centre for Climate Change Research, University of Bern, Switzerland and ⁶Exeter college, University of Oxford, United Kingdom

Abstract

We present the ‘SISAL webApp’—a web-based tool to query the Speleothem Isotope Synthesis and AnaLysis (SISAL) database. The software provides an easy-to-use front-end interface to mine data from the SISAL database while providing the SQL code alongside as a learning tool. It allows for simple and increasingly complex querying of the SISAL database based on various data and metadata fields. The SISAL webApp version currently hosts SISALv2 of the database with 691 records from 294 sites, 512 of which have standardized chronologies. The SISAL webApp has sufficient flexibility to host future versions of the SISAL database, which may include allied speleothem information such as trace elements and cave-monitoring records. The SISAL webApp will increase accessibility to the SISAL database while also functioning as a learning tool for more advanced ways of querying paleoclimate databases. The SISAL webApp is available at http://geochem.hu/SISAL_webApp.

Keywords: Speleothems, Stable isotopes, SQL, Database, Online application

(Received 9 February 2023; accepted 17 July 2023)

INTRODUCTION

Natural sedimentary archives are globally distributed and are valuable for local- to continental-scale assessment of past climate and environmental changes because they record the environmental conditions that prevailed during their formation (Williamson et al., 2009; Birks et al., 2014). Speleothems are natural sedimentary archives formed of calcium carbonate and preserved in caves. Speleothems often have excellent age control (Fairchild and Baker, 2012), are widely distributed in terrestrial regions around the world, and provide high-resolution records of past changes in climate and environment that are encoded mainly in carbon and oxygen isotopes (McDermott, 2004; Wong and Breecker, 2015) and trace elements (Fairchild and Treble, 2009). SISAL (Speleothem Isotope Synthesis and AnaLysis) is a working group of the Past Global Changes (PAGES) project with the goal to provide a comprehensive compilation of speleothem records for climate reconstruction and model evaluation (Comas-Bru and Harrison, 2019).

The first version of the SISAL database, SISALv1 (Atsawawaranant et al., 2018), contained 381 speleothem records from 174 cave sites and was supported by publications on ways to

use these data for data-model comparisons (Comas-Bru and Harrison, 2019) and regional interpretations of the isotopic records in the database (Kaushal et al., 2018; Lechleitner et al., 2018; Braun et al., 2019; Burstyn et al., 2019; Deininger et al., 2019; Kern et al., 2019; Oster et al., 2019). The last published version of the database, SISALv2, encompassed 691 speleothem records from 294 sites (Comas-Bru et al., 2020a) and provided additional standardized chronologies that are essential for better age control, which is required for analysis based on multiple speleothem records (Bühler et al., 2022). Regional and/or temporal subsets of the SISALv2 database provided essential data for evaluating environmental response to climate events in time (Kukla et al., 2021; Parker and Harrison, 2022), evolution of climate phenomena in a spatial domain (Parker et al., 2021; Gorenstein et al., 2022), data-model comparisons (Bühler et al., 2021, 2022), improved interpretations of speleothem data (Treble et al., 2022), or improved assessment of the robustness of spectral analysis of unequally spaced sedimentary proxies with chronological uncertainty (Hatvani et al., 2022). A third version of the database, SISALv3, is currently being compiled and will be made available to the public in 2023. The new database will also contain Mg/Ca, Sr/Ca, Ba/Ca, U/Ca, P/Ca, and Sr isotope records (Kaushal et al., 2023). With this new version of the database, it will be possible to explore the global significance of trace-element signatures in speleothems systematically, and to refine climatic interpretations gained from the stable isotope records.

While the SISAL database is clearly useful for the climate community, additional applications and new scientific results can be

*Corresponding author: Nikita Kaushal; Email: nikita.kaushal@exeter.ox.ac.uk

[†]Authors have contributed equally to the work.

Cite this article: Hatvani IG, Kern Z, Tanos P, Wilhelm M, Lechleitner FA, Kaushal N (2024). The SISAL webApp: exploring the speleothem climate and environmental archives of the world. *Quaternary Research* 118, 211–217. <https://doi.org/10.1017/qua.2023.39>

achieved by effectively facilitating broader access to the database (Kaushal et al., 2021). At the moment, the SISAL database is hosted as an SQL file or multiple *.csv files that are linked together by identification numbers. These formats require the use of software such as MySQL, R, Python, or MATLAB to query the database. In addition, there has been a tremendous increase in the number of paleoclimate databases and in their use for research purposes over the last 10 years (Sundqvist et al., 2014; PAGES2k Consortium, 2017; Konecky et al., 2020; Kukla et al., 2021). Because researchers need to learn how to use such databases to address scientific questions at hand, and further how to appropriately query databases to get accurate datasets for analysis, we created the SISAL webApp in order to fill these knowledge gaps.

The paper aims to (1) describe the architecture of the SISAL webApp, (2) provide instructions on the logic by which databases can be mined for required data, and (3) provide accompanying SQL output code the user can build on the basic functionalities of the SISAL webApp by using a tool like MySQL to directly mine the database.

DESIGN

The SISAL webApp's architecture is based on the SISALv2 SQL database (Comas-Bru et al., 2020b). The SISAL Query Server, written in JavaScript using the node.js code library, is responsible for the database operations. The web interface accessible to users is served by the SISAL Query Client, which also is written in JavaScript and relies on the React.js code library. This architecture allows users to query the SISALv2 database using the most popular web browsers supporting ES5 methods (e.g., Google Chrome, Safari, Microsoft Edge, Mozilla Firefox) by specifying a few parameters. The graphical user interface created allows this without the user having to generate SQL code (Fig. 1). The SISALv2 database, the SISAL Query Server, and the SISAL Query Client are hosted on the servers of the Research Centre for Astronomy and Earth Sciences, Eötvös Loránd Research Network, Budapest, Hungary. We plan an update of the SISAL webApp to the SISALv3 database after its release.

FEATURES

The SISAL webApp offers two types of querying. Basic querying, in which the SISALv2 database can be explored based on site name, geographical information, and/or temporal constraints.

Basic querying provides the user with the metadata of the queried speleothem records, their sample data ($\delta^{18}\text{O}$ and $\delta^{13}\text{C}$), and dating information, both the original author-generated chronology and seven SISAL-generated standardized chronologies (*lin_interp_age*, *lin_reg_age*, *Bchron_age*, *Bacon_age*, *OxCal_age*, *copRa_age*, *StalAge_age*) derived by different approaches for age–depth modeling (Amirnezhad-Mozhdehi and Comas-Bru, 2019; Comas-Bru et al., 2019; Rehfeld et al., 2020). The second type of querying supported by the SISAL webApp is the Advanced querying option through which database information can be extracted based on number of available radiometric ages and sample data resolution.

As an additional feature, the SISAL webApp provides SQL codes to help the user get a deeper insight into how the database is queried. It is our intention to make the webApp a steppingstone in the usage of the SISAL database and other databases like it.

Basic querying

Step 1: determining the basic spatiotemporal constraints

Basic querying provides the tool to extract SISAL database information based on the most fundamental filters. After providing an email address (recommended for query logging purposes) the user can choose to query based on the name of the cave site or within spatial (e.g., latitude and longitude limits) and/or within temporal constraints (*interp_age*). At least one of the following “Filter types” must be correctly filled out.

Filter type 1. Site name (*site_name*).

Filter type 2. Latitude and longitude (from–to; default is global coverage values from -90° to 90° and from -180° to 180°). In the first column, the southern and western boundaries should be provided for latitude and longitude, respectively. Latitude in degrees decimal (N: +; S: –) and longitude in degrees decimal (E: +; W: –). Other formats are not accepted, in which case the SISAL webApp will return no results. If asking for global coverage, only the metadata are made available through the SISAL webApp. A secure feature does not allow the user to overload the server, therefore download is limited to 30,000 rows.

Filter type 3. *interp_age* (interpolated age from younger–older) according to the original author-generated age model expressed in years BP, where BP refers to “before present,” (present = AD 1950). For details see Atsawawaranunt et al. (2018, table S9) and the SISAL repository at University of Reading.

This first querying step will return a list of speleothem records, their site and entity metadata fulfilling the query criteria provided.




Figure 1. Architecture of the SISAL webApp.

Email
info@paleoclimate_2ka.hu

Filter type 1 (Site)

site_name

Filter type 2 (Lat-Lon)

Latitude from -90°
-10

Latitude to 90°
10

Longitude from -180°
-50

Longitude to 180°
15

Filter type 3 (interp_age)

Usage is mandatory for advanced querying!

interp_age from (years BP)

interp_age to (years BP)

Included speleothem_type(s)

At least one of the checkboxes below must be checked!

Non-Composite Composite

[Get entity list](#)

Figure 2. Basic querying first step illustrating the query options. Only Filter type 2 is used determining the geographical constraints as Lat: -10° to 10° and Lon: -50° to 15° . Note that temporal constraints (Filter type 3) must be used if one aims to use the advanced query options later.

The sites with the queried entities are shown on a map (OpenStreetMap contributors, 2023); clicking on the markers shows information on *site_name*, *site_id*, *geology*, and *rock_age*.

In addition, the user can choose from querying (1) only non-composite records (tick ‘Non-Composite’ checkbox), (2) only composite records (tick ‘Composite’ check-box), or (3) both non-composite and composite records (tick both ‘Non-Composite’ and ‘Composite’ check-boxes). The default is set to only non-composite records.

Step 2: selection based on the metadata

All records fulfilling the criteria set in Step 1 can subsequently be selected by checking the box next to the *doi* column header. Alternatively, the user can specify a subset of data to be extracted, based on criteria specified in the metadata (e.g., mineralogy; Fig. 3).

The original author-generated chronology is a default output, and the user of the SISAL webApp has to choose at least one SISAL chronology to be extracted for the queried record(s) under the ‘Select chronos’ section (Fig. 3). The alternative age-depth models (SISAL chronologies with corresponding uncertainties) were provided by SISAL (Amirnezhad-Mozhdehi and Comas-Bru, 2019; Rehfeld et al., 2020) for records that are not composites (i.e., time series based on more than one speleothem record) and which are $^{230}\text{Th}/\text{U}$ dated (see Comas-Bru et al., 2020b). All the SISAL chronologies can be selected with the ‘Select all’ checkbox, or in any combination, for example asking only for the *Bacon_age* SISAL chronology (Fig. 3).

Step 3: data extraction

By pressing the blue download buttons at the bottom of the page the user can download the (1) metadata of the selected records (called EntityList.xlsx), (2) their dating information (called

DatingInfo.xlsx), and/or (3) selected chronologies and the sample data (called SampleData.xlsx) in three separate files. In addition, the SQL codes are provided in a worksheet called ‘SQL query’ in each output file to help the user get a deeper insight into how the database is queried.

The extracted sample data are trimmed according to the temporal constraints if applied in ‘Filter type 3 (*interp_age*)’, but the complete dating information table is given for the selected record (s). Note that *interp_age* (original author-generated age model expressed in years BP, where BP refers to “before present” with the present being AD 1950) is provided by default in the fifth column of the output file SampleData.xlsx.

Advanced querying

Advanced querying provides tools to extract SISAL database information based on the number of available radiometric ages and sample data resolution. In descriptive words, this option would be suitable for a query like “extract all data from Asia, covering the last 2,000 years BP, where each record has at least two radiometric age measurements over the 2,000 year period, and the sampling interval for isotopes is less than ten years between successive samples.” This option is available after Basic filtering when a corresponding list of records are received and selected. Note that if no temporal constraint is applied in the Basic querying, the Advanced query will search within full temporal coverage of the selected records.

In advanced querying, two filters can be applied and combined: (1) minimum number of radiometric ages for the chosen record(s) regarding the whole available time interval, or shorter if a filter is applied in the Basic querying part: first step (note that ages excluded by the original authors to develop the age-depth models [i.e., where *date_used* = *no* in the database] are

Filtered metadata

<input type="checkbox"/>	doi	site_name	site_id	entity_id	entity_name	entity_status	corresponding_current	depth_ref	cover_thickness	distance_entranc
<input checked="" type="checkbox"/>	10.1038/ngeo444	Abissal cave	18	79	Abissal	current		from top		
<input type="checkbox"/>	10.1038/ngeo444	Abissal cave	18	80	Ale-1	current		from top		
<input type="checkbox"/>	10.1038/ngeo444	Rainha cave	111	219	RN1	current		from top		
<input checked="" type="checkbox"/>	10.1038/ngeo444	Rainha cave	111	220	RN4	current		from top		

Select chronos

- Select all
 lin_interp_age
 lin_reg_age
 Bchron_age
 Bacon_age
 OxCal_age
 copRa_age
 StalAge_age

[Download meta data](#)
[Download dating information](#)
[Download Sample data \(chosen chronology\)](#)

[Advanced Query Parameters](#)

Figure 3. Basic querying output information and selection of SISAL chronology/chronologies. The example shows the output using the setting shown in Figure 2. The *entity_name* Abissal (*entity_id* = 79) and the *entity_name* RN4 (*entity_id* = 220) speleothem records are selected with the *Bacon_age* SISAL chronology.

not considered); and (2) maximum allowed ‘age gap’ given as number of years in the original chronology (*interp_age*), or in another chosen *sisal_chronology*, considering the whole available time interval, or shorter if a filter is applied in the Basic querying part: first step (i.e., a large age gap means either the sample resolution is coarse and/or the estimated duration of any hiatus in the record exceeds the given maximum allowed ‘age gap’ in the queried interval).

In an example, the Middle Holocene (5500–6500 yr BP) was queried globally (Fig. 4A), which provided 178 records. When the ‘Advanced query filter 1’ was chosen and at least three radiometric ages were required (Fig. 4B) from each entity from within the Middle Holocene, the number of records decreased to 30, with 6065 lines of sample data altogether. This querying took 23 seconds for the server to finish. When the advanced constraints were made stricter with only a maximum of 100 years allowed between consecutive sample dates in *interp_age* (interpolated age based on *original_chronology*) to exclude the coarse resolution records (Fig. 4C), the number of obtained records dropped to 15, with the querying taking 24 seconds. When only ‘Advanced query filter 2’ is used (Fig. 4D) the output is 66 entities with 6332 lines of sample data in 25 seconds.

In all cases the output table (called *advancedRes.xlsx*) consists of six worksheets: (1) *reportInfo* with the SQL code of the selection of the records (Step 2; Fig. 3); (2) *entityAdvFiltered* with the list of records meeting the advanced criteria provided; (3) *datingAdvFiltered* with the corresponding radiometric ages; (4) *chronoAdvFiltered* with the sample data and the

chosen SISAL chronology (in this case *copRa_age*; Fig. 4B–D); (5) *SQLsAdvFiltered* with the SQL code of the advanced querying; and (6) ‘entity,’ ‘dating,’ ‘chrono,’ and ‘SQLs’ worksheets with the list of entities, radiometric ages, sample data, and SQL code, respectively, provided by the basic querying (Fig. 4A).

COMMON ERROR MESSAGES AND THEIR BACKGROUND/SOLUTION

Most common errors are associated with basic querying. If no field is completed, the SISAL webApp will return: “None of the query’s filter parameters are specified correctly! Please specify the *site_name* and/or Lat-Lon coordinates and/or *interp_age* interval and try again!”.

In general, the user should pay attention to using the proper formatting of the spatial constraints and use decimal degree units, otherwise the SISAL webApp will return the message: “The coordinates are incorrect or some are missing! Please revise the coordinates, and try again! Default is global coverage from –90° to 90° and from –180° to 180°.”

Secondly, the temporal constraints should follow the instructions given in this paper and the user manual (http://geochem.hu/SISAL_webApp). If the *interp_age* is provided incorrectly, the following error message is given: “The *interp_age* interval is incorrect or incomplete! Please revise the beginning (younger) and end (older) of the interval, and try again!”.

Advanced querying does not work without the Basic querying part being used and at least one entity selected in Step 2;

Filter type 3 (interp_age)

Usage is mandatory for advanced querying!

A)

interp_age from (years BP)
5500interp_age to (years BP)
6500[Advanced Query Parameters](#)

Advanced query filter 1

B)

At least n radiometric dates in each and every entity (numeric)
3

NOTE: Leave this field blank if you do not want to use this filter!

Advanced query filter 2

Max gap in the chosen chronology (numeric)

copRa_age ▶

NOTE: Leave this field blank if you do not want to use this filter!

Download advanced result

Advanced query filter 1

C)

At least n radiometric dates in each and every entity (numeric)
3

NOTE: Leave this field blank if you do not want to use this filter!

Advanced query filter 2

Max gap in the chosen chronology (numeric)
100

copRa_age ▶

NOTE: Leave this field blank if you do not want to use this filter!

Download advanced result

Advanced query filter 1

D)

At least n radiometric dates in each and every entity (numeric)

NOTE: Leave this field blank if you do not want to use this filter!

Advanced query filter 2

Max gap in the chosen chronology (numeric)
100

copRa_age ▶

NOTE: Leave this field blank if you do not want to use this filter!

Download advanced result

Figure 4. Advanced querying examples. Temporal constraints are applied in the basic querying (A). Advanced query filter 1 (B), filter 2 (C) and both advanced query filters (D) are applied with the *copRa_age* SISAL chronology chosen to accompany the sample data in the advanced query output.

otherwise, the following error message is returned: “Download request denied! Please select at least one entity!”.

Please note, in case of large output tables (maximum 30,000 rows), querying may take up to minutes.

SOFTWARE AVAILABILITY, FUTURE UPDATES, AND TERMS OF USE

The SISAL webApp is available at the Research Centre for Astronomy and Earth Sciences (http://geochem.hu/SISAL_webApp), and has been tested to work in all major

browsers (e.g., Google Chrome, Safari, Microsoft Edge, Mozilla Firefox). SISAL is continuing to expand the global database by including new records and extended sets of data and metadata information. The SISAL webApp is intended to be able to perform more advanced querying on the most updated version of the SISAL database.

Planned updates include (1) the option to allow the user to pick the annual laminated samples; (2) replacing the current map output with an interactive alternative, that would allow the selection of the area of interest, while showing the cave sites and records; and (3) making the SISAL webApp capable of

providing information on the mean sample resolution and mean chronological uncertainty, providing input data for immediate verification of inherent constraints for spectral analyses (e.g., CUSP (<https://geochem.hu/CUSP/>); Hatvani et al., 2022) of the considered speleothem record(s).

The PAGES-SISAL project is a community-lead effort. This dataset is licensed by the rights-holder(s) under a Creative Commons Attribution 4.0 International License: <https://creativecommons.org/licenses/by/4.0/>. In order to assure traceability, any presentation, report, or publication that uses the SISALv2 database should cite the dataset (<https://doi.org/10.17864/1947.256>) along with the following publications: Atsawaranunt et al. (2018), Comas-Bru et al. (2019), and Comas-Bru et al. (2020a). If the SISAL webApp is used for data extraction from the SISAL database, it is required to cite the webApp itself (http://geochem.hu/SISAL_webApp) and this paper (Hatvani et al., 2023; doi: 10.1017/qua.2023.39).

If using individual sites, the literature citations for published work provided in the database also should be cited. Contact information of data contributors of unpublished data is also provided, and these data contributors should be contacted when unpublished records are used on an individual basis. In addition, users are advised to verify which queried reference reports the particular record downloaded, because for example, oxygen and carbon stable isotope records from the same stalagmite may be published in different papers.

Acknowledgments. This study was undertaken by SISAL (Speleothem Isotopes Synthesis and Analysis), a working group of the Past Global Changes (PAGES) project, which in turn receives support from the Swiss Academy of Sciences and the Chinese Academy of Sciences. The PAGES Data Stewardship Scholarship (DSS_108) also supported this work. We thank SISAL members who contributed their published data to the database and provided additional information when necessary. Authors also thank the IT team of the RCAES, namely Evelin Bányai for installing the app on servers. We are also grateful to the SISAL Phase 2 Steering Committee for help in debugging the app. This is contribution No. 85 of the 2ka Paleoclimatology Research Group. The first author dedicates this paper to the memory of his beloved grandmother, Marta Chiovini (1928–2023).

Author contributions. I.G.H conceived the idea and designed the app with P.T. P.T. developed the app with input from Z.K. and I.G.H. I.G.H. produced the figures. I.G.H. and Z.K. wrote the paper, with contributions from N.K., T.P., F.L., and M.W. All authors have read and agreed to the published version of the manuscript and helped in the debugging and final development of the app.

REFERENCES

- Amirnezhad-Mozhdehi, S., Comas-Bru, L., 2019. MATLAB scripts to produce OxCal chronologies for SISAL database (scripts V1) (1.0). *Zenodo*. <https://doi.org/10.5281/zenodo.3586280>.
- Atsawaranunt, K., Comas-Bru, L., Amirnezhad Mozhdehi, S., Deininger, M., Harrison, S.P., Baker, A., Boyd, M., et al., 2018. The SISAL database: a global resource to document oxygen and carbon isotope records from speleothems. *Earth System Science Data* **10**, 1687–1713.
- Birks, J., Battarbee, R., Mackay, A., Oldfield, F. (Eds.), 2014. *Global Change in the Holocene*. Routledge, New York.
- Braun, K., Nehme, C., Pickering, R., Rogerson, M., Scroxton, N., 2019. A window into Africa's past hydroclimates: the SISAL_v1 database contribution. *Quaternary* **2**, 4. <https://doi.org/10.3390/quat2010004>.
- Bühler, J.C., Axelsson, J., Lechleitner, F.A., Fohlmeister, J., LeGrande, A.N., Midhun, M., Sjolte, J., Werner, M., Yoshimura, K., Rehfeld, K., 2022. Investigating stable oxygen and carbon isotopic variability in speleothem records over the last millennium using multiple isotope-enabled climate models. *Climate of the Past* **18**, 1625–1654.
- Bühler, J.C., Roesch, C., Kirschner, M., Sime, L., Holloway, M.D., Rehfeld, K., 2021. Comparison of the oxygen isotope signatures in speleothem records and iHadCM3 model simulations for the last millennium. *Climate of the Past* **17**, 985–1004.
- Burstyn, Y., Martrat, B., Lopez, J.F., Iriarte, E., Jacobson, M.J., Lone, M.A., Deininger, M., 2019. Speleothems from the Middle East: an example of water limited environments in the SISAL database. *Quaternary* **2**, 16. <https://doi.org/10.3390/quat2020016>.
- Comas-Bru, L., Atsawaranunt, K., Harrison, S., SISAL Working Group Members, 2020b. SISAL (Speleothem Isotopes Synthesis and Analysis Working Group) database version 2.0. University of Reading, Dataset. <http://dx.doi.org/10.17864/1947.256>.
- Comas-Bru, L., Harrison, S.P., 2019. SISAL: Bringing added value to speleothem research. *Quaternary* **2**, 7. <https://doi.org/10.3390/quat2010007>.
- Comas-Bru, L., Harrison, S.P., Werner, M., Rehfeld, K., Scroxton, N., Veiga-Pires, C., SISAL Working Group Members, 2019. Evaluating model outputs using integrated global speleothem records of climate change since the last glacial. *Climate of the Past* **15**, 1557–1579.
- Comas-Bru, L., Rehfeld, K., Roesch, C., Amirnezhad-Mozhdehi, S., Harrison, S.P., Atsawaranunt, K., Ahmad, S.M., et al., 2020a. SISALv2: A comprehensive speleothem isotope database with multiple age-depth models. *Earth System Science Data* **12**, 2579–2606.
- Deininger, M., Ward, B.M., Novello, V.F., Cruz, F.W., 2019. Late Quaternary variations in the South American Monsoon System as inferred by speleothems—new perspectives using the SISAL Database. *Quaternary* **2**, 6. <https://doi.org/10.3390/quat2010006>.
- Fairchild, I.J., Baker, A., 2012. Geochemistry of speleothems. In: Fairchild, I.J., Baker, A., *Speleothem Science*. John Wiley & Sons, Ltd, pp. 245–289.
- Fairchild, I.J., Treble, P.C., 2009. Trace elements in speleothems as recorders of environmental change. *Quaternary Science Reviews* **28**, 449–468.
- Gorenstein, I., Prado, L.F., Bianchini, P.R., Wainer, I., Griffiths, M.L., Pausata, F.S.R., Yokoyama, E., 2022. A fully calibrated and updated mid-Holocene climate reconstruction for eastern South America. *Quaternary Science Reviews* **292**, 107646. <https://doi.org/10.1016/j.quascirev.2022.107646>.
- Hatvani, I.G., Tanos, P., Mudelsee, M., Kern, Z., 2022. Robust periodic signals in proxy records with chronological uncertainty and variable temporal resolution. *Quaternary Science Reviews* **276**, 107294. <https://doi.org/10.1016/j.quascirev.2021.107294>.
- Kaushal, N., Breitenbach, S.F.M., Lechleitner, F.A., Sinha, A., Tewari, V.C., Ahmad, S.M., Berkelhammer, M., et al., 2018. The Indian Summer Monsoon from a speleothem $\delta^{18}\text{O}$ perspective—a review. *Quaternary* **1**, 29. <https://doi.org/10.3390/quat1030029>.
- Kaushal, N., Comas-Bru, L., Lechleitner, F., Hatvani, I.G., Kern, Z., 2021. Improving access to paleoclimate data. *Eos* **102**, 5. <https://doi.org/10.1029/2021EO155315>.
- Kaushal, N., Wilhelm, M., Lechleitner, F., Braun, K., Rehfeld, K., Gabor Hatvani, I., Tanos, P., et al., 2023. Update to the SISAL speleothem database—links to monitoring data, additional palaeoenvironmental proxies and enhanced accessibility. *EGU General Assembly 2023, Vienna, Austria*, 24–28 Apr 2023, EGU23-688. <https://doi.org/10.5194/egusphere-egu23-688>.
- Kern, Z., Demény, A., Perşoiu, A., Hatvani, I.G., 2019. Speleothem records from the eastern part of Europe and Turkey—discussion on stable oxygen and carbon isotopes. *Quaternary* **2**, 31. <https://doi.org/10.3390/quat2030031>.
- Konecky, B.L., McKay, N.P., Churakova, O.V., Comas-Bru, L., Dassié, E.P., DeLong, K.L., Falster, G.M., et al., 2020. The Iso2k database: a global compilation of paleo- $\delta^{18}\text{O}$ and $\delta^2\text{H}$ records to aid understanding of Common Era climate. *Earth System Science Data* **12**, 2261–2288.
- Kukla, T., Ahlström, A., Maezumi, S.Y., Chevalier, M., Lu, Z., Winnick, M.J., Chamberlain, C.P., 2021. The resilience of Amazon tree cover to past and present drying. *Global and Planetary Change* **202**, 103520. <https://doi.org/10.1016/j.gloplacha.2021.103520>.
- Lechleitner, F.A., Amirnezhad-Mozhdehi, S., Columbu, A., Comas-Bru, L., Labuhn, I., Pérez-Mejías, C., Rehfeld, K., 2018. The potential of speleothems from Western Europe as recorders of regional climate: a critical assessment of the SISAL Database. *Quaternary* **1**, 30. <https://doi.org/10.3390/quat1030030>.

- McDermott, F., 2004. Palaeo-climate reconstruction from stable isotope variations in speleothems: a review. *Quaternary Science Reviews* **23**, 901–918.
- OpenStreetMap contributors, 2023. <https://planet.openstreetmap.org> [20 May 2023].
- Oster, J.L., Warken, S.F., Sekhon, N., Arienzo, M.M., Lachniet, M., 2019. Speleothem paleoclimatology for the Caribbean, Central America, and North America. *Quaternary* **2**, 5. <https://doi.org/10.3390/quat2010005>.
- PAGES2k Consortium, 2017. A global multiproxy database for temperature reconstructions of the Common Era. *Scientific Data* **4**, 170088. <https://doi.org/10.1038/sdata.2017.88>.
- Parker, S.E., Harrison, S.P., 2022. The timing, duration and magnitude of the 8.2 ka event in global speleothem records. *Scientific Reports* **12**, 10542. <https://doi.org/10.1038/s41598-022-14684-y>.
- Parker, S.E., Harrison, S.P., Comas-Bru, L., Kaushal, N., LeGrande, A.N., Werner, M., 2021. A data–model approach to interpreting speleothem oxygen isotope records from monsoon regions. *Climate of the Past* **17**, 1119–1138.
- Rehfeld, K., Roesch, C., Comas-Bru, L., Amirnezhad-Mozhdehi, S., 2020. Age-depth model ensembles for SISAL v2 speleothem records, (1.0) [Data set]. *Zenodo*. <https://doi.org/10.5281/zenodo.3816804>.
- Sundqvist, H.S., Kaufman, D.S., McKay, N.P., Balascio, N.L., Briner, J.P., Cwynar, L.C., Sejrup, H.P., *et al.*, 2014. Arctic Holocene proxy climate database—new approaches to assessing geochronological accuracy and encoding climate variables. *Climate of the Past* **10**, 1605–1631.
- Treble, P.C., Baker, A., Abram, N.J., Hellstrom, J.C., Crawford, J., Gagan, M.K., Borsato, A., *et al.*, 2022. Ubiquitous karst hydrological control on speleothem oxygen isotope variability in a global study. *Communications Earth & Environment* **3**, 29. <https://doi.org/10.1038/s43247-022-00347-3>.
- Williamson, C.E., Saros, J.E., Vincent, W.F., Smol, J.P., 2009. Lakes and reservoirs as sentinels, integrators, and regulators of climate change. *Limnology and Oceanography* **54**, 2273–2282.
- Wong, C.I., Brecker, D.O., 2015. Advancements in the use of speleothems as climate archives. *Quaternary Science Reviews* **127**, 1–18.