

Invariant Keisler measures for ω -categorical structures

Paolo Marimon

May 16, 2024

Abstract

A recent article of Chernikov, Hrushovski, Kruckman, Krupinski, Moconja, Pillay and Ramsey finds the first examples of simple structures with formulas which do not fork over the empty set but are universally measure zero. In this article we give the first known simple ω -categorical counterexamples. These happen to be various ω -categorical Hrushovski constructions. Using a probabilistic independence theorem from Jahel and Tsankov, we show how simple ω -categorical structures where a formula forks over \emptyset if and only if it is universally measure zero must satisfy a stronger version of the independence theorem.

1 Introduction

Keisler measures yield a natural notion of smallness for a definable subset of a structure: a subset X of M^x defined by $\phi(x, a)$ is **universally measure zero** if it is assigned measure zero by every invariant Keisler measure on M . A more classical model theoretic notion of smallness is forking: dividing captures the idea that a small subset of a model can be "moved enough" by automorphisms so as to not overlap with itself. We are interested in comparing these notions of smallness in a sufficiently saturated and strongly homogeneous model.

In stable theories, a definable set is universally measure zero if and only if it forks over \emptyset [3]. This is also the case in ω -categorical NIP theories [2]. Until recently, it was unknown whether the two notions coincided in simple theories. The first counterexample, showing that in simple theories there can be non-forking formulas which are universally measure zero is given in [3]. Using the same technique, the authors also give examples of simple groups which are not definably amenable and prove some positive results in the context of small theories. Neither of the counterexamples given is ω -categorical. The theory of the first counterexample contains as a reduct the theory of the free action of the free group F_5 on an infinite set, which has infinitely many 1-types over a single parameter. Meanwhile, on the definably amenable group case, all groups definable in an ω -categorical simple theory are definably amenable [3, Corollary 4.14]. Indeed, ω -categorical supersimple groups are actually amenable being finite-by-abelian-by-finite [10].

It is natural to ask whether adding the assumption of ω -categoricity we can prove that being universally measure zero is the same as forking. Firstly, the known counterexample makes heavy use of a construction which is inherently not ω -categorical. Secondly, ω -categoricity implies that any invariant Keisler measure is also definable, i.e. the set of parameters a for which $\mu(\phi(x, a)) = \alpha$ is \emptyset -definable. In general, this is a substantially stronger assumption than invariance.

Another motivation for an ω -categorical counterexample comes from the study of ω -categorical MS -measurable structures [16]. Until recently [8, 18], it was an open question whether all supersimple ω -categorical structures of finite SU -rank are MS -measurable. Indeed, this question of Elwes and Macpherson [7] was the initial motivation for our study

of Keisler measures in the context of ω -categorical Hrushovski constructions. An MS -measurable structure [16] has a dimension-measure function which is definable, finite and such that the dimension and measures satisfy Fubini's theorem. As we shall see in Lemma 4.8, forking and being universally measure zero yield the same notion of smallness in an MS -measurable context.

In this article, we show how for various classes of supersimple ω -categorical Hrushovski constructions we have formulas that do not fork over \emptyset but are universally measure zero:

Theorem 5.4. *There are ω -categorical supersimple theories T of finite SU -rank with a formula $\phi(x, a)$ which does not fork over the empty set, but which is universally measure zero.*

More generally, in Theorem 4.4 we show that if forking and being universally measure zero agree in a simple ω -categorical structure, then it must satisfy a stronger version of the independence theorem. It is easy to build ω -categorical Hrushovski constructions for which this fails. These structures are extremely amenable in the sense of [12], which implies the existence of invariant types, and so invariant Keisler measures, in each variable.

We begin with Section 2, where we introduce ergodic measures and Keisler measures. Ergodic measures simplify our study since any measure can be considered as an "integral average" of them. Meanwhile, Keisler measures are the natural notion of a measure on a first-order structure. In Section 3, we study the L^2 -spaces associated with an invariant Keisler measure. From some results of Jahel and Tsankov [13, Theorem 3.2, Corollary 3.5], we know that in ω -categorical structures, ergodic measures are better behaved and a weak form of algebraic independence implies a form of probabilistic independence in the measure (Corollary 3.8). In Section 4, we show how in simple ω -categorical structures with forking and being universally measure zero agreeing we have a stronger version of the independence theorem (Theorem 4.4). We conclude the section with some implications for MS -measurable structures. Finally, in Section 5, we give the example of an ω -categorical structure, supersimple of finite SU -rank with a formula which does not fork over the empty set but which is universally measure zero.

This article requires some knowledge in model theory. Chapters 1-4 of Tent & Ziegler's book [22] should be sufficient, together with Ch.16 of [20] for understanding imaginaries and weak elimination of imaginaries. On simple theories, Chapters 2 and 3 of [14] cover the relevant definitions and results, including the definition of SU -rank and the independence theorem. We also require some basic knowledge of Hilbert spaces and L^2 -spaces, such as Chapter 1 of [5]. In Section 2 we give a self-contained introduction to ergodic measures and Keisler measures. The reader interested in MS -measurable structures should consult [16] or [7]. This article does not require knowledge of Hrushovski constructions since all of the properties that we use are listed in Theorem 5.1. Further details about ω -categorical Hrushovski constructions can be found in [24, §6.2.1], while [18], especially in the appendix, provides the details on the specific construction we give as an example in Theorem 5.4.

To conclude, we provide some notation and conventions. Firstly, we work with a complete countable first-order \mathcal{L} -theory T . From Section 3, by \mathcal{M} we denote the countable model of an ω -categorical theory. We write M for when we consider \mathcal{M} as a set. We use greek letters ϕ, ψ, χ, \dots to refer to formulas. The letters x, y, z, \dots indicate variables, and may also indicate a finite tuple of variables. Similarly, the lowercase letters a, b, c, \dots indicate parameters from M , and may also indicate a finite tuple. Meanwhile, we indicate subsets of M by the uppercase A, B, C, \dots . For a, a' tuples from \mathcal{M} and $A \subseteq M$, we write $a \equiv_A a'$ to say that a and a' have the same type over A .

Acknowledgements: This article was written during my PhD at Imperial College London under the supervision of David Evans and Charlotte Kestner, both of whom I thank for their support and help. Nicholas Ramsey gave an excellent talk at Imperial about [3],

which motivated me to look at whether ω -categorical Hrushovski constructions may also have been a counterexample. I am also grateful to my friend Matteo Tabaro for helpful discussions during my learning of ergodic theory. Finally, some of the ideas behind this article are greatly indebted to Ehud Hrushovski and his unpublished article [11], which he kindly shared with me.

Funding: the author's PhD was supported by an Admin Roth Scholarship funded by the Department of Mathematics of Imperial College London. This research is also funded by the European Union (ERC, POCOCOP, 101071674). Views and opinions expressed are however those of the author only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

2 Keisler measures and ergodic measures

We begin with a brief and self-contained introduction to ergodic measures and Keisler measures. Firstly, we introduce ergodic measures in a general context. Then, we explain their utility for the study of Keisler measures.

2.1 Ergodic measures

Ergodic measures are an essential tool in our paper. They are better behaved than invariant measures and any invariant measure can be decomposed as an integral average of ergodic measures. Here we briefly introduce these measures and mention some basic results about them. Chapter 12 of Phelps' book [19] covers the theory we discuss at the adequate level of generality for our purposes.

We work in the context of a topological group G acting on a topological space X via a continuous action $\cdot : G \times X \rightarrow X$. When X is compact and Hausdorff, we call this action a **G-flow**. Let $\mathcal{B}(X)$ be the set of Borel subsets of X , and let $\mu : \mathcal{B}(X) \rightarrow [0, 1]$ be a Borel probability measure. We say that μ is **G-invariant** if for any $\tau \in G$ and any $A \in \mathcal{B}(X)$, we have that $\tau \cdot A \in \mathcal{B}(X)$ and $\mu(\tau \cdot A) = \mu(A)$.

Definition 2.1. We say that a G -invariant Borel probability measure μ is **G-ergodic** if for all $A \in \mathcal{B}(X)$, we have that if for all $\tau \in G$,

$$\mu(A \Delta \tau \cdot A) = 0,$$

then either $\mu(A) = 0$ or $\mu(A) = 1$.

An alternative definition tells us that any invariant function is constant [6, Prop.2.14]:

Proposition 2.2. Let (X, \mathcal{B}, μ) be a probability space and G be a group acting on X such that μ is G -invariant. Then, the following are equivalent:

1. the measure μ is G -ergodic; and
2. any measurable function $f : X \rightarrow \mathbb{C}$, which is G -invariant almost everywhere (i.e. for any $\tau \in G$, $f \circ \tau = f$ a.e.) is constant almost everywhere.

We are interested in studying G -ergodic measures since, when X is metrizable, they yield an ergodic decomposition of any G -invariant measure. Hence, their study is essential to the understanding of the G -invariant measures on X . Below, for $f \in C(X, \mathbb{C})$, we write $\mu(f) = \int_X f d\mu$. From [19, p.77] we have:

Theorem 2.3 (Ergodic decomposition). Let X be a compact metrizable space with a group G acting continuously on it. Let μ be a G -invariant Borel probability measure on X . Then, the space $\mathfrak{M}(X)$ of G -invariant Borel probability measures on X is also metrizable and the set of G -ergodic

measures $\text{Erg}(X)$ is a Borel subset of $\mathfrak{M}(X)$. Furthermore, there is a unique Borel probability measure \mathfrak{m} on $\mathfrak{M}(X)$ such that for any $f \in C(X, \mathbb{C})$,

$$\mu(f) = \int_{\text{Erg}(X)} v(f) \, d\mathfrak{m}(v).$$

2.2 Keisler measures

Keisler measures are finitely additive probability measures on the space of definable subsets of a structure. Chapter 7 of Pierre Simon's book [21] is a good introduction to the subject. We give a brief self-contained discussion and note the importance of ergodic measures in the study of invariant Keisler measures.

Let T be a complete \mathcal{L} -theory, where \mathcal{L} is a countable first-order language and $\mathcal{M} \models T$. Let $\text{Def}_x(M)$ denote the Boolean algebra of $\mathcal{L}(M)$ -formulas in the free variable x up to $\text{Th}(\mathcal{M}_M)$ -equivalence, where \mathcal{M}_M is the expansion of \mathcal{M} by constant symbols for each element of M . Let $S_x(M)$ be the Stone space of types over M in the variable x with the usual topology. We write $[\phi(x, a)]$ for the clopen set of types containing the formula $\phi(x, a)$. This space is always compact and Hausdorff. Moreover, when M is countable, it is also metrizable. This can be seen by taking an enumeration of the $\mathcal{L}(M)$ -formulas ϕ_1, ϕ_2, \dots and then for $p_1, p_2 \in S_x(M)$, letting $d(p_1, p_2) = \frac{1}{2^n}$, where n is the smallest natural number such that ϕ_n is not contained in both p_1 and p_2 . The automorphism group $\text{Aut}(M)$ naturally acts on $\text{Def}_x(M)$ and on $S_x(M)$, where for $\tau \in \text{Aut}(M)$, $\tau \cdot \phi(x, a) = \phi(x, \tau(a))$ and for $p \in S_x(M)$,

$$\tau \cdot p = \{\phi(x, \tau(a)) : \phi(x, a) \in p\}.$$

Definition 2.4. Let \mathcal{M} be an \mathcal{L} -structure. We say that $\mu : \text{Def}_x(M) \rightarrow [0, 1]$ is a **Keisler measure** if it is a finitely additive measure such that $\mu(x = x) = 1$. We say that μ is **Aut(M)-invariant** if for any $\tau \in \text{Aut}(M)$,

$$\mu(\tau \cdot \phi(x, a)) = \mu(\phi(x, a)).$$

For an infinite cardinal κ , we say that \mathcal{M} is **strongly κ -homogeneous** if whenever a and a' are tuples of cardinality $< \kappa$ such that $a \equiv a'$, there is $\tau \in \text{Aut}(M)$ such that $\tau(a) = a'$. If \mathcal{M} is strongly ω -homogeneous, for a Keisler measure to be $\text{Aut}(M)$ -invariant is equivalent to saying that, if $a \equiv a'$, then $\mu(\phi(x, a)) = \mu(\phi(x, a'))$.

As noted in the introduction, Keisler measures give us a natural notion of smallness for a definable set.

Definition 2.5. Let \mathcal{M} be ω -saturated and strongly ω -homogeneous. We say that $\phi(x, b) \in \text{Def}_x(M)$ is **universally measure zero** if $\mu(\phi(x, b)) = 0$ for every $\text{Aut}(M)$ -invariant Keisler measure. Similarly, let $A \subseteq M$ and let \mathcal{M} be κ -saturated and strongly κ -homogeneous for $\kappa = \max\{|A|^+, \omega\}$. We say that $\phi(x, b) \in \text{Def}_x(M)$ is **universally measure zero over A** if it is assigned measure zero by every $\text{Aut}(M/A)$ -invariant Keisler measure. We call $\mathcal{O}(A)$ the set of formulas in $\text{Def}_x(M)$ which are universally measure zero over A .

The set $\mathcal{O}_x(A) = \mathcal{O}(A) \cap \text{Def}_x(M)$ forms an ideal in the Boolean algebra of $\text{Def}_x(M)$ [3]. Similarly, $F_x(A)$, the set of formulas in the variable x with parameters from M forking over A , also forms an ideal. Our main result is that there are ω -categorical supersimple theories for which $\mathcal{O}(\emptyset)$ strictly contains the set of formulas forking over \emptyset , $F(\emptyset)$.

Remark 2.6. In other articles [3], being universally measure zero is defined in the monster model for the theory. In this article, since we work over finite sets, no generality is lost by focusing on an ω -saturated strongly ω -homogeneous model. In fact, if $F(\emptyset) = \mathcal{O}(\emptyset)$ in an ω -saturated strongly ω -homogeneous model of T , this is the case in all such models (cf. [12, Proposition 2.3]). In particular, we may focus on invariant Keisler measures on the countable model of the ω -categorical theory we are studying. More generally, whenever we compare the sizes of $F(A)$ and $\mathcal{O}(A)$ in this article we always work over a sufficiently saturated and strongly homogeneous model.

A Keisler measure μ can be extended uniquely to a regular σ -additive Borel probability measure $\mu : \mathcal{B}_x(M) \rightarrow [0, 1]$, where $\mathcal{B}_x(M)$ is the set of Borel subsets of $S_x(M)$ [21, §7.1]. Conversely, any regular Borel probability measure on $S_x(M)$ induces a Keisler measure by considering its restriction to clopen sets. This correspondence still holds between $\text{Aut}(M)$ -invariant Keisler measures in the variable x and $\text{Aut}(M)$ -invariant regular Borel probability measures on $S_x(M)$. From here, we shall speak interchangeably of the two.

Remark 2.7. An $\text{Aut}(M^{eq})$ -invariant Keisler measure μ^{eq} on \mathcal{M}^{eq} in the real variable x is entirely determined by its restriction to its induced $\text{Aut}(M)$ -invariant Keisler measure μ on \mathcal{M} . Furthermore, μ^{eq} is ergodic if and only if μ is.

If we are interested in studying the universally measure zero formulas for a countable structure, it is helpful to study $\text{Aut}(M)$ -ergodic measures on $S_x(M)$. From the ergodic decomposition 2.3, we get:

Corollary 2.8. *Let \mathcal{M} be a countable structure and let μ be an $\text{Aut}(M)$ -invariant Borel probability measure on $S_x(M)$. Let $\mathfrak{M}_x(M)$ be the space of $\text{Aut}(M)$ -invariant Borel probability measures on $S_x(M)$. Then, there is a unique Borel probability measure \mathfrak{m} on $\mathfrak{M}_x(M)$ such that for any $\mathcal{L}(M)$ -formula $\phi(x, a)$,*

$$\mu([\phi(x, a)]) = \int_{\text{Erg}_x(M)} \nu([\phi(x, a)]) d\mathfrak{m}(\nu),$$

where $\text{Erg}_x(M)$ is the space of $\text{Aut}(M)$ -ergodic measures.

Remark 2.9. When M is countable, every Borel measure μ on $S_x(M)$ is regular [1, Theorem 7.1.7]. From [9, Corollary 1.2], we actually know that when \mathcal{M} is the countable model of an ω -categorical theory, $\text{Erg}_x(M)$ is closed in the space $\mathfrak{M}_x(M)$.

For conciseness of notation, in subsequent sections we shall generally refer to μ as an ergodic measure on \mathcal{M}^{eq} (in the variable x). By this we mean that μ is a Borel probability measure on $S_x(M^{eq})$ invariant under the action of $\text{Aut}(M^{eq})$, which is also $\text{Aut}(M^{eq})$ -ergodic. As noted above, when x is a variable in the real sort, there is a one-to-one correspondence between these ergodic measures and the ergodic $\text{Aut}(M)$ -invariant Borel probability measures on $S_x(M)$. More generally, given the various correspondences explained in this section we can safely use μ to denote both the Keisler measure and the corresponding Borel probability measures on $S_x(M)$ and $S_x(M^{eq})$.

3 Weak algebraic independence and probabilistic independence

Given the Borel probability space $(S_x(M^{eq}), \mathcal{B}_x(M^{eq}), \mu)$ induced by an invariant Keisler measure on M in the variable x , we wish to study the associated Hilbert space $L^2(\mu)$. In this section we introduce some of the relevant tools for this, following the discussion of [13] and [23]. Furthermore, we shall show how various results of [13] yield that a weak form of algebraic independence between parameters implies a form of probabilistic independence in an ergodic measure. This result is Corollary 3.8.

Let \mathcal{M} be a countable ω -categorical structure. We have that $\text{Aut}(M)$ is a Polish group, that is, a separable completely metrizable topological group. In particular, in the topology of $G = \text{Aut}(M)$, the pointwise stabilizers of finite sets $G_A = \text{Aut}(M/A)$ for $A \subset M$ finite, are neighbourhoods of the identity and the set of cosets of these pointwise stabilizers form a basis of open sets for the topology [15, §4.1].

Definition 3.1. Let \mathcal{H} be a complex Hilbert space and $U(\mathcal{H})$ be its unitary group. A **unitary representation** of a topological group G is a continuous action of G on \mathcal{H} by unitary operators. Equivalently, we may say that it is a homomorphism $\pi : G \rightarrow U(\mathcal{H})$ such that for each $f \in \mathcal{H}$ the map $\tau \mapsto \pi(\tau) \cdot f$ is continuous.

As noted in the previous section, a Keisler measure μ on \mathcal{M} in the variable x induces a Borel probability space $(S_x(M^{eq}), \mathcal{B}_x(M^{eq}), \mu)$. We are interested in studying the complex L^2 -space $L^2(S_x(M^{eq}), \mathcal{B}_x(M^{eq}), \mu)$, which we abbreviate $L^2(\mu)$.

The action of $\text{Aut}(M)$ on M naturally induces an action $\lambda : \text{Aut}(M) \times L^2(\mu) \rightarrow L^2(\mu)$, where for $\sigma \in \text{Aut}(M), f \in L^2(\mu)$,

$$\lambda(\sigma, f)(p) = f(\sigma^{-1}(p)) \text{ for all } p \in S_x(M).$$

This action is well defined and preserves integrals. For each $\sigma \in \text{Aut}(M)$, the map $\Lambda_\sigma : L^2(\mu) \rightarrow L^2(\mu)$ given by $f \mapsto \lambda(\sigma, f)$ is a unitary operator. Moreover, the map $\pi : \text{Aut}(M) \rightarrow U(L^2(\mu))$ from the group of automorphisms of M to the group of unitary operators of $L^2(\mu)$ given by $\sigma \mapsto \Lambda_\sigma$ is a unitary representation (cf. §3 of [23]).

Let \mathcal{H} be a complex Hilbert space and the action of $G = \text{Aut}(M)$ on \mathcal{H} be a **unitary representation**. For $A \subseteq M^{eq}$ we write, following [13],

$$\mathcal{H}_A = \overline{\{f \in \mathcal{H} \mid G_{A'} \cdot f = f \text{ for some finite } A' \subseteq A\}},$$

where for $S \subset \mathcal{H}$, \bar{S} is its closure.

Definition 3.2. Let \mathcal{H} be a Hilbert space and $\mathcal{K} \subseteq \mathcal{H}$ be a subspace. We denote the orthogonal complement of \mathcal{K} in \mathcal{H} (often denoted as \mathcal{K}^\perp) as $\mathcal{H} \ominus \mathcal{K}$. When \mathcal{K} is a closed subspace, we have that $\mathcal{H} = \mathcal{K} \oplus (\mathcal{H} \ominus \mathcal{K})$. Given subspaces $\mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2 \subseteq \mathcal{H}$, and $\mathcal{H}_0 \subseteq \mathcal{H}_i$ for $i \in \{1, 2\}$ we say that \mathcal{H}_1 and \mathcal{H}_2 are **orthogonal** over \mathcal{H}_0 and write $\mathcal{H}_1 \perp_{\mathcal{H}_0} \mathcal{H}_2$ if $(\mathcal{H}_1 \ominus \mathcal{H}_0) \perp (\mathcal{H}_2 \ominus \mathcal{H}_0)$.

Note that when the \mathcal{H}_i are all closed subspaces for $i \in \{1, 2, 3\}$ with $\mathcal{H}_1 \perp_{\mathcal{H}_0} \mathcal{H}_2$, we have that for $f \in \mathcal{H}_1$ we can decompose f into $f = f_1 + f_0$, where $f_1 \in \mathcal{H}_1 \ominus \mathcal{H}_0$ and $f_0 \in \mathcal{H}_0$. Similarly, for $g \in \mathcal{H}_2$ we can decompose it as $g = g_2 + g_0$. Now, by the orthogonalities $(\mathcal{H}_i \ominus \mathcal{H}_0) \perp \mathcal{H}_0$ for $i \in \{1, 2\}$ and $(\mathcal{H}_1 \ominus \mathcal{H}_0) \perp (\mathcal{H}_2 \ominus \mathcal{H}_0)$, we have that

$$\langle f, g \rangle = \langle f_0, g_0 \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the inner product on \mathcal{H} .

In an ω -categorical structure, we have the following theorem of Jahel and Tsankov, which translates weak algebraic independence into orthogonality of the associated Hilbert spaces:

Theorem 3.3. [13, Theorem 3.2] *Let M be ω -categorical and $G = \text{Aut}(M)$. Let $A, B \subseteq M^{eq}$ be algebraically closed with respect to acl^{eq} . Then, $\mathcal{H}_A \perp_{\mathcal{H}_{A \cap B}} \mathcal{H}_B$.*

Remark 3.4. Recently, [4] develops some results of this kind outside the context of ω -categorical structures.

Note that the space of constant functions is closed in $L^2(\mu)$, hence, from Proposition 2.2 we get:

Corollary 3.5. *Let μ be an ergodic measure on \mathcal{M}^{eq} in the variable x . Then, $L^2(\mu)_{\text{dcl}^{eq}(\emptyset)}$ is generated by the constant indicator function $\mathbb{1}$.*

Proof. Suppose $f \in L^2(\mu)_{\text{dcl}^{eq}(\emptyset)}$. Then, for some finite $C \subseteq \text{dcl}^{eq}(\emptyset)$, f is G_C -invariant. However, since any element in $\text{dcl}^{eq}(\emptyset)$ is fixed by $\text{Aut}(M^{eq})$ -automorphisms, $G_C = G_\emptyset$. This means that f is invariant almost everywhere, and therefore constant by ergodicity of the measure. \square

An ergodic measure also concentrates on an orbit. So, for ergodic μ , in $L^2(\mu)$ the constant function $\mathbb{1}$ will be in the same equivalence class as $\mathbb{1}_\phi$, where ϕ isolates one of the finitely many types over the empty set, by Ryll-Nardzewski.

Definition 3.6. Let $A, B, C \subseteq \mathcal{M}^{eq}$. Then, A and C are **weakly algebraically independent** over B , written $A \perp_B^a C$ if $\text{acl}^{eq}(AB) \cap \text{acl}^{eq}(BC) = \text{acl}^{eq}(B)$.

Theorem 3.3 yields very powerful results when considering an ergodic measure and the inner product on $L^2(\mu)$.

Theorem 3.7. *Let \mathcal{M} be an ω -categorical countable structure with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Consider the complex Hilbert space $\mathcal{H} = L^2(\mu)$, where μ is an ergodic measure. Suppose that $A, B \subseteq M^{eq}$ are algebraically closed and that $A \perp^a B$. Let $f \in \mathcal{H}_A, g \in \mathcal{H}_B$. Then,*

$$\langle f, g \rangle = \langle f, \mathbb{1} \rangle \overline{\langle g, \mathbb{1} \rangle}.$$

Proof. Let f_0 and g_0 be the projections of f and g respectively on \mathcal{H}_\emptyset . Now, since $f \perp_{\mathcal{H}_\emptyset} g$, $\langle f, g \rangle = \langle f_0, g_0 \rangle$. Since $f_0, g_0 \in \mathcal{H}_\emptyset$, by Lemma 3.5, $f_0 = \alpha \mathbb{1}, g_0 = \beta \mathbb{1}$ for $\alpha, \beta \in \mathbb{C}$. Now, this yields that

$$\langle f, g \rangle = \langle f_0, g_0 \rangle = \langle \alpha \mathbb{1}, \beta \mathbb{1} \rangle = \alpha \bar{\beta} \langle \mathbb{1}, \mathbb{1} \rangle.$$

Being in a probability space, $\langle \mathbb{1}, \mathbb{1} \rangle = 1$. However, since $\langle f - f_0, f_0 \rangle = 0$, we obtain that $\langle f, \mathbb{1} \rangle = \alpha$, and similarly, $\langle g, \mathbb{1} \rangle = \beta$. This yields the desired result. \square

This is already substantially observed in Corollary 3.5 and Remark 3.6 of [13]. In particular, since we are interested in the measures of formulas, we have that

Corollary 3.8. *Let \mathcal{M} be ω -categorical. Let μ be an ergodic measure on M^{eq} in the variable x . Suppose that $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Let a, b be tuples from M^{eq} such that $a \perp^a b$. Then, for any \mathcal{L}^{eq} -formulas $\phi(x, y), \psi(x, z)$,*

$$\mu(\phi(x, a) \wedge \psi(x, b)) = \mu(\phi(x, a))\mu(\psi(x, b)).$$

Proof. Note that the indicator functions $\mathbb{1}_{\phi(x, a)}$ and $\mathbb{1}_{\psi(x, b)}$ are in $\mathcal{H}_{\text{acl}^{eq}(a)}$ and $\mathcal{H}_{\text{acl}^{eq}(b)}$ respectively, and that

$$\begin{aligned} \mu(\phi(x, a) \cap \psi(x, b)) &= \int_{S_x(M)} \mathbb{1}_{\phi(x, a) \cap \psi(x, b)} d\mu \\ &= \int_{S_x(M)} \mathbb{1}_{\phi(x, a)} \cdot \mathbb{1}_{\psi(x, b)} d\mu \\ &= \left(\int_{S_x(M)} \mathbb{1}_{\phi(x, a)} d\mu \right) \overline{\left(\int_{S_x(M)} \mathbb{1}_{\psi(x, b)} d\mu \right)} \\ &= \mu(\phi(x, a)) \overline{\mu(\psi(x, b))} \\ &= \mu(\phi(x, a))\mu(\psi(x, b)). \end{aligned}$$

Here the third equality holds by Theorem 3.7 and the last one follows since our measure is real-valued. Hence, the result follows. \square

Remark 3.9. This corollary has a partial converse which is well known in ergodic theory [13, Proposition 4.8]. Fix an invariant Keisler measure μ and suppose that for any formula $\phi(x, a)$ there is an automorphism $\sigma \in \text{Aut}(M)$ such that

$$\mu(\phi(x, a) \wedge \phi(x, \sigma \cdot a)) = \mu(\phi(x, a))^2.$$

Then, μ is ergodic.

We conclude this section with a brief discussion of the assumption of $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$ in Theorem 3.7 and Corollary 3.8. The assumption is needed in order to have the equality $L^2(\mu)_{\text{acl}^{eq}(\emptyset)} = L^2(\mu)_{\text{dcl}^{eq}(\emptyset)}$. The latter then yields the desired results since we know from Corollary 3.5 that $L^2(\mu)_{\text{dcl}^{eq}(\emptyset)}$ is generated by the constant indicator function. However, we can also obtain the relevant independence results when $\text{acl}^{eq}(\emptyset) \supseteq \text{dcl}^{eq}(\emptyset)$. The following lemma is common knowledge:

Lemma 3.10. *Let \mathcal{M} be ω -categorical. Let $A \subseteq M$ be finite. Fix the variable x in the home sort. Then, there is $A_0 \subseteq \text{acl}^{eq}(A)$ finite such that for every b , a tuple from \mathcal{M} in the variable x ,*

$$\text{tp}(b/A_0) \vdash \text{tp}(b/\text{acl}^{eq}(A)). \quad (1)$$

By this we mean that if b and b' in the variable x have the same type over A_0 , then they have the same type over $\text{acl}^{eq}(A)$ in M^{eq} .

Proof. This is substantially Lemma 2.4 in [9]. □

A useful consequences of the lemma is:

Corollary 3.11. *Let \mathcal{M} be ω -categorical. Let μ be an invariant Keisler measure on \mathcal{M}^{eq} in the variable x . Then, there is finite $A_0 \subseteq \text{acl}^{eq}(\emptyset)$ such that*

$$L^2(\mu)_{A_0} = L^2(\mu)_{\text{acl}^{eq}(\emptyset)}.$$

Proof. Let $A_0 \subseteq \text{acl}^{eq}(\emptyset)$ be finite such that **1** from Lemma 3.10 holds. Suppose that $f \in L^2(\mu)$ is G_A -invariant for $G = \text{Aut}(M^{eq})$ and A a finite subset of $\text{acl}^{eq}(\emptyset)$. Then, f is also $G_{A'}$ -invariant for $A' = A \cup A_0$. But then, by choice of A_0 (with respect to the variable x), the $G_{A'}$ -orbit of any type in $S_x(M^{eq})$ is the same as its G_{A_0} -orbit. Hence, f is G_{A_0} -invariant. □

For an ω -categorical structure \mathcal{M} and given the variable x and A_0 as above, there are finitely many types over A_0 in the variable x , isolated by formulas $\chi_1(x), \dots, \chi_m(x)$. By additivity, any invariant Keisler measure μ can be written as a weighted sum of measures μ_{χ_i} for $1 \leq i \leq m' \leq m$, where, for $\mu(\chi_i(x)) > 0$, μ_{χ_i} is the $\text{Aut}(M/A_0)$ -invariant Keisler measure induced from μ by

$$\mu_{\chi_i}(\phi(x, a)) = \frac{\mu(\phi(x, a) \wedge \chi_i(x))}{\mu(\chi_i(x))}.$$

From Corollary 3.11, we know that for any $\text{Aut}(M/A_0)$ -ergodic measure ν we have that for $a \perp^a b$, and \mathcal{L}^{eq} -formulas $\phi(x, y), \psi(x, z)$,

$$\nu(\phi(x, a) \wedge \psi(x, b)) = \nu(\phi(x, a))\nu(\psi(x, b)).$$

Hence, in the context of $\text{acl}^{eq}(\emptyset) \supseteq \text{dcl}^{eq}(\emptyset)$, in order to study invariant Keisler measures on an ω -categorical structure \mathcal{M} we may naturally move to study the $\text{Aut}(M/A_0)$ -invariant Keisler measures.

4 A strong independence theorem

In this section, we prove that in a simple ω -categorical structure \mathcal{M} with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$ if forking over the emptyset is the same as being universally measure zero, then \mathcal{M} satisfies a stronger version of the independence theorem over finite algebraically closed sets. We conclude with some consequences for ω -categorical MS-measurable structures. Unless specified otherwise, \mathcal{M} denotes the countable model of an ω -categorical theory.

Firstly, we note how the measure $\mu(\phi(x, a) \wedge \psi(x, b))$ only depends on the types of the individual parameters when a and b are weakly algebraically independent.

Corollary 4.1. *Let \mathcal{M} be an ω -categorical structure with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Suppose that $a, b \in \mathcal{M}^{eq}$ are such that $a \perp^a b$. Let $\phi(x, y), \psi(x, z)$ be \mathcal{L}^{eq} -formulas. Then, for an arbitrary $\text{Aut}(M)$ -invariant Keisler measure $\mu : \text{Def}_x(M) \rightarrow [0, 1]$,*

$$\mu(\phi(x, a) \wedge \psi(x, b))$$

only depends on $\text{tp}(a)$ and $\text{tp}(b)$.

Proof. Let $a' \perp^a b'$ be such that $a' \equiv a$ and $b' \equiv b$. Then, by Corollary 3.8, we get that for any ergodic measure ν ,

$$\nu(\phi(x, a) \wedge \psi(x, b)) = \nu(\phi(x, a))\nu(\psi(x, b)) = \nu(\phi(x, a'))\nu(\psi(x, b')) = \nu(\phi(x, a') \wedge \psi(x, b')).$$

But then, by the ergodic decomposition, Corollary 2.8, we have that for any $\text{Aut}(M)$ -invariant Keisler measure

$$\mu(\phi(x, a) \wedge \psi(x, b)) = \mu(\phi(x, a') \wedge \psi(x, b')).$$

□

Definition 4.2. Let \mathcal{M} be an \mathcal{L} -structure and $A \subseteq M$. We write $F(A)$ for the set of formulas with parameters from \mathcal{M} forking over A .

Following Remark 2.6, we always compare the forking and universally measure zero ideals in sufficiently saturated and strongly homogeneous models. As noted in the introduction, from [3] we know that $F(\emptyset) \subseteq \mathcal{O}(\emptyset)$. We are interested in studying what happens when $F(\emptyset) = \mathcal{O}(\emptyset)$ in order to find a structure where the two sets differ.

Definition 4.3. Let \mathcal{M} be an \mathcal{L} -structure. We say that \mathcal{M} **satisfies the strong independence theorem** over $A \subseteq M^{eq}$ if the following holds:

Let $a, b, c_0, c_1 \in M^{eq}$ be such that $a \downarrow_A^a b$, $c_0 \equiv_A c_1$ and $c_0 \downarrow_A a$, $c_1 \downarrow_A b$. Then, there is $c^* \in M^{eq}$ such that $c^* \equiv_{Aa} c_0$, $c^* \equiv_{Ab} c_1$, and $c^* \downarrow_A ab$.

Here, the relation \downarrow denotes non-forking independence. From Corollary 4.1 we obtain that simple ω -categorical structures where forking coincides with being universally measure zero satisfy the strong independence theorem over the empty set.

Theorem 4.4. Let \mathcal{M} be a simple ω -categorical structure with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Suppose that $F(\emptyset) = \mathcal{O}(\emptyset)$, i.e. a formula forks over the empty set if and only if it is universally measure zero. Then, \mathcal{M} satisfies the strong independence theorem over the empty set.

Proof. Suppose that there are $a, b, c_0, c_1 \in M^{eq}$ as in Definition 4.3. Let $\phi(x, a)$ and $\psi(x, b)$ isolate $\text{tp}(c_0/a)$ and $\text{tp}(c_1/b)$. By the existence property of non-forking independence, there is $b' \equiv b$ such that $b' \downarrow a$. By Corollary 4.1, for any Keisler measure

$$\mu(\phi(x, a) \wedge \psi(x, b)) = \mu(\phi(x, a) \wedge \psi(x, b')).$$

By simplicity, $\phi(x, a) \wedge \psi(x, b')$ does not fork over the empty set since the independence theorem holds over \emptyset . Hence, by $F(\emptyset) = \mathcal{O}(\emptyset)$, $\phi(x, a) \wedge \psi(x, b)$ is not universally measure zero, and so is non-forking over the empty set. This proves the strong independence theorem over the empty-set for \mathcal{M} . \square

Remark 4.5. In general, a simple ω -categorical theory with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$ satisfies the independence theorem over the empty set (see, for example, [18]). However, the condition of $F(\emptyset) = \mathcal{O}(\emptyset)$ yields a strengthening of the independence theorem, where the "base" of the amalgamation is weakly algebraically independent rather than non-forking independent. In fact, while non-forking independence implies weak algebraic independence, the converse does not hold in general.

Definition 4.6. We say that an ω -categorical simple structure \mathcal{M} is **one-based** if given $A, B \subseteq M^{eq}$ algebraically closed, then,

$$A \downarrow_{A \cap B} B.$$

If \mathcal{M} is one-based, for $A \subseteq M$ we have that

$$b \downarrow_A^a c \text{ if and only if } b \downarrow_A c.$$

In particular, in a one-based structure, satisfying the independence theorem over A is equivalent to satisfying the strong independence theorem over A .

However, there are not one-based simple ω -categorical structures. The only known example of this are ω -categorical Hrushovski constructions. For these, to satisfy the strong independence theorem is a genuinely stronger requirement than satisfying the independence theorem.

We conclude this section with some consequences for MS -measurable structures in an ω -categorical context. For a general introduction to MS -measurable structures we suggest [7] or the original article [16]. In [18], I discuss in detail ω -categorical MS -measurable structures, and we direct the reader to that article for the relevant definitions and some basic

results. The general idea is that MS -measurable structures have an associated dimension-measure function assigning each definable set a dimension and a measure. The dimension-measure function is invariant (and definable), the measure always takes strictly positive values and the dimension and the measure satisfy Fubini's theorem. Being MS -measurable is a property of a theory. An important feature proved in [16] is that \mathcal{M} is MS -measurable if and only if \mathcal{M}^{eq} is.

Remark 4.7. The dimension d of the dimension-measure function in an MS -measurable theory induces a natural notion of independence: for tuples b, c and small A , we write

$$b \underset{A}{\perp}^d c \text{ if and only if } d(b/Ac) = d(b/A),$$

where for a tuple b and A small,

$$d(b/A) = \min\{d(\phi(x, a)) \mid \phi(x, a) \in \text{tp}(b/A)\}.$$

A folklore result attributed to Ben-Yaacov in [7] is that this notion of independence corresponds to non-forking independence. See [18] for a proof of this. Using this fact, we can prove that in MS -measurable theories forking coincides with being universally measure zero (over any small set).

Lemma 4.8. *Let \mathbb{M} be the monster model of an MS -measurable theory. Then, $F(A) = \mathcal{O}(A)$ for any small $A \subseteq \mathbb{M}$.*

Proof. Suppose there is a formula $\phi(x, b)$ which does not fork over A , but has measure zero for every $\text{Aut}(\mathbb{M}/A)$ -invariant Keisler measure. Let $c \models \phi(x, b)$ be such that $c \underset{A}{\perp} b$. Since dimension-independence corresponds to non-forking independence, as noted in Remark 4.7, $d(c/Ab) = d(c/A)$. Let $\psi(x, a) \in \text{tp}(c/A)$ be such that $d(\psi(x, a)) = d(c/A)$. By MS -measurability [16, Proposition 5.3], we have an induced $\text{Aut}(\mathbb{M}/A)$ -invariant Keisler measure μ_ψ on $\text{Def}_x(\mathbb{M})$ given by

$$\mu_\psi(\chi(x, d)) = \begin{cases} \frac{\mu(\chi(x, d) \wedge \psi(x, a))}{\mu(\psi(x, a))} & \text{for } d(\chi(x, d) \wedge \psi(x, a)) = d(\psi(x, a)), \\ 0 & \text{otherwise,} \end{cases}$$

where μ is the measure in the dimension-measure. Being universally measure zero (over A), we have that $\mu_\psi(\phi(x, b)) = \mu(\phi(x, b) \wedge \psi(x, a)) = 0$, which contradicts positivity of the measure. \square

From Corollary 4.1 we obtain that ω -categorical MS -measurable structures satisfy the strong independence theorem over the algebraic closures of finite sets. Indeed, we also get the corresponding probabilistic independence statement. Recall from Lemma 3.10 that for $A \subseteq M^{eq}$ the algebraic closure of a finite set, and fixing the variable x , there is $A_0 \subset A$ finite such that types in the variable x over A are isolated by \mathcal{L}^{eq} -formulas with parameters from A_0 .

Theorem 4.9. *Let \mathcal{M} be ω -categorical and MS -measurable. Then, it satisfies the strong independence theorem over the algebraic closures of finite sets:*

Let $A \subseteq M^{eq}$ be the algebraic closure of a finite set, a_0, a_1, b, c tuples from M^{eq} . Suppose $a_0 \equiv_A a_1$, $b \underset{A}{\perp}^a c$ and that $a_0 \underset{A}{\perp} b$, $a_0 \underset{A}{\perp} c$. Then, there is a^ such that $a^* \equiv_{Ab} a_0$, $a^* \equiv_{Ac} a_1$, and $a^* \underset{A}{\perp} bc$. Moreover,*

$$\mu(\text{tp}(a_0/Ab) \cup \text{tp}(a_1/Ac)) = \frac{\mu(a_0/Ab)\mu(a_1/Ac)}{\mu(a_0/A)}.$$

Proof. In [18] we proved a version of this statement where $b \underset{A}{\perp} c$ (instead of $b \underset{A}{\perp}^a c$) simplifying some more general results in [11]. We shall use this and Corollary 4.1 to deduce the theorem. Let $\phi(x, b)$ and $\psi(x, c)$ isolate $\text{tp}(a_0/Ab)$ and $\text{tp}(a_1/Ac)$ respectively. Let $\chi(x)$ isolate $\text{tp}(a_i/A)$, and let μ_χ be the $\text{Aut}(M^{eq}/A)$ invariant Keisler measure induced by $\chi(x)$.

We consider $\mu_\chi(\phi(x, b) \wedge \psi(x, c))$. By extension we can find $c' \equiv_A c$ such that $c' \perp_A b$. Since non-forking independence implies weak algebraic independence, by Corollary 4.1

$$\mu_\chi(\phi(x, b) \wedge \psi(x, c')) = \mu_\chi(\phi(x, b) \wedge \psi(x, c)). \quad (2)$$

But the former has positive measure by the independence theorem over algebraically closed sets (and d -independence being the same as non-forking independence). Hence, $\phi(x, b) \wedge \psi(x, c)$ must have a realisation which is independent from bc over A . From [18], for $b \perp_A c'$ we have that

$$\mu(\phi(x, b) \wedge \psi(x, c')) = \frac{\mu(\phi(x, b))\mu(\psi(x, c))}{\mu(\chi(x))}.$$

However, multiplying both sides of 2 by $\mu(\chi(x))$,

$$\mu(\phi(x, b) \wedge \psi(x, c')) = \mu(\phi(x, b) \wedge \psi(x, c)).$$

This yields the desired equation. \square

Remark 4.10. By the proof above and Remark 3.9, an MS -measurable ω -categorical structure induces various ergodic Keisler measures. Let $A \subseteq M^{eq}$ be the algebraic closure of a finite set and $\chi(x)$ isolate a type over A . Then, the induced $\text{Aut}(M/A)$ -invariant Keisler measure μ_χ is $\text{Aut}(M/A)$ -ergodic. We can also prove this using [18] and Remark 3.9.

5 The counterexample

We are now ready to introduce our example of a simple ω -categorical structure with $F(\emptyset) \subsetneq \mathcal{O}(\emptyset)$. It will be an ω -categorical Hrushovski construction. For details on Hrushovski constructions the reader may refer to [24, § 6.2.1]. This construction is also used in [18] as an example of an ω -categorical supersimple structure which is not MS -measurable, so the reader may refer to that article for details on this particular construction. The basic idea is that (non-trivial) simple ω -categorical Hrushovski constructions are not one-based. Hence, there are pairs which are weakly algebraic independent, but forking-dependent. This allows us to build Hrushovski constructions which are simple but which do not satisfy the strong independence theorem.

Theorem 5.1. *There is an ω -categorical graph \mathcal{M} supersimple of SU -rank 2 with the following properties:*

- $\text{Aut}(M)$ acts transitively on the vertices of M .
- Points are algebraically closed;
- Edges are algebraically closed, but for $b \in M$ the formula $E(x, b)$ asserting that x has an edge with b forks over \emptyset ;
- For $a, b \in M$ with no edge between them, $\text{acl}(ab) = ab$ or $\text{acl}(ab) = abc$, where abc is a path of length two with endpoints a and b . In either case, $a \perp b$;
- The smallest k -cycle in \mathcal{M} is a 6-cycle;
- The structure \mathcal{M} has weak elimination of imaginaries;

Furthermore, we can choose \mathcal{M} to satisfy independent n -amalgamation over the algebraic closures of finite sets for any $n \in \mathbb{N}$, and also for all $n \in \mathbb{N}$.

Proof. The structure is an ω -categorical Hrushovski construction. It is the same construction as in [18, Construction 5.1]. In the appendix of that article we prove supersimplicity, weak elimination of imaginaries and note how higher independent amalgamation can be obtained. The other properties also follow by construction and basic calculations with the dimension, recalling that in a Hrushovski construction, the Hrushovski dimension corresponds to SU -rank [24, Corollary 6.2.26]. \square

Remark 5.2. The graph \mathcal{M} in Theorem 5.1 is also **extremely amenable** in the sense of [12]. That is, every type in $S(\emptyset)$ extends to an $\text{Aut}(M)$ -invariant type over M . Since invariant types can be considered as invariant Keisler measures taking only the values 0 and 1, this guarantees that there are some invariant Keisler measures on \mathcal{M} in each variable. To see that these structures are extremely amenable, for a finite tuple \bar{a} from M , consider the type $p(\bar{x})$ given by

$$\bigcup_{B \subseteq M \text{ finite}} \{ \text{tp}(\bar{a}'/B) \mid \bar{a}' \equiv \bar{a}, \text{acl}(\bar{a}B) = \text{acl}(\bar{a}) \cup \text{acl}(B), \neg E(c, b) \text{ for } c \in \text{acl}(\bar{a}), b \in \text{acl}(B) \}.$$

Substantially, $p(\bar{x})$ asserts that \bar{x} has no relation to \mathcal{M} and is weakly algebraically independent from it. From [18, Section 4] we can conclude that this type is consistent, complete and invariant by the extension property and because types of finite tuples are determined by the quantifier-free types of their algebraic closures.

It is commonly known that in a structure with weak elimination of imaginaries, for $A, B \subseteq M$ algebraically closed in \mathcal{M} , we have that

$$\text{acl}^{\text{eq}}(A) \cap \text{acl}^{\text{eq}}(B) = \text{acl}^{\text{eq}}(A \cap B).$$

Hence, for a and b sharing an edge, $a \downarrow^a b$, but $\text{tp}(a/b)$ forks over \emptyset . Meanwhile, for a and c at distance two from each other, $a \downarrow c$. Hence, for \mathcal{M} to satisfy the strong independence theorem over the empty set, \mathcal{M}_f should include 5-cycles. But we have built \mathcal{M}_f to exclude these. This implies the existence of a formula which does not fork over the empty-set, but is universally measure zero. We can give this formula explicitly:

Theorem 5.3. *Let \mathcal{M} be the ω -categorical structure described in Theorem 5.1. Let $\phi(x, y)$ be the formula stating that the points x and y are exactly at distance two from each other. Then, for $a \in M$, the formula $\phi(x, a)$ is universally measure zero but does not fork over the emptyset.*

Proof. Let $a, b \in M$ share an edge. Now, $\phi(x, a) \wedge \phi(x, b)$ is inconsistent since \mathcal{M} avoids 5-cycles. Meanwhile, $\phi(x, a)$ does not fork over the empty set since $c \downarrow a$ for c at distance two from a . However, since $a \downarrow^a b$, for an ergodic measure,

$$0 = \mu(\phi(x, a) \wedge \phi(x, b)) = \mu(\phi(x, a))\mu(\phi(x, b)) = \mu(\phi(x, a))^2,$$

where the second equality follows by Corollary 3.8 and the last equality by transitivity of the action of $\text{Aut}(M)$. From the calculation above we get that $\mu(\phi(x, a)) = 0$ for any ergodic measure. By Corollary 2.8, $\phi(x, a)$ is universally measure zero. \square

Hence, we get the desired counterexample:

Theorem 5.4. *There are ω -categorical supersimple theories T of finite SU -rank with a formula $\phi(x, a)$ which does not fork over the empty set, but which is universally measure zero.*

Furthermore, by Lemma 4.8, we get a counterexample to the question of Elwes and Macpherson [7]:

Corollary 5.5. *There are ω -categorical supersimple theories T of finite SU -rank which are not MS -measurable.*

The above was also answered in [8] and [18], with ω -categorical Hrushovski constructions as counterexamples.

Remark 5.6. The point at the heart of our proof is that it is possible to build simple ω -categorical Hrushovski constructions which do not satisfy the strong independence theorem. This can be done in different relational languages and with different ranks for the final structure. For example, using a 3-hypergraph, we can build a simple ω -categorical Hrushovski construction of SU -rank 1 with non-forking formulas which are universally measure zero.

Remark 5.7. As noted after Definition 4.6, for a simple ω -categorical structure with $\text{acl}^{eq}(A) = \text{dcl}^{eq}(A)$, being one-based implies satisfying the strong independence theorem over A . However, satisfying the strong independence theorem over the algebraic closures of finite sets does not imply being one-based: the standard example of a not one-based supersimple ω -categorical Hrushovski construction [24, Example 6.2.27] does satisfy the strong independence theorem over the algebraic closures of finite sets. See [17] for details. At the moment of writing, we do not know whether satisfying $F(A) = \mathcal{O}(A)$ for all finite sets implies being one-based.

5.1 Conclusions

Our result makes substantial use of the fact we work with not one-based ω -categorical structures. In fact, our counterexample relies on our simple structure not satisfying the strong independence theorem. As we noted in Remark 5.7, for this to be possible we must work with a not one-based structure. This raises two natural questions:

Question 5.8. Suppose that \mathcal{M} is a simple ω -categorical structure satisfying the strong independence theorem over the algebraic closures of finite sets. Does this imply that $F(\emptyset) = \mathcal{O}(\emptyset)$?

Question 5.9. Suppose that \mathcal{M} is a simple one-based ω -categorical structure. Does this imply that $F(\emptyset) = \mathcal{O}(\emptyset)$?

Regarding Question 5.8, we might also ask whether satisfying some sufficiently strong higher amalgamation property implies $F(\emptyset) = \mathcal{O}(\emptyset)$. In this article we have shown that satisfying independent n -amalgamation for all $n \in \mathbb{N}$ over finite algebraically closed sets is not sufficient, but an adequate generalisation of the strong independence theorem might work.

For both questions we may also ask whether the hypotheses mentioned (substituting "simple" by "supersimple") imply MS -measurability. The finite rank assumption is not needed since supersimple ω -categorical one-based structures are of finite rank [10]. In this context, the author's PhD thesis [17] provides negative answers to both questions. In particular, we know of an example of a supersimple ω -categorical Hrushovski construction of SU -rank 1 which is not MS -measurable in spite of satisfying the strong independence theorem over the algebraic closures of finite sets and independent n -amalgamation for all n [17, Ch. 7.5]. We strongly suspect that some ω -categorical Hrushovski construction will yield an example of a structure with $F(\emptyset) \subsetneq \mathcal{O}(\emptyset)$ while satisfying the strong independence theorem (and arbitrarily strong higher amalgamation properties). Meanwhile, the universal homogeneous tetrahedron-free 3-hypergraph is not MS -measurable in spite of being ω -categorical supersimple and one-based [17, Ch. 7.3]. If one could prove that $F(\emptyset) \subsetneq \mathcal{O}(\emptyset)$ for this structure we would answer negatively both questions.

On the other hand, we do not know whether being ω -categorical and MS -measurable implies being one-based. Indeed, this question was also asked by Elwes and Macpherson in [7]. In a more general context, it is unclear whether one might be able to show that forking and being universally measure zero disagree for not one-based supersimple ω -categorical structures:

Question 5.10. Suppose that \mathcal{M} is supersimple ω -categorical with $F(A) = \mathcal{O}(A)$ for all finite $A \subset M$. Is \mathcal{M} one-based?

By Lemma 4.8, a positive answer to this question would imply that all ω -categorical MS -measurable structures are one-based, answering the question of Elwes and Macpherson.

References

- [1] Vladimir I. Bogachev. *Measure Theory*. 1st ed. 2007. Measure Theory; Berlin, Heidelberg: Springer Berlin Heidelberg, 2007. ISBN: 1-280-74570-3.

- [2] Samuel Braunfeld and Marimon Paolo. *Invariant measures in ω -categorical NIP theories*. Ongoing work. 2022.
- [3] Artem Chernikov, Ehud Hrushovski, Alex Kruckman, Krzysztof Krupiński, Slavko Moconja, Anand Pillay, and Nicholas Ramsey. “Invariant measures in simple and in small theories”. In: *Journal of Mathematical Logic* 23.02 (2023), p. 2250025. DOI: [10.1142/S0219061322500258](https://doi.org/10.1142/S0219061322500258). eprint: <https://doi.org/10.1142/S0219061322500258>. URL: <https://doi.org/10.1142/S0219061322500258>.
- [4] Alexis Chevalier and Ehud Hrushovski. *Piecewise Interpretable Hilbert Spaces*. 2021. DOI: [10.48550/ARXIV.2110.05142](https://arxiv.org/abs/2110.05142). URL: <https://arxiv.org/abs/2110.05142>.
- [5] John B. Conway. *A course in functional analysis*. 2nd ed. Graduate texts in mathematics ; 96. New York: Springer-Verlag, 1990. ISBN: 0387972455.
- [6] Manfred Einsiedler. *Ergodic Theory with a view towards Number Theory*. 1st ed. 2011. Graduate Texts in Mathematics, 259. London: Springer London, 2011. ISBN: 0-85729-021-5.
- [7] Richard Elwes and Dugald Macpherson. “A survey of asymptotic classes and measurable structures”. In: *Model Theory with Applications to Algebra and Analysis*. Ed. by Z. Chatzidakis, D. Macpherson, A. Pillay, and A. Wilkie. Vol. 2. London Mathematical Society Lecture Note Series. Cambridge University Press, 2008, pp. 125–160. DOI: [10.1017/CB09780511735219.004](https://doi.org/10.1017/CB09780511735219.004).
- [8] David M. Evans. *Higher amalgamation properties in measured structures*. To appear in *Model Theory*. 2022. DOI: [10.48550/ARXIV.2202.10183](https://arxiv.org/abs/2202.10183). URL: <https://arxiv.org/abs/2202.10183>.
- [9] David M. Evans and Todor Tsankov. “Free actions of free groups on countable structures and property (T)”. In: *Fundamenta Mathematicae* 232 (2016), pp. 49–63. DOI: [10.4064/fm232-1-4](https://doi.org/10.4064/fm232-1-4). URL: <http://dx.doi.org/10.4064/fm232-1-4>.
- [10] David M. Evans and Frank O. Wagner. “Supersimple ω -Categorical Groups and Theories”. In: *The Journal of Symbolic Logic* 65.2 (2000), pp. 767–776. ISSN: 00224812. URL: <http://www.jstor.org/stable/2586568> (visited on 07/12/2022).
- [11] Ehud Hrushovski. “Approximate Equivalence Relations”. Unpublished notes. 2015.
- [12] Ehud Hrushovski, Krzysztof Krupiński, and Anand Pillay. *On first order amenability*. 2020. DOI: [10.48550/ARXIV.2004.08306](https://arxiv.org/abs/2004.08306). URL: <https://arxiv.org/abs/2004.08306>.
- [13] Colin Jahel and Todor Tsankov. “Invariant measures on products and on the space of linear orders”. In: *Journal de l’École polytechnique — Mathématiques* 9 (2022), pp. 155–176.
- [14] Byunghan Kim. *Simplicity theory*. Oxford University Press, 2014.
- [15] Dugald Macpherson. “A survey of homogeneous structures”. In: *Discrete Mathematics* 311.15 (2011). Infinite Graphs: Introductions, Connections, Surveys, pp. 1599–1634. ISSN: 0012-365X. DOI: <https://doi.org/10.1016/j.disc.2011.01.024>. URL: <https://www.sciencedirect.com/science/article/pii/S0012365X11000422>.
- [16] Dugald Macpherson and Charles Steinhorn. “One-dimensional asymptotic classes of finite structures”. In: *Transactions of the American Mathematical Society* 360.1 (2008), pp. 411–448. ISSN: 0002-9947.
- [17] Paolo Marimon. “Measures and amalgamation properties in ω -categorical structures”. PhD thesis. Department of Mathematics, Imperial College London, 2023.
- [18] Paolo Marimon. *On the non-measurability of ω -categorical Hrushovski constructions*. 2022. DOI: [10.48550/ARXIV.2208.06323](https://arxiv.org/abs/2208.06323). URL: <https://arxiv.org/abs/2208.06323>.
- [19] Robert R. Phelps. *Lectures on Choquet’s Theorem*. 2nd ed. 2001. Lecture Notes in Mathematics, 1757. Berlin, Heidelberg: Springer Berlin Heidelberg, 2001. ISBN: 9783540487197.
- [20] Bruno Poizat. *A course in model theory: an introduction to contemporary mathematical logic*. Universitext. New York ; London: Springer, 2000. ISBN: 9780387986555.
- [21] Pierre Simon. *A Guide to NIP Theories*. Lecture Notes in Logic. Cambridge University Press, 2015. DOI: [10.1017/CB09781107415133](https://doi.org/10.1017/CB09781107415133).
- [22] Katrin Tent and Martin Ziegler. *A Course in Model Theory*. Lecture Notes in Logic. Cambridge University Press, 2012. DOI: [10.1017/CB09781139015417](https://doi.org/10.1017/CB09781139015417).
- [23] Todor Tsankov. “Unitary Representations of Oligomorphic Groups”. In: *Geometric and functional analysis* 22.2 (2012), pp. 528–555. ISSN: 1016-443X.
- [24] Frank O. Wagner. “Relational structures and dimensions”. In: *Automorphisms of first-order structures*. Ed. by R. Kaye and D. Macpherson. Clarendon Press, 1994, pp. 153–180.