

greater use that is being made to treat COVID-19 and the higher doses used in severe cases (up to 120 mg).

Conclusions: To conclude, we need to know characteristics of these episodes in order to be able to prevent and treat them properly (minimum effective dose and less time), since they will probably occur more frequently at this time.

Keywords: SARS-CoV-2; corticosteroids; Psychiatric symptoms; steroid-induced psychosis

EPP0467

Post-traumatic stress disorder among tunisian healthcare professionals facing the pandemic coronavirus (COVID-19)

L. Aribi¹, R. Jbir^{2*}, N. Messedi², D. Gdoura², W. Bouattour², N. Kolsi², F. Charfeddine² and J. Aloulou³

¹Psychiatry, Hedi Chaker Sfax, Sfax, Tunisia; ²Psychiatry, Hedi chaker Sfax, sfax, Tunisia and ³Psychiatry (b), Hedi Chaker University hospital, sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.835

Introduction: The new coronavirus has spread rapidly across the planet confining entire populations, filling hospitals overwhelmed by massive arrivals of patients This new health situation was traumatic especially for health-professionals

Objectives: To study the prevalence and predictors of post-traumatic stress disorder, among health-workers.

Methods: Our study was descriptive and analytical cross-sectional, between May until June 2020. An anonymous online-survey was sent to collect those parameters Sociodemographic-information Physical symptoms The existence of contact with a suspected case The need for quarantine The stressful event The state of mental health, using: PCL-5: 20 items which measures the 20 symptoms of post-traumatic stress-disorder according to DSM-5. PSQI: 9 questions to see the existence or not of a disturbance in sleep

Results: 125 participants: 28 university-hospital doctors, 55 residents, 5 interns, 4 specialist-doctors, 2 general-practitioners, 14 nurses, 14 senior-technicians, 2 midwives and a pharmacist. The average seniority at the job was 6 years. Two factors were the most stressful: The characteristics of this pandemic 37.6% The fear of caching the virus and transmit it to their families: 37.6%. 42.4% of participants presented a post-traumatic stress disorder. 3 parameters were correlated with post-traumatic stress disorder: young age, having children ($p = 0.007$) and fewer years of professional-experience. This pandemic altered the quality of sleep of caregivers, 62.4% of them had a bad quality of sleep. The bad sleepers developed more post-traumatic stress disorder

Conclusions: This health crisis had a major impact on the mental health of our heroes that is why we should provide them with the necessary psychological support.

Keywords: Post traumatic Stress Disorder; COVID-19; health professionals

EPP0470

Resilient care in times of covid: The stress buddy

N. Rius-Ottenheim^{1*}, E. Vermetten¹, E. Giltay¹, M. Boeschoten², N. De Bles¹, N. Van Der Wee¹ and A. Van Hemert¹

¹Psychiatry, LUMC, Leiden, Netherlands and ²Psychotrauma, Arq Foundation, Diemen, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.836

Introduction: The COVID-19 outbreak poses a challenge for health care professionals due to a surge in care demands, overwork, fear of contagion and concerns on the availability of protective equipment, and coping with distress of patients and their families. Although there is emerging evidence on prevalence of stress and its predictors, less is known on the trajectory of stress symptoms and the differences between cohorts of health care professionals.

Objectives: To sustain and restore health care professionals the Leiden University Medical Center has launched the Digital Stress Buddy, a mobile app, to assess psychological stress, depressive symptoms, anxiety and posttraumatic stress symptoms.

Methods: Participants fill in a 14-item questionnaire on stress and resilience resources, followed by a COVID-related questionnaire and finally a set of validated questionnaires on depression and anxiety (DASS-21), posttraumatic stress-symptoms (IES-R), burn-out (CBI) and resilience (RES).

Results: To date, 959 health care workers have completed the stress monitor, of whom 223 (23%) showed relevant stress levels. Within this group, anxiety and posttraumatic symptoms were most prevalent (45%), followed by depressive symptoms (15%). Predictors of stress were being female, coping with distress of patients and their families, teleworking, and overwork.

Conclusions: By identifying vulnerabilities and resilience for psychological distress, we are able to tailor the support interventions for health care workers within our hospital. This is an ongoing study and future follow-up during the second wave of the pandemic will provide more insight on the trajectories of stress-related symptoms.

Conflict of interest: No significant relationships.

EPP0471

Psychosocial consequences of the COVID-19 social isolation in the italian general population: Preliminary results

A. Ciaramella^{1,2*}, M. Rossi^{1,2}, M. Jarego³ and A. Ferreira-Valente³

¹Gift Institute Of Integrative Medicine, Psychosomatic Center, Pisa, Italy; ²Education Programme Partner With University Of Pisa, Florence, Padua, Miur, Italy, Aplysia APS, Pisa, Italy and ³Health Psychology, ISPA, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.837

Introduction: Although some philosophers recognize in the loneliness an evolutionary existential process, a 2019 declaration of World Health Organization underlines the major health problem in the worldwide is the perception of state of loneliness. The feeling of loneliness linked to the social isolation (SI) or a lack of social opportunity activate a stressful condition associated to an increase of social dependence. This 'learned social helplessness' can be dangerous so that it is associated with an increased prevalence of suicides (Cacioppo and Cacioppo, 2018; Bzdok and Dunbar, 2020). Considering the impact of loneliness on the mental health we can assume that the COVID-19 forced SI affects the state of health and psychosocial well-being.

Objectives: To evaluate the psychosocial impact of the SI in Italy.