

impedance analysis (TANITA-BC-418MA). A validated physical activity questionnaire was used to ascertain their PE dislikes/barriers, and their answers were compiled into different categories, according to the core themes that were identified during thematic data analysis.

*Results:* 19.5% ( $n = 74$ ) of the subjects were classified as either obese/overweight. 14.2% of the subjects reported having no dislikes/barriers with regard to PE and among this population, only a small proportion (20.4%) of the subjects were among those classified as overweight/obese. This indicates that overweight/obese pupils are more likely to have PE dislikes/barriers. The most common barrier reported among the overweight/obese subjects, was a

dislike of running/sprinting activities (23.2%). Other barriers reported include: unenjoyable/boring activities, tiring and physically unpleasant activities, lack of an adequate range of activities, too competitive PE environments, dislike of PE assessment and dressing out procedures, being dominated by skilled pupils or put into low-ability PE groups and fear of weight-related teasing/ridicule by peers.

*Conclusions:* Physical educators should find innovative ways to address the PE barriers/concerns of overweight/obese pupils, provide them a supportive PE environment to build their confidence and reduce the risk of embarrassment by designing creative activities that are appropriate to their ability.

doi:10.1017/S1368980012002145

## 40 – Association between adiposity and physical activity levels of schoolchildren during physical education lessons

ME Jacobson, CJ Laws and JA Potter

University of Chichester, West Sussex, England

*Introduction:* It is increasingly acknowledged that school physical education (PE) is some children's only experience of physical activity (PA) and therefore has an important role to play in the fight against childhood obesity. The present study explored the relationship between the PA variables (percentage of PE lesson time engaged in (i) moderate-to-vigorous physical activity (MVPA) and (ii) sedentary activity) and the body composition variables (i) BMI; (ii) waist circumference (WC) and (iii) waist-to-height ratio (WHtR)), in 158 pupils (82 boys and 76 girls), aged 11–13 years, from five secondary schools in the South of England.

*Method:* Anthropometry (height, body mass and BMI), and PA data (using Actigraph-GT1M accelerometers) of the pupils were measured using standardized methods.

*Results:* A significant negative correlation was observed between the percentage of PE lesson time engaged in MVPA and the body composition variables of WHtR ( $r = -0.32$ ) and WC ( $r = -0.30$  amongst boys ( $P < 0.01$ ) and WC among girls ( $r = -0.25$ ,  $P < 0.05$ ). There was a small but significant positive correlation ( $r = 0.25$ ) between the percentage of PE lesson time engaged in sedentary activities and WHtR ( $P < 0.05$ ) amongst boys. No significant associations were found between the BMI and any of the PA variables ( $P > 0.05$ ) for either gender. The study suggests that obese pupils are less active and more likely to be sedentary than non-obese pupils, during PE lessons.

*Conclusions:* Physical educators should find innovative ways to increase MVPA participation in overweight/obese pupils. WC and WHtR may be better proxy indices of the lack of PA than BMI.

doi:10.1017/S1368980012002157

## 41 – The influence of a comprehensive programme for treatment of overweight and obesity on physical fitness of children and adolescents in health-resort conditions

Renata Archacka<sup>1</sup>, Ewa Mojs<sup>2</sup>, Artur Cieślak<sup>1</sup>, Grzegorz Ferdynus<sup>1</sup> and Włodzimierz Samoborski<sup>3</sup>

<sup>1</sup>The Klodzko Health Resorts Group Joint Stock Co., Poland; <sup>2</sup>Clinical Psychology Department of the Faculty of Health Sciences of the Poznań University of Medical Sciences, Poland; <sup>3</sup>Department and Clinic of Physiotherapy, Rheumatology and Rehabilitation of the Poznań University of Medical Sciences, Poland