

Nutrition Society Congress 2024, 2–5 July 2024

## Vitamin D intakes and food sources in breastfeeding women in Ireland: findings from the WellFed study

M. Ali<sup>1</sup>, B. Murphy<sup>1</sup>, F. O'Dwyer<sup>1</sup>, M. Sanecka<sup>1</sup>, S. O'Reilly<sup>1</sup> and A. O'Sullivan<sup>1</sup>  
<sup>1</sup>UCD Institute of Food & Health, University College Dublin, Belfield, Dublin 4, Ireland

Limited sunshine and dietary vitamin D make vitamin D deficiency very common in Ireland. Breastfeeding is the universal recommendation and supplies the best combination of nutrients and bioactive factors for all infants<sup>(1)</sup>. However, maternal diet and nutrition status impacts concentrations of certain nutrients in breastmilk and therefore vitamin D concentrations can be lower than recommended for infants<sup>(2,3)</sup>. This study aimed to determine the vitamin D intakes and food sources of breastfeeding mothers in Ireland.

Data was collected as part of the WellFed study, a 4-week randomised control trial investigating the effectiveness of a food supplement containing a protein hydrolysate and beta glucan (Wellmune<sup>®</sup>) during lactation on immune and gut health (LS-23-07-O'Sullivan). Participant vitamin D intakes were assessed using FoodBook24 self-completed 24-hour recalls. Foods were grouped into food groups to determine their contribution to intake as well as the contribution from natural and added vitamin D foods. Data was analysed using IBM SPSS Statistics (version 27).

Dietary analysis of 45 participants indicated that maternal vitamin D intakes were low ( $6.4 \pm 4.6 \mu\text{g}/\text{day}$ ). Most participants (96%) did not meet the  $15 \mu\text{g}/\text{day}$  recommendation, despite 56% of participants reporting they consumed some form of vitamin D supplement. Natural vitamin D foods like eggs and egg dishes, fish and fish products, meat and meat products and foods with added vitamin D including breakfast cereals, beverages, and milk and yoghurt were the key food groups contributing to intake, with the majority coming from natural food sources.

Low dietary vitamin D intake is common among breastfeeding women and diet alone is insufficient to achieve the daily vitamin D requirements even when supplements are consumed. As natural food sources of vitamin D are limited and infrequently consumed, recommendations to consume more vitamin D foods may not be successful as a standalone strategy to improve vitamin D status. A combination of approaches including the promotion of vitamin D rich food consumption, supplementation strategies, and the introduction of food fortification policies may be a feasible method for improving vitamin D across the population.

### Acknowledgments

The authors would like to acknowledge the funders Food for Health Ireland, Grant no: TC201800025, and all participants who took part in the WellFed study.

### References

1. Ballard O, Morrow AL. (2013) *Pediatr Clin North Am* **60**, 49–74.
2. Dawodu A, Zalla L, Woo JG *et al.* (2014) *Maternal Child Nutr* **10**, 383–397.
3. Hollis BW, Roos BA, Draper H *et al.* (1981) *J Nutr* **111**, 1240–1248.