

Tues-P111 COMPLEX THERAPY IN SEX REASSIGNMENT PROCESS

L.M. Vasilenko¹. ¹*Moscow Research Institute of Psychiatry, Moscow, Russia*

The aim of the investigation was to develop complex therapy methods for transsexual persons subjected to plastic surgery on genitals. Clinical assessment, psychopharmacotherapy and psychotherapy have been accomplished before and after the plastic surgery. 125 transsexual patients in age from 18 to 42 years have been investigated. Special tests showed that before the surgery most patients knew almost nothing about the procedure itself, were quite unaware of its possible complications, and had unreasonable expectations and impractical plans concerning their future. All the patients had mild mental disorders as following: (1) anxious-phobic disturbances (34 patients); depressive disturbances (68); mixed depressive and phobic disturbances (26).

Complex therapy included medicinal treatment in combination with psychotherapy. Medicinal treatment was directed to affective disturbances reduction. Psychotherapy method was chosen with account to the patient's personality. Besides cognitive therapy together with educational programme were conducted for all the patients.

The results of this therapy approach showed different positive changes in patients' clinical status and their attitudes toward the surgery, in particular reduction of mood disturbances, realistic predispositions toward the surgery, priority of health value. Some patients chose partial surgery of the genitals for the purpose of possible complications reduction. Expressed positive effect has been evident in 68% of the patients; 23% of the patients showed mild positive effect and 9% showed only slight positive alterations.

Thus this complex therapy showed significant efficacy and can be recommended for transsexual persons treatment.

Tues-P112 CLINICAL PRESENTATION AND THERAPY OF SEXUAL SOMATIZATION DISORDERS IN MALES

I. Kan¹. ¹*Moscow Research institute of Psychiatry, Moscow, Russia*

Clinical particularities and variants were studied in groups of 60 male patients suffering from sexual somatization disorders (age limits from 17 up to 35). The diagnosis of somatization sexual disorder requires that a patient have a specific number of medically unexplained somatic sexual symptoms. The predispositions are represented by particularities of personality, annoying states (social and sexual fears), hysterical reactions and weak variant of sexual constitution. The sexual somatization is accompanied by symptoms of depression and anxiety, what is especially actual for men because of social and psychological importance of male sexual function.

On the basis of psychopathological mechanisms and particularities of personality in structure of sexual somatization disorders we choose certain methods of therapy which include the treatment of anxiety, anxiety associated with depression and neurotic depression by anxiolytics and antidepressants such as Xanax (alprazolam tablets), Coaxil (tianeptin) and others in combination with psychotherapy (rational psychotherapy, psychotherapy in the state of hypnotic suggestion, hypnotherapy, autogenous training, suggestion in the state of walking, sex therapy and others). The satisfactory results were reached in 65–70% of describing cases.

Tues-P113 SEXUAL ANGEDONIA IN WOMEN

I. Airiants^{1*}, N. Kibrik¹. ¹*Moscow Research Institute of Psychiatry, Moscow, Russia*

In our practice, we very often meet such psychiatric and sexual disharmony, as sexual anhedonia. It may accompany wide specter of sexual problems, for example: anorgasmia, dispareunia, marital disharmony and other. At the same time, sexual anhedonia is one of marks of anhedonia in depressive state (in our patients we usually can find subdepression, masked depression, or neurotic depression).

In clinical presentation if this syndrome we see compliance's to absence of sensation (emotional and physical) during sexual foreplay and intercourse itself, lack of enjoyment in sexual communication, sometimes to unpleasant and even disgusting feelings during sexual act (in some cases even more hard during foreplay, deep kisses). Some of our patients have travesty, artificially increased estimate of sexual pleasure, sexual norms, orgasm sensations (for example, they suggest, that orgasm must be accompanied with swoon, or partial consciousness, or be prolonged about 5–10 minutes at least, or they must feel "incredible strong muscle spasms" and so on...). Very often such patients really have orgasmic experience, but just don't want to admit it.

In treatment of our patients we starting with salvation of affective, psychiatric problems - we use antidepressants with tranquilizing effects, such as Xanax, Coaxil and others, and adaptogenes, vitamins (to combat psychastenia). We also employ different methods of psychotherapy (individual, family, rational, suggestive and others). All this measures in complex with special sex-therapy give us possibility to improve sexual life and "quality of life" in common in our patients.

Tues-P114 DIFFERENT FACTORS INFLUENCING THE DECISION TO CHANGE THE GENDER

A.A. Dolnykova¹. ¹*Moscow Research Institute of Psychiatry, Moscow, Russia*

All people who want to change their biological and social gender must pass through many expertises and tests to get the permission to do it. One of them is psychodiagnostics. In our laboratory of psychology and psychotherapy we do psychodiagnostics for all of them. All these patients wanted to have a formal conclusion and were not asking for any help. We tested some features of their mentality, their personality and the gender identification. We define three groups of factors making people to wish to change the gender. The first one is transsexualism, gender identification does not correspond with the biological and social gender a person has. Some patients have some special mental features or thinking decrease which make us supposing some personal or mental disorders. And the last group of factors were some psychological problems, especially problems of the early childhood and relationship with their parents. According to this factors we divide people who want to change their biological and social gender to different groups. The biggest group is the group of "real", "veritable" transsexuals. They do have some psychological problem, but transsexualism is not a consequence of them, rather they are consequence of transsexualism. The other group consists of people with some personal and mental deceases and their desire to change the gender is one of the symptoms they have. The third group is the group of people with psychological problems, they are not aware of them and the problems are represented at the conscience level as