

Identifying dietary patterns in Irish schoolchildren and their association with nutritional knowledge and markers of health before and after intervention

Alison Merrotsy^{1*}, Aoife L. McCarthy², Sean Lacey³ and Tara Coppinger¹

¹Department of Sport, Leisure and Childhood Studies, Cork Institute of Technology, Cork T12 P928, Republic of Ireland

²Department of Biological Sciences, Cork Institute of Technology, Cork T12 P928, Republic of Ireland

³Department of Mathematics, Cork Institute of Technology, Cork T12 P928, Republic of Ireland

(Submitted 29 September 2019 – Final revision received 31 August 2020 – Accepted 7 October 2020 – First published online 15 October 2020)

Abstract

The aim of the study was to identify dietary patterns (DP) and examine differences in anthropometric measures, blood pressure (BP), cardiorespiratory fitness and nutritional knowledge of 6- and 10-year-old children at baseline and following a nutrition and physical activity intervention, with respect to DP and treatment group. This is a longitudinal study. Food diary, nutritional knowledge questionnaire and 550-m walk/run test measured dietary intake, nutritional knowledge and cardiorespiratory fitness, respectively. BP, weight, height and waist circumference were also measured and BMI and waist-to-height ratio (WtHR) were derived. All measurements were performed at baseline and following intervention. Two primary schools (one intervention, one control) in Cork, Ireland, were selected. Participants were 6- (n 39, age 5.9 (SD 0.6) years) and 10- (n 49, age 9.8 (SD 0.5) years)-year-olds. Two DP were identified, using k -means cluster analysis, for both 6- (unhealthy and nutrient-dense) and 10-year-olds (processed and Western diet) at baseline. DP derived post-intervention were (1) plant-based and (2) processed foods for 6-year-olds and (1) nutrient-dense and (2) unhealthy for 10-year-olds. There was no statistically significant difference in DP for 6- and 10-year-olds at baseline and post-intervention ($P > 0.05$). Following the intervention, a multivariate ANOVA showed there were no statistically significant differences in nutritional knowledge, BMI, WtHR, cardiorespiratory fitness and BP based on DP and intervention/control group for both age groups ($P > 0.05$). Three out of four dietary patterns identified for 6- and 10-year-olds were unfavourable. While no statistically significant evidence of intervention impact was found on DP, a positive trend was emerging among 10-year-olds.

Key words: Nutrition: Dietary patterns: Dietary intake: Nutritional knowledge: Cardiorespiratory fitness and health markers of Cork children

Examining dietary patterns, rather than individual foods, provides a more accurate account of dietary intake^(1,2). The study of dietary patterns facilitates the study of the whole diet, examining the interactive effect of individual nutrients, interactions often not detected in isolated analyses of nutrients⁽¹⁾. The increasing prevalence of overweight, low levels of fitness and elevated blood pressure (BP) in Irish children is a growing health concern⁽³⁾ and there is strong evidence to suggest that 'unhealthy' dietary patterns are associated with CVD risk factors^(4,5). Since dietary habits established in childhood are known to track into later life⁽⁶⁾, the early establishment of healthy dietary patterns is important for reducing future cardiometabolic risk⁽⁷⁾.

Several international studies have examined dietary patterns in children^(6,8–12), describing children's diets as 'healthy/health conscious'^(6,10,11) as well as 'traditional'^(6,10,11,13). While a small number of studies have identified dietary patterns in Irish children^(14,15), none have examined their relationship with health

markers, particularly over time. O'Brien *et al.*⁽¹⁴⁾ derived two clusters, 'more healthful' (higher intake of fruits and vegetables, rice, pasta, starches and grains, wholemeal bread and products, breakfast cereals, low-fat milk, fish and fresh poultry) and 'less healthful' (higher intake of white bread and products, butters and spreads, chips and processed potato products, processed red meats, sugars and confectioneries and high-energy beverages). Walton *et al.*⁽¹⁵⁾ explored dietary patterns in relation to change in dietary energy density with age and found that children moved from a less energy-dense diet in pre-school to a more energy-dense diet (consuming less fruit and vegetables, and more fat, confectionery, snacks and sugar-sweetened beverages) as they got older.

Identifying dietary patterns and their association with markers of health may provide a better prediction of diet–disease relationships⁽¹⁶⁾. Shang *et al.*⁽¹⁷⁾ found that children with 'healthy' dietary patterns (high in fruits, vegetables and white meat) had a lower prevalence of obesity, and Smith *et al.*⁽¹⁸⁾

* Corresponding author: Alison Merrotsy, email alison.merrotsy@cit.ie



established that a 'Health Aware' dietary pattern (high in fibre, fruit and vegetables) was linked with decreased fat mass gain in girls aged 9–11 years. However, evidence is not yet conclusive, as Saeedi *et al.*⁽¹⁹⁾ found no significant associations between dietary patterns and markers of CVD, including cardiorespiratory fitness, hand grip strength, fat mass index and arterial stiffness in their study of New Zealand children aged 9–11 years.

There is evidence that school-based obesity prevention interventions have some success in improving dietary intake by increasing fruit and vegetable consumption^(20–23) and reducing fat intake^(24,25). An example of one such intervention in Ireland, Project Spraoi⁽²⁶⁾, has shown success in increasing fibre and protein intake in children, while also improving nutritional knowledge, waist-to-height ratio (WHtR) and BP⁽²⁷⁾. Yet, to date, no Irish study has examined differences in nutritional knowledge and markers of health with respect to dietary patterns or investigated the effect of a school-based obesity prevention intervention on overall dietary patterns in children.

The present study aims to identify dietary patterns in 6- and 10-year-old primary schoolchildren from Cork, Ireland, attending one intervention school and to examine any associations they may have with nutritional knowledge, cardiorespiratory fitness, BP, BMI and WHtR over a 2-year period. The ability of the Project Spraoi (<http://www.isrctn.com/ISRCTN92611015>) intervention to change dietary patterns over time will also be examined.

Methods

Study design and subject participation

Project Spraoi (<http://www.cit.ie/projectspraoi>) is a primary school health promotion intervention that aims to promote physical activity (PA), improve dietary intake and increase nutritional knowledge of Irish schoolchildren and has been described in detail elsewhere^(26,27). Briefly, 20-min targeted nutrition and PA classes that included healthy lunch box ideas, healthy food-tasting sessions and education sessions on sugary beverages, takeaways, healthy breakfasts and food groups as well as 'huff and puff' games and other PA-centred activities were delivered by an 'Energizer' from October 2014 to June 2016. Two schools from Cork were selected via convenience sampling to participate and gain an initial insight into dietary patterns. The design of the present study was exploratory, to inform the direction of a larger study. Inclusion criteria for schools included mixed sex, middle/high socio-economic status, urban/rural, medium sized, proximity of 20 km to the research institute, willingness to implement the 'Project Spraoi' intervention and not currently participating in another PA and/or healthy eating intervention in their school. All students from the intervention school (*n* 274) received the PA and nutrition intervention; children in the control school (*n* 567) did not receive any of the intervention components. The evaluation of dietary intake, nutritional knowledge, BP, anthropometric and cardiorespiratory fitness measurements were carried out on 6- and 10-year-old children only.

A total of eighty-eight children (6-year-olds, *n* 39; males = 22, females = 17; 10-year-olds, *n* 49; male = 23, females = 26)

participated. The mean age was 5.9 (SD 0.6) years for 6-year-olds and 9.8 (SD 0.5) years for 10-year-olds. Data were collected over 4 d (two weekdays and two weekend days) at two time points (baseline, October 2014 and post-intervention, June 2016) by trained postgraduate researchers and staff of Cork Institute of Technology (Cork, Ireland). By the end of the study, following the 2-year intervention, the 6-year-olds were aged 8 and the 10-year-olds were aged 12 years. The same children were assessed at baseline and post-intervention; non-response rate for completed food diaries increased from 25% (*n* 25) at baseline to 36% (*n* 36) post-intervention. Informed consent and parental consent were required from all children and their parents/guardians before participation. Ethical approval was obtained from Cork Institute of Technology Research Ethics Committee in September 2013.

Dietary analysis

Dietary intake was examined using a 4-d estimated food diary (two weekdays and two weekend days), a validated measure in children⁽²⁸⁾. The diary was adapted from the Cork Children's Lifestyle Study (CCLaS) 3-d food diary⁽²⁹⁾. Written instructions to complete the food diary, including food photographs from the Young Person's Food Atlas Primary⁽³⁰⁾ to aid portion size estimation, were provided at the beginning of the food diary. The researcher also provided verbal instruction to participants in the classroom on each day of diary completion. Each day of the food diary was broken into six meal sections: breakfast, morning snack, lunch, afternoon snack, dinner and evening snack. Parents of the 6-year-old children were asked to complete the food diary for their child (with the child also present), while the 10-year-old children were advised to seek help with the completion of their diaries from parents/guardian and teachers. The Irish Food Portion Sizes Database⁽³¹⁾ was used to assign an age-appropriate median portion size, where no estimation was given and also to estimate the number of unhealthy snacks (sweets, crisps, chocolate, ice cream, cake, biscuits, bars and pastries) consumed.

All foods and beverages were entered into the Dietplan 7 (Forestfield Software Ltd, 2015) software package. Output measures included fruit and vegetable intake, number of unhealthy snacks per day and macro- (protein, fat, saturated fat and carbohydrate) and micro- (Na, Ca and Fe) nutrient intake.

Nutrition knowledge questionnaire

Participant's knowledge of healthy eating was assessed using the validated questionnaire 'Fit Kids "r" Healthy Kids'⁽³²⁾. It was first piloted by the researcher in a primary school in Cork (*n* 23) to ensure its relevance in an Irish setting and the Irish food pyramid⁽³³⁾ replaced the United States Department of Agriculture (USDA) food pyramid⁽³⁴⁾ used in the original questionnaire. Fifteen multiple-choice questions assessed knowledge on food groups, healthful foods and food functions.

Anthropometric and blood pressure measurements

Height, body mass, BP, heart rate and waist circumference were measured by trained researchers using standard procedures⁽³⁵⁾.



Height and body mass values were used to derive BMI and BMI Z scores^(36,37). Waist circumference and height measurements were used to calculate WHtR and children were classified into two categories (WHtR < 0.50 and WHtR ≥ 0.50), where a WHtR greater than or equal to 0.50 was considered to be an indicator of central obesity⁽³⁸⁾.

Cardiorespiratory fitness

A validated 550-m walk/run test^(39,40) was used to measure cardiorespiratory fitness. The run was performed outside on grass, where a 110-m rope was arranged in an oval shape. Children were instructed to run/walk as fast as they could for the five laps of the test. Completion time was recorded using a stopwatch to the nearest minute and second. Run scores were then classified into fast (≤50th centile) and slow (>50th centile) categories based on the Waikato 2011 centile charts for time to complete 550 m⁽⁴¹⁾.

Dietary patterns

Dietary patterns can be derived using two common approaches: *a priori* and *a posteriori*. *A priori* methods assess diet based on prior knowledge and scientific evidence⁽⁴²⁾, while *a posteriori* methods use statistical approaches to provide information about existing dietary patterns within the population⁽⁴²⁾. Cluster analysis is one of the most commonly used methods for deriving dietary patterns, which combines individuals into non-overlapping groups based on similarity of dietary intakes^(43,44). Cluster analysis also provides a simple way of examining changes in dietary patterns over time⁽¹⁰⁾ and allows for the identification of dietary patterns without any pre-defined criteria⁽⁴⁵⁾. Cluster analysis, in comparison with the *a priori* approach, has previously been used to identify relationships between dietary patterns and central obesity⁽⁴⁶⁾ and was thus the method of choice for this research. Dietary patterns were determined using *k*-means cluster analysis⁽⁴⁴⁾. Fibre, Na, Ca, Fe, portions of fruit, portions of vegetables, number of unhealthy snacks, fat percentage, carbohydrate percentage, protein percentage, sugar percentage, starch percentage and saturated fat percentage were selected as cluster variables. Energy was adjusted prior to analysis, and all macronutrients were expressed as a percentage of total energy consumed⁽²⁶⁾.

Statistical analysis

Data were analysed using IBM SPSS (version 22.0 for Windows). Children absent from school on the day of testing were recorded as missing values (*n* 15) and excluded from the analysis. A *k*-means cluster analysis derived two dietary patterns for each age group. Variables were standardised prior to the *k*-means cluster analysis being performed and analyses were run for 2–4 clusters. The *k*-means algorithm converged after three and four iterations for 6-year-olds and seven and three iterations for 10-year-olds. The best cluster solution was chosen based on the size of the clusters. ANOVA was then used to explore the differences in nutritional knowledge, BP, BMI, WHtR and cardiorespiratory fitness with respect to dietary patterns.

Changes in dietary patterns over the 2-year period were analysed with χ^2 analyses.

A multivariate ANOVA was conducted to explore differences in results of BMI, WHtR, BP, cardiorespiratory fitness and nutritional knowledge from baseline to post-intervention with respect to dietary patterns and treatment groups (intervention or control). The results are complemented by corresponding measures of effect sizes (standardised mean difference) using the partial eta-squared (η^2) classification (small $0.01 \leq \eta^2 < 0.06$; medium $0.06 \leq \eta^2 < 0.14$; large $\eta^2 \geq 0.14$)⁽⁴⁷⁾ and observed power. Shapiro–Wilk test, Box’s test and Levene’s test were used as criteria for satisfying the assumptions of measurements being normally distributed, covariance and homogeneity of variances, respectively. All statistical testing was performed using a 5% level of significance.

Results

Baseline anthropometric and dietary intake data

Descriptive statistics of baseline data has previously been reported⁽²⁷⁾ and revealed that 84.2% of children were normal weight, 10.9% were overweight, 4% were obese and 1% were underweight. A total of 69% of 6-year-old children had normal BP, 18% had high-normal and 12% had high BP, respectively, whereas 56.9% of 10-year-olds had normal BP, 27.5% had high-normal and 15.7% had high BP. Intakes of Ca, Fe and fibre were below recommended levels and Na and saturated fat intakes were higher than recommended⁽²⁷⁾. As previous research⁽²⁷⁾ found no statistical differences between sexes, results for boys only and girls only were not included in this manuscript’s statistical inference.

Baseline dietary patterns and differences

K-means clustering derived clusters based on the selected cluster variables. Due to the small sample size, the two-cluster solution was found to give the best fit. At baseline, these clusters for 6-year-olds were labelled as: (1) unhealthy (*n* 19) and (2) nutrient-dense (*n* 12) (see Fig. 1). Fig. 2 shows dietary patterns at baseline for 10-year-olds, which were labelled as: (1) processed (*n* 22) and (2) Western diet (*n* 19).

For 6-year-olds, the ‘unhealthy’ cluster had higher intakes of fat and saturated fat, while the ‘nutrient-dense’ cluster had higher intakes of fruit, vegetables and fibre. For 10-year-olds, the ‘processed’ cluster had higher intakes of carbohydrates and starch, while the ‘Western diet’ had higher intakes of fat, saturated fat and Na.

A one-way multivariate ANOVA showed there were no statistical differences for nutritional knowledge, BMI, WHtR, cardiorespiratory fitness and BP combined with respect to dietary patterns for 6-year-olds ($P=0.924$) and 10-year-olds ($P=0.997$). Tables 1 and 2 present the results to the one-way ANOVAs undertaken for the individual variables (nutritional knowledge, BMI, WHtR, cardiorespiratory fitness and BP) for 6- and 10-year-olds, respectively, and there were no statistical differences ($P>0.05$). The corresponding effect size and observed power statistics to the mean differences are also presented in Tables 1 and 2.






Fig. 1. Baseline dietary patterns (DP) for 6-year-olds. Dietary intake: ■ fibre; ■ sodium; ■ calcium; ■ iron; ■ portions of fruits; ■ portions of vegetables; ■ total fruits and vegetables; ■ number of unhealthy snacks; ■ fat; ■ cholesterol; ■ protein; ■ sugars; ■ starch; ■ saturated fat.




Fig. 2. Baseline dietary patterns (DP) for 10-year-olds. Dietary intake: ■ fibre; ■ sodium; ■ calcium; ■ iron; ■ portions of fruit; ■ portions of vegetables; ■ total fruits and vegetables; ■ number of unhealthy snacks; ■ fat; ■ cholesterol; ■ protein; ■ sugars; ■ starch; ■ saturated fat.

Post-intervention dietary patterns and differences

At post-intervention, clusters were labelled for 6-year-olds as: (1) plant-based and (2) processed foods (see Fig. 3) and for 10-year-olds as: (1) nutrient-dense and (2) unhealthy (see Fig. 4). For 6-year-olds, the ‘plant-based’ cluster had higher intakes of fruit and vegetables, while the ‘processed’ cluster had higher intakes of carbohydrates and starch. For 10-year-olds, the ‘nutrient-dense’ cluster had higher intakes of fibre, fruit and vegetables, while the ‘unhealthy’ cluster had higher intakes of fat, saturated fat protein and Na.

A two-way multivariate ANOVA showed there were no statistical differences for mean nutritional knowledge, BMI, WHtR, cardiorespiratory fitness and BP combined based on the interaction effect between dietary patterns and intervention/control group for 6-year-olds ($P=0.162$) and 10-year-olds ($P=0.746$). Tables 3 and 4 present the results to the following two-way ANOVAs for the individual variables (nutritional knowledge, BMI, WHtR, cardiorespiratory fitness and BP) for 6- and 10-year-olds, respectively. The corresponding effect size and observed power statistics to the mean differences are also presented in Tables 3 and 4.

Changes in dietary patterns following intervention

A χ^2 analysis revealed there was no significant association in dietary patterns for 6- ($P=0.661$) and 10-year-olds ($P=0.738$) at baseline and post-intervention (see Table 5). Baseline and percentage change (baseline to post-intervention) descriptive statistics on individual foods included in the *k*-means cluster analysis are presented in Tables 6 and 7.

Discussion

Identification of the overall combination of foods consumed is essential to understanding the relationships between diet and disease⁽⁴⁸⁾. The aim of the present study was to identify dietary patterns and investigate their relationship with nutritional knowledge, cardiorespiratory fitness, BP, BMI and WHtR and to examine the effect of the intervention on dietary patterns.



The present study identified two dietary patterns at baseline for 6-year-old (unhealthy and nutrient-dense) and 10-year-old (processed and Western) Irish children. Dietary patterns derived post-intervention were (1) plant-based and (2) processed foods for 6-year-olds and (1) nutrient-dense and (2) unhealthy for 10-year-olds. Although not statistically significant and the effect sizes being low, the positive change in dietary patterns towards

Table 1. Baseline differences for 6-year-olds in markers of health and nutritional knowledge of Irish primary schoolchildren with respect to dietary patterns (*n* 34)

Marker of health	Unhealthy (<i>n</i> 22)	Nutrient-dense (<i>n</i> 12)	<i>P</i>	Effect size (η^2)	95 % CI	Power
BMI (kg/m ²)	16.4	16.0	0.411	0.023	0.000, 0.201	0.127
Systolic blood pressure (mmHg)	102.5	101.6	0.780	0.003	0.000, 0.099	0.059
Diastolic blood pressure (mmHg)	62.0	58.9	0.401	0.024	0.000, 0.204	0.131
Waist-to-height ratio (cm/cm)	0.5	0.4	0.349	0.030	0.000, 0.216	0.151
Run time (s)	202.6	207.8	0.553	0.012	0.000, 0.173	0.089
Nutritional knowledge	9.1	9.6	0.561	0.012	0.000, 0.172	0.088

Table 2. Baseline differences for 10-year-olds in markers of health and nutritional knowledge of Irish primary schoolchildren with respect to dietary patterns (n46)

Marker of health	Processed (n 26)	Western diet (n 20)	P	Effect size (η^2)	95 % CI	Power
BMI (kg/m ²)	17.5	17.3	0.844	0.001	0.000, 0.039	0.054
Systolic blood pressure (mmHg)	105.9	105.9	0.992	<0.0005	0.000, 0.000	0.050
Diastolic blood pressure (mmHg)	64.6	64.4	0.944	<0.0005	0.000, 0.005	0.051
Waist-to-height ratio (cm/cm)	0.4	0.4	0.751	0.003	0.000, 0.097	0.061
Run time (s)	171.0	165.9	0.496	0.012	0.000, 0.145	0.103
Nutritional knowledge	12.5	12.5	0.956	<0.000	0.000, 0.003	0.050


Fig. 3. Post-intervention dietary patterns (DP) for 6-year-olds. Dietary intake: ■, fibre; ■, sodium; ■, calcium; ■, iron; ■, fruits; ■, vegetables; ■, fruits and vegetables; ■, unhealthy snacks; ■, fat; ■, cholesterol; ■, protein; ■, sugars; ■, starch; ■, saturated fat.

Fig. 4. Post-intervention dietary patterns (DP) for 10-year-olds. Dietary intake: ■, fibre; ■, sodium; ■, calcium; ■, iron; ■, fruits; ■, vegetables; ■, fruits and vegetables; ■, unhealthy snacks; ■, fat; ■, cholesterol; ■, protein; ■, sugars; ■, starch; ■, saturated fat.

more nutrient-dense diets for 10-year-olds may be indicative of a healthier trend emerging, which could have substantial clinical implications if proved significant in a larger sample size and/or longer intervention delivery time.

Previous research examining the overall diets of children identified 'unhealthy' dietary patterns to consist of high intakes of processed foods, low intakes of fibre and high intakes of Na, fat and refined carbohydrates^(17,49–51). Contrastingly, 'healthy' dietary patterns in the literature are characterised by high intakes of fruit and vegetables, milk/yogurt, eggs and legumes^(17,51,52). O'Brien *et al.*⁽¹⁴⁾ identified two patterns in Irish children aged 7–13 years and termed these as 'more healthful' or 'less healthful'. These clusters are similar to the two clusters identified for 6-year-olds in the present study.

Some evidence suggests that 'unhealthy' dietary patterns are associated with CVD risk factors^(4,5), but contrary to these findings, the present study found no significant relationship between dietary pattern and nutritional knowledge, BMI, BP, WHtR or cardiorespiratory fitness. However, evidence for any association between dietary pattern and markers of health is mixed^(4,11,18,19,53). In a study of children from the UK, a 'Health Aware' dietary pattern was linked to decreased fat mass gain in girls between the ages of 9 and 11 years⁽¹⁸⁾. Craig *et al.*⁽¹¹⁾ also found that 'healthier' dietary patterns were associated with less screen time in Scottish for 5- to 11-year-olds. This is supported by a more recent review by Akbarzadeh *et al.*⁽⁵³⁾ who found diets with a higher healthy eating index to be inversely associated with the metabolic syndrome in youth, whereas Western-type diets (diets that score lower on the healthy eating index) were positively associated with the condition. A systematic review by Rocha *et al.*⁽⁴⁾ also found a positive association between 'unhealthy' dietary patterns and cardiometabolic alterations in children and adolescents. Similar to the present study's findings, however, Craig *et al.*⁽¹¹⁾ found little association between dietary patterns and overweight and obesity, and the National Children's Food Survey⁽⁵⁴⁾ found no evidence that overweight or obese children have unhealthy dietary patterns. Saeedi *et al.*⁽¹⁹⁾ also found no significant associations between dietary patterns and markers of cardiovascular health in 9-year-old New Zealand children. The conflicting evidence may be due to differences in sample size, methods used to analyse dietary patterns, duration of the study and the age of participants, which may all effect the ability to detect an association between dietary patterns and health markers. Furthermore, associations between dietary patterns in childhood and health markers may not become clear until later in life⁽⁵⁵⁾.



Table 3. Intervention effect on dietary pattern for 6-year-olds in relation to nutritional knowledge and markers of health (*n* 31)

Marker of health	Plant-based (<i>n</i> 10)		Processed foods (<i>n</i> 21)		<i>P</i>	Effect size (η^2)	95 % CI	Power
	Intervention	Control	Intervention	Control				
BMI (kg/m ²)	16.1	15.7	17.1	18.0	0.500	0.023	0.000, 0.241	0.100
Systolic blood pressure (mmHg)	105.2	105.1	104.5	95.3	0.181	0.088	0.000, 0.341	0.262
Diastolic blood pressure (mmHg)	56.6	60.4	56.8	60.3	0.966	<0.0005	0.000, 0.004	0.050
Waist-to-height ratio (cm/cm)	0.4	0.4	0.4	0.4	0.688	0.008	0.000, 0.195	0.067
Run time (s)	162.8	178.2	180.9	175.3	0.342	0.045	0.000, 0.282	0.153
Nutritional knowledge	12.8	11.4	12.0	12.0	0.412	0.034	0.000, 0.263	0.126

Table 4. Intervention effect on dietary pattern for 10-year-olds in relation to nutritional knowledge and markers of health (*n* 38)

Marker of health	Nutrient-dense (<i>n</i> 20)		Unhealthy (<i>n</i> 18)		<i>P</i>	Effect size (η^2)	95 % CI	Power
	Intervention	Control	Intervention	Control				
BMI (kg/m ²)	16.0	18.1	18.2	20.5	0.916	<0.0005	0.000, 0.015	0.051
Systolic blood pressure (mmHg)	110.3	103.8	112.8	105.9	0.945	<0.0005	0.000, 0.007	0.051
Diastolic blood pressure (mmHg)	64.9	60.5	63.4	58.1	0.985	<0.0005	0.000, 0.000	0.050
Waist-to-height ratio (cm/cm)	0.4	0.4	0.4	0.4	0.252	0.044	0.000, 0.236	0.205
Run time (s)	148.3	160.2	153.0	164.0	0.952	<0.0005	0.000, 0.005	0.05
Nutritional knowledge	14.0	13.4	13.6	13.0	0.950	<0.0005	0.000, 0.005	0.05

Table 5. Cluster solution for 6- and 10-year-olds, baseline and post-intervention*

Six-year-olds	Cluster post-intervention		Total
	Plant-based	Processed foods	
Cluster baseline			
Unhealthy		13	17
Nutrient dense	3	6	9
Total	7	19	26
Ten-year-olds	Cluster post-intervention		Total
	Nutrient-dense	Unhealthy	
Cluster baseline			
Processed	10	7	17
Western diet	9	9	18
Total	19	16	35

* Table includes children who were at both baseline and post-intervention, that is, paired data. Tables 1–4 are presenting results from baseline and post-intervention independently.

Strengths and limitations

A significant strength of the present study is the longitudinal design and the use of a 4-d food diary to investigate dietary intake. Furthermore, examining dietary patterns, as opposed to individual nutrients, is a useful method for examining diet and the association with health markers. This was the first study in Ireland known to investigate the longitudinal association between dietary patterns and obesity-related outcomes in children and to utilise dietary pattern analysis to assess a health promotion intervention in children.

Limitations of the present study include subjective decision-making in determination of the number of clusters and how the clusters were labelled. The dietary patterns identified in the present study, however, were similar to previous research⁽¹⁴⁾.

In addition, the food diaries were self-reported; therefore, some misreporting and non-reporting are likely to have occurred⁽⁵⁶⁾. As reporting error is a common limitation associated with the direct measurement of dietary intake in children⁽⁵⁷⁾, data should be evaluated with caution. Debriefing with the researcher at the end of diary completion may have also introduced reporting bias of dietary intake⁽⁵⁸⁾. The small sample size also made it more difficult to definitively assess dietary patterns and their association with health markers. As the intervention was only implemented in one intervention school, and only 6- and 10-year-olds were evaluated, the clinical implications of the results are limited. However, the study's purpose was exploratory in nature to help inform further research, and the 6- and 10-year-old age groups were selected on the basis that they mark sensitive periods of growth for the child (mid-childhood and early adolescence)^(59,60). A potential limitation of the study could be that children may have changed their dietary intake as a result of increasing independence as they near adolescence. However, as these children were all still in primary school at the time of follow-up, these changes may not be substantial, since increasing independence usually coincides with the commencement of secondary school in Ireland.

Public health implications

Project Spraoi is based on Project Energize, which has been implemented in New Zealand since 2004, and proven to be successful and cost-effective⁽⁶¹⁾. Initial findings from the Project Spraoi research⁽²⁷⁾ indicate that the intervention has the potential to improve the health of children in Ireland, which supports recent policy⁽⁶²⁾. Preventing children from becoming overweight and obese is the best approach and this is the focus of Project Spraoi, which aims to make healthy eating and increased PA a daily habit. Although some of the findings were not statistically

Table 6. Baseline measurements and percentage change (baseline to post-intervention) in individual foods for 6-year-olds (Mean values and standard deviations)

Individual foods	Intervention (n 23 at baseline, n 18 at post-intervention)		Control (n 11 at baseline, n 8 at post-intervention)	
	Mean	SD	Mean	SD
Fibre (g)				
Baseline	11.17	3.68	11.35	3.03
% Change	-25.1	42.1	5.62	28.26
Na (mg)				
Baseline	1446.43	333.09	1353.82	392.23
% Change	-0.92	29.93	-5.33	31.94
Ca (mg)				
Baseline	716.35	197.77	753.82	278.98
% Change	-10.99	41.03	-8.49	25.25
Fe (mg)				
Baseline	6.9	1.77	7.33	1.83
% Change	-21.48	37.12	-20.94	48.91
Fruit (servings/d)				
Baseline	1.47	1.01	1.11	0.42
% Change	-27.48	102.26	-6.64	90.49
Vegetables (servings/d)				
Baseline	1.11	0.8	0.78	0.40
% Change	29.35	58.78	-63.29	92.58
Total fruits and vegetables (servings/d)				
Baseline	2.58	1.69	1.99	0.74
% Change	0.15	65.07	-22.06	51.53
No. of unhealthy snacks (servings/d)				
Baseline	1.24	0.72	1.22	0.72
% Change	-32.98	107.83	1.03	41.49
Fat (% total energy)				
Baseline	34.47	4.51	35.85	4.15
% Change	5.88	13.21	1.96	16.57
Carbohydrate (% total energy)				
Baseline	48.79	4.64	46.52	5.52
% Change	-3.54	9.39	-3.42	19.37
Protein (% total energy)				
Baseline	15.05	2.03	15.92	2.57
% Change	-5.99	12.63	-2.49	19.86
Sugars (% total energy)				
Baseline	21.88	5.87	18.65	4.11
% Change	11.43	35.12	17.2	32.03
Starch (% total energy)				
Baseline	26.9	3.9	27.85	5.53
% Change	-19.97	22.63	-6.63	22.7
Saturated fat (% total energy)				
Baseline	16.34	2.69	15.37	3.19
% Change	11.79	20.72	-3.59	24.13

Table 7. Baseline measurements and percentage change (baseline to post-intervention) in individual foods for 10-year-olds (Mean values and standard deviations)

Individual foods	Intervention (n 22 at baseline, n 18 at post-intervention)		Control (n 24 at baseline, n 18 at post-intervention)	
	Mean	SD	Mean	SD
Fibre (g)				
Baseline	10.88	2.34	11.05	2.83
% Change	-29.83	39.95	-20.9	38.87
Na (mg)				
Baseline	1605.59	461.15	1338.54	305.14
% Change	-7.97	20.64	-26.81	78.3
Ca (mg)				
Baseline	660.68	260.29	576.79	145.87
% Change	-34.82	58.57	-8.72	30.75
Fe (mg)				
Baseline	7.18	1.52	6.45	2.00
% Change	-26.16	45.94	-23.24	30.39
Fruits (servings/d)				
Baseline	1.03	0.91	1.04	0.61
% Change	-90.95	177.35	-43.04	89.69
Vegetables (servings/d)				
Baseline	0.82	0.56	0.63	0.39
% Change	-39.01	192.07	-43.85	137.80
Total fruits and vegetables (servings/d)				
Baseline	1.91	1.18	1.68	0.72
% Change	-133.45	435.94	-44.42	60.62
No. of unhealthy snacks (servings/d)				
Baseline	1.62	1.01	1.45	0.82
% Change	-18.65	124.75	-26.29	164.68
Fat (% total energy)				
Baseline	36.44	3.68	34.32	4.41
% Change	3.74	12.01	1.69	9.86
Carbohydrate (% total energy)				
Baseline	48.1	4.64	48.08	4.08
% Change	-0.65	12.24	-1.93	7.66
Protein (% total energy)				
Baseline	13.99	2.1	15.89	2.37
% Change	-11.7	16.6	-2.06	18.21
Sugars (% total energy)				
Baseline	19.65	5.98	19.27	3.26
% Change	-9.64	43.27	6.31	27.14
Starch (% total energy)				
Baseline	28.38	4.38	28.78	3.71
% Change	-0.71	20.28	-6.75	17.09
Saturated fat (% total energy)				
Baseline	15.92	2.17	14.53	3.09
% Change	1.88	23.9	0.01	22.55

significant, there were positive trends emerging for the 10-year-olds and these findings are important given that small cumulative differences may ultimately contribute to significant improvements in dietary intake and markers of health in a public health context. It is widely recognised that childhood obesity is a multi-dimensional issue; however, the WHO⁽⁶³⁾ has highlighted that schools are one of the most important sectors to address obesity. Yet, Ireland has very few school-based preventive strategies or interventions that target overweight and obesity in children. These interventions are important as part of the whole school approach recommended by the Irish Obesity Policy and Action Plan⁽⁶²⁾. From a public health context, therefore, if the small cumulative differences found in this school-based intervention study were explored in more schools, over a longer

period of time, significant improvements in dietary intake and markers of health may be found.

Conclusion

This is the first study in Ireland to examine the relationship between dietary patterns and markers of health in children and to assess the impact of a health promotion intervention on dietary patterns over time. Despite finding no statistically significant evidence to support the relationship between dietary patterns and health markers, previous research has shown that dietary patterns established during childhood may track into adulthood and are therefore important to monitor for health outcomes later in life. While no statistically significant evidence of intervention impact was found on dietary patterns, a positive

trend was emerging among 10-year-olds in the present study. This exploratory study will inform a future study, which aims to have a larger sample size and be delivered to more schools. Additional studies with larger sample sizes that deliver interventions over a longer period of time are warranted to further clarify the role of dietary patterns with markers of health and inform future interventions.

Acknowledgements

The authors would sincerely like to thank the schools and participants for their involvement in and contribution to the present study.

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

T. C., A. L. M. and A. M. designed the research project. A. M. conducted the research and T. C. assisted with data collection. A. M. analysed the data and performed the statistical analysis under the guidance of S. L. The lead author was A. M.; however, all authors contributed to the writing of the article. All authors then reviewed the final article for content.

There are no conflicts of interest.

References

- Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
- Ritchie L, Woodward-Lopez G, Gerstein D, *et al.* (2007) Preventing obesity: what *should* we eat? *Calif Agric* **61**, 112–118.
- Childhood Obesity Research Bulletin. Children's Health, Issue 1. <https://childhoodobesityresearchnetwork.org/files/ChildhoodObesityBulletin2017.docx>
- Rocha NP, Milagres LC, Longo GZ, *et al.* (2017) Association between dietary pattern and cardiometabolic risk in children and adolescents: a systematic review. *J Pediatr (Rio J)* **93**, 214–222.
- Funtikova AN, Benítez-Arciniega AA, Fitó M, *et al.* (2015) Modest validity and fair reproducibility of dietary patterns derived by cluster analysis. *Nutr Res* **35**, 265–268.
- Mikkilä V, Räsänen L, Raitakari OT, *et al.* (2005) Consistent dietary patterns identified from childhood to adulthood: the Cardiovascular Risk in Young Finns Study. *Br J Nutr* **93**, 923–931.
- Ambrosini GL, Huang RC, Mori TA, *et al.* (2010) Dietary patterns and markers for the metabolic syndrome in Australian adolescents. *Nutr Metab Cardiovasc Dis* **20**, 274–283.
- North K & Emmett P (2000) Multivariate analysis of diet among three-year-old children and associations with socio-demographic characteristics. The Avon Longitudinal Study of Pregnancy and Childhood (ALSPAC) Study Team. *Eur J Clin Nutr* **54**, 73–80.
- Rasanen M, Lehtinen JC, Niinikoshi H, *et al.* (2002) Dietary patterns and nutrient intakes of 7-year-old children taking part in an atherosclerosis prevention project in Finland. *J Am Diet Assoc* **4**, 518–524.
- Northstone K, Smith AD, Newby PK, *et al.* (2013) Longitudinal comparisons of dietary patterns derived by cluster analysis in 7- to 13-year-old children. *Br J Nutr* **109**, 2050–2058.
- Craig LC, McNeill G, Macdiarmid JI, *et al.* (2010) Dietary patterns of school-age children in Scotland: association with socio-economic indicators, physical activity and obesity. *Br J Nutr* **103**, 319–334.
- Gubbels J, Kremers S, Goldbohm A, *et al.* (2012) Energy balance-related behavioral patterns in 5-year-old children and the longitudinal association with weight status development in early childhood. *Public Health Nutr* **15**, 1402–1410.
- Shin KO, Oh SY & Park HS (2007) Empirically derived major dietary patterns and their associations with overweight in Korean preschool children. *Br J Nutr* **98**, 416–421.
- O'Brien SA, Feeney EL, Scannell AG, *et al.* (2013) Bitter taste perception and dietary intake patterns in Irish children. *J Nutrigenet Nutrigenomics* **6**, 43–58.
- Walton J, McNulty BA, Nugent AP, *et al.* (2014) Diet, lifestyle and body weight in Irish children: findings from Irish Universities Nutrition Alliance national surveys. *Proc Nutr Soc* **73**, 190–200.
- Slattery ML (2010) Analysis of dietary patterns in epidemiological research. *Appl Physiol Nutr Metab* **35**, 207–210.
- Shang X, Li Y, Liu A, *et al.* (2012) Dietary pattern and its association with the prevalence of obesity and related cardiometabolic risk factors among Chinese children. *PLOS ONE* **7**, e43183.
- Smith AD, Emmett PM, Newby PK, *et al.* (2014) Dietary patterns obtained through principal components analysis: the effect of input variable quantification. *Br J Nutr* **109**, 1881–1891.
- Saeedi P (2017) Dietary patterns, physical fitness, and markers of cardiovascular health in 9–11 year-old Dunedin children. Thesis, Doctor of Philosophy, University of Otago. <http://hdl.handle.net/10523/7720>
- Anderson AS, Porteous L, Foster E, *et al.* (2005) The impact of a school-based nutrition education intervention on dietary intake and cognitive and attitudinal variables relating to fruits and vegetables. *Public Health Nutr* **8**, 650–656.
- Fung C, Kuhlle S, Lu C, *et al.* (2012) From 'best practice' to 'next practice': the effectiveness of school-based health promotion in improving healthy eating and physical activity and preventing childhood obesity. *Int J Behav Nutr Phys Act* **9**, 27.
- Laurence S, Peterken R, Burns C (2007) Fresh Kids: the efficacy of a Health Promoting Schools approach to increasing consumption of fruit and water in Australia. *Health Promot Int* **22**, 218–236.
- Te Velde SJ, Brug J, Wind M, *et al.* (2008) Effects of a comprehensive fruit- and vegetable promoting school-based intervention in three European countries: the Pro Children Study. *Br J Nutr* **99**, 893–903.
- Luepker RV, Perry CL, McKinlay SM, *et al.* (1996) Outcomes of a field trial to improve children's dietary patterns and physical activity: the Child and Adolescent Trial for Cardiovascular Health (CATCH). *JAMA* **275**, 768–776.
- Caballero B, Clay T, Davis SM, *et al.* (2003) Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. *Am J Clin Nutr* **78**, 1030–1038.
- Coppinger T, Lacey S, O'Neill C, *et al.* (2016) 'Project Spraoi': a randomized control trial to improve nutrition and physical activity in school children. *Contemp Clin Trials Commun* **3**, 94–101.
- Merrotsy A, McCarthy AL, Flack J, *et al.* (2018) Project Spraoi: dietary intake, nutritional knowledge, cardiorespiratory fitness and health markers of Irish primary school children. *Int J Child Health Nutr* **7**, 63–73.
- Yang YJ, Kim MK, Hwang SH, *et al.* (2010) Relative validities of 3-day food records and the food frequency questionnaire. *Nutr Res Pract* **4**, 142–148.
- Keane E, Kearney PM, Perry IJ, *et al.* (2014) Diet, physical activity, lifestyle behaviors, and prevalence of childhood obesity in Irish children: the Cork Children's Lifestyle Study Protocol. *JMIR Res Protoc* **3**, e44.
- Foster E, Hawkins A & Adamson A (2010) Young person's food atlas primary. London: Food Standards Agency.



- https://www.fuse.ac.uk/uploads/doc/vid_6325_foodatlasprimary0310.pdf
31. Lyons J & Giltinan M (2013) The Irish Food Portion Sizes Database, First Edition. <http://www.iuna.net>
 32. Gower J, Laurie MS, Moyer-Mileur J, *et al.* (2010) Validity and Reliability of a nutrition knowledge survey for assessment in elementary school children. *J Am Diet Assoc* **110**, 452–456.
 33. Food Safety Authority (2011) Scientific Recommendations for Healthy Eating Guidelines in Ireland. <https://www.fsai.ie/recommendationsforhealthyeatingguidelinesinireland.html>
 34. US Department of Agriculture. MyPyramid for Kids. MyPyramid.gov Web site. <http://www.mypyramid.gov/kids/index.html>
 35. Graham D, Appleton S, Rush E, *et al.* (2008) Increasing activity and improving nutrition through a schools-based programme: Project Energize. 1. design, programme, randomisation and evaluation methodology. *Public Health Nutr* **11**, 1076–1084.
 36. Cole TJ, Freeman JV & Preece MA (1995) Body mass index reference curves for the UK, 1990. *Arch Dis Child* **73**, 25–29.
 37. Cole TJ & Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* **7**, 284–294.
 38. Ashwell M (2009) Obesity risk: importance of the waist-to-height ratio. *Nurs Stand* **23**, 49–54.
 39. Albon HM, Hamlin MJ & Ross JJ (2008) Secular trends and distributional changes in health and fitness performance variables of 10–14-year-old children in New Zealand between 1991 and 2003. *Br J Sports Med* **44**, 263–269.
 40. Hamlin MJ, Fraser M, Lizamore C, *et al.* (2014) Measurement of cardiorespiratory fitness in children from two commonly used field tests after accounting for body fatness and maturity. *J Hum Kinet* **40**, 83–92.
 41. Rush E & Obolonkin V (2014) Waikato 2011 centile charts for assessment of time to run 550 m [Microsoft Excel work-book]. <http://aut.researchgateway.ac.nz/handle/10292/8235>
 42. Ocké MC (2013) Evaluation of methodologies for assessing the overall diet: dietary quality scores and dietary pattern analysis. *Proc Nutr Soc* **72**, 191–199.
 43. Moeller SM, Reedy J, Millen AE, *et al.* (2007) Dietary patterns: challenges and opportunities in dietary patterns research. *J Am Diet Assoc* **107**, 1233–1239.
 44. Newby PK & Tucker KL (2004) Empirically derived eating patterns using factor or cluster analysis: a review. *Nutr Rev* **62**, 177–203.
 45. Pérez-Rodrigo C, Gil Á, González-Gross M, *et al.* (2016) Clustering of dietary patterns, lifestyles, and overweight among Spanish children and adolescents in the ANIBES study. *Nutrients* **8**, 11.
 46. Wadolowska L, Kowalkowska J, Czarnocinska J, *et al.* (2017) Comparing dietary patterns derived by two methods and their associations with obesity in Polish girls aged 13–21 years: The cross-sectional GEBaHealth study. *Perspect Public Health* **137**, 182–189.
 47. Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences*, 2nd ed. Hillside, NJ: Lawrence Erlbaum Associates.
 48. Harrington J, Perry I, Lutomski J, *et al.* (2008) *SLÁN 2007: Survey of Lifestyle, Attitudes and Nutrition in Ireland. Dietary Habits of the Irish Population, Department of Health and Children*. Dublin: The Stationery Office.
 49. Karatzi K, Moschonis G, Barouti AA, *et al.* (2014) Dietary patterns and breakfast consumption in relation to insulin resistance in children. The Healthy Growth Study. *Public Health Nutr* **17**, 2790–2797.
 50. Park SJ, Lee SM, Kim SM, *et al.* (2013) Gender specific effect of major dietary patterns on the metabolic syndrome risk in Korean pre-pubertal children. *Nutr Res Pract* **7**, 139–145.
 51. Romero-Polvo A, Denova-Gutiérrez E & Rivera-Paredes B, *et al.* (2012) Association between dietary patterns and insulin resistance in Mexican children and adolescents. *Ann Nutr Metab* **61**, 142–150.
 52. Mikkilä V, Räsänen L, Raitakari O, *et al.* (2007) Major dietary patterns and cardiovascular risk factors from childhood to adulthood. The Cardiovascular Risk in Young Finns Study. *Br J Nutr* **98**, 218–225.
 53. Akbarzadeh Z, Nourian M, Hovsepian S, *et al.* (2018) Dietary patterns and metabolic syndrome in children and adolescents: a systematic review. *J Pediatr Rev* **6**, 2–13.
 54. Safefood (2011) Body weight and eating habits in 5–12 year old Irish children: the National Children's Food Survey: summary report. Cork: Safefood. <http://www.safefood.eu/SafeFood/media/SafeFoodLibrary/Documents/Publications/Research%20Reports/Bodyweight-pdf.pdf>
 55. Bull CJ & Northstone K (2016) Childhood dietary patterns and cardiovascular risk factors in adolescence: results from the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort. *Public Health Nutr* **19**, 3369–3377.
 56. Poslusna K, Ruprich J, de Vries JHM, *et al.* (2009) Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *Br J Nutr* **101**, S73–S85.
 57. Livingstone M, Robson P & Wallace J (2004) Issues in dietary intake assessment of children and adolescents. *Br J Nutr* **92**, S213–S222.
 58. Bingham S, Cassidy A, Cole T, *et al.* (1995) Validation of weighed records and other methods of dietary assessment using the 24 hour urine nitrogen technique and other biological markers. *Br J Nutr* **73**, 531–550.
 59. Cameron N & Demerath EW (2002) Critical periods in human growth and their relationship to diseases of aging. *Am J Phys Anthropol Suppl* **119**, 159–184.
 60. Graham D, Appleton S, Rush E, *et al.* (2008) Increasing activity and improving nutrition through a schools-based programme: Project Energize. 1. design, programme, randomisation and evaluation methodology. *Public Health Nutr* **11**, 1076–1084.
 61. Rush E, Cairncross C, Williams MH, *et al.* (2016) Project Energize: intervention development and 10 years of progress in preventing childhood obesity. *BMC Res Notes* **9**, 44.
 62. Department of Health (2016) *A Healthy Weight for Ireland 2016–2025, Obesity Policy and Action Plan*. Dublin: The Stationery Office.
 63. World Health Organization (2015) *Draft Final Report of the Commission on Ending Childhood Obesity*. <https://www.who.int/end-childhood-obesity/commission-ending-childhood-obesity-draft-final-report-en.pdf>

