

Transplantation-free Survival after Norwood surgery for Hypoplastic Left Heart Syndrome with Aortic Atresia. A Swedish National Cohort Study. – ERRATUM

Erratum

Cite this article: Öhman A, El-Segaier M, Bergman G, Hanseus K, Malm T, Nilsson B, Pivodic A, Rydberg A, Sonesson S-E, and Mellander M (2020) Transplantation-free Survival after Norwood surgery for Hypoplastic Left Heart Syndrome with Aortic Atresia. A Swedish National Cohort Study. – ERRATUM. *Cardiology in the Young* 30: 361. doi: [10.1017/S1047951120000190](https://doi.org/10.1017/S1047951120000190)

First published online: 4 February 2020

Annika Öhman, Milad El-Segaier, Gunnar Bergman, Katarina Hanseus, Torsten Malm, Boris Nilsson, Aldina Pivodic, Annika Rydberg, Sven-Erik Sonesson and Mats Mellander

DOI: <https://doi.org/10.1017/S1047951119003263> Published online by Cambridge University Press: 10 January 2020

The Publisher apologises that upon publication of this paper the p-values in the upper half of Table 4 have a minus sign which should not have been there. The corrected table is as below

Table 4. The results of the univariable Cox regression analysis to investigate risk factors for death or heart transplantation in the 121 patients who underwent Norwood surgery in the period 1993–2010

Variable	Missing data	Value	Events n (%)	Univariable analysis	
				Hazard Ratio (95% CI)	P-value
Categorical					
Gender	0	Male	35 (42.2)	1.00 (Ref)	0.0018
		Female	26 (68.4)	2.25 (1.35–3.76)	
Birth period	0	1993–2000 (early)	38 (63.3)	1.00 (Ref)	0.010
		2001–2010 (late)	23 (37.7)	0.50 (0.30–0.85)	
Prenatal diagnosis	0	No	51 (51.5)	1.00 (Ref)	0.62
		Yes	10 (45.5)	0.84 (0.43–1.66)	
Premature	2	No	59 (51.8)	1.00 (Ref)	0.70
		Yes	2 (40.0)	0.76 (0.19–3.11)	
Surgical center	0	Gothenburg	37 (53.6)	1.00 (Ref)	0.33
		Lund	24 (46.2)	0.78 (0.46–1.30)	
Extra-cardiac malformation	0	No	58 (50.4)	1.00 (Ref)	0.96
		Yes	3 (50.0)	1.03 (0.32–3.29)	
				The HR (95% CI) for each unit increase of the continuous variable	
Continuous		Unit			P-value
Age at stage I (days)	0	1 day	61 (50.4)	1.01 (0.96–1.06)	0.76
GA (weeks)	2	1 week	61 (51.3)	0.96 (0.83–1.11)	0.61
BW (kg)	1	0.5 kg	61 (50.8)	0.70 (0.55–0.89)	0.0044
		0.5 kg for values ≤ 2.5 kg	61 (50.8)	0.60 (0.41–0.88)	0.0088
		0.5 kg for values > 2.5 kg		0.92 (0.52–1.62)	0.77
BW SD score	1	0.5 SD-score	61 (50.8)	0.82 (0.72–0.94)	0.0033
		0.5 SD-score for values ≤ –0.46	61 (50.8)	0.73 (0.60–0.89)	0.0019
		0.5 SD-score for values > –0.46		1.04 (0.76–1.42)	0.80
BHC (cm)	38	1 cm	46 (55.4)	0.75 (0.61–0.92)	0.0068
BHC SD score	38	0.5 SD-score	46 (55.4)	0.90 (0.78–1.03)	0.13
BHC/BW ratio	38	1 unit in ratio	46 (55.4)	1.38 (1.14–1.66)	0.0010

BHC = birth head circumference; BW = birth weight; SD score = standard deviation score.

© Cambridge University Press 2020.

Reference

Öhman A, El-Segaier M, Bergman G, Hanseus K, Malm T, Nilsson B, . . . Mellander, M. (n.d.). Transplantation-free survival after Norwood surgery for hypoplastic left heart syndrome with aortic atresia: A Swedish national cohort study. *Cardiology in the Young*, 1–8. doi: [10.1017/S1047951119003263](https://doi.org/10.1017/S1047951119003263)