

DISTRIBUTION-FREE CALCULATION OF THE STANDARD ERROR OF CHAIN LADDER RESERVE ESTIMATES

BY THOMAS MACK

Munich Re, Munich

ABSTRACT

A distribution-free formula for the standard error of chain ladder reserve estimates is derived and compared to the results of some parametric methods using a numerical example.

KEYWORDS

Claims reserving; chain ladder; standard error.

1. INTRODUCTION

The chain ladder method is probably the most popular method for estimating IBNR claims reserves. The main reason for this is its simplicity and the fact that it is distribution-free, i.e. that it seems to work with almost no assumptions. On the other hand, it is well-known that chain ladder reserve estimates for the most recent accident years are very sensitive to variations in the data observed. Moreover, in recent years many other claims reserving procedures have been proposed and the results of all these procedures vary widely and also differ more or less from the chain ladder result. Therefore it would be very helpful to know the standard error of the chain ladder reserve estimates as a measure of the uncertainty contained in the data and in order to see whether the difference between the results of the chain ladder method and any other method is significant or not.

Up to now only a few papers on claims reserving have been published which also consider the calculation of the standard error of the reserve estimate: In the papers by TAYLOR/ASHE 1983, ZEHNWIRTH 1985, RENSHAW 1989, CHRISTOFIDES 1990, VERRALL 1990, VERRALL 1991 essentially the same method for the calculation of the standard error is used, namely a least squares regression approach which (with the exception of Taylor/Ashe) is applied to the logarithms of the incremental claims amounts (i.e. assuming a lognormal distribution). Slightly different approaches have been proposed by WRIGHT (1990, via a generalized linear model and the method of scoring) and MACK (1991, using a gamma distribution and maximum likelihood estimation). All methods cited require a rather high amount of programming in order to calculate the many covariances between the parameter estimators.

In the present paper, a very simple formula for the standard error of chain ladder reserve estimates is developed. The decisive step towards this formula was made by SCHNIEPER (1991). In order to calculate the rate for a casualty excess of loss cover he used a mixture of the Bornhuetter-Ferguson technique and the chain ladder method. Within this model he developed an approximation to the standard error of the estimated premium rate using a Taylor series approximation.

The present paper adapts Schnieper's idea to the claims reserving situation and contains the following additional points:

1. The model is specialized for the pure chain ladder case. This makes things easier and also makes it possible to replace the Taylor series approximation with a more exact procedure.
2. An estimate of the process variance is additionally included in the standard error of the reserve estimate. This is necessary here because the claims reserve is a random variable and not a parameter like the net premium (= expected value).
3. Schnieper intuitively claimed that the chain ladder development factors were "not strongly correlated". We prove that they are in fact uncorrelated and that therefore the reserve estimate is unbiased.
4. Besides the standard error for each accident year, a formula for the standard error of the overall reserve estimator is given, too, which takes the correlations between the estimates for the individual accident years into account.

Finally, two numerical examples are given and the results are compared to the results obtained by the aforementioned methods of Taylor/Ashe, Zehn-wirth, Renshaw/Christofides, Verrall and Mack.

2. NOTATIONS AND BASIC RESULTS

Let C_{ik} denote the accumulated total claims amount of accident year i , $1 \leq i \leq I$, either paid or incurred up to development year k , $1 \leq k \leq I$. We consider C_{ik} a random variable of which we have an observation if $i+k \leq I+1$ (run-off triangle). The aim is to estimate the ultimate claims amount C_{iI} and the outstanding claims reserve

$$R_i = C_{iI} - C_{i, I+1-i}$$

for accident year $i = 2, \dots, I$.

The basic chain ladder assumption is that there are development factors $f_1, \dots, f_{I-1} > 0$ with

$$(1) \quad E(C_{i, k+1} | C_{i1}, \dots, C_{ik}) = C_{ik} f_k, \quad 1 \leq i \leq I, \quad 1 \leq k \leq I-1.$$

The chain ladder method consists of estimating the f_k by

$$\hat{f}_k = \sum_{j=1}^{I-k} C_{j, k+1} \bigg/ \sum_{j=1}^{I-k} C_{jk}, \quad 1 \leq k \leq I-1,$$

and the ultimate claims amount C_{iI} by

$$\hat{C}_{iI} = C_{i, I+1-i} \cdot \hat{f}_{I+1-i} \cdots \hat{f}_{I-1},$$

or equivalently the reserve R_i by

$$\hat{R}_i = C_{i, I+1-i} (\hat{f}_{I+1-i} \cdots \hat{f}_{I-1} - 1).$$

Because the chain ladder algorithm does not take into account any dependencies between accident years, we can additionally assume that the variables C_{ik} of different accident years, i.e.

(2) $\{C_{i1}, \dots, C_{iI}\}, \{C_{j1}, \dots, C_{jI}\}, i \neq j$, are independent.

This must be regarded as a further implicit assumption of the chain ladder method. In practise, the independence of the accident years can be distorted by certain calendar year effects like major changes in claims handling or in case reserving.

The following theorem makes it clear that (1) and (2) are indeed the implicit assumptions of the chain ladder method.

Theorem 1: Let $D = \{C_{ik} | i+k \leq I+1\}$ be the set of all data observed so far. Under the assumptions (1) and (2) we have

$$E(C_{iI} | D) = C_{i, I+1-i} f_{I+1-i} \cdots f_{I-1}.$$

Proof: We use the abbreviation

$$E_i(X) = E(X | C_{i1}, \dots, C_{i, I+1-i}).$$

Then (2) and repeated application of (1) yield

$$\begin{aligned} E(C_{iI} | D) &= E_i(C_{iI}) \\ &= E_i(E(C_{iI} | C_{i1}, \dots, C_{i, I-1})) \\ &= E_i(C_{i, I-1} f_{I-1}) \\ &= E_i(C_{i, I-1}) f_{I-1} \\ &= \text{etc.} \\ &= E_i(C_{i, I+1-i}) f_{I+1-i} \cdots f_{I-1} \\ &= C_{i, I+1-i} f_{I+1-i} \cdots f_{I-1}. \end{aligned} \quad \square$$

This theorem shows that the estimator \hat{C}_{iI} has the same form as $E(C_{iI} | D)$ which is the best forecast of C_{iI} based on the observations D . The next theorem shows that estimating $f_{I+1-i} \cdots f_{I-1}$ by $\hat{f}_{I+1-i} \cdots \hat{f}_{I-1}$ is indeed a reasonable procedure.

Theorem 2: Under the assumptions (1) and (2) the estimators $\hat{f}_k, 1 \leq k \leq I-1$, are unbiased and uncorrelated.

Proof: Let

$$B_k = \{C_{ij} | j \leq k, i + j \leq I + 1\}, 1 \leq k \leq I.$$

Then (2) and (1) yield

$$E(C_{i, k+1} | B_k) = E(C_{i, k+1} | C_{i1}, \dots, C_{ik}) = C_{ik} f_k.$$

We therefore have

$$E(\hat{f}_k | B_k) = \sum_{j=1}^{I-k} E(C_{j, k+1} | B_k) \bigg/ \sum_{j=1}^{I-k} C_{jk} = f_k,$$

which immediately gives the unbiasedness

$$E(\hat{f}_k) = E(E(\hat{f}_k | B_k)) = f_k, 1 \leq k \leq I - 1,$$

of the parameter estimates. Also, the \hat{f}_k are uncorrelated because for $j < k$

$$\begin{aligned} E(\hat{f}_j \hat{f}_k) &= E(E(\hat{f}_j \hat{f}_k | B_k)) \\ &= E(\hat{f}_j E(\hat{f}_k | B_k)) \\ &= E(\hat{f}_j) f_k \\ &= E(\hat{f}_j) E(\hat{f}_k). \end{aligned}$$

□

The uncorrelatedness of the \hat{f}_k 's is surprising because \hat{f}_{k-1} and \hat{f}_k depend on the same data $C_{1k} + \dots + C_{I-k, k}$. The foregoing proof of the uncorrelatedness easily extends to arbitrary products of pairwise different \hat{f}_k , i.e. we have

$$E(\hat{f}_{I+1-i} \dots \hat{f}_{I-1}) = f_{I+1-i} \dots f_{I-1},$$

which shows that $\hat{C}_{ii} = C_{i, I+1-i} \hat{f}_{I+1-i} \dots \hat{f}_{I-1}$ is an unbiased estimator of $E(C_{ii} | D) = C_{i, I+1-i} f_{I+1-i} \dots f_{I-1}$. In the same way, the reserve estimator $\hat{R}_i = \hat{C}_{ii} - C_{i, I+1-i}$ is an unbiased estimator of the true reserve $R_i = C_{ii} - C_{i, I+1-i}$.

3. CALCULATION OF MEAN SQUARED ERROR AND STANDARD ERROR

The mean squared error $mse(\hat{C}_{ii})$ of the estimator \hat{C}_{ii} of C_{ii} is defined to be

$$mse(\hat{C}_{ii}) = E((\hat{C}_{ii} - C_{ii})^2 | D)$$

where $D = \{C_{ik} | i + k \leq I + 1\}$ is the set of all data observed so far. Note that we are not using the unconditional mean squared error $E((\hat{C}_{ii} - C_{ii})^2) = E(E((\hat{C}_{ii} - C_{ii})^2 | D))$ as this averages over all possible data D from the underlying distribution. Instead, in practise, we are more interested in the conditional mean squared error of the particular estimated amount \hat{C}_{ii} based on the specific data set D observed and therefore have to use $E((\hat{C}_{ii} - C_{ii})^2 | D)$ which just gives us the average deviation between \hat{C}_{ii} and C_{ii} due to future randomness only.

First, we see that

$$mse(\hat{R}_i) = E((\hat{R}_i - R_i)^2 | D) = E((\hat{C}_{ii} - C_{ii})^2 | D) = mse(\hat{C}_{ii}).$$

Next, because of the general rule $E(X-a)^2 = \text{Var}(X) + (E(X)-a)^2$ we have

$$mse(\hat{C}_{il}) = \text{Var}(C_{il}|D) + (E(C_{il}|D) - \hat{C}_{il})^2$$

which shows that the mean squared error is the sum of the stochastic error (process variance) and of the estimation error.

In order to further calculate the *mse* we need a formula for the variance of C_{ik} . From the fact that \hat{f}_k is the C_{ik} -weighted mean of the individual development factors $C_{i,k+1}/C_{ik}$, $1 \leq i \leq I-k$, we can induce that $\text{Var}(C_{i,k+1}/C_{ik}|C_{i1}, \dots, C_{ik})$ should be inversely proportional to C_{ik} , or equivalently

$$(3) \quad \text{Var}(C_{i,k+1}|C_{i1}, \dots, C_{ik}) = C_{ik} \sigma_k^2, \quad 1 \leq i \leq I, \quad 1 \leq k \leq I-1,$$

with unknown parameters σ_k^2 , $1 \leq k \leq I-1$. This is the variance assumption which is implicitly underlying the chain ladder method.

Later on, we will need an estimator for σ_k^2 . Similarly as for \hat{f}_k it can be shown that

$$\hat{\sigma}_k^2 = \frac{1}{I-k-1} \sum_{i=1}^{I-k} C_{ik} \left(\frac{C_{i,k+1}}{C_{ik}} - \hat{f}_k \right)^2, \quad 1 \leq k \leq I-2.$$

is an unbiased estimator of σ_k^2 , $1 \leq k \leq I-2$. We still lack an estimator for σ_{I-1} . If $\hat{f}_{I-1} = 1$ and if the claims development is believed to be finished after $I-1$ years, we can put $\hat{\sigma}_{I-1} = 0$. If not, we extrapolate the usually exponentially decreasing series $\hat{\sigma}_1, \dots, \hat{\sigma}_{I-3}, \hat{\sigma}_{I-2}$ by one additional member, for instance by loglinear regression or more simply by requiring that

$$\hat{\sigma}_{I-3}/\hat{\sigma}_{I-2} = \hat{\sigma}_{I-2}/\hat{\sigma}_{I-1}$$

holds at least as long as $\hat{\sigma}_{I-3} > \hat{\sigma}_{I-2}$. This last possibility leads to

$$\hat{\sigma}_{I-1}^2 = \min(\hat{\sigma}_{I-2}^4/\hat{\sigma}_{I-3}^2, \min(\hat{\sigma}_{I-3}^2, \hat{\sigma}_{I-2}^2))$$

which has been used in the examples.

Now, we are able to state and prove the main result

Theorem 3: Under the assumptions (1), (2) and (3) the mean squared error $mse(\hat{R}_i)$ can be estimated by

$$\widehat{mse}(\hat{R}_i) = \hat{C}_{il}^2 \sum_{k=I+1-i}^{I-1} \frac{\hat{\sigma}_k^2}{\hat{f}_k^2} \left(\frac{1}{\hat{C}_{ik}} + \frac{1}{\sum_{j=1}^{I-k} C_{jk}} \right)$$

where $\hat{C}_{ik} = C_{i,I+1-i} \hat{f}_{I+1-i} \dots \hat{f}_{k-1}$, $k > I+1-i$, are the estimated values of the future C_{ik} and $\hat{C}_{i,I+1-i} = C_{i,I+1-i}$.

Proof: We use the abbreviations

$$E_i(X) = E(X|C_{i1}, \dots, C_{i, I+1-i}),$$

$$\text{Var}_i(X) = \text{Var}(X|C_{i1}, \dots, C_{i, I+1-i}).$$

We start from

$$\text{mse}(\hat{R}_i) = \text{Var}(C_{i1}|D) + (E(C_{i1}|D) - \hat{C}_{i1})^2.$$

Repeated application of the basic chain ladder assumption (1) and of the above variance assumption (3) yields for the first term of $\text{mse}(\hat{R}_i)$

$$\begin{aligned} \text{Var}(C_{i1}|D) &= \text{Var}_i(C_{i1}) \\ &= E_i(\text{Var}(C_{i1}|C_{i1}, \dots, C_{i, I-1})) + \\ &\quad + \text{Var}_i(E(C_{i1}|C_{i1}, \dots, C_{i, I-1})) \\ &= E_i(C_{i, I-1})\sigma_{I-1}^2 + \text{Var}_i(C_{i, I-1})f_{I-1}^2 \\ &= E_i(C_{i, I-2})f_{I-2}\sigma_{I-1}^2 + E_i(C_{i, I-2})\sigma_{I-2}^2 f_{I-1}^2 + \\ &\quad + \text{Var}_i(C_{i, I-2})f_{I-2}^2 f_{I-1}^2 \\ &= \text{etc.} \\ &= C_{i, I+1-i} \sum_{k=I+1-i}^{I-1} f_{I+1-i} \dots f_{k-1} \sigma_k^2 f_{k+1}^2 \dots f_{I-1}^2 \end{aligned}$$

because of $\text{Var}_i(C_{i, I+1-i}) = 0$.

Due to Theorem 1 we obtain for the second term of $\text{mse}(\hat{R}_i)$

$$(*) \quad (E(C_{i1}|D) - \hat{C}_{i1})^2 = C_{i, I+1-i}^2 (f_{I+1-i} \dots f_{I-1} - \hat{f}_{I+1-i} \dots \hat{f}_{I-1})^2.$$

In practice, we must find estimators for these two terms of $\text{mse}(\hat{R}_i)$. For the first term this will be done by replacing the unknown parameters f_k and σ_k^2 with their estimators \hat{f}_k and $\hat{\sigma}_k^2$, i.e. we estimate $\text{Var}(C_{i1}|D)$ by

$$\begin{aligned} &C_{i, I+1-i} \left(\sum_{k=I+1-i}^{I-1} \hat{f}_{I+1-i} \dots \hat{f}_{k-1} \sigma_k^2 \hat{f}_{k+1}^2 \dots \hat{f}_{I-1}^2 \right) \\ &= \hat{C}_{i1}^2 \sum_{k=I+1-i}^{I-1} \frac{\hat{\sigma}_k^2 \hat{f}_k^2}{\hat{C}_{ik}} \end{aligned}$$

where we have used the notation \hat{C}_{ik} introduced in the theorem.

But in the second term (*) of $\text{mse}(\hat{R}_i)$ we can not simply replace f_k with \hat{f}_k because this would yield 0. We therefore use a different approach. We can write

$$\begin{aligned} F &= f_{I+1-i} \dots f_{I-1} - \hat{f}_{I+1-i} \dots \hat{f}_{I-1} \\ &= S_{I+1-i} + \dots + S_{I-1} \end{aligned}$$

with

$$S_k = \hat{f}_{I+1-i} \dots \hat{f}_{k-1} (f_k - \hat{f}_k) f_{k+1} \dots f_{I-1}$$

and therefore

$$\begin{aligned}
 F^2 &= (S_{I+1-i} + \dots + S_{I-1})^2 \\
 &= \sum_{k=I+1-i}^{I-1} S_k^2 + 2 \sum_{j < k} S_j S_k.
 \end{aligned}$$

Now we replace S_k^2 with $E(S_k^2|B_k)$ and $S_j S_k, j < k$, with $E(S_j S_k|B_k)$. This means that we approximate S_k^2 and $S_j S_k$ by averaging over as little data as possible such that as many values C_{ik} as possible from the observed data are kept fixed. Because of $E(f_k - \hat{f}_k|B_k) = 0$ (see the proof of Theorem 2) we obtain $E(S_j S_k|B_k) = 0$ for $j < k$. Because of

$$\begin{aligned}
 E((f_k - \hat{f}_k)^2|B_k) &= \text{Var}(\hat{f}_k|B_k) \\
 &= \sum_{j=1}^{I-k} \text{Var}(C_{j, k+1}|B_k) \left| \left(\sum_{j=1}^{I-k} C_{jk} \right)^2 \right. \\
 &= \sigma_k^2 \left| \sum_{j=1}^{I-k} C_{jk} \right.
 \end{aligned}$$

we obtain

$$E(S_k^2|B_k) = \hat{f}_{I+1-i}^2 \cdot \dots \cdot \hat{f}_{k-1}^2 \cdot \sigma_k^2 \cdot f_{k+1}^2 \cdot \dots \cdot f_{I-1}^2 \left| \sum_{i=1}^{I-k} C_{ik} \right.$$

Taken together, we replace $F^2 = (\sum S_k)^2$ with $\sum_k E(S_k^2|B_k)$ and because all terms of this sum are positive we now can replace all unknown parameters f_k, σ_k^2 with their unbiased estimators $\hat{f}_k, \hat{\sigma}_k^2$. Altogether, we estimate $F^2 = (f_{I+1-i} \cdot \dots \cdot f_{I-1} - \hat{f}_{I+1-i} \cdot \dots \cdot \hat{f}_{I-1})^2$ by

$$\begin{aligned}
 &\sum_{k=I+1-i}^{I-1} \left(\hat{f}_{I+1-i}^2 \cdot \dots \cdot \hat{f}_{k-1}^2 \cdot \hat{\sigma}_k^2 \cdot \hat{f}_{k+1}^2 \cdot \dots \cdot \hat{f}_{I-1}^2 \left| \sum_{j=1}^{I-k} C_{jk} \right. \right) \\
 &= \hat{f}_{I+1-i}^2 \cdot \dots \cdot \hat{f}_{I-1}^2 \sum_{k=I+1-i}^{I-1} \frac{\hat{\sigma}_k^2 |\hat{f}_k^2}{\sum_{j=1}^{I-k} C_{jk}}
 \end{aligned}$$

This finally leads to the estimator stated in the theorem. □

The square root s.e. (\hat{R}_i) of an estimator of the mean squared error is defined to be the standard error of \hat{R}_i .

Often the standard error of the overall reserve estimate $\hat{R} = \hat{R}_2 + \dots + \hat{R}_I$ is of interest, too. In this case we cannot simply add together the values of (s.e. (\hat{R}_i))², $2 \leq i \leq I$, because they are correlated via the common estimators \hat{f}_k and $\hat{\sigma}_k$. We therefore proceed as before and obtain:

Corollary: With the assumptions and notations of Theorem 3 the mean squared error of the overall reserve estimate $\hat{R} = \hat{R}_2 + \dots + \hat{R}_I$ can be estimated by

$$\widehat{mse}(\hat{R}) = \sum_{i=2}^I \left\{ (\text{s.e.}(\hat{R}_i))^2 + \hat{C}_{iI} \left(\sum_{j=i+1}^I \hat{C}_{jI} \right) \sum_{k=I+1-i}^{I-1} \frac{2\hat{\sigma}_k^2 / \hat{f}_k^2}{\sum_{n=1}^{I-k} C_{nk}} \right\}$$

Proof: We have

$$\begin{aligned} mse \left(\sum_{i=2}^I \hat{R}_i \right) &= E \left(\left(\sum_{i=2}^I \hat{R}_i - \sum_{i=2}^I R_i \right)^2 \middle| D \right) \\ &= E \left(\left(\sum_{i=2}^I \hat{C}_{iI} - \sum_{i=2}^I C_{iI} \right)^2 \middle| D \right) \\ &= \text{Var} \left(\sum_{i=2}^I C_{iI} \middle| D \right) + \left(E \left(\sum_{i=2}^I C_{iI} \middle| D \right) - \sum_{i=2}^I \hat{C}_{iI} \right)^2. \end{aligned}$$

The independence of the accident years yields

$$\text{Var} \left(\sum_{i=2}^I C_{iI} \middle| D \right) = \sum_{i=2}^I \text{Var} (C_{iI} \middle| D),$$

whose summands have already been calculated in the proof of Theorem 3. Furthermore

$$\begin{aligned} \left(E \left(\sum_{i=2}^I C_{iI} \middle| D \right) - \sum_{i=2}^I \hat{C}_{iI} \right)^2 &= \left(\sum_{i=2}^I (E(C_{iI} \middle| D) - \hat{C}_{iI}) \right)^2 \\ &= \sum_{i,j} (E(C_{iI} \middle| D) - \hat{C}_{iI}) \cdot (E(C_{jI} \middle| D) - \hat{C}_{jI}) \\ &= \sum_{i,j} C_{i,I+1-i} C_{j,I+1-j} F_i F_j \end{aligned}$$

with

$$F_i = f_{I+1-i} \cdots f_{I-1} - \hat{f}_{I+1-i} \cdots \hat{f}_{I-1}.$$

Observing

$$mse(\hat{R}_i) = \text{Var} (C_{iI} \middle| D) + (C_{i,I+1-i} F_i)^2$$

(cf. (*) in the proof of theorem 3) we see that

$$mse \left(\sum_{i=2}^I \hat{R}_i \right) = \sum_{i=2}^I mse(\hat{R}_i) + \sum_{2 \leq i < j \leq I} 2 \cdot C_{i,I+1-i} C_{j,I+1-j} F_i F_j.$$

An analogous procedure as for F^2 in the above proof yields for $F_i F_j$, $i < j$, the estimator

$$\sum_{k=i+1}^{j-1} \hat{f}_{i+1-j} \cdots \hat{f}_{i-i} \hat{f}_{i+1-i}^2 \cdots \hat{f}_{k-1}^2 \hat{\sigma}_k^2 \hat{f}_{k+1}^2 \cdots \hat{f}_{j-1}^2 \left| \sum_{n=1}^{j-k} C_{nk} \right.$$

This completes the proof. □

4. EXAMPLES

In the first example we use the TAYLOR/ASHE (1983) data, which were also used by VERRALL (1990, 1991).

TABLE 1
RUN-OFF TRIANGLE (ACCUMULATED FIGURES)

<i>i</i>	C_{i1}	C_{i2}	C_{i3}	C_{i4}	C_{i5}	C_{i6}	C_{i7}	C_{i8}	C_{i9}	C_{i10}
1	357848	1124788	1735330	2218270	2745596	3319994	3466336	3606286	3833515	3901463
2	352118	1236139	2170033	3353322	3799067	4120063	4647867	4914039	5339085	
3	290507	1292306	2218525	3235179	3985995	4132918	4628910	4909315		
4	310608	1418858	2195047	3757447	4029929	4381982	4588268			
5	443160	1136350	2128333	2897821	3402672	3873311				
6	396132	1333217	2180715	2985752	3691712					
7	440832	1288463	2419861	3483130						
8	359480	1421128	2864498							
9	376686	1363294								
10	344014									

This yields the following parameter estimates ($k = 1, \dots, 9$):

\hat{f}_k : 3.49, 1.75, 1.46, 1.174, 1.104, 1.086, 1.054, 1.077, 1.018
 $\hat{\sigma}_k^2/1000$: 160, 37.7, 42.0, 15.2, 13.7, 8.19, 0.447, 1.15, 0.477

TABLE 2
ESTIMATED RESERVES \hat{R}_i IN 1000 s

	Chain ladder	Verrall 1991	Renshaw Christofides	Zehnwirth	Mack 1991	Taylor Ashe
<i>i</i> = 2	95	96	111	109	93	298
<i>i</i> = 3	470	439	482	473	447	600
<i>i</i> = 4	710	608	661	648	611	745
<i>i</i> = 5	985	1011	1091	1069	992	1077
<i>i</i> = 6	1419	1423	1531	1500	1453	1788
<i>i</i> = 7	2178	2150	2311	2265	2186	2879
<i>i</i> = 8	3920	3529	3807	3731	3665	4221
<i>i</i> = 9	4279	4056	4452	4364	4122	4866
<i>i</i> = 10	4626	4340	5066	4965	4516	5827
overall	18681	16652	19512	19124	18085	22301

TABLE 3
STANDARD ERROR IN % OF \hat{R}_i

	Chain ladder	Verrall 1991	Renshaw Christofides	Zehnwirth	Mack 1991	Taylor Ashe
$i = 2$	80 %	49 %	54 %	49 %	40 %	27 %
$i = 3$	26 %	37 %	39 %	35 %	30 %	20 %
$i = 4$	19 %	30 %	32 %	29 %	24 %	18 %
$i = 5$	27 %	27 %	28 %	25 %	21 %	16 %
$i = 6$	29 %	25 %	26 %	24 %	20 %	16 %
$i = 7$	26 %	25 %	26 %	24 %	20 %	14 %
$i = 8$	22 %	27 %	28 %	26 %	21 %	14 %
$i = 9$	23 %	30 %	31 %	30 %	24 %	14 %
$i = 10$	29 %	38 %	40 %	39 %	31 %	14 %
overall	13 %	15 %	16 %	16 %		9 %

Comments:

The results for 'Taylor/Ashe' and 'Verrall 1991' have been taken from these papers. Taylor/Ashe produced much lower standard errors than the other methods. This is due to the fact that their reserve estimates employed only 6 parameters (as compared to 19 of the other methods) and that they additionally used the information on the numbers of claims finalized.

Renshaw and Christofides describe the same loglinear regression method which is also identical to Verrall's (1990) Bayesian approach without any prior information. Therefore the results for 'Renshaw/Christofides' have been taken from VERRALL (1990), Table 2.

The results for 'Zehnwirth' have been obtained by using his ICRFS software package version 6.1 employing one of his fixed parameter development factor models which he calls 'chain ladder model'. We have used it without any further adjustment. It should be remarked that this is not what Zehnwirth intends, as his software package is a modelling framework and any initial model should be further adjusted interactively with the help of the indications and plots given by the program. Without any further adjustment this 'chain ladder model' is identical to the Renshaw/Christofides model, i.e. it is a loglinearized approximation of the usual chain ladder model. The fact that it leads to slightly lower results is attributable to using a different estimator for the model variance.

The results for 'Mack 1991' have been obtained according to a previous paper (MACK (1991)) of the author but additionally an estimate of the process variance has been included, as this is the case with all the other methods.

The estimated reserves of all methods except 'Taylor/Ashe' differ by less than 20% and are therefore according to Table 3 within one standard error. For the chain ladder method neither the reserve estimates nor the standard errors are systematically higher or lower than for the other methods (except 'Taylor/Ashe'). The reason for the comparatively high chain ladder standard

error of 80% for accident year 2 is the fact that the reserve \hat{R}_2 itself is very low in comparison to the other reserves $\hat{R}_3, \dots, \hat{R}_{10}$: If we look at the sequence $\hat{R}_{10}, \hat{R}_9, \dots, \hat{R}_4, \hat{R}_3$ we see that \hat{R}_{i-1} is always greater than $\hat{R}_i/2$ but \hat{R}_2 is smaller than $\hat{R}_3/4$. This fact is very well reflected by the high standard error of 80%.

A closer look at the Taylor/Ashe data shows that the individual development factors $C_{i, k+1}/C_{ik}, 1 \leq i \leq I-k$, do not fluctuate much around their mean value \hat{f}_k so that the whole triangle can be considered as relatively regular. Therefore Taylor/Ashe were able to dispense with taking logarithms and thus avoided the problem of transforming back the result into the original data space. We therefore give a second example, which is less regular and where the claims amounts of the most recent accident years are much lower than in the previous years. These data (mortgage guarantee business) were compiled from a competition presented by SANDERS (1990).

TABLE 4
RUN-OFF TRIANGLE (ACCUMULATED FIGURES)

<i>i</i>	C_{i1}	C_{i2}	C_{i3}	C_{i4}	C_{i5}	C_{i6}	C_{i7}	C_{i8}	C_{i9}
1	58046	127970	476599	1027692	1360489	1647310	1819179	1906852	1950105
2	24492	141767	984288	2142656	2961978	3683940	4048898	4115760	
3	32848	274682	1522637	3203427	4445927	5158781	5342585		
4	21439	529828	2900301	4999019	6460112	6853904			
5	40397	763394	2920745	4989572	5648563				
6	90748	951994	4210640	5866482					
7	62096	868480	1954797						
8	24983	284441							
9	13121								

Parameter estimates ($k = 1, \dots, 8$):

\hat{f}_k : 11.1, 4.09, 1.71, 1.28, 1.14, 1.069, 1.026, 1.023

$\hat{\sigma}_k^2/1000$: 1787, 977, 194, 42.8, 27.0, 5.57, 1.26, 0.285

TABLE 5
ESTIMATED RESERVES \hat{R}_i IN 1000 s

	Chain ladder	Renshaw Christofides	Zehnwirth	Mack 1991
$i = 2$	93	91	87	62
$i = 3$	265	275	262	199
$i = 4$	834	818	778	682
$i = 5$	1568	1979	1884	1639
$i = 6$	3696	5497	5231	4420
$i = 7$	3487	6650	6328	5378
$i = 8$	2956	4331	4122	3143
$i = 9$	1647	2339	2226	1555
overall	14547	21980	20919	17078

TABLE 6
STANDARD ERROR IN % OF \hat{R}_i

	Chain ladder	Renshaw Christofides	Zehnwirth	Mack 1991
$i = 2$	65 %	90 %	80 %	60 %
$i = 3$	53 %	60 %	53 %	41 %
$i = 4$	38 %	51 %	45 %	37 %
$i = 5$	38 %	48 %	42 %	35 %
$i = 6$	28 %	46 %	41 %	33 %
$i = 7$	37 %	47 %	42 %	34 %
$i = 8$	61 %	50 %	47 %	36 %
$i = 9$	133 %	66 %	64 %	47 %
overall	26 %		24 %	

Here all results have been calculated by the author. In comparison with the standard errors of the first example, the chain ladder standard errors now reflect very well the generally higher uncertainty of this second triangle and especially the uncertainty of the last two accident years where the relative standard errors are very high because the reserve estimates are comparatively low. The most extreme deviation between the reserve estimates of the different methods is for accident year 7 where the 'Renshaw/Christofides' reserve exceeds the chain ladder reserve by 2.5 standard errors.

Altogether, if the impressions of these two examples can be taken as typical, we can conclude that the standard errors are of about the same size for the chain ladder as with the other methods, although they do not show such a smooth pattern as these because the other methods use only one σ^2 parameter as compared to $I-1$ of chain ladder. But this could also be achieved for the chain ladder method by smoothing out the $\hat{\sigma}_k^2$'s by means of an exponential function $\exp(a-bk)$.

Finally, we must bear in mind that these standard errors can only reflect the estimation error and the statistical error, but not the specification error, i.e. the fact that the model chosen can be wrong or that the future development may not be in accordance with past experience.

ACKNOWLEDGEMENT

I am indebted to ALOIS GISLER for pointing out the correct definition of the mean squared error and some further useful remarks.

REFERENCES

- CHRISTOFIDES, S. (1990) Regression Models Based on Logincremental Payments. In: *Claims Reserving Manual*, Vol. 2, Institute of Actuaries, London.
- MACK, Th. (1991) A Simple Parametric Model for Rating Automobile Insurance or Estimating IBNR Claims Reserves. *ASTIN Bulletin* 21, 93-109.
- RENSHAW, A. (1989) Chain Ladder and Interactive Modelling. *Journal of the Institute of Actuaries* 116, 559-587.

- SANDERS, D. E. A. (1990) Competition Presented at a London Market Actuaries Dinner.
- SCHNIEPER, R. (1991) Separating True IBNR and IBNER Claims. *ASTIN Bulletin* **21**, 111–127.
- TAYLOR, G. C. and ASHE, F. R. (1983) Second Moments of Estimates of Outstanding Claims. *Journal of Econometrics* **23**, 37–61.
- VERRALL, R. J. (1990) Bayes and Empirical Bayes Estimation for the Chain Ladder Model. *ASTIN-Bulletin* **20**, 217–243.
- VERRALL, R. J. (1991) On the Estimation of Reserves from Loglinear Models. *Insurance: Mathematics and Economics* **10**, 75–80.
- WRIGHT, T. S. (1990) A Stochastic Method for Claims Reserving in General Insurance. *Journal of the Institute of Actuaries* **117**, 677–731.
- ZEHNWIRTH, B. (≥ 1985) Interactive Claims Reserving Forecasting System. Insuware P/L, E. St. Kilda Vic 3183, Australia.

THOMAS MACK

*Münchener Rückversicherungs-Gesellschaft, Königinstrasse 107,
D-80791 München.*