INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by all the authors and should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size ar illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made some indication of how they may be grouped to make a corporate plate without unduc wastage of space, should be indicated. Some indication of scale (preferably a scale bar should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and car have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: Ecology and Physiology of parasites (editor, A. M. Fallis) pp. 213-222. Adam Hilger Ltd. London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains o Schistosoma haematobium. Journal of Helminthology, 47, 49-59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
The source, release and specificity of proteolytic enzyme activity produced by <i>Anisakis simplex</i> larvae (Nematoda: Ascaridida) <i>in vitro.</i> B. E. MATTHEWS	175–185
Echinostoma revolutum (Froelich, 1802) Looss, 1899 and Isthmiophora melis (Schrank, 1788) Lühe, 1909 (Echinostomatinae, Digenea): Scanning electron microscopy of the tegumental surfaces. L. R. SMALES and H. D. BLANKESPOOR	187–195
Uptake of diethylcarbamazine by microfilariae and adults of <i>Litomosoides carinii</i> and <i>Dipetalonema viteae</i> . (Research Note). V. M. L. SRIVASTAVA, D. P. SINGH, R. K. CHATTERJEE and A. B. SEN	197–199
Histochemical localization and functional significance of some enzymes in various tissues of Ceylonocotyle scoliocoelium cultured in vitro. A. N. SHARMA	201–206
Activation in vitro of some biological systems by extracts of adult worms and microfilariae of Dirofilaria immitis. P. F. L. BOREHAM	207–212
Studies on some monogenean parasites of Haemulon sciurus (Shaw, 1803) (Pomadasyidae). A. KOHN, B. ABRAMSON and B. MACEDO	213–218
Parasitological and pathological observations on schistosomiasis in sheep in Senegal. (Research Note). J. VERCRUYSSE and P. SCHANDEVYL	219–220
Nematospiroides dubius in the jird, Meriones unguiculatus: stimulation and expression of acquired immunity. J. M. BEHNKE and J. HANNAH	221–229
The presence of Angiostrongylus cantonensis in Viti Levu, Fiji. R. UCHIKAWA, M. TAKAGI, S. MATAYOSHI and A. SATO	231–234
Stage-specific immunogens of Hymenolepis nana in mice. (Research Note). A. ITO	235–238
The possible role of slugs (Agriolimax spp.) in the transmission of infective larvae of Cooperia oncophora. (Research Note). J. GRØNVOLD and P. NANSEN	239–240
Infectivity, growth and development of Echinostoma revolutum (Trematoda) in the domestic chick. B. FRIED	241-244
Induction of host resistance to <i>Brugia pahangi</i> in jirds (<i>Meriones unguiculatus</i>) protected by chemoprophylaxis. W. CHUSATTAYANOND and D. A. DENHAM	245–249
Studies of potential filaricides: Part 14—Activity of 1-iso-butoxycarbonyl-4-methyl-piperazine against experimental filariasis. R. RASTOGI, S. SHARMA, N. ANAND, A. DUTTA, N. FATMA, R. K. CHATTERJEE and A. B. SEN	251–254
Studies of Angiostrongylus cantonensis in two experimental hosts. K. UCHIDA, K. YOSHIMURA, K. SATO and T. H. YOSIDA	255–258
Detection of circulating antigen in bancroftian filariasis by sandwich ELISA using filarial serum IgG. M. V. R. REDDY, A. MALHOTRA and B. C. HARINATH	259–262
Intestinal absorption of glucose in rats after continuous challenge with larvae of Nippostrongylus brasiliensis. K. J. CHEEMA and A. M. SCOFIELD	263–266
Observations on the infectivity and fecundity of a Sudanese isolate of <i>Schistosoma bovis</i> in albino mice. A. M. SAAD and M. F. HUSSEIN	267–270
Announcements 186	, 200, 250
	-