

Method. This retrospective cohort study was carried out at the Phoenix Care Centre Dublin, Ireland. Informed consent was not sought as this was a retrospective chart study involving anonymised clinical data which was collected as part of routine clinical care and no items of information were reported that would enable the identification of any subject. We described primary outcomes using frequencies, percentages, mean and standard deviations, median and interquartile ranges (IQR). Between groups comparisons were made using χ^2 tests for categorical variables; t-tests, ANOVA tests, or Kruskal-Wallis tests, for continuous variables; All analyses were two-tailed, and a P-value ≤ 0.05 was considered statistically significant

Result. Over the study period from Jan 2014 to Jan 2017 inclusive, there were 96 admission episodes to the PICU. The mean age of admitted cases was 37.1 (SD = 11.3) years (range 18–63 years). The mean length of stay (LOS) was 59.3 (SD = 61.0) days (median 39.5 days). All patients were admitted under the Mental Health Act legislation. We identified assault as the primary risk factor for pre-admission 62% (n = 62) to the PICU. Antipsychotic polypharmacy was used in 61% (n = 55) of the admission. The mean daily antipsychotic dosage was 139.4 % (SD = 65.1) of BNF maximum daily dose. A diagnosis of acute psychotic disorder (B = -1.027, p = 0.003, 95% CI: -1.691 to -0.363) was associated with reduced LOS in PICU. Majority of admissions 43% (n = 39) had a diagnosis of schizophrenia, followed by Bipolar affective disorder BPAD 21% (n = 21), schizoaffective disorder 18% (n = 18), and acute psychotic disorder 9% (n = 9).

Conclusion. Psychiatric Intensive Care Unit is an essential service for the severely ill psychiatric patients and is a progressively developing sub-speciality. An important finding from our study describes the cohort of patients admitted being predominantly male, younger-aged, single, with a diagnosis of schizophrenia, legally detained, and from an Irish background. The primary indication for a referral is the risk of assault, showing the need for the intensive and secure treatment model that a PICU can provide.

Prevalence and correlates of depressive symptoms among professional drivers in Saudi Arabia: a cross-sectional study

Adnan Raed Alnaser*, Osama A. Zitoun, Tawfik Rajab, Abdullah Khojah, Juliann Saquib and Nazmus Saquib

College of Medicine, Sulaiman Al Rajhi University

*Corresponding author.

doi: 10.1192/bjo.2021.751

Aims. Due to the nature of their work, professional drivers face a considerable risk of developing depression and other mental illnesses. We sought to assess the prevalence and the factors influencing depressive symptoms among professional drivers in Saudi Arabia.

Method. Using convenience sampling, we have conducted an interviewer-administered survey on 324 professional drivers in Qassim Region in Saudi Arabia using Depression subscale from the Depression, Anxiety and Stress Scale 21 (DASS-21). Participants were interviewed in their native language, and responses were outlined directly into an online form in English. Data were then extracted and analyzed using SPSS software.

Result. Participants' mean age was 38.6 years, and mean driving hours per day were 9.86 hours/day. The mean DASS-21 depression score among the professional drivers was 2.88. Overall, 21.9% of the included drivers had variable degrees of depressive symptoms, with 7.4% suffered from extremely severe symptoms. Depressive symptoms were influenced by the driver's nationality, educational level, vehicle type, driving years, BMI, presence of chronic medical conditions, physical activity, and sexual activity. Moreover, poor sleep quality increased the risk of developing

depressive symptoms among the drivers by 31.9 times (OR: 31.9, CI: 9.03–112.63, P < 0.001).

Conclusion. Nearly one-fifth of professional drivers in Saudi Arabia (Qassim region) suffer from depressive symptoms. Unhealthy life-style practices (i.e. being obese and physically inactive) have been closely related to depressive symptoms. Education, sexual activity, type of driven vehicle, and the number of chronic conditions were also associated with depressive symptoms. Also, poor and fair sleep quality was strongly associated with the development of depressive symptoms as compared with excellent sleep quality. As drivers are always on the move and hardly reachable, we would propose psychological support and counseling to be administered via telemedicine services. Future research is needed to better comprehend the needs of this vulnerable population.

Can probiotics benefit young people with autism spectrum disorders?

Pooja Ramani^{1*} and Regina Sala²

¹Great Ormond Street Hospital and ²East London NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.752

Aims. The aims are to evaluate the effectiveness of Probiotics on young people with Autism Spectrum Disorder.

We hypothesized that there will be an improvement of the comorbid gastrointestinal symptoms that can accompany Autism Spectrum Disorder.

We believe that the use of probiotics can exert bidirectional effects on the gut-brain axis which may result in improvements in core Autism symptoms.

Method. A literature search was performed in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. We used databases including OVID MEDLINE, Pubmed, EMBASE, AMED and the Cochrane register of controlled trials. Studies using Probiotics as a treatment for children with ASD were identified by key search terms; Child*, young person*, adoles*, teenagers, ASD, Autism Spectrum Disorder, Autism, Pervasive developmental disorder, PDD, Probiotics, Supplements, Lactobacillus, and Bifidobacterium. Inclusion criteria: Children of age range 2-18 with a diagnosis of ASD and having at least one gastrointestinal symptom were included. Exclusion criteria: The following were excluded: studies looking at Autism with interventions aside from Probiotics; studies where Probiotics were tested in conjunction with other interventions; studies where there were additional neurodevelopmental disorders.

Result. Twelve studies identified all utilized probiotics. This included 7 Randomised Control Trials, 2 Open-Label studies, 1 pre and post-intervention design and 1 Case study. All RCTs gave probiotics or placebo to children.

Ten studies showed an improvement in gastrointestinal symptoms. Six studies showed improvements in various behavioral measures. Four studies showed improvements in core autism symptoms. However, the sample sizes in these studies were not large enough to prove statistical significance.

Conclusion. No studies showed an adverse reaction which indicates probiotics can be considered a safe treatment.

The improvements in a variety of parameters imply probiotics a suitable adjunctive intervention that may help improve ASD core symptoms in young people as well as improving physical and behavioural comorbidities which in some cases was noted by parents.

However, due to high dropout rates and generally small sample sizes, larger-scale trials are needed to critically confirm the efficacy of probiotics for children with ASD.

Comparison of hospitalizations in patients on first generation versus second generation long acting injectable (LAI) antipsychotics

Seemab Rasool* and Paster Venan

Essex Partnership University NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.753

Aims. There is limited data on the comparison of efficacy between first and second antipsychotic LAIs. One good indicator of efficacy is the rates of hospitalization. Some studies have shown that second generation depot antipsychotics, significantly reduce hospitalizations as compared to conventional depots.

Our aim was to compare hospitalizations in patients on first and second generation LAI antipsychotics.

Method. A retrospective observational study was done by reviewing the records of all the depot clinics in South Essex, United Kingdom.

A list of patients enrolled and receiving LAI antipsychotics was obtained from the 6 depot clinics. Data were collected by going through the electronic records of the patients on the depot clinic lists and taking down the demographics, diagnosis and the hospital admissions. Other variables like comorbid drug abuse were also recorded.

Result. Amongst a total of 346 patients 223 (64 %) were males and 123 (36%) were females. Average age was 50.3 (range 21 to 88 years) and 290 (83%) patients were single. An overwhelming majority of patients 299 (87 %) were not in employment. Regarding the diagnosis, the majority, 237 patients were diagnosed with Paranoid Schizophrenia, 49 patients were diagnosed with Schizoaffective disorder, 38 patients were diagnosed with Bipolar affective disorder, 20 patients had a diagnosis of Delusional disorder and only 2 patients had a primary diagnosis of Mental and Behavioral disorders due to substance abuse. Of the total 346 only 17 patients were on a Community treatment Order.

Risperidone was the most commonly used second generation LAI at 26%, Aripiprazole in 10% and Paliperidone was used in 5% patients. Olanzapine LAI was only used in 2 patients. Amongst first generation LAIs Zuclopentixol, Fluclopentixol were both used in 24%, and Haloperidol in 10% patients. 21 % of patients were reported to be actively abusing drugs.

65 (32.6%) of the total 200 patients on 1st Generation LAIs had hospital admissions

55 (39.8%) of the total 138 patients on 2nd Generation LAIs had hospital admissions

This difference was not statistically significant (Z test)- P value of 0.082427

Conclusion. The results in our observational study are equivocal, both LAIs providing equitable decrease in the hospital admissions albeit with a slightly favourable outcome (not statistically significant though) attributable to the first generation LAIs. There was a high incidence of unemployment and drug abuse in our cohort of patients, thus targeted interventions can be established in rehabilitation of such individuals.

Quantifying the disorganization and the core deficit in classical schizophrenia

Mohan Rathnaiah¹*, Elizabeth B Liddle², Lauren Gascoyne², Jyothika Kumar², Mohammad Zia Ul-Haq Katshu², Catherine Faruqui³, Christina Kelly³, Malkeet Gill³, Sian Robson², Peter Morris², Mathew Brookes² and Peter Liddle²

¹Institute of Mental Health, University of Nottingham; ²University of Nottingham and ³Nottinghamshire Healthcare NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.754

Aims. To derive scores for mental disorganization and impoverishment from commonly used rating scales, and test the hypothesis that disorganization and impoverishment, along with impaired cognition and role-function reflect a latent variable that is a plausible candidate for the putative core deficit.

Background. For more than 100 years, disorganization and impoverishment of mental activity have been recognised as fundamental symptoms of schizophrenia. These symptoms may reflect a core brain process underlying persisting disability. Delusions and hallucinations have been regarded as accessory features. The psychopathological processes predisposing to persisting disability in schizophrenia are poorly understood. The delineation of a core deficit underlying persisting disability would be potentially of great value in predicting outcome and developing improved treatment.

Method. Patients aged 18–55 years were included if: they satisfied DSM IV criteria for schizophrenia or schizoaffective disorder. Healthy controls were recruited by public advertisement and selected to match the patient group in age and sex. Study sample included 39 participants with schizophrenia, 1 with schizoaffective disorder and 44 matched healthy controls. We derived disorganization and impoverishment scores from three symptom scales: PANSS, SSPI and CASH. We computed composite scores for disorganization and for impoverishment and employed Confirmatory Factor Analysis to test the hypothesis that a single factor accounts for the relationships between disorganization, impoverishment, cognitive impairment and impaired role function. We assessed the relationship between this latent “core deficit” and diminished Post Movement Beta Rebound (PMBR), an electrophysiological measure from Magnetoencephalography (MEG), associated with persisting brain disorders.

Result. Fit indices for the single factor model from CFA indicated a good fit: $\chi^2(2) = 1.817$, $p = .403$; RMSEA $<.001$ GFI = .979. PMBR was significantly reduced in the schizophrenia group compared to healthy controls, $t(68) = 3.55$, $p < .001$. Within the patient group, PMBR was significantly and negatively correlated with the CFA factor scores representing the Core Deficit score, $r = -.543$, $p < .01$, indicating that high core deficit scores were associated with reduced PMBR. PMBR was significantly correlated with the composite Disorganization score, $r = -.521$, $p < .001$.

Conclusion. Our findings demonstrate that the shared variance between impoverishment (psychomotor poverty); disorganization; cognitive impairment; and impaired role function can be accounted for by a latent variable that can reasonably be described as the core deficit of classical schizophrenia. The demonstration that the severity of the putative core deficit is correlated with the reduction in PMBR provides evidence that the core deficit is associated with an identifiable abnormality of brain dysfunction.

Prevalence of mental disorders in prisons in the UK: a systematic review and meta-analysis

Nivedita Rebbapragada*, Vivek Furtado and George William Hawker-Bond

Warwick Medical School

*Corresponding author.

doi: 10.1192/bjo.2021.755

Aims. To report pooled prevalence of all mental disorders among the general prison population in the United Kingdom (UK). This includes individuals in Young Offender Institutions (YOI), youth custody and adult prisons across all categories. A secondary aim explores possible sources of heterogeneity by performing