

new therapist, in this context intensive follow-up is carried out in the event of the appearance of pharmacological secondary effects, pharmacological readjustment is carried out with good results.

During the joint follow-up with nursing, the cessation of secondaryisms is confirmed and we are informed of the gradual appearance of overvalued ideas in relation to the new therapist, which are gradually structured in the form of erotomanic delirium that coincides with the cessation of follow-up by said therapist. Consultations in the emergency room occur on a couple of occasions due to mild behavioral alterations secondary to messages and communications that he reports receiving where said love is confirmed. Despite readjustments, there continues to be an increase in clinical symptoms due to abandonment of medication, finally producing serious alterations aimed at the search for said therapist, finally culminating in admission to the acute care unit for containment of said condition.

Objectives: The objectives is the diferencial diagnosis, in this case symptoms could be classified as positive symptoms of schizophrenia, although it is its own nosological entity.

Methods: .

Results: .

Conclusions: This patient represents a classic example of De Clerambault syndrome and is a faithful expression of the recurrent syndrome associated with delusions of grandeur, eroticism and jealousy. There have also been ideas of reference and agitated behavior associated with his delusional process.

Disclosure of Interest: None Declared

EPV0975

The Course of Schizophrenia Spectrum Disorders With Episodes of Catatonic Depressions

A. Barkhatova*, M. Bolgov and A. Smolnikova

Department of endogenous mental disorders and affective states, Mental health research center, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1563

Introduction: Mood symptoms, especially depressive ones, occur in the majority of patients with schizophrenia spectrum disorders (SSD). Therefore, depression is often identified as one of the symptomatological dimensions of schizophrenia. Catatonia is also considered by some researchers as one of the dimensions of schizophrenia, or as an independent transnosological formation. Catatonia in SSD may be associated with affective dysregulation and is often accompanied by depression. Although the clinical course of SSD has been well studied previously, its relationship with psychopathological structure of episodes of SSD remains not entirely clear.

Objectives: To determine the impact of episodes of catatonic depression on the course and prognosis of SSD.

Methods: A sample of 60 patients with episodic course of SSD who met the criteria for catatonia according to the Bush-Francis Catatonia Screening Instrument (BFCSI) and for depression according to the Calgary depression schizophrenia scale (CDSS) was analyzed. An analysis of the clinical course of SSD was carried out on the basis of the medical history of all patients in the study sample and follow-up observation of 42 patients for 5 years. Global

Assessment of Functioning Scale (GAF) was used to assess the prognosis of SSD.

Results: Patients were divided into two groups depending on the period of manifestation of catatonia in the clinical course of SSD: during the first episode or during subsequent episodes. The sample of patients with the first episode (n=43, 71.7 %) was divided into three subgroups. A relatively favorable course of SSD was observed only in 13 patients (30.2 %; 21.7 % of SSD sample). The course of disorder was characterized by similar episodes with a high proportion of affective symptoms, long-term remissions and minimally expressed negative symptoms (GAF score=75.2±5.82). A relatively unfavorable course of SSD was observed in 15 patients (34.9 %; 25.0 % of SSD sample). It was characterized by moderate negative and chronic subdepressive symptoms with low frequency of catatonic and psychotic relapses (GAF score=62.3). An unfavorable course of SSD was also observed in 15 patients (34.9 %; 25.0 % of SSD sample). It was characterized by a high frequency of relapses with a tendency to form a chronic conditions with residual catatonic signs and psychotic symptoms (GAF score=50.1). In the sample of patients with manifestation of catatonia in the second or subsequent episodes (n=17; 28.3 %), the clinical course of SSD was unfavorable. It was characterized by a rapid augmenting of negative symptoms with the formation of psychomotor poverty syndrome with residual catatonic symptoms (GAF score=52.7).

Conclusions: Our study shows that the occurrence of catatonic depressive episodes in the clinical course of SSD in most cases is an unfavorable prognostic factor.

Disclosure of Interest: None Declared

EPV0976

Efficacy and tolerability Aripiprazole once-monthly long-acting injectable in schizophrenia. Two-injection start regimen. A 24 months follow-up and mirror image study

S. L. Romero Guillena^{1*}, F. Gotor Sanchez-Luengo² and B. O. Plasencia Garcia de Diego³

¹Psychiatry, UGC Mental Health Virgen Macarena; ²Psychiatry, UGC Mental Health Virgen Rocio and ³Psychiatry, UGC Mental Health Virgen Rocio, Seville, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1564

Introduction: Relapse prevention is crucial in patients with schizophrenia, as repeated episodes can worsen psychopathology and functionality. There is strong evidence of antipsychotics efficacy in preventing relapse; however, non-compliance rates in patients with schizophrenia are very high. Long-acting injectable antipsychotics (LAIs) are an important treatment option but remain underutilized.

Aripiprazole once-monthly is a long-acting intramuscular injectable formulation of aripiprazole indicated for the maintenance treatment of schizophrenia in adult patients stabilized on oral aripiprazole.

If one injection start regimen is adopted, on the day of initiation, an injection of 400mg Aripiprazole once monthly should be administered accompanied by 10mg to 20mg of oral aripiprazole per day for the successive 14 days New treatment regimen: On the day it begins,