

DIETARY SURVEYS AND NUTRITIONAL EPIDEMIOLOGY

Plasma 25-hydroxyvitamin D₃, folate and vitamin B₁₂ biomarkers among international colorectal cancer patients: a pilot study

Cornelia M. Ulrich^{1,2*}, Adetunji T. Toriola³, Erin M. Siegel⁴, Hermann Brenner⁵, Jenny Chang-Claude⁶, Clare Abbenhardt¹, Jana Kotzmann¹, Xiaoling Song², Robert W. Owen¹, Michael Hoffmeister⁵, Heiko Becher⁷, David Shibata⁴, Kathy Vickers², Shannon K. Rush², Karen Makar², Gerd Würtele¹, Roswitha Haubner¹, Thomas A. Sellers⁴ and William Grady on behalf of the ColoCare Consortium²

¹Division of Preventive Oncology, National Center for Tumor Diseases (NCT) and German Cancer Research Center (DKFZ), Heidelberg, Germany

²Fred Hutchinson Cancer Research Center, Seattle, WA, USA

³Division of Public Health Sciences, Department of Surgery, Washington University School of Medicine and Siteman Cancer Center, St Louis, MO, USA

⁴H. Lee Moffitt Cancer Center, Tampa, FL, USA

⁵Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany

⁶Department of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

⁷University of Heidelberg, Heidelberg, Germany

(Received 23 March 2012 – Final revision received 14 September 2012 – Accepted 29 October 2012)

Journal of Nutritional Science (2013), vol. 2, e9, page 1 of 6

doi:10.1017/jns.2012.28

Abstract

Vitamin D and folate are associated with decreased colorectal cancer risk and their association with colorectal cancer prognosis is under investigation. We assessed the levels of plasma 25-hydroxyvitamin D₃ (25(OH)D₃), folate and vitamin B₁₂ in an international pilot study in order to determine variability of these biomarkers based on geographical location. Plasma 25(OH)D₃, folate and vitamin B₁₂ concentrations were measured in 149 invasive, newly diagnosed colorectal cancer cases from Heidelberg (Germany), Seattle (WA, USA), and Tampa (FL, USA) and in ninety-one age- and sex-matched controls. Their associations with potential predictors were assessed using multivariate linear regression analyses. Plasma 25(OH)D₃, folate and vitamin B₁₂ concentrations differed by location. Other predictors were season for 25(OH)D₃ and tumour stage (vitamin B₁₂). Season-corrected average 25(OH)D₃ concentrations were higher in Heidelberg (31.7 ng/ml; range 11.0–83.0 ng/ml) than in Seattle (23.3 ng/ml; range 4.0–80.0 ng/ml) and Tampa (21.1 ng/ml; range 4.6–51.6 ng/ml). In Heidelberg, a strong seasonal variation was observed. Folate (11.1 ng/ml) and vitamin B₁₂ (395 pg/ml) concentrations in Heidelberg were lower than those in Seattle (25.3 ng/ml and 740 pg/ml, respectively) and Tampa (23.8 ng/ml and 522 pg/ml, respectively). Differences in plasma 25(OH)D₃ and folate concentrations between Heidelberg and the US sites were observed, probably reflecting variation in outdoor activities and sun-avoidance behaviour during summer as well as in folic acid fortification and supplement use. Intra-site differences at each study location were greater than between-location variability, suggesting that individual health behaviours play a significant role. Nevertheless, the intra-site differences we observed may be due to chance because of the limited sample size. Our pilot study illustrates the value of an international cohort in studying colorectal cancer prognosis to discern geographical differences in a broad range of exposures.

Key words: Colorectal cancer: Survival: Vitamin D: Vitamin B₁₂: Folate

Vitamin D and folate are critical for cell signalling and functioning and can influence processes related to cancer risk

and prognosis^(1,2). The anti-carcinogenic effects of vitamin D are mediated through its diverse effects on cellular

Abbreviations: DACHS, Darmkrebs Chancen der Verhütung durch Screening; 25(OH)D₃, 25-hydroxyvitamin D₃.

* **Corresponding author:** Cornelia M. Ulrich, email neli.ulrich@nct-heidelberg.de



differentiation, proliferation, apoptosis and angiogenesis⁽³⁾. Folate and other B vitamins such as vitamin B₁₂ play important roles as co-factors in one-carbon metabolism, which is essential for nucleotide synthesis and DNA methylation^(1,2). Disruption of these processes leads to abnormalities in DNA methylation and imbalance of DNA precursors, leading to aberrant DNA synthesis and repair^(1,2).

Globally, colorectal cancer is the second most commonly diagnosed cancer in women and the third one in men⁽⁴⁾. Studies have demonstrated that a higher pre-diagnostic vitamin D status is associated with a decreased colorectal cancer risk^(5–7) and there is emerging evidence that higher plasma vitamin D concentrations can also improve survival in patients with colorectal cancer^(8,9). On the other hand, folate appears to have a dual effect on colorectal carcinogenesis, depending on the timing and dosage^(10–13). In normal colorectal epithelial cells, folate deficiency predisposes to colorectal carcinogenesis whereas in established neoplasms, folate deficiency appears to have an inhibitory effect on colorectal carcinogenesis and high folate levels may promote the growth of an existing cancer^(10,11,13,14). Vitamin B₁₂ is an important co-factor in folate metabolism but studies relating plasma vitamin B₁₂ to colorectal cancer risk and prognosis are relatively very few and results equivocal^(15,16). Because there is paucity of information on how these nutritional biomarkers impact on colorectal cancer survival, more studies are needed to discern their effects.

Plasma vitamin biomarker levels may vary based on dietary and other health-related practices in different countries. However, studies comparing this wide range of vitamin biomarker levels across geographical regions are scarce. A characterisation of this variability may be important for studying the association of these biomarkers with colorectal cancer prognosis. In order to discern geographical differences in plasma 25-hydroxyvitamin D₃ (25(OH)D₃), folate and vitamin B₁₂, we evaluated levels and predictors among colorectal cancer cases and controls in an international pilot study with participants from Heidelberg (Germany), Seattle (WA, USA), and Tampa (FL, USA).

Methods

Study groups

The study participants from the USA are members of a colorectal cancer prognosis study (ColoCare), with sites at the Fred Hutchinson Cancer Research Center, Seattle (WA, USA), H. Lee Moffitt Cancer Center and Research Institute, Tampa (FL, USA), and the National Center for Tumor Diseases, German Cancer Research Center, Heidelberg (Germany). The objective of our consortium is to identify the factors that determine short-term survival in a prospective cohort of colorectal cancer patients. These include inherited genetic and tumour characteristics that affect early detection, treatment response and prognosis as well as health behaviours that patients adopt that may have health outcomes. German samples included in this pilot study were obtained from participants of the Darmkrebs Chancen der Verhütung durch Screening (DACHS) study at the German Cancer Research

Center in Heidelberg (Germany)⁽¹⁷⁾. Samples included from Tampa were collected as part of Moffitt's Total Cancer Care Program⁽¹⁸⁾.

Eligible for the study were colorectal cancer patients, at least 18 years of age, with a first diagnosis of invasive colon or rectal cancer (stages I–IV) receiving care at any of the ColoCare consortium/DACHS sites. This pilot study involved 149 colorectal cancer cases and ninety-one age- and sex-matched controls (Heidelberg, forty cases and forty controls; Seattle, ninety-one cases and thirty-three frequency-matched controls; Tampa, eighteen cases and eighteen controls), recruited between the years 2006 and 2010 and whose biospecimens were obtained within the same time frame with identical blood-draw protocols at each of the sites. Blood samples were obtained at the time of recruitment into the study. For the cases, the samples were pre-treatment samples obtained at the time of surgery. In the DACHS study, control subjects were community-based and randomly selected from population registries. They were frequency-matched to cases on age, sex and county of residence. We excluded controls with a history of colorectal cancer; otherwise eligibility criteria were the same as for the cases. Controls were contacted through mail and follow-up calls, and interviews were conducted at their homes. The participation rate among eligible control participants was a little higher than 50 %⁽¹⁷⁾. The controls in Moffitt, Tampa were frequency matched to the cases on age (5-year intervals), sex and race (white non-Hispanic). Controls in Seattle were drawn from participants in a screening and surveillance colonoscopy programme who had no evidence of pathology within the colon and were matched to the cases for age, sex and race.

This study was conducted according to the guidelines laid down by the Declaration of Helsinki and all procedures involving human subjects were approved by the Institutional Review Boards of the German Cancer Research Center (DKFZ), the Fred Hutchinson Cancer Research Center, Seattle (WA, USA) and H. Lee Moffitt Cancer Center and Research Institute, Tampa (FL, USA). Written informed consent was obtained from all the subjects.

Laboratory analysis

Plasma 25(OH)D₃ was measured by reverse-phase HPLC in the Biochemistry and Biomarkers Laboratory of the Division of Preventive Oncology, German Cancer Research Center, DKFZ (Dr Owen). The intra-assay and inter-assay CV of 25(OH)D₃ at a concentration of 50 ng/ml were 2.3 and 3.1 %, respectively. Plasma folate and vitamin B₁₂ were quantified using SimulTRAC[®] B₁₂/FOLATE – SNB 57Co/¹²⁵I Radioassay Kit (MP Biomedicals) in the Biomarker Laboratory (Dr Song) at the Fred Hutchinson Cancer Research Center with intra-assay CV of 3.1 and 3.5 % and inter-assay CV of 7.0 and 5.1 % for folate and vitamin B₁₂, respectively.

Statistical analysis

To account for skewed distributions, Wilcoxon or Kruskal–Wallis tests were used to test for the differences in biomarker concentration by age group (18–59, 60–69 and 70–80 years),



case status, tumour stage (I–IV), study site (Heidelberg, Seattle and Tampa), and season of blood draw (spring: March–May; summer: June–August; autumn: September–November; winter: December–February). Because the samples were unequally distributed across seasons at the three study sites, we computed season-corrected levels for the descriptive analyses of 25(OH)D₃ in order to give equal weight to all four seasons. Linear regression analysis (using both forward stepwise and enter methods) with log-transformed outcome parameters was used for multivariate analyses. All statistical analyses were conducted using SAS (V9.1.2, SAS Institute Inc.). Statistical significance was defined as $P < 0.05$ and all statistical tests were two-sided.

Results

Cases and controls were, on average, aged 60.0 (SD 13.3) and 60.9 (SD 13.1) years, respectively. The majority of tumours were of stages III (39.6%) and IV (26.2%). Among the cases, the mean 25(OH)D₃, folate and vitamin B₁₂ concentrations were 25.6 (SD 13.5) ng/ml, 21.6 (SD 14.0) ng/ml, and 638 (SD 597) pg/ml, respectively, whereas among the controls, they were 25.9 (SD 14.0) ng/ml, 18.0 (SD 13.7) ng/ml, and 515 (SD 241) pg/ml (Table 1).

In multivariate modelling, significant predictors for the biomarkers included season for 25(OH)D₃, study site for 25(OH)D₃, folate, vitamin B₁₂, and tumour stage for vitamin B₁₂ (Table 2). Probably because of the limited sample size of this pilot study, case status was not statistically significant as a predictor; hence, cases and controls were combined in subsequent analyses. Also, the results from models excluding case-status were identical to those including case-status in the model. The association from controls only was similar, yet did not reach statistical significance. Plasma 25(OH)D₃ concentrations were, on average, higher in Heidelberg (31.7 ng/ml; range 11.0–83.0 ng/ml) than in Seattle (23.3 ng/ml; range 4.0–80.0 ng/ml) and Tampa (21.1 ng/ml; range 4.6–51.6 ng/ml) (Table 3). Hypovitaminosis D (plasma 25(OH)D₃ <20 ng/ml) was more prevalent in Tampa (54%) and Seattle (50%) than in Heidelberg (23%) (data not shown). On the other hand, plasma folate concentrations were, on average, lower in Heidelberg (11.1 ng/ml; range 0.3–48.2 ng/ml) than in Seattle (25.3 ng/ml; range 3.5–70.3 ng/ml) and Tampa (23.8 ng/ml; range 6.8–69.7 ng/ml). No participants from Seattle and Tampa had folate deficiency (folate <3 ng/ml), while 10% of those from Heidelberg had folate deficiency. Plasma vitamin B₁₂ concentrations followed the same pattern as folate with lower average values recorded in Heidelberg (395 pg/ml; range 96–1641 pg/ml) than in Seattle (740 pg/ml; range 183–3750 pg/ml) and Tampa (522 pg/ml; range 94–1085 pg/ml).

Seasonal differences in plasma 25(OH)D₃ concentrations were evident in Heidelberg, where the highest plasma 25(OH)D₃ concentrations were observed in summer (37.0 ng/ml) and the lowest in spring (21.3 ng/ml), but no such seasonal variation was noted in Seattle and Tampa (Fig. 1).

Table 1. Baseline characteristics of colorectal cancer cases and controls

Characteristic	Cases (n 149)		Controls (n 91)	
	n	%	n	%
Study site				
Heidelberg	40	26.8	40	44.0
Seattle	91	61.1	33	36.3
Tampa	18	12.1	18	19.8
Age (years)				
Mean		60		61
SD		13		13
18–59	64	43	36	40
60–69	47	32	32	35
70–90	38	26	23	25
Sex				
Male	80	54	48	53
Female	69	46	43	47
Race				
White, non-Hispanic	137	92	84	92
Non-white or other	12	8	7	8
Tumour stage				
I (I, IA, IB and IC)	20	13.4	–	–
II (II, IIA, IIB and IIC)	30	20.1	–	–
III (III, IIIA, IIIB and IIIC)	59	39.6	–	–
IV	39	26.2	–	–
Unknown	1	0.7	–	–
Season of blood draw				
Spring (March–May)	27	18.1	23	25.3
Summer (June–August)	32	21.5	25	27.5
Autumn (September–November)	44	29.5	23	25.3
Winter (December–February)	46	30.9	20	22.0
Nutritional biomarker				
25(OH)D ₃ (ng/ml)				
Mean		25.6		25.9
SD		13.5		14.0
Range		4.6–80.0		4.0–83.0
Folate (ng/ml)				
Mean		21.6		18.0
SD		14.0		13.7
Range		0.3–70.3		1.4–62.5
Vitamin B ₁₂ (pg/ml)				
Mean		638		515
SD		597		241
Range		96–3750		94–1352

25(OH)D₃, 25-hydroxyvitamin D₃.

Discussion

To the best of our knowledge, this is the first study to compare a range of nutritional biomarkers (25(OH)D₃, folate and vitamin B₁₂) among colorectal cancer patients and controls in different geographical regions. Our international pilot study

Table 2. Predictors of plasma 25-hydroxyvitamin D₃ (25(OH)D₃), vitamin B₁₂ and folate among colorectal cancer cases and controls†

Biomarker	Predictor	Direction	Significance
25(OH)D ₃	Heidelberg	+	**
	Summer season	+	**
	Autumn season	+	*
Vitamin B ₁₂	Heidelberg	–	**
	Stage IV	+	**
Folate	Heidelberg	–	**
	Age 60–69 years	+	*

* $P < 0.10$, ** $P < 0.05$.

† Cases and controls were combined because case status was not statistically significant.



Table 3. Nutritional biomarker concentrations among colorectal cancer cases and controls in the three study locations (Mean values and ranges)

	Heidelberg		Seattle		Tampa	
	Mean	Range	Mean	Range	Mean	Range
25(OH)D ₃ (ng/ml)	31.7	11.0, 83.0	23.3	4.0, 80.0	21.1	4.6, 51.6
Folate (ng/ml)	11.1	0.3, 48.2	25.3	3.5, 70.3	23.8	6.8, 69.7
Vitamin B ₁₂ (pg/ml)	395	96, 1641	740	183, 3750	522	94, 1085

25(OH)D₃, 25-hydroxyvitamin D₃.

revealed differences in biomarker concentrations between the US sites and Germany.

Because dietary sources of vitamin D are very limited, cutaneous synthesis of vitamin D (within the UVB wavelength of 290–315 nm) is a very important source of plasma 25(OH)D₃⁽¹⁹⁾. The UV radiation required for the synthesis of vitamin D in the skin is more abundant and more functional during summer than during winter^(19,20). Hence, among people who do not use adequate vitamin D supplements or consume vitamin D-fortified foods, season is one of the most important determinants of plasma 25(OH)D₃ concentrations.

Thus, the seasonal variations in plasma 25(OH)D₃ observed in Heidelberg are in line with expectations. Somewhat surprising is the lack of seasonal variation and the low plasma 25(OH)D₃

concentrations observed in Seattle and Tampa. Of the three sites, Tampa has the highest daytime temperatures during summer but the participants from Tampa had the lowest plasma 25(OH)D₃ concentrations during summer. The differences in plasma 25(OH)D₃ concentration during summer among participants in our study could reflect the extent of outdoor activities and possible sun-avoidance behaviours (including sunscreen use) in the three cities. In Heidelberg, people are more likely to commute with bicycles or walk than use cars, whereas people are more likely to practise sun-avoidance in Florida⁽²¹⁾.

Few studies have examined the impact of season on 25(OH)D₃ concentrations in lower altitude regions in the USA. Although a previous study in South Florida reported a seasonal variation in serum 25(OH)D₃ concentrations⁽²¹⁾, the




Fig. 1. (a) Plasma vitamin D (25-hydroxyvitamin D; 25(OH)D) concentration by study site. (b) Plasma folate concentration by study site. (c) Plasma vitamin B₁₂ concentration by study site. The horizontal bars represent medians and the circles represent potential outliers in the data. (d) Seasonal plasma 25(OH)D by study site: □, spring; ■, summer; ■, autumn; ■, winter. Values are means.



magnitude (14 %) is much smaller than what is observed (38 %) among residents of northern latitudes in the USA^(21,22). Compared with the previous study from South Florida, the winter 25(OH)D₃ concentrations among participants from Tampa in our study are identical (23.3 v. 23.1 ng/ml, respectively) but the summer concentrations were lower (17.7 v. 26.8 ng/ml). Thus, the low summer 25(OH)D₃ concentrations in Tampa could be a chance finding due to the low number of participants from Tampa in our study.

Studies have revealed that current food fortification practices in the USA may not be adequate to prevent vitamin D deficiency among vulnerable people during winter^(23,24). Likewise, it has been shown that although supplement use increases serum 25(OH)D₃ concentrations, it does not make up for insufficient sun exposure⁽²³⁾, and current supplement recommendations are not adequate to prevent vitamin D deficiency⁽²⁵⁾. As vitamin D supplementation is discussed in the context of improving survival in colorectal cancer patients^(8,26), our study shows that high plasma vitamin D concentrations can be achieved in populations with substantial outdoor activity during summer (such as Heidelberg). Nevertheless, vitamin D fortification in the USA did not prevent low levels of serum 25(OH)D₃ in Tampa or Seattle, cities that practise sun-avoidance or reduced sun exposure.

The lower plasma folate concentrations among participants from Heidelberg compared with those from the USA are not unexpected considering the fact that mandatory fortification of staple foods with folic acid commenced in the USA in 1998⁽²⁷⁾ while fortification is not practised in Germany⁽²⁸⁾. The mandatory fortification programme has resulted in a substantial increase in plasma folate concentrations in the USA, where the prevalence estimates for low serum folate concentrations (<3 ng/ml) declined from 20.6 % in the period 1988–1994 to 0.6 % in 2004⁽²⁹⁾.

However, we also observed large intra-site differences in plasma folate and vitamin B₁₂ concentrations at each study site. This may be due mostly to the use of folic acid-containing supplements. It has been shown that supplement and multivitamin use is higher among cancer patients compared with the general population⁽³⁰⁾. Between 64 and 81 % of cancer patients use any vitamin or mineral supplement, compared with 50 % in the general population⁽³⁰⁾. In a large study in Washington state, USA, colorectal cancer patients were 33 % more likely to use folic acid supplements compared with cancer-free controls⁽³¹⁾ and such a supplement use is associated with higher plasma folate and vitamin B₁₂ concentrations⁽³²⁾. Therefore, the use of folic acid-containing supplements is an important determinant of plasma folate concentrations in colorectal cancer patients. This may be of potential concern considering the fact that very high plasma folate concentrations that can be achieved with supplement use may accelerate the growth of already existing colorectal cancers⁽¹⁰⁾. In a randomised controlled trial, folic acid given at a dosage of 1 mg/d was associated with a 1.7-fold increased risk of advanced colorectal cancers⁽¹²⁾, thus excluding its utility and safety as a secondary or tertiary prevention agent for colorectal cancer. However, this is an area of ongoing research and further studies are needed to elucidate the relationship between folate and survival in patients with colorectal cancer.

A particular strength of our study is the comparison of biomarkers across international study sites with identical laboratories for assays and consistent blood-draw protocols. However, our sample size as well as variable set was limited and we had no information on supplement use; hence we could not investigate the association of supplement use with nutritional biomarker concentrations.

In conclusion, we observed substantial differences in nutritional biomarker concentrations across international study sites, probably reflecting dissimilarities in supplement use, food fortification, and sun-seeking behaviour, among other health practices. However, intra-site differences at each study location were larger than between-location variability, suggesting that individual health behaviours play a significant role. As we aim to understand predictors of colorectal cancer prognosis, there is much to learn from international cohorts, which offer a broad range of exposure and health behaviours that cannot be obtained at a single site.

Acknowledgements

We thank all the participants of the DACHS study and the interviewers, physicians and recruiting hospitals who have been involved in the study. We also acknowledge the excellent technical assistance rendered by Ute Handte-Daub, Renate Hettler-Jensen, Belinda Su Kapereit and Muhabbet Celik.

The DACHS study was supported by grants from the German Research Council (Deutsche Forschungsgemeinschaft, grant numbers BR 1704/6-1, BR 1704/6-3, BR 1704/6-4 and CH 390 117/1-1), and the German Federal Ministry of Education and Research (grant numbers 01KH0404 and 01ER0814).

The study was conceived of and designed by C. M. U., E. S., H. B., J. C. C., C. A., M. H., K. M., T. A. S. and W. G. Funding was obtained by C. M. U., E. S., H. B., J. C. C., D. S., K. M., T. A. S. and W. G. Data were acquired by C. M. U., E. S., H. B., J. C. C., C. A., X. S., R. W. O., M. H., D. S., K. V., S. K. R., G. W. and R. H. In charge of quality control of data and algorithms were H. B., J. C. C., C. A., X. S., R. W. O., M. H., K. V., S. K. R., G. W. and R. H. Data analysis and interpretation were performed by C. M. U., A. T. T., E. S., H. B., J. C. C., M. H., H. B., K. M., T. A. S. and W. G. Statistical analysis was done by J. K. and H. B. The manuscript was prepared by A. T. T., C. A., J. K. and S. K. R. The manuscript was edited by C. M. U., E. S., X. S., M. H., H. B., D. S., K. V., K. M., T. A. S. and W. G. The manuscript was reviewed by C. M. U., E. S., H. B., J. C. C., R. W. O., G. W. and R. H.

There is no conflict of interest.

References

1. Ulrich CM (2005) Nutrigenetics in cancer research – folate metabolism and colorectal cancer. *J Nutr* **135**, 2698–2702.
2. Lamprecht SA & Lipkin M (2003) Chemoprevention of colon cancer by calcium, vitamin D and folate: molecular mechanisms. *Nat Rev Cancer* **3**, 601–614.



3. Deeb KK, Trump DL & Johnson CS (2007) Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat Rev Cancer* **7**, 684–700.
4. Jemal A, Bray F, Center MM, *et al.* (2011) Global cancer statistics. *CA Cancer J Clin* **61**, 69–90.
5. Jenab M, Bueno-de-Mesquita HB, Ferrari P, *et al.* (2010) Association between pre-diagnostic circulating vitamin D concentration and risk of colorectal cancer in European populations: a nested case–control study. *Br Med J* **340**, b5500.
6. Gandini S, Boniol M, Haukka J, *et al.* (2011) Meta-analysis of observational studies of serum 25-hydroxyvitamin D levels and colorectal, breast and prostate cancer and colorectal adenoma. *Int J Cancer* **128**, 1414–1424.
7. Yin L, Grandi N, Raum E, *et al.* (2009) Meta-analysis: longitudinal studies of serum vitamin D and colorectal cancer risk. *Aliment Pharmacol Ther* **30**, 113–125.
8. Ng K, Meyerhardt JA, Wu K, *et al.* (2008) Circulating 25-hydroxyvitamin D levels and survival in patients with colorectal cancer. *J Clin Oncol* **26**, 2984–2991.
9. Ng K, Sargent DJ, Goldberg RM, *et al.* (2011) Vitamin D status in patients with stage IV colorectal cancer: findings from Intergroup trial N9741. *J Clin Oncol* **29**, 1599–1606.
10. Kim YI (2006) Folate: a magic bullet or a double edged sword for colorectal cancer prevention? *Int J Gastroenterol Hepatol* **55**, 1387–1389.
11. Ulrich CM & Potter JD (2006) Folate supplementation: too much of a good thing? *Cancer Epidemiol Biomarkers Prev* **15**, 189–193.
12. Cole BF, Baron JA, Sandler RS, *et al.* (2007) Folic acid for the prevention of colorectal adenomas: a randomized clinical trial. *J Am Med Assoc* **297**, 2351–2359.
13. Song J, Medline A, Mason JB, *et al.* (2000) Effects of dietary folate on intestinal tumorigenesis in the apcMin mouse. *Cancer Res* **60**, 5434–5440.
14. Ulrich CM & Potter JD (2007) Folate and cancer – timing is everything. *J Am Med Assoc* **297**, 2408–2409.
15. Dahlin AM, Van Guelpen B, Hultdin J, *et al.* (2008) Plasma vitamin B₁₂ concentrations and the risk of colorectal cancer: a nested case–referent study. *Int J Cancer* **122**, 2057–2061.
16. Eussen SJ, Vollset SE, Hustad S, *et al.* (2010) Plasma vitamins B₂, B₆, and B₁₂, and related genetic variants as predictors of colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* **19**, 2549–2561.
17. Brenner H, Chang-Claude J, Seiler CM, *et al.* (2011) Protection from colorectal cancer after colonoscopy: a population-based, case–control study. *Ann Intern Med* **154**, 22–30.
18. Fenstermacher DA, Wenham RM, Rollison DE, *et al.* (2011) Implementing personalized medicine in a cancer center. *Cancer J* **17**, 528–536.
19. IARC Working Group on Vitamin D (2008) *Vitamin D and Cancer*. Lyon, France: IARC Press.
20. Kimlin MG (2008) Geographic location and vitamin D synthesis. *Mol Aspects Med* **29**, 453–461.
21. Levis S, Gomez A, Jimenez C, *et al.* (2005) Vitamin D deficiency and seasonal variation in an adult South Florida population. *J Clin Endocrinol Metab* **90**, 1557–1562.
22. Woitge HW, Knothe A, Witte K, *et al.* (2000) Circaannual rhythms and interactions of vitamin D metabolites, parathyroid hormone, and biochemical markers of skeletal homeostasis: a prospective study. *J Bone Miner Res* **15**, 2443–2450.
23. Yetley EA (2008) Assessing the vitamin D status of the US population. *Am J Clin Nutr* **88**, 558S–564S.
24. Calvo MS, Whiting SJ & Barton CN (2004) Vitamin D fortification in the United States and Canada: current status and data needs. *Am J Clin Nutr* **80**, 6 Suppl., 1710S–1716S.
25. Ginde AA, Liu MC & Camargo CA Jr (2009) Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* **169**, 626–632.
26. Ulrich CM & Holmes RS (2008) Shedding light on colorectal cancer prognosis: vitamin D and beyond. *J Clin Oncol* **26**, 2937–2939.
27. Aufreiter S, Gregory JF III, Pfeiffer CM, *et al.* (2009) Folate is absorbed across the colon of adults: evidence from cecal infusion of ¹³C-labeled [6S]-5-formyltetrahydrofolic acid. *Am J Clin Nutr* **90**, 116–123.
28. Herrmann W & Obeid R (2011) The mandatory fortification of staple foods with folic acid: a current controversy in Germany. *Dtsch Arztebl Int* **108**, 249–254.
29. Pfeiffer CM, Johnson CL, Jain RB, *et al.* (2007) Trends in blood folate and vitamin B₁₂ concentrations in the United States, 1988–2004. *Am J Clin Nutr* **86**, 718–727.
30. Velicer CM & Ulrich CM (2008) Vitamin and mineral supplement use among US adults after cancer diagnosis: a systematic review. *J Clin Oncol* **26**, 665–673.
31. Greenlee H, White E, Patterson RE, *et al.* (2004) Supplement use among cancer survivors in the vitamins and lifestyle (VITAL) study cohort. *J Altern Complement Med* **10**, 660–666.
32. Yang Q, Cogswell ME, Hamner HC, *et al.* (2010) Folic acid source, usual intake, and folate and vitamin B₁₂ status in US adults: National Health and Nutrition Examination Survey (NHANES) 2003–2006. *Am J Clin Nutr* **91**, 64–72.