

COFINITENESS OF GENERALIZED LOCAL COHOMOLOGY MODULES

KEIVAN BORNA , PARVIZ SAHANDI and SIAMAK YASSEMI

(Received 6 February 2010)

Abstract

Let \mathfrak{a} be an ideal of a Noetherian ring R . Let s be a nonnegative integer and let M and N be two R -modules such that $\text{Ext}_R^j(M/\mathfrak{a}M, H_{\mathfrak{a}}^i(N))$ is finite for all $i < s$ and all $j \geq 0$. We show that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite provided $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is a finite R -module. In addition, for finite R -modules M and N , we prove that if $H_{\mathfrak{a}}^i(M, N)$ is minimax for all $i < s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite. These are two generalizations of the result of Brodmann and Lashgari [‘A finiteness result for associated primes of local cohomology modules’, *Proc. Amer. Math. Soc.* **128** (2000), 2851–2853] and a recent result due to Chu [‘Cofiniteness and finiteness of generalized local cohomology modules’, *Bull. Aust. Math. Soc.* **80** (2009), 244–250]. We also introduce a generalization of the concept of cofiniteness and recover some results for it.

2010 *Mathematics subject classification*: primary 13D45; secondary 13E10.

Keywords and phrases: cofiniteness, generalized local cohomology, local cohomology, minimaxness.

1. Introduction

Throughout this paper R is a commutative Noetherian ring, \mathfrak{a} is an ideal and M and N are two R -modules. Let $H_{\mathfrak{a}}^i(M)$ be the i th local cohomology module of M with respect to \mathfrak{a} . In [10] Grothendieck conjectured that ‘for any finite R -module M , $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^i(M))$ is finite for all i ’. Although Hartshorne disproved Grothendieck’s conjecture (see [11]), there are some partial positive answers to it. For example, in [3, Theorem 4.1] the authors showed that for a nonnegative integer s if M is a finite R -module such that $H_{\mathfrak{a}}^i(M)$ is finite for all $i < s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M))$ is finite. Recall that an R -module M is called \mathfrak{a} -cofinite if $\text{Supp}(M) \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(R/\mathfrak{a}, M)$ is a finite R -module for each i . In [9, Theorem 2.1] the authors improved [3, Theorem 4.1] by showing that for a nonnegative integer s , if M is a finite R -module such that $H_{\mathfrak{a}}^j(M)$ is \mathfrak{a} -cofinite for all $j \leq s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M))$ is finite.

The research of K. Borna was in part supported by a grant from IPM (No. 89130049). The research of P. Sahandi was in part supported by a grant from IPM (No. 89130051). The research of S. Yassemi was in part supported by a grant from IPM (No. 89130213).

© 2011 Australian Mathematical Publishing Association Inc. 0004-9727/2011 \$16.00

Let M and N be two R -modules. The generalized local cohomology module

$$H_{\mathfrak{a}}^i(M, N) = \varinjlim_n \text{Ext}_R^i(M/\mathfrak{a}^n M, N)$$

was introduced by Herzog in [12] and studied further in [17, 18]. Note that $H_{\mathfrak{a}}^i(R, N) = H_{\mathfrak{a}}^i(N)$. Now it is natural to think about Grothendieck's conjecture for the generalized local cohomology modules.

QUESTION 1.1. Let M and N be two finite R -modules. When is

$$\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^i(M, N))$$

finite?

In [2, Theorem 1.2] it is shown that, for a nonnegative integer s , if M and N are two finite R -modules such that $H_{\mathfrak{a}}^i(M, N)$ is finite for all $i < s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite. Recently, in [8] Chu has shown that for a nonnegative integer s , if M and N are two finite R -modules such that $H_{\mathfrak{a}}^i(M, N)$ is Artinian for all $i < s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite.

Let M and N be two R -modules. Then we say that N is (\mathfrak{a}, M) -cofinite if $\text{Supp}(N) \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(M/\mathfrak{a}M, N)$ is a finite R -module for all i . Note that (\mathfrak{a}, R) -cofinite is the classical \mathfrak{a} -cofinite.

In this paper we give some answers to Question 1.1 which improve on some previous results. Our first main result shows that for two R -modules M and N such that $H_{\mathfrak{a}}^i(N)$ is (\mathfrak{a}, M) -cofinite for all $i < s$, the R -module $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite provided $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is finite.

Recall that an R -module M is called minimax if there is a finite submodule N of M such that M/N is Artinian; see [19]. The class of minimax modules includes all finite and all Artinian modules. We show that for finite R -modules M and N if $H_{\mathfrak{a}}^i(M, N)$ is a minimax module for all $i < s$, then for all finite R -submodules L of $H_{\mathfrak{a}}^s(M, N)$, $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L)$ is finite and in particular, $H_{\mathfrak{a}}^s(M, N)$ has finitely many associated primes. This result improves on some previous ones, for example [6, Theorem 2.2], [3, Theorem 4.1], [9, Theorem 2.1], [13, Theorem 2.1], [2, Theorem 1.2] and [8, Theorem 2.5]. We use the terminology and notation of [7].

2. (\mathfrak{a}, M) -cofiniteness

DEFINITION 2.1. Let M and N be two R -modules. We say that N is (\mathfrak{a}, M) -cofinite if $\text{Supp}(N) \subseteq V(\mathfrak{a})$ and $\text{Ext}_R^i(M/\mathfrak{a}M, N)$ is a finite R -module for all i .

Note that (\mathfrak{a}, R) -cofinite is the classical \mathfrak{a} -cofinite. In the following we establish some results on (\mathfrak{a}, M) -cofinite modules. The proofs of the following two results are classical.

LEMMA 2.2. *In a short exact sequence, if two modules in the sequence are (\mathfrak{a}, N) -cofinite, then so is the third.*

COROLLARY 2.3. *If $f : H \rightarrow K$ is a homomorphism of (\mathfrak{a}, N) -cofinite modules and one of $\ker f$, $\text{im } f$ and $\text{coker } f$ is (\mathfrak{a}, N) -cofinite, then all three are (\mathfrak{a}, N) -cofinite.*

THEOREM 2.4. *Let (R, \mathfrak{m}, k) be a local ring. If M is (\mathfrak{m}, N) -cofinite, then $\text{Hom}_R(N, M)$ is Artinian.*

PROOF. The result follows clearly if $\text{Hom}_R(N, M) = 0$. It is well known that a nonzero module M is Artinian if and only if $\text{Supp}_R(M) = \{\mathfrak{m}\}$ and $\text{Hom}(k, M)$ is finite. Note that

$$\text{Hom}_R(N/\mathfrak{m}N, M) \cong \text{Hom}_R(R/\mathfrak{m} \otimes_R N, M) \cong \text{Hom}(R/\mathfrak{m}, \text{Hom}_R(N, M)).$$

Since $\text{Hom}(N/\mathfrak{m}N, M)$ is finite, $\text{Hom}_R(R/\mathfrak{m}, \text{Hom}_R(N, M))$ is finite too. Further $\text{Supp}_R(\text{Hom}_R(N, M)) \subseteq \text{Supp}_R\{M\} \subseteq \{\mathfrak{m}\}$ and $\text{Hom}_R(k, \text{Hom}_R(N, M))$ is finite. \square

LEMMA 2.5. *Let M be a minimax module and N be \mathfrak{a} -cofinite. Then $\text{Ext}_R^i(N/\mathfrak{a}N, M)$ is minimax for all i .*

PROOF. It is well known that in an exact sequence $A \rightarrow B \rightarrow C$ of R -modules and R -homomorphisms, if A and C are minimax, then B is minimax too; see [4, Lemma 2.1]. Then one can deduce that $\text{Hom}_R(M, N)$ is minimax whenever M is finite and N is minimax. Hence for such M and N we have that $\text{Ext}_R^i(M, N)$ is minimax, as it can be seen using a projective resolution for M . Now let $\mathfrak{a} = (x_1, \dots, x_n)$. Then $N/\mathfrak{a}N \cong H^n(x_1, \dots, x_n, N)$. As N is \mathfrak{a} -cofinite, all its Koszul cohomology modules are finite. In particular, $N/\mathfrak{a}N$ is finite. Now apply the argument for the finite case. \square

One can replace ‘minimax’ with ‘finite’ in Lemma 2.5 to deduce the following.

COROLLARY 2.6. *Let M be a finite module and let N be \mathfrak{a} -cofinite. Then $\text{Ext}_R^i(N/\mathfrak{a}N, M)$ is finite for all i . In particular, if $\text{Supp}_R(M) \subseteq V(\mathfrak{a})$, then M is (\mathfrak{a}, N) -cofinite.*

THEOREM 2.7. *Let M be a (\mathfrak{a}, N) -cofinite R -module. Then $\text{Ass}_R(\text{Hom}_R(N, M))$ is finite.*

PROOF. Set $P := \text{Hom}_R(N, M)$. Then $\text{Hom}_R(N/\mathfrak{a}N, M) \cong \text{Hom}_R(R/\mathfrak{a}, P) \cong 0 :_P \mathfrak{a}$ is finite. The essence of [14, Proposition 1.3] is that $0 :_P \mathfrak{a}$ is an essential submodule of P . For this let $0 \neq x \in P$. Since $\text{Supp}_R(P) \subseteq V(\mathfrak{a})$, there is a natural number n such that $\mathfrak{a}^n x = 0$ but $\mathfrak{a}^{n-1} x \neq 0$. Thus $0 \neq \mathfrak{a}^{n-1} x \subseteq Ax \cap 0 :_P \mathfrak{a}$. Hence each submodule of P has a nonzero intersection with $0 :_P \mathfrak{a}$. That is, $0 :_P \mathfrak{a}$ is an essential submodule of P . In other words, P has finite Goldie dimension. Hence $\text{Ass}_R(P)$ is finite. \square

The following result is the counterpart for the change of rings principle for (\mathfrak{a}, P) -cofinite modules, where P is a finite flat module; see [14, Proposition 1.5].

THEOREM 2.8. *Let $f : A \rightarrow B$ be a homomorphism of Noetherian rings. Let \mathfrak{a} be an ideal of A , M an A -module and P a finite flat A -module.*

- (a) *If f is flat, then $M \otimes_A B$ is $(\mathfrak{a}B, P \otimes_A B)$ -cofinite whenever M is (\mathfrak{a}, P) -cofinite.*
- (b) *If f is faithfully flat, the converse of (a) holds as well.*

PROOF. Note that $\text{Ext}_A^i(P/\mathfrak{a}P, M) \otimes_A B \cong \text{Ext}_B^i(P \otimes_A B/\mathfrak{a}B, M \otimes_A B)$; see [16, Proposition 7.39]. Since P is a flat A -module, $P \otimes_A B/\mathfrak{a}B \cong P \otimes_A B/P \otimes_A \mathfrak{a}B \cong (P \otimes_A B)/\mathfrak{a}(P \otimes_A B)$. Hence, $\text{Ext}_A^i(P/\mathfrak{a}P, M) \otimes_A B \cong \text{Ext}_B^i((P \otimes_A B)/\mathfrak{a}(P \otimes_A B), M \otimes_A B)$. \square

3. Cofiniteness and minimaxness

There are several papers devoted to partially answering Question 1.1 in more general situations, for example [3, 6, 8, 9, 13]. The following theorem is the first main result of this paper in this vein.

THEOREM 3.1. *Let s be a nonnegative integer. Let M and N be two R -modules such that $H_{\mathfrak{a}}^i(N)$ is (\mathfrak{a}, M) -cofinite for all $i < s$. If $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is a finite (respectively minimax) R -module, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite (respectively minimax).*

PROOF. We proceed by induction on s . If $s = 0$, then $H_{\mathfrak{a}}^0(M, N) \cong \Gamma_{\mathfrak{a}}(\text{Hom}(M, N))$ and $\text{Hom}_R(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(\text{Hom}_R(M, N)))$ is equal to the finite (respectively minimax) R -module

$$\text{Hom}_R(R/\mathfrak{a}, \text{Hom}_R(M, N)) \cong \text{Hom}_R(M/\mathfrak{a}M, N).$$

Suppose that $s > 0$ and that the case $s - 1$ is settled. We have that $\text{Ext}_R^j(M/\mathfrak{a}M, \Gamma_{\mathfrak{a}}(N))$ is finite (respectively minimax) for all $j \geq 0$. Using the exact sequence $0 \rightarrow \Gamma_{\mathfrak{a}}(N) \rightarrow N \rightarrow N/\Gamma_{\mathfrak{a}}(N) \rightarrow 0$, we get that $\text{Ext}_R^s(M/\mathfrak{a}M, N/\Gamma_{\mathfrak{a}}(N))$ is finite (respectively minimax). On the other hand, $H_{\mathfrak{a}}^0(M, N/\Gamma_{\mathfrak{a}}(N)) = 0$ and for all $i > 0$ and all $j \geq 0$,

$$\text{Ext}_R^j(M/\mathfrak{a}M, H_{\mathfrak{a}}^i(N/\Gamma_{\mathfrak{a}}(N))) \cong \text{Ext}_R^j(M/\mathfrak{a}M, H_{\mathfrak{a}}^i(N)).$$

Thus we may assume that $\Gamma_{\mathfrak{a}}(N) = 0$. Let E be an injective hull of N and put $T = E/N$. Then $\Gamma_{\mathfrak{a}}(E) = 0$ and $\text{Hom}_R(M/\mathfrak{a}M, E) = 0$. Consequently, $\text{Ext}_R^i(M/\mathfrak{a}M, T) \cong \text{Ext}_R^{i+1}(M/\mathfrak{a}M, N)$ and $H_{\mathfrak{a}}^i(M, T) \cong H_{\mathfrak{a}}^{i+1}(M, N)$ for all $i \geq 0$. Now the induction hypothesis yields that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{s-1}(M, T))$ is finite (respectively minimax) and hence $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite (respectively minimax). \square

COROLLARY 3.2. *Let s be a nonnegative integer. Let \mathfrak{a} be an ideal of R and let M and N be two R -modules. Let L be a submodule of $H_{\mathfrak{a}}^s(M, N)$ such that $\text{Ext}_R^1(R/\mathfrak{a}, L)$ is a finite (respectively minimax) R -module. If $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is a finite (respectively minimax) R -module and $H_{\mathfrak{a}}^i(N)$ is (\mathfrak{a}, M) -cofinite for all $i < s$, then the module $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L)$ is finite (respectively minimax). In particular, $H_{\mathfrak{a}}^s(M, N)/L$ has finitely many associated primes.*

PROOF. Let L be a submodule of $H_{\mathfrak{a}}^s(M, N)$ such that $\text{Ext}_R^1(R/\mathfrak{a}, L)$ is a finite (respectively minimax) R -module. The short exact sequence $0 \rightarrow L \rightarrow H_{\mathfrak{a}}^s(M, N) \rightarrow H_{\mathfrak{a}}^s(M, N)/L \rightarrow 0$ induces the exact sequence

$$\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)) \rightarrow \text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, L).$$

Since by Theorem 3.1 the left-hand term and by hypothesis the right-hand term are finite (respectively minimax), we have that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L)$ is finite (respectively minimax). For the last statement note that

$$\text{Supp}(H_{\mathfrak{a}}^s(M, N)/L) \subseteq \text{Supp}(H_{\mathfrak{a}}^s(M, N)) \subseteq V(\mathfrak{a}).$$

Therefore $\text{Ass}(\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L)) = \text{Ass}(H_{\mathfrak{a}}^s(M, N)/L)$ is a finite set. \square

The following corollary is the main result of Brodmann and Lashgari [6].

COROLLARY 3.3. *Let M be a finite R -module. Let s be a nonnegative integer such that $H_{\mathfrak{a}}^i(M)$ is finite for each $i < s$. Then for any finite submodule N of $H_{\mathfrak{a}}^s(M)$, the set $\text{Ass}(H_{\mathfrak{a}}^s(M)/N)$ has finitely many elements.*

One can define the term (\mathfrak{a}, M) -coartinian by replacing ‘Artinian’ with ‘finite’ in our definition of (\mathfrak{a}, M) -cofinite. Then a similar proof as for Theorem 3.1 implies the following.

THEOREM 3.4. *Let s be a nonnegative integer. Let M and N be two R -modules such that $H_{\mathfrak{a}}^i(N)$ is (\mathfrak{a}, M) -coartinian for all $i < s$. If $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is an Artinian R -module, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is Artinian.*

Using [1, Theorem 2.9], we are able to express this result in several equivalent situations. Note that by [1, Example 2.4(b)] the class of Artinian R -modules is closed under taking submodules, quotients, extensions and injective hulls. Hence it satisfies condition C_a in the notation of [1, Theorem 2.9].

COROLLARY 3.5. *Let \mathfrak{a} be an ideal of a Noetherian ring R . Let s be a nonnegative integer. Let M and N be two R -modules such that $M/\mathfrak{a}M$ is finite and $\text{Ext}_R^s(M/\mathfrak{a}M, N)$ is Artinian. Then in any of the following cases, $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is Artinian:*

- (a) $H_{\mathfrak{a}}^i(N)$ is Artinian for all $i < s$;
- (b) $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is Artinian for all $i < s$;
- (c) $\text{Ext}_R^i(T, N)$ is Artinian for all $i < s$ and each finite module T such that $\text{Supp}_R(T) \subseteq V(\mathfrak{a})$;
- (d) there is a finite R -module T with $\text{Supp}_R(T) = V(\mathfrak{a})$ such that $\text{Ext}_R^i(T, N)$ is Artinian for all $i < s$;
- (e) $H^i(x_1, \dots, x_r, N)$ is Artinian for all $i < s$ where x_1, \dots, x_r generate \mathfrak{a} ;
- (f) $H_{\mathfrak{a}}^i(T, N)$ is Artinian for each finite R -module T and for all $i < s$.

The following theorem is the second main result of this paper.

THEOREM 3.6. *Let M and N be finite R -modules. If $H_{\mathfrak{a}}^i(M, N)$ is a minimax module for all $i < s$, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite.*

PROOF. First we prove the theorem under the additional assumption that $H_{\mathfrak{a}}^i(M, N)$ is \mathfrak{a} -cofinite module for all $i < s$. If $s = 0$, then $H_{\mathfrak{a}}^0(M, N) \cong \Gamma_{\mathfrak{a}}(\text{Hom}_R(M, N))$ is a

finite R -module. Now suppose that $s > 0$. The short exact sequence $0 \rightarrow \Gamma_{\mathfrak{a}}(N) \rightarrow N \rightarrow N/\Gamma_{\mathfrak{a}}(N) \rightarrow 0$ induces the exact sequence

$$H_{\mathfrak{a}}^t(M, N) \rightarrow H_{\mathfrak{a}}^t(M, N/\Gamma_{\mathfrak{a}}(N)) \rightarrow H_{\mathfrak{a}}^{t+1}(M, \Gamma_{\mathfrak{a}}(N)).$$

For $t < s$ the R -module $H_{\mathfrak{a}}^t(M, N)$ is \mathfrak{a} -cofinite and minimax and by [13, Lemma 1.1] the R -module $H_{\mathfrak{a}}^{t+1}(M, \Gamma_{\mathfrak{a}}(N))$ is finite. Thus by [15, Corollary 4.4] we have that $H_{\mathfrak{a}}^t(M, N/\Gamma_{\mathfrak{a}}(N))$ is an \mathfrak{a} -cofinite and minimax R -module. Thus without loss of generality we can assume that $\Gamma_{\mathfrak{a}}(N) = 0$. Choosing an arbitrary N -regular element x in \mathfrak{a} and using an argument similar to the proof of [4, Lemma 2.2], we obtain that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N))$ is finite.

Next suppose that $H_{\mathfrak{a}}^i(M, N)$ is minimax module for all $i < s$. In view of the first part of the proof, it is enough to show that $H_{\mathfrak{a}}^i(M, N)$ is \mathfrak{a} -cofinite for all $i < s$. We proceed by induction on i . The case $i = 0$ is obvious as $H_{\mathfrak{a}}^0(M, N)$ is finite. Thus let $i > 0$, and assume the result has been proved for smaller values of i . By the inductive hypothesis, $H_{\mathfrak{a}}^j(M, N)$ is \mathfrak{a} -cofinite for $j < i$. Thus by the first part, $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^i(M, N))$ is finite. Therefore by [15, Proposition 4.3], $H_{\mathfrak{a}}^i(M, N)$ is \mathfrak{a} -cofinite. Hence $H_{\mathfrak{a}}^i(M, N)$ is \mathfrak{a} -cofinite minimax for all $i < s$. Now the assertion follows from the first part. \square

By an argument similar to the proof of Corollary 3.2 we have the following corollary.

COROLLARY 3.7. *Let s be a nonnegative integer. Let \mathfrak{a} be an ideal of R and let M and N be two finite R -modules. Let L be a submodule of $H_{\mathfrak{a}}^s(M, N)$ such that $\text{Ext}_R^1(R/\mathfrak{a}, L)$ is a finite R -module. If $H_{\mathfrak{a}}^i(M, N)$ is minimax for all $i < s$, then the module $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M, N)/L)$ is finite. In particular, $H_{\mathfrak{a}}^s(M, N)/L$ has finitely many associated primes.*

Applying Theorem 3.6, we have the following result; see [5, Theorem 2.2].

COROLLARY 3.8. *Let \mathfrak{a} be an ideal of R and let s be a nonnegative integer. Let N be an R -module such that $\text{Ext}_R^s(R/\mathfrak{a}, N)$ is a finite R -module. If $H_{\mathfrak{a}}^i(N)$ is minimax for all $i < s$ then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(N))$ is a minimax module. Furthermore, if L is a finite R -module such that $\text{Supp}(L) \subseteq V(\mathfrak{a})$, then $\text{Hom}_R(L, H_{\mathfrak{a}}^s(N))$ is a minimax module.*

In the same way we can apply Theorem 3.6 to deduce the following result; see [4, Theorem 2.3].

COROLLARY 3.9. *Let R be a Noetherian ring, M a nonzero finite R -module and \mathfrak{a} an ideal of R . Let s be a nonnegative integer such that $H_{\mathfrak{a}}^i(M)$ is minimax for all $i < s$. Then the R -module $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^s(M))$ is finite. In particular, $\text{Ass}_R(H_{\mathfrak{a}}^s(M))$ is finite.*

References

- [1] M. Aghapournahr and L. Melkerson, 'Local cohomology and Serre subcategories', *J. Algebra* **320** (2008), 1275–1287.
- [2] J. Asadollahi, K. Khashyarmansh and Sh. Salarian, 'On the finiteness properties of the generalized local cohomology modules', *Comm. Algebra* **30** (2002), 859–867.

- [3] J. Asadollahi, K. Khashyarmanesh and Sh. Salarian, 'A generalization of the cofiniteness problem in local cohomology modules', *J. Aust. Math. Soc.* **75** (2003), 313–324.
- [4] K. Bahmanpour and R. Naghipour, 'On the cofiniteness of local cohomology modules', *Proc. Amer. Math. Soc.* **136**(7) (2008), 2359–2363.
- [5] K. Borna, P. Sahandi and S. Yassemi, 'Artinian local cohomology modules', *Canad. Bull. Math.* **50**(4) (2007), 598–602.
- [6] M. P. Brodmann and A. Lashgari, 'A finiteness result for associated primes of local cohomology modules', *Proc. Amer. Math. Soc.* **128** (2000), 2851–2853.
- [7] M. P. Brodmann and R. Y. Sharp, *Local Cohomology: an Algebraic Introduction with Geometric Applications*, Cambridge Studies in Advanced Mathematics, 60 (Cambridge University Press, Cambridge, 1998).
- [8] L. Chu, 'Cofiniteness and finiteness of generalized local cohomology modules', *Bull. Aust. Math. Soc.* **80** (2009), 244–250.
- [9] M. T. Dibaei and S. Yassemi, 'Associated primes and cofiniteness of local cohomology modules', *Manuscripta Math.* **117** (2005), 199–205.
- [10] A. Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Advanced Studies in Pure Mathematics, 2 (North-Holland, Amsterdam, 1968).
- [11] R. Hartshorne, 'Affine duality and cofiniteness', *Invent. Math.* **9** (1970), 145–164.
- [12] J. Herzog, 'Komplexe, Auflösungen und Dualität in der lokalen Algebra', Preprint, University of Regensburg, 1974.
- [13] L. Khatami, T. Sharif and S. Yassemi, 'Associated primes of generalized local cohomology modules', *Comm. Algebra* **30** (2002), 327–330.
- [14] L. Melkersson, 'Properties of cofinite modules and applications to local cohomology', *Math. Proc. Cambridge Philos. Soc.* **125** (1999), 417–423.
- [15] L. Melkersson, 'Modules cofinite with respect to an ideal', *J. Algebra* **285** (2005), 649–668.
- [16] J. Rotman, *Introduction to Homological Algebra* (Academic Press, New York, 1979).
- [17] N. Suzuki, 'On the generalized local cohomology and its duality', *J. Math. Kyoto Univ.* **18** (1978), 71–85.
- [18] S. Yassemi, 'Generalized section functor', *J. Pure Appl. Algebra* **95** (1994), 103–119.
- [19] H. Zöschinger, 'Minimax-Moduln', *J. Algebra* **102** (1986), 1–32.

KEIVAN BORNA, Faculty of Mathematical Sciences and Computer,
Tarbiat Moallem University, Tehran, Iran
and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM),
Tehran, Iran
e-mail: borna@ipm.ir

PARVIZ SAHANDI, Department of Mathematics, University of Tabriz, Tabriz, Iran
and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM),
Tehran, Iran
e-mail: sahandi@ipm.ir

SIAMAK YASSEMI, Department of Mathematics, University of Tehran, Tehran, Iran
and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM),
Tehran, Iran
e-mail: yassemi@ipm.ir