

Executive function and in-patient violence in forensic patients with schizophrenia

Rachael S. Fullam and Mairead C. Dolan

Background

The literature on the association between neuropsychological deficits and in-patient violence in schizophrenia is limited and the findings inconsistent.

Aims

To examine the role of executive function deficits in in-patient violence using measures of dorsolateral (DLPFC) and ventrolateral prefrontal cortical (VLPFC) function.

Methods

Thirty-three violent and forty-nine non-violent male forensic in-patients with schizophrenia were assessed using neuropsychological tasks probing DLPFC and VLPFC function and on measures of symptoms and psychopathy.

Results

There were no significant group differences in

neuropsychological task performance. Higher rates of violence were significantly associated with lower current IQ scores and higher excitement symptom scores. The 'violent' group had significantly higher interpersonal and antisocial domain psychopathy scores. In a logistic regression analysis, IQ and the interpersonal domain of psychopathy were significant discriminators of violent v. non-violent status.

Conclusions

Personality factors rather than symptoms and neuropsychological function may be important in understanding in-patient violence in forensic patients with schizophrenia.

Declaration of interest

None. Funding detailed in Acknowledgements.

Reviews of the literature on the association between executive function and community or in-patient violence in schizophrenia have produced inconsistent findings.¹ Some,^{2–6} but not all studies,^{7–10} report links between violence and impairment in executive function. The inconsistent findings may reflect methodological variation between studies including: definitions of violence; sample heterogeneity; acute v. chronic illness; comorbidity with antisocial personality/psychopathy; variation in the neuropsychological test batteries used; and a lack of differentiation between tasks sensitive to dorsolateral and ventrolateral prefrontal cortical (DLPFC and VLPFC respectively) function. The majority of studies focus on community violence with only four studies to date examining in-patient violence.^{2,3,4,7} In-patient violence is not necessarily predicted by previous community violence¹¹ and may therefore show a different relationship to executive function in people with schizophrenia. In addition, no previous study in the current literature has adequately examined the interactive role of personality factors such as psychopathy in the relationship between executive function and violence in a large sample of people with schizophrenia. The aim of the present study was to examine the contribution of executive function to in-patient violence in schizophrenia by utilising a large forensic sample of individuals and looking at neuropsychological tasks specific to DLPFC and VLPFC function. The study also examined the role of psychopathy and symptoms in the discrimination of violent v. non-violent status.

majority ($n=64$, 78%) of the sample were White. The remainder were of Black African–Caribbean origin. Potential participants were identified by core clinical teams using the following criteria:

- inclusion: currently clinically stable on medication, able to give informed consent;
- exclusion: a history of organic brain syndrome, head injury or recent history of electroconvulsive therapy.

Symptom and psychopathy assessment

Symptom severity was assessed using the Positive and Negative Syndrome Scale (PANSS).¹³ The five-factor structure for the PANSS formulated by Lindenmayer *et al*¹⁴ was computed. The latter model uses 25 of the 30 items on the PANSS to form five factors: positive, negative, cognitive, excitement and depression.

Psychopathy was assessed based on interview and file review using the Psychopathy Checklist – Screening Version (PCL–SV).¹⁵ Factor 1 of the PCL–SV reflects affective/interpersonal traits and factor 2 reflects the behavioural/social deviance components of psychopathy. Data were analysed using both the two-factor and more recent four-facet (interpersonal, affective, lifestyle and antisocial) models of psychopathy.^{16,17} As there are no established UK cut-off scores for psychopathy using the PCL–SV, the UK cut-off score for psychopathy on the Psychopathy Checklist Revised¹⁸ was translated across to the PCL–SV using percentile points from the development samples. This resulted in cut-offs of ≥ 17 for psychopathy and < 11 for non-psychopathy. Interrater reliability checks in ten cases resulted in an intraclass correlation of 0.93 for total score.

Method

Participants

A sample of 82 male in-patients meeting the DSM–IV¹² criteria for schizophrenia (diagnosis made by each participant's responsible medical officer) were recruited (approximate recruitment rate was one in three) from medium- ($n=53$) and high- ($n=29$) security forensic hospitals in the North-West of England. The

Assessment of in-patient violence

An independent researcher masked to scores on the psychometric and neuropsychological measures reviewed computerised official incident records within the hospital. An incident was considered violent if the individual was the clear instigator or co-aggressor,

and if the incident involved physical aggression to staff, in-patients or property. The median number of violent incidents across the sample was 0. Based on this we assigned groups into non-violent (non-violent=0 incidents) and violent (violent \geq 1 incident) groups. This generated 49 individuals that had not been physically violent since admission and 33 individuals that had been involved in at least one physically aggressive incident within the institution since admission. Of the 33 violent participants, 11 had been involved in one incident since admission, 8 had been involved in between two and five incidents, 7 had been involved in between six and ten incidents, and 7 had been involved in ten or more incidents. Rate of violent incidents per year since admission were also calculated for each participant.

Neuropsychological assessments

Premorbid intellectual function was assessed using the National Adult Reading Test (NART).¹⁹ Current IQ was assessed using sub-tests (vocabulary and matrix reasoning) of the Wechsler Abbreviated Scale of Intelligence (WASI).²⁰

The Cambridge Automated Neuropsychological Test Battery (CANTAB-2)²¹ was used to assess spatial planning ability (Stockings of Cambridge) and cognitive set shifting (intra-/extra-dimensional set shifting). The CANTAB-2 is a culture-free visual computerised assessment battery that overcomes assessment problems resulting from poor reading ability. Participants were also tested on the Stop Task behavioural inhibition task that was developed by Rubia *et al*²² as an adaptation of the Schacher & Logan²³ task (a description of each task is detailed in online Table DS1).

Data analysis

All data were analysed using SPSS version 12. As a large proportion of the sample had failed to complete one or more tasks in the battery ($n=42$), we were unable to use a multivariate analysis of variance to examine all neuropsychological variables across tasks simultaneously. Between-group differences were examined using independent *t*-tests and chi-squared tests where appropriate. In order to examine the violence data dimensionally and to account for possible differences in length of admission, we used Spearman's correlation coefficient to investigate the relationship between the rate of violent incidents per year since admission and the neuropsychological, symptomatological and neuropsychological variables of interest. Data that were not normally distributed were transformed using square root or log base 10 transformations to reduce skew. The categorical outputs from

the intra-/extra-dimensional set shifting task were analysed using the likelihood ratio method with the resulting statistic $2i$ being distributed as χ^2 . In order to control for multiple statistical comparisons, Bonferroni corrections were applied to acceptable probability levels for each set of analyses.

A binary logistic regression using the enter method was used to examine the prediction of non-violent *v.* violent group status. Only the symptomatological, personality and neuropsychological variables that were significantly different in the univariate non-violent *v.* violent group comparisons were entered as predictors.

Results

Group characteristics

The violent group had a longer mean current admission length than the non-violent group, although this difference failed to reach significance following Bonferroni corrections. There were no significant group differences between the violent and non-violent groups in terms of mean age, years of education, medication dose (converted to chlorpromazine equivalents using the British National Formulary),²⁴ age of first criminal offence, or number of previous criminal offences. Similarly, these variables did not show significant relationships with the rate of violent incidents per year since admission (Table 1). There were no significant differences in the proportion of each group who had a previous history of violent offending (non-violent 34.0% *v.* violent 53.1%; $\chi^2=2.85$, not significant), or those who had a history of substance misuse (drug or alcohol) (non-violent 31.3% *v.* violent 42.4%; $\chi^2=0.70$, not significant).

Symptoms

The violent group had a higher PANSS excitement scale score than the non-violent group, although this group difference failed to reach significance following Bonferroni correction. However, there was a significant positive correlation between PANSS excitement scale score and rate of violent incidents per year since admission ($r=0.35$, $P=0.001$). There were no significant differences between the violent and non-violent groups on any of the remaining PANSS symptom scales, and scores on these scales showed no dimensional relationship to violence (Table 2).

Psychopathy

Overall, 7 (14.3%) of the non-violent group and 13 (39.4%) of the violent group met the UK criteria for a diagnosis of psychopathy

Table 1 General characteristics of the non-violent and violent groups

	Non-violent group ($n=49$)		Violent group ($n=33$)		<i>t</i> (d.f.=80)	<i>P</i>	Cohen's <i>d</i>
	Mean	(s.d.)	Mean	(s.d.)			
Age, years	36.84	(9.36)	35.52	(9.61)	0.62	ns	0.14
Years in education ^a	11.24	(1.90)	10.97	(1.65)	0.68	ns	0.15
Length of current admission, years (min. 0.17, max. 25.80) ^a	3.40	(4.44)	6.48	(6.55)	-2.35	0.022 ^b	-0.55
Illness duration since first psychiatric admission, years ^a	11.24	(9.12)	12.58	(9.20)	-0.60	ns	-0.15
Age at first criminal offence, years	18.69 ^c	(6.43)	17.47 ^d	(4.52)	0.68	ns	0.22
Previous criminal offences, n^a	8.76	(12.13)	13.19 ^e	(15.11)	-1.49	ns	-0.32
Medication, mg/day ^f	461.84	(362.75)	572.99	(454.53)	-0.91	ns	-0.27

ns, not significant.

a. Performed using log transformed data

b. ns after Bonferroni corrections.

c. $n=48$.

d. $n=30$.

e. $n=31$.

f. Converted into standard chlorpromazine units. Analysis performed using square root transformations.

Table 2 The mean Positive and Negative Syndrome Scale (PANSS) scores for non-violent and violent groups

	Non-violent group (n=49)		Violent group (n=33)		t (d.f.=80)	P	Cohen's d
	Mean	(s.d.)	Mean	(s.d.)			
Positive scale ^a	9.22	(5.40)	10.52	(4.88)	-1.39	ns	-0.25
Negative scale ^a	13.35	(6.59)	10.73	(4.63)	1.85	ns	0.46
Cognitive scale ^a	7.49	(2.27)	8.18	(2.52)	-1.31	ns	-0.29
Excitement scale ^a	5.69	(2.62)	7.09	(3.21)	-2.33	0.022 ^b	-0.48
Depression scale ^a	8.33	(2.87)	8.00	(3.00)	0.59	ns	0.11
Total score	52.53	(17.02)	53.63	(12.12)	-0.32	ns	-0.07

a. Analysis performed using log transformed data.
b. ns after Bonferroni corrections.

on the PCL-SV ($\chi^2=12.17$, $P<0.001$). The violent group had significantly higher total psychopathy and traditional sub-factor scores than the non-violent group. In addition, rate of violent incidents per year since admission showed a significant positive correlation with PCL-SV total score ($r=0.41$, $P=0.001$), factor 1 ($r=0.30$, $P=0.006$), and factor 2 scores ($r=0.36$, $P=0.001$). Analysis of the four-facet model revealed that the violent group had significantly higher scores on the interpersonal and antisocial factors. The violent group also showed higher scores on the lifestyle facet, although this difference failed to reach significance following Bonferroni corrections. Similarly, the correlational analysis revealed significant positive correlations between rate of violent incidents per year since admission and scores on the interpersonal ($r=0.34$, $P=0.002$) and antisocial facets ($r=0.39$, $P=0.001$). The correlation between rate of violence and scores on the lifestyle facet failed to reach significance following correction ($r=0.25$, $P=0.022$). There were no significant group differences found for the affective facet, and scores on this facet did not show a significant relationship to rate of violent incidents (Table 3).

Neuropsychological function

The violent group had a lower mean WASI IQ, although this group difference failed to reach significance following Bonferroni correction. However, there was a significant negative correlation between WASI IQ score and rate of violent incidents per year since admission ($r=-0.32$, $P=0.004$). The non-violent group had higher mean scores on the WASI vocabulary sub-test and scores on this scale showed a negative correlation with rate of violent incidents ($r=-0.23$, $P=0.046$), but neither result reached significance following Bonferroni correction. Similarly, mean NART IQ score was also higher in the non-violent group, and NART IQ scores showed a negative correlation with rate of violence ($r=-0.26$, $P=0.041$), but again these results did not reach significance following Bonferroni correction (Table 4).

There were no significant differences between the violent and non-violent groups on the Stockings of Cambridge, intra-/extra-dimensional set shifting, or Stop tests using either the categorical or dimensional analyses of outputs from these tasks (Table 4 and Fig.1). Similarly, scores on these assessments showed no significant relationship with rate of violent incidents per year since admission.

Predicting violent/non-violent group status

A logistic regression was carried out using the enter method for the prediction of non-violent *v.* violent group status. Current IQ, PANSS excitement scale score and PCL-SV interpersonal and antisocial facet scores were entered as independent variables into the regression in one block. A total of 82 cases were included in the regression analysis. The Hosmer and Lemeshow Test indicated reasonable goodness of fit ($\chi^2=8.85$, d.f.=8, $P=0.36$). The Nagelkerke R^2 was 0.33. The model was significant ($\chi^2=22.84$, d.f.=4, $P=0.001$), overall correct classification was 74.4%. As can be seen in Table 5, current IQ score and PCL-SV interpersonal facet score contributed significantly to the equation. The PANSS excitement scale score and PCL-SV antisocial facet score did not contribute significantly.

Discussion

This study investigated the role of neuropsychological factors, particularly executive function deficits, in in-patient violence in schizophrenia. In order to control for methodological problems with previous studies in this field we ensured that our groups were large and well-matched on potential confounding variables such as demography, clinical and criminal history profile.

This is one of the few studies comparing violent and non-violent forensic in-patients with schizophrenia on measures of neuropsychological function using a well-validated culture-free

Table 3 The mean psychopathy scores for the non-violent and violent groups

PCL-SV scores	Non-violent group (n=49)		Violent group (n=33)		t-value (d.f.=80)	P	Cohen's d
	Mean	(s.d.)	Mean	(s.d.)			
Total score	10.80	(4.92)	14.82	(4.75)	-3.68	0.001	-0.83
Factor 1	5.12	(2.79)	6.97	(3.15)	-2.79	0.007	-0.62
Factor 2	5.67	(3.26)	7.85	(2.71)	-3.17	0.002	-0.73
Interpersonal	1.41	(1.41)	2.70	(1.98)	-3.23	0.002	-0.75
Affective	3.69	(1.88)	4.24	(1.70)	-1.35	ns	-0.31
Lifestyle	2.80	(1.63)	3.52	(1.50)	-2.02	0.047 ^a	-0.46
Antisocial	2.94	(1.93)	4.27	(1.44)	-3.38	0.001	-0.78

PCL-SV, Psychopathy Checklist – Screening Version.
a. ns after Bonferroni corrections.

Table 4 The neuropsychological assessment scores for each group comparison

	Non-violent group (n=49)		Violent group (n=33)		t (d.f.=80)	P	Cohen's d
	Mean	(s.d.)	Mean	(s.d.)			
NART IQ ^a	105.61	(10.53)	98.91	(14.55)	2.11	0.04 ^g	0.53
WASI IQ	94.67	(16.33)	83.85	(16.68)	2.92	0.005 ^g	0.66
WASI Vocabulary score	43.85	(12.97)	37.50	(12.32)	2.18	0.032 ^g	0.50
WASI Matrix Reasoning score	42.98	(16.52)	39.41	(14.80)	0.98	ns	0.23
Stockings of Cambridge ^b							
Mean number of moves: 4 move problems	6.03	(1.23)	6.09	(0.98)	-0.21	ns	-0.05
Mean number of moves: 5 move problems	7.74	(1.75)	7.82	(1.59)	-0.21	ns	-0.05
Initial thinking time: 4 move problems ^c (sec)	11.25	(7.47)	10.19	(7.05)	0.73	ns	0.15
Initial thinking time: 5 move problems ^c	12.82	(14.91)	12.61	(22.48)	1.08	ns	0.01
Intra-extra-dimensional set shift ^d							
Extra-dimensional shift stage errors	17.09	(10.79)	17.72	(11.32)	-0.25	ns	-0.06
Reversal stage errors	5.10	(2.83)	5.85	(3.12)	-1.08	ns	-0.25
DMTS ^e							
Total correct	10.97	(2.42)	10.92	(3.49)	0.49	ns	0.02
Number correct – long delay	3.61	(1.09)	3.68	(1.28)	0.16	ns	-0.06
Number correct – medium delay	3.42	(1.35)	3.60	(1.53)	-0.10	ns	-0.12
Number correct – short delay	3.94	(1.06)	3.68	(1.49)	1.06	ns	0.20
Stop probability of inhibition ^f	65.85	(25.91)	55.90	(24.49)	1.52	ns	0.39

DMTS, Delayed Matching to Sample; NART, National Adult Reading Test; ns, not significant; WASI, Wechsler Abbreviated Scales of Intelligence.

a. Non-violent n=41, violent n=22.
b. Non-violent n=46, violent n=31.
c. Analysis performed using log transformed data.
d. Non-violent n=47, violent n=29.
e. Non-violent n=34, violent n=25.
f. Non-violent n=41, violent n=24.
g. ns after Bonferroni corrections.




Fig. 1 The proportion of non-violent and violent groups reaching criterion at each stage of the intra-/extra-dimensional set shift task.

Extra-dimensional shift (EDS) stage likelihood ratio=0.17, d.f.=1, not significant; extra-dimensional reversal (EDR) stage likelihood ratio=0.34, not significant. CD, compound discrimination; CR, compound reversal; IDS, intra-dimensional shift; IDR, intra-dimensional reversal.

computerised test battery that distinguishes putative DLPFC and VLPFC functions. We also compared the groups on measures of symptoms and measures of psychopathy as these variables have also been reported to be associated with violence in schizophrenia (for example, symptoms,^{25,26} psychopathy^{27–29}).

Neuropsychological function

We found that higher rates of in-patient violence were associated with lower current IQ scores, and that current IQ was a significant predictor of violent *v.* non-violent status. The evidence for an association between IQ and violence is contradictory with some,² but not all,⁷ studies reporting lower IQ in violent compared with

non-violent in-patients with schizophrenia. In the general violence literature, low IQ has been associated with an increased risk for violence,^{30,31} and there is evidence to suggest that low IQ combined with psychopathy presents with an additive risk for increased violence.³² As IQ shows a strong association with number of years in education,³³ it is possible that educational factors may account for the discrepant findings across studies. In addition, although this comparison was not possible in the present study, the findings of studies focusing on community violence³⁴ would suggest that verbal IQ may be more strongly associated with in-patient violence than performance IQ.

To date, there is a lack of well-powered studies examining the relationship between in-patient violence and specific neuropsychological deficits in schizophrenia. None the less, based on Naudts & Hodgins¹ review on neuropsychological function and community violence in schizophrenia, we had postulated that our violent in-patients would have greater deficits in executive function and behavioural inhibition than those who were not violent. However, we did not find this to be the case. Our findings concur with previous reports that DLPFC function is not specifically associated with violence in individuals with schizophrenia.^{4–6,9,10}

Our finding that the violent group did not have a specific impairment in putative VLPFC function on the Stop Task generally fits with previous reports on community violence in schizophrenia.⁸ However, Rasmussen *et al*⁵ noted that forensic patients with schizophrenia showed poorer performance on another type of behavioural inhibition task (Go/NoGo task) than non-forensic patients, suggesting that criminality (rather than violence *per se*) may be associated with poor behavioural inhibition. As violence can be characterised in the instrumental *v.* reactive domain, it is possible that future studies may find a more specific association between impairments in behavioural inhibition and reactive rather than instrumental aggression and violence.

Table 5 Logistic regression for the prediction of non-violent *v.* violent group status

	Wald	P	Exp(B)	95% CI for Exp(B)	
				Lower	Upper
PANSS excitement scale score	0.07	0.798	1.03	0.84	1.25
WASI IQ	4.90	0.027	0.96	0.93	0.99
PCL–SV Interpersonal score	6.01	0.014	1.49	1.08	2.04
PCL–SV Antisocial score	2.60	0.107	1.32	0.94	1.85

PANSS, Positive and Negative Syndrome Scale; PCL–SV, Psychopathy Checklist – Screening Version; WASI, Wechsler Abbreviated Scale of Intelligence.

In this study we used the Stop Task as a putative measure of ventrolateral prefrontal function, and the intra-/extra-dimensional set shifting and Stockings of Cambridge tasks as putative measures of DLPFC. It is possible that tasks assessing orbitofrontal function rather than ventrolateral prefrontal function, such as smell discrimination/identification tests,³⁵ may be able to differentiate violent from non-violent groups. Given the literature suggesting that people with schizophrenia who engage in community violence show impairments on tasks assessing social cognition (i.e. theory of mind³⁶ and face expression recognition³⁷), future studies should also explore the utility of these tasks in distinguishing those who engage in in-patient violence.

It is important to note that although specific neuropsychological tasks are thought to probe specific brain regions, it is increasingly recognised that they actually activate integrated neural circuits that include both frontal and limbic brain regions. Future studies should use functional magnetic resonance imaging techniques to examine subtle dysfunction in neurocircuitry that may contribute to community and in-patient violence in people with schizophrenia.

It is possible that illness-related reductions in IQ may have masked subtle group differences in executive function. In the present study, due to small group sizes, we were unable to examine the relationship between in-patient violence and executive function in those whose IQ had remained stable with illness onset. Future studies may want to address this issue, although, Elliott *et al.*³⁸ have demonstrated that neuropsychological dysfunction is present and detectable in people with schizophrenia regardless of whether or not their IQ remains stable.

Symptoms

Overall, we did not find that the violent patients had higher positive (hallucinations and delusions) symptom scores on the PANSS. However, higher scores on the PANSS excitement scale (which contains items such as hostility and poor impulse control) were associated with higher rates of in-patient violence. Previous studies have reported an association between high PANSS positive scores and high rates of in-patient aggression,^{39,40} or higher levels of positive symptoms in violent compared with non-violent individuals.^{25,41} The lack of an observed association between the traditional positive symptoms of schizophrenia and violence in this study probably reflects the clinically stable nature of this sample. In line with our findings, others^{41,42} have reported an association between aggression and PANSS hostility and impulsivity scores. Similar to other studies, we did not observe an association between violence and negative symptoms.^{40–42}

Psychopathy

In line with previous studies we found an association between violence and psychopathy.⁴³ A novel aspect of this study is our analysis based on the newer four-facet model of psychopathy¹⁶ which indicated that the violent group had significantly higher

scores on the antisocial and interpersonal factors, but not on the affective or lifestyle factors. However, our regression analysis revealed that the interpersonal factor was the most significant discriminator of violent *v.* non-violent status. Our work partly confirms previous reports of an association between the antisocial components of psychopathy and violence.^{44–46} However, the lack of an association between violence and the affective components of psychopathy contrasts with Vitacco *et al.*⁴⁷ who found that both the affective and antisocial components of the four-facet model of psychopathy were associated with community violence in civil psychiatric patients. The discrepancy may reflect differences in samples (civil *v.* forensic) and differences in the context in which violence occurs (e.g. in-patient *v.* out-patient). However, given that in the present study the PCL–SV interpersonal factor and IQ were both significant discriminators of violent *v.* non-violent status, the interaction between psychopathy, IQ and violence in both in-patient and community settings in those with schizophrenia is an area worthy of further research.

Implications

The findings from this study suggest that, in clinically and demographically well-matched individuals with schizophrenia in forensic settings, violent and non-violent in-patients are best distinguished on the basis of key personality traits such as psychopathy rather than specific deficits in neuropsychological function. Given the significance of personality factors in distinguishing violent from non-violent individuals with schizophrenia in forensic settings, future studies should take account of the high levels of comorbid antisocial and psychopathic personality disorder pathology, and examine the relative role of personality factors in in-patient violence risk. As the non-psychotic literature suggests that psychopathy may be associated with specific deficits in VLPFC function⁴⁸ and DSM–IV antisocial personality disorder may be associated with a broader range of DLPFC and VLPFC deficits,^{49,50} future studies need to look at the impact of these comorbid personality pathologies on both neuropsychological function and violence in in-patient samples with schizophrenia.

Limitations

The focus on recruitment in secure forensic settings that have high base rates of in-patient violence but have highly controlled environments may influence the findings. Further work is needed in less secure in-patient settings. As environmental factors can influence institutional violence⁵¹ a measure of institutional environment should assist in understanding the complex array of factors associated with in-patient violence in people with schizophrenia. Our participants were assessed when clinically stable so the findings cannot be generalised to more acutely ill people. Given that the sample was purely male, it is difficult to comment on the applicability of the present findings to a mixed or solely female population. Future studies should examine the

contribution of gender differences. As this is a cohort rather than prospective study, no causal associations between predictors and outcome measures can be established. Future studies examining the relationship between symptoms, neuropsychological function and personality traits should use a prospective study design with an emphasis on cohorts with their first episode of psychosis. For studies specifically looking at incarcerated samples the potential moderator effects of environment must be considered as the latter factors may attenuate or exaggerate the person-specific risks of in-patient violence.

Rachael S. Fullam, PhD, University of Manchester, Manchester, and **Mairead C. Dolan**, MRCPsych, PhD, University of Manchester and Bolton Salford and Trafford NHS Mental Health Trust, Manchester, UK

Correspondence: Rachael Fullam, Centre for Forensic Behavioural Science, School of Psychology, Psychiatry and Psychological Medicine, Monash University, Victorian Institute for Forensic Mental Health, Locked Bag 10, Fairfield, VIC 3078, Australia. Email: rachel.fullam@forensicare.vic.gov.au

First received 14 May 2007, final revision 19 Mar 2008, accepted 31 Mar 2008

Acknowledgements

This study was supported by grants from the National Alliance for Research on Schizophrenia and Depression (NARSAD), the National Forensic Mental Health R&D Programme and Merseycare NHS Trust. We thank the staff and participants at key institutions.

References

- Naudts K, Hodgins H. Neurobiological correlates of violent behaviour among persons with schizophrenia. *Schizophr Bull* 2006; **32**: 562–72.
- Krakowski M, Convit A, Jaeger J, Lin S, Volavka J. Neurological impairment in violent schizophrenic inpatients. *Am J Psychiatry* 1989; **146**: 849–53.
- Adams JJ, Meloy JR, Moritz S. Neuropsychological deficits and violent behaviour in incarcerated schizophrenics. *J Nerv Ment Dis* 1990; **178**: 253–6.
- Forster HG, Hillbrand M, Silverstein M. Neuropsychological deficit and aggressive behaviour: a prospective study. *Prog Neuropsychopharmacol Biol Psychiatry* 1993; **17**: 939–46.
- Rasmussen K, Levander S, Sletvold H. Aggressive and non-aggressive schizophrenics: symptom profile and neuropsychological differences. *Psychol Crime Law* 1995; **2**: 119–29.
- Barkataki I, Kumari V, Das M, Morris R, O'Connell P, Taylor P, Sharma T. A neuropsychological investigation into violence and mental illness. *Schizophr Res* 2005; **74**, 1–13.
- Roy S, Herrera J, Parent M, Costa J. Violent and non-violent schizophrenic patients: Clinical and developmental characteristics. *Psychol Rep* 1987; **61**: 855–61.
- Lapierre D, Braun CM, Hodgins S, Toupin J, Leveillee S, Constantineau C. Neuropsychological correlates of violence in schizophrenia. *Schizophr Bull* 1995; **21**: 253–62.
- Lafayette MD, Frankle WG, Pollock A, Dyer K, Goff DC. Clinical characteristics, cognitive functioning and criminal histories of outpatients with schizophrenia. *Psychiatr Serv* 2003; **54**: 1635–40.
- Silver H, Goodman C, Knoll G, Isakov V, Modai I. Schizophrenia patients with a history of severe violence differ from non-violent schizophrenia patients in perception of emotions but not cognitive function. *J Clin Psychiatry* 2005; **66**: 300–8.
- Dinakar H, Sobel RN. Violence in the community as a predictor of violence in the hospital. *Psychiatr Serv* 2001; **52**: 240–1.
- American Psychiatric Association. *Diagnostic Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. APA, 1994.
- Kay SR, Opler LA, Fiszbein A. *The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia: Manual*. Multi-Health Systems, 1992.
- Lindenmayer JP, Grochowski S, Bark N. The five factor model of schizophrenia: replication across samples. *Schizophr Res* 1995; **14**: 229–34.
- Hart SD, Cox DN, Hare RD. *The Psychopathy Checklist – Screening Version (PCL-SV)*. Multi-Health Systems, 1995.
- Hare RD, Neumann CS. Structural models of psychopathy. *Curr Psychiatry Rep* 2005; **7**: 57–64.
- Hill CD, Neumann CS, Rogers R. Confirmatory factor analysis of the Psychopathy Checklist: screening version. *Psychol Ass* 2003; **15**: 41–5.
- Hare RD. *The Psychopathy Checklist-Revised (PCL-R)*. Multi-Health Systems, 1991.
- Nelson H. *National Adult Reading Test Manual*. nferNelson, 1991.
- The Psychological Corporation. *Wechsler Abbreviated Scale of Intelligence*. Harcourt, Brace & Co., 1999.
- Fray PJ, Robbins TW. CANTAB battery: proposed utility in neurotoxicology. *Neurotoxicol Teratol* 1996; **18**: 499–504.
- Rubia K, Russell T, Overmeyer S, Bullmore ET, Sharma T, Simmons A, Williams SC, Giampietro V, Andrew CM, Taylor E. Mapping motor inhibition: conjunctive brain activations across different versions of go/no-go and stop tasks. *Neuroimage* 2001; **13**: 250–61.
- Schacher R, Logan GD. Impulsivity and inhibitory control in normal development and childhood psychopathology. *Dev Psychol* 1990; **26**: 710–20.
- British Medical Association & Royal Pharmaceutical Society. *British National Formulary*. British Medical Journal & Pharmaceutical Press, 2005.
- Cheung P, Schweitzer I, Crowley K, Tuckwell V. Violence in schizophrenia: role of hallucinations and delusions. *Schizophr Res* 1997; **26**: 181–90.
- Taylor PJ, Leese M, Williams D, Butwell M, Daly R, Larkin E. Mental disorder and violence. A special (high security) hospital study. *Br J Psychiatry* 1998; **172**: 218–26.
- Tengström A, Grann M, Långström N, Kullgren G. Psychopathy (PCL-R) as a predictor of violent recidivism among criminal offenders with schizophrenia. *Law Hum Behav* 2000; **24**: 45–58.
- Hodgins S, Hiscoke UL, Freese MD. The antecedents of aggressive behaviour among men with schizophrenia: a prospective investigation of patients in community treatment. *Behav Sci Law* 2003; **21**: 523–46.
- Dolan M, Davies G. Psychopathy and institutional outcome in patients with schizophrenia in forensic settings in the UK. *Schizophr Res* 2006; **81**: 277–81.
- Raine A. *The Psychopathology of Crime: Criminal Behavior as a Clinical Disorder*. Academic Press, 1997.
- Gendreau P, Goggin C, Little T. *Predicting Adult Offender Recidivism: What Works!* Public Works and Government Services Canada, 1997.
- Walsh Z, Swogger MT, Kosson DS. Psychopathy, IQ, and violence in European American and African American county jail inmates. *J Consult Clin Psychol* 2004; **72**: 1165–69.
- Matarazzo JD, Herman DO. Relationship of education and IQ in the WAIS-R standardization sample. *J Consult Clin Psychol* 1984; **52**: 631–4.
- Elliott AK, Mirsky AF. Cognitive antecedents of violence and aggression. In *The Neurobiology of Criminal Behaviour* (ed J Glicksohn): 111–36. Kluwer Academic Publishers, 2002.
- Sabri M, Radnovich AJ, Li TQ, Kareken DA. Neural correlates of olfactory change detection. *Neuroimage* 2005; **25**: 969–74.
- Abu-Akel A, Abushua'leh K. 'Theory of mind' in violent and nonviolent patients with paranoid schizophrenia. *Schizophr Res* 2004; **69**: 45–53.
- Silver H, Feldman P, Bilker W, Gur RC. Working memory as a core neuropsychological dysfunction in schizophrenia. *Am J Psychiatry* 2003; **160**: 1809–16.
- Elliott R, McKenna PJ, Robbins TW, Sahakian BI. Specific neuropsychological deficits in schizophrenic patients with preserved intellectual function. *Cog Neuropsychiatry* 1998; **3**: 45–70.
- Bowie CR, Moriarty PJ, Harvey PD, Parrella M, White L, Davis KL. Aggression in elderly schizophrenia patients: a comparison of nursing home and state hospital residents. *J Neuropsychiatry Clin Neurosci* 2001; **13**: 357–66.
- Fresán A, Apiquian R, de la Fuente-Sandoval C, Löyzaga C, Garcia-Anaya M, Meyenberg N, Nicolini H. Violent behavior in schizophrenia patients: Relationship with clinical symptoms. *Aggress Behav* 2005; **31**: 511–20.
- Arango C, Calcedo-Barba A, González-Salvador T, Calcedo-Ordóñez A. Violence in inpatients with schizophrenia: a prospective study. *Schizophr Bull* 1999; **25**: 493–503.
- Nolan KA, Volavka J, Czober P, Sheitman B, Lindenmayer JP, Citrome LL, McEvoy J, Lieberman J.A. Aggression and psychopathology in treatment resistant inpatients with schizophrenia and schizoaffective disorder. *J Psychiatr Res* 2005; **39**: 109–15.
- Hare RD. Psychopathy: a clinical and forensic overview. *Psychiatr Clin North Am* 2006; **29**: 709–24.
- Skeem JL, Mulvey EP. Psychopathy and community violence among civil psychiatric patients. Results from the MacArthur Violence Risk Assessment Study. *J Consult Clin Psychol* 2001; **69**: 358–74.
- Skeem JL, Mulvey EP, Grisso T. Applicability of traditional and revised models of psychopathy to the Psychopathy Checklist: screening version. *Psychol Assess* 2003; **15**: 41–55.

- 46 Skeem JL, Miller JD, Mulvey E, Tiemann J, Monahan J. Using a five-factor lens to explore the relation between personality traits and violence in psychiatric patients. *J Consult Clin Psychol* 2005; **73**: 454–65.
- 47 Vitacco MJ, Neumann CS, Jackson RL. Testing a four-factor model of psychopathy and its association with ethnicity, gender, intelligence, and violence. *J Consult Clin Psychol* 2005; **73**: 466–76.
- 48 Blair KS, Newman C, Mitchell DG, Richell RA, Leonard A, Morton J, Blair RJ. Differentiating among prefrontal substrates in psychopathy: neuropsychological test findings. *Neuropsychology* 2006; **20**: 153–65.
- 49 Morgan AB, Lilienfeld SO. A meta-analytic review of the relation between antisocial behavior and neuropsychological measures of executive function. *Clin Psychol Rev* 2000; **20**: 113–36.
- 50 Dolan M, Park I. The neuropsychology of antisocial personality disorder. *Psychol Med* 2002; **32**: 417–27.
- 51 Gadon L, Johnstone L, Cooke D. Situational variables and institutional violence: a systematic review of the literature. *Clin Psychol Rev* 2006; **26**: 515–34.



Psychiatry in the Old Testament

Did Ezekiel have catatonia?

George Stein

3:25 'As for you mortal, cords shall be placed on you with them so that you cannot go out among the people.
26 And I will make your tongue cling to the roof of your mouth so that you shall be speechless.'

The picture here is of cords being placed upon Ezekiel so that he cannot move, but we do not know whether these were real, as in the case of Jeremiah, or allegorical cords. The phrase 'you will be speechless' is suggestive of mutism. The traditional rabbinical explanation for these verses is that Ezekiel was such a hypercritical prophet who was constantly berating the people for being rebellious that he had to be silenced and confined to his house. However, his period of silence and immobilisation went on for a long time.

4:4 'Then lie on your left side and place the punishment of the house of Israel upon it; you shall bear the punishment for the number of days you lie there
5 for I assign to you a number of days, 390 days and so you shall bear the punishment of the house of Israel.
6 When you have completed these, you shall lie down a second time, but on your right side and bear the punishment of the house of Judah: 40 days I assign you.
7 You shall set your face towards the siege of Jerusalem, and with your arm you shall prophesy against it.
8 See I am putting cords on you so that you cannot turn from one side to another until you have completed the days of your siege.'

In this passage Ezekiel must prophesy with his arm because he cannot speak and he cannot move, even to roll over. This lasts for a very long time, more than a year for the first episode and 40 days for the second episode. The most common and possibly the only non-fatal human condition lasting for such a long time where a person moves very little, does not speak and from which there may be complete recovery is catatonia.

Ezekiel ate very poorly and drank very little in his period of immobility and it seems he had to cook his food using human faeces (possibly his own): 4:12 'You shall eat it as barley cake, baking it in their sight on human dung'. The phrase 'in their sight' suggests that the local people must have witnessed this. The taboos about handling human excrement or defilement were thought to be very strong in ancient Israel, particularly among the priesthood (Ezekiel was an important priest). However, such taboos are sometimes lost in people with serious mental illness, especially schizophrenia. Previous psychiatric commentators, including Jaspers, in *Aneignung und Polemik*, have also suggested that Ezekiel may have suffered from catatonic schizophrenia.

The British Journal of Psychiatry (2008)
193, 253. doi: 10.1192/bjp.193.3.253